

Prevalence of post intensive care syndrome among Japanese intensive care unit patients: A prospective, multicenter, observational J-PICS study

Daisuke Kawakami (✉ dsk_kwkm_n9s@hotmail.co.jp)

Kobe City Medical Center General Hospital <https://orcid.org/0000-0001-7325-5512>

Shigeki Fujitani

Saint Marianna University School of Medicine: Sei Marianna Ika Daigaku

Takeshi Morimoto

Hyogo College of Medicine: Hyogo Ika Daigaku

Hisashi Dote

Seirei Hamamatsu Hospital: Seirei Hamamatsu Byoin

Mumon Takita

Saint Marianna University School of Medicine: Sei Marianna Ika Daigaku

Akihiro Takaba

JA Hiroshima General Hospital

Masaaki Hino

Kurashiki Central Hospital: Kurashiki Chuo Byoin

Michitaka Nakamura

Nara Prefecture General Medical Center: Nara-ken Sogo Iryo Center

Hiromasa Irie

Kurashiki Central Hospital: Kurashiki Chuo Byoin

Tomohiro Adachi

Tokyo Women's Medical University Medical Center East: Tokyo Joshi Ika Daigaku Higashi Iryo Center

Mami Shibata

Wakayama Medical University

Jun Kataoka

Tokyo Bay Urayasu Ichikawa Medical Center

Akira Korenaga

Japan Red Cross Wakayama Medical Center: Nihon Sekijujisha Wakayama Iryo Center

Tomoya Yamashita

Osaka City General Hospital: Osaka Shiritsu Sogo Iryo Center

Tomoya Okazaki

Kagawa University Hospital: Kagawa Daigaku Igakubu Fuzoku Byoin

Masatoshi Okumura

Aichi Medical University Hospital: Aichi Ika Daigaku Byoin

Takefumi Tsunemitsu

Hyogo Prefectural Amagasaki General Medical Center: Hyogo Kenritsu Amagasaki Sogo Iryo Center

Research

Keywords: Post-intensive care syndrome, Health-related Quality of Life, SF-36, Disability, Critical care, Mechanical ventilation, Intensive care unit

Posted Date: February 12th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-76221/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 16th, 2021. See the published version at <https://doi.org/10.1186/s13054-021-03501-z>.

Abstract

Background Many studies have compared quality of life of post-intensive care syndrome (PICS) patients with age-matched population-based controls. Many studies on PICS used the 36-item Short Form (SF-36) health survey questionnaire version 2 but lack the data for SF-36 values before and after intensive care unit (ICU) admission. Thus, clinically important changes in the parameters of SF-36 are unknown. Therefore, we determined the frequency of co-occurrence of PICS impairments at 6 months after ICU admission. We also evaluated the changes in SF-36 subscales and interpreted the patients' subjective significance of impairment.

Methods A prospective, multicenter, observational cohort study was conducted in 16 ICUs across 14 hospitals in Japan. Adult ICU patients expected to receive mechanical ventilation for >48 hours were enrolled, and their 6-month outcome was assessed using the questionnaires. PICS definition was based on the physical status, indicated by the change in SF-36 physical component score (PCS) ≥ 10 points; mental status, indicated by the change in SF-36 mental component score (MCS) ≥ 10 points; and cognitive function, indicated by the worsening of Short-Memory Questionnaire (SMQ) score and SMQ score at 6 months <40. Multivariate logistic regression model was used to identify the factors associated with PICS occurrence. The patients' subjective significance of physical and mental symptoms was assessed using the 7-scale Global Assessment Rating to evaluate minimal clinically important difference (MCID).

Results Among 192 patients, 48 (25%) died at 6 months. Among the survivors at 6 months, 96 patients responded to the questionnaire; ≥ 1 PICS impairment occurred in 61 (63.5%) patients, and ≥ 2 occurred in 17 (17.8%) patients. Physical, mental, and cognitive impairments occurred in 32.3%, 14.6%, and 37.5% patients, respectively. Population with only mandatory education was associated with PICS occurrence (odds ratio: 4.0, 95% confidence interval: 1.1–18.8, $P=0.029$). The MCID of PCS and MCS scores was 6.5 and 8.0, respectively.

Conclusions Among the survivors who received mechanical ventilation, 64% had PICS at 6 months; co-occurrence of PICS impairments occurred in 20%. PICS was associated with population with only mandatory education. Future studies elucidating the MCID of SF-36 scores among ICU patients and standardizing the PICS definition are required.

Trial registration UMIN000034072

Background

Recent improvement in intensive care unit (ICU) mortality rate has resulted in numerous ICU survivors. Most ICU survivors experience long-term impairment in their quality of life, known as post-intensive care syndrome (PICS). PICS is defined as a new or worsening impairment in physical, cognitive, or mental health status arising and persisting after hospitalization for critical illness [1]. Globally, many observational studies have been conducted for each PICS parameter among ICU patients [2–5]. These

studies have several limitations. First, most studies focused on only one or two aspects of PICS (i.e., only cognitive function, or only physical and mental status). Only a few studies focusing on the co-occurrence of PICS symptoms in individual patients have been conducted [6,7]. Second, limited data regarding PICS after implementation of PICS prophylaxis strategy are available, especially for Asian populations with different cultural backgrounds and races from Western countries [8–11]. Many guidelines and bundle strategies aimed at growing awareness about PICS have been published and implemented to decrease the occurrence of PICS. These include clinical practice guidelines for the prevention and management of pain, agitation, sedation, delirium, immobility, and sleep disruption in adult patients in the ICU [12]; guidelines for family centered care in the neonatal, pediatric, and adult ICU [13]; and ABCDEFGH bundle [14]. ABCDEFGH bundle comprises airway management; breathing trials; coordination of care and communication; delirium assessment; early mobility; family involvement, follow-up referrals, and functional reconciliation; good handoff communication; and handout materials on PICS and PICS-family. The Japanese PICS survey conducted in 2019 revealed that more than 60% medical staff recognized PICS, and early rehabilitation was provided in more than 90% cases [15]. Third, ICU patients often report lower health-related quality of life (HRQOL) than the general population before ICU admission. However, many studies have compared quality of life of PICS patients with age-matched population-based controls, instead of baseline quality of life [16]. Furthermore, many studies on PICS used the 36-item Short Form (SF-36) health survey questionnaire version 2, which is the most popular HRQOL scale for the outcome measure, among ICU survivors [17]; however, these studies lack the data for SF-36 values before and after ICU admission. Thus, the clinically important changes in the parameters of SF-36 in these studies are unknown.

Therefore, the Japanese PICS (J-PICS) study aimed to evaluate the co-occurrence of PICS symptoms, along with the assessment of baseline HRQOL, in Japanese patients admitted to multicenter ICUs, after publication of several guidelines and bundles about PICS prophylaxis. Additionally, we evaluated the subjective significance of changes in SF-36 parameters before and after ICU admission among the patients.

Methods

Study design and setting

The J-PICS study was a prospective, multicenter, observational cohort study of mechanically ventilated patients, conducted in 16 ICUs across 14 hospitals of Japan. Five of the 16 ICUs were university-affiliated hospitals, and the others were tertiary teaching hospitals. The median number of ICU beds was 10 [interquartile range (IQR), 8–15]. All ICUs were mixed ICUs and were managed by the intensivists. The study protocol was approved by the Kobe City Medical Center General Hospital (KCGH) ethics committee and the ethics committees of all participating hospitals (KCGH approval number: Zn181008). Written informed consent was obtained from all patients or authorized surrogates. The J-PICS study has been registered at University Hospital Medical Information Network Clinical Trials Registry (registration number: UMIN000034072).

Study population and eligibility criteria

All consecutive adult ICU patients who were expected to receive mechanical ventilation for more than 48 hours between April 01, 2019 and September 30, 2019 were recruited in the study. The patients who received noninvasive mechanical ventilation were also enrolled. Eligibility criteria were assessed next morning at 8:00 am after admission in the ICUs. The patients with the following conditions were excluded from the study: 1) patients who had primary brain injury that was likely to result in conscious or cognitive disorder (e.g., traumatic brain injury, subarachnoid hemorrhage, acute stroke, post cardiac arrest, meningitis, and encephalitis); 2) patients with pre-admission diagnosis of dementia; 3) patients who received home ventilation prior to admission; 4) patients with end-stage cancer; 5) patients with withdraw/withhold status; 6) expected death within 24 h; 7) second or subsequent admission to ICU during the study period; 8) patients who had no family members; 9) patients who did not speak Japanese; and 10) patients who could not be followed-up (e.g., did not live in Japan and/or were homeless).

Variables and measurements

The following demographic and hospital data of the patients were collected: age, sex, body mass index, Charlson comorbidity index (CCI) [18], clinical frailty scale [19], do not attempt resuscitation code status at the time of ICU admission, educational level, employment status, marital status, patient's residential living status before admission, history of treatment with benzodiazepines and steroids, source of admission to ICU, and primary diagnosis at the time of ICU admission. We chose 9 years as the cut-off value for educational level because elementary and junior high school are mandatory for everyone, which usually last for 9 years in Japan, following which, many people go to high school for three years, followed by universities and graduate schools. The number of patients with sepsis and acute respiratory distress syndrome (ARDS) was also recorded. Sepsis and ARDS were diagnosed on the basis of Sepsis-3 [20] and Berlin definition [21], respectively. The severity of illness was measured using the Acute Physiology and Chronic Health Evaluation (APACHE II) score and Sequential Organ Failure Assessment (SOFA) score [22] within 24 hours of ICU admission. The management data in the ICU, including the use of inotropes or vasopressors, paralytic agents except during intubation, renal replacement therapy, extracorporeal membrane oxygenation (ECMO), intra-aortic balloon pump, and tracheostomy were also collected. The data for the use of inotropes, vasopressors, and paralytic agents were collected during the first four days of ICU admission. The data on patient outcomes, including ICU mortality; in-hospital mortality; length of stay in ICU; length of stay in hospital; duration of mechanical ventilation; occurrence of delirium, diagnosed by the Confusion Assessment Method for the Intensive Care Unit [23], during the first four days; and discharge status among survivors were collected.

Patient-reported outcomes

Six months after ICU admission, the authors sent the questionnaires by post to all patients except those who died. Patient-reported data were collected centrally. If the participants did not revert, the lead author asked each participating institution to attempt a contact via telephone to return the questionnaire.

The patient-reported outcome survey evaluated the physical and mental functions of patients through the assessment of HRQOL and cognitive functions. HRQOL was assessed using the SF-36 questionnaire [24–26], which is available in the Japanese language [27,28]. At the time of enrolment in the study after ICU admission, the patients' baseline SF-36 questionnaire was completed by a proxy (4-week recall assessment before the patients' current acute illness). At 6 months after ICU admission, SF-36 questionnaire was obtained by mail from the patient or proxy. The SF-36 questionnaire has established acceptability, reliability, and validity in critically ill patient populations and as a surrogate-completed proxy measure [29–36]. SF-36 questionnaire is a comprehensive 36-item survey of HRQOL with two summary scales, physical component scale (PCS) and mental component scale (MCS), with scores ranging from 0 to 100. A higher score indicates better physical and mental functions. Both scales were transformed to a normalized scale using norm-based scoring (NBS) with 50 as the population mean and 10 points representing one standard deviation. The scores were calculated based on the standard methods [37]. The missing data were treated using the standard methods. If a patient answered more than half of the items on the subscale, the missing data were replaced with the mean of the subscale. In contrast, if a patient answered less than half items on the subscale, the data of the answered questions were excluded [37].

Cognitive functioning was assessed using the Short-Memory Questionnaire (SMQ) [38]. The SMQ was to be completed by a proxy, similar to the Informant Questionnaire on Cognitive Decline in the Elderly [39]. The SMQ is the only questionnaire filled by a proxy that has been translated and established in the Japanese language [40]. The SMQ is a 12-item questionnaire with scores ranging from 4 to 46. A score less than 40 indicates cognitive dysfunction.

The primary outcome was the occurrence of PICS 6 months after ICU admission. The definition of PICS was any of the following criteria: 1) decline in physical status, indicated by PCS score decrease of ≥ 10 points; 2) decline in mental status, indicated by MCS score decrease of ≥ 10 points; or 3) cognitive function impairment, indicated by decline in SMQ score and SMQ score < 40 at 6 months after ICU admission. A 10-point change was considered clinically important in a previous study in the ICU setting [36]. For the assessment of subjective significance of SF-36 score change, participants also answered anchor questions on whether their physical and mental status had improved, worsened, or remained unchanged on a 7-point Likert scale (from large negative change to large positive change) [41–43] at 6 months. The questions related to patients' employment status at 6 months were also asked.

Statistical analyses

Since the analyses focused on patient-reported outcomes at 6 months, participants who were lost to follow-up and who died before 6 months were excluded from the final analysis. Sample size was based on the total number of patients meeting the inclusion criteria because this prospective study was descriptive and hypothesis generating by nature. Continuous variables are presented as medians and interquartile range. Categorical variables are presented as absolute values and percentages. A univariate analysis was performed using the Wilcoxon rank-sum test for continuous variables, and the chi-squared

test or Fisher's exact test was used for categorical variables, if the number was less than 10. Multivariate logistic regression model was used to identify the factors associated with the occurrence of PICS. Multivariate analysis included the following variables as confounding factors: age, APACHE II score, CCI, and educational status [7, 44–46]. Age, APACHE II score, and CCI were log-transformed. The risk of PICS is expressed as odds ratio (OR) with 95% confidence interval (CI). The number of missing data have been reported, and no assumptions have been made except for SF-36 questionnaire data. The changes in SF-36 PCS, MCS, and SMQ scores between baseline and 6 months after ICU admission were compared using the Wilcoxon signed-rank test.

The subjective significance of change in SF-36 PCS and MCS was assessed by an anchor-based question with a Global Assessment Rating as large negative change, moderate negative change, small negative change, no change, small positive change, moderate positive change, and large positive change in physical and mental status. The change in each Global Assessment Rating for SF-36 PCS and MCS scores is expressed as mean and standard deviation (SD). Further, we evaluated the responsiveness of SF-36 PCS and MCS. Responsiveness is the instrument's ability to detect change over time in the construct to be measured [47,48]. Responsiveness of SF-36 PCS and MCS scores to negative change or no change in Global Assessment Rating at 6 months was calculated using the effect size index [49,50]. Cohen's d effect size was used to evaluate the changes as small (0.2–0.49), moderate (0.5–0.79), and large (> 0.80) [47,51]. We also assessed interpretability, which is the degree to which one can assign qualitative meaning to the quantitative scores or change in scores. We also evaluated the floor effect and ceiling effect of SF-36 PCS and MCS scores. Floor and ceiling effects were calculated as the percentage of participants scoring their lowest and highest (0 and 100) score, respectively. The floor and ceiling effects were considered relevant at 15% [52]. The minimal clinically important difference (MCID) of SF-36 PCS and MCS scores was calculated as the mean change score for no change patients minus the mean change score for small negative change patients [53,54].

The differences with P value less than 0.05 (two-sided) were considered statistically significant. All data were analyzed using JMP 15.1.0 (SAS Institute Inc., Cary, NC, USA).

Results

Patients' characteristics

During the study period, 947 consecutive patients were registered in the J-PICS study, among whom, 755 patients were excluded from the study based on the exclusion criteria. Finally, 192 patients were included in the study. Among 192 patients, 30 (15.6%) patients did not return the questionnaire survey, 48 (25.0%) patients died before 6 months of follow-up, and 18 (9.4%) patients did not complete the questionnaires. Thus, in total, 96 (50.0%) patients were enrolled for the assessment of 6-month outcome (Fig. 1).

Overall characteristics of the patients, along with stratification based on the presence and absence of PICS, are shown in Table 1. In total, 61 (63.5%) patients were in the PICS group and 35 (36.5%) in the non-PICS group. The mean age of the patients was 74 [64–81] years, the proportion of men was 65%, mean

APACHE II score was 23 [18–28], and mean SOFA score was 8 [6–11]. The main cause of ICU admission was acute respiratory failure. There were 35% and 41% patients with ARDS and sepsis, respectively. Population with only mandatory education was associated with PICS occurrence.

Hospital outcomes

Among the patients included in the study, ICU mortality, hospital mortality, and 6-month mortality were observed in 15 (7.8%), 33 (17.2%), and 48 (29.6%) patients, respectively. Table 2 shows the hospital outcomes and 6-month outcomes among the patients who were enrolled for assessment at 6 months after ICU admission. The median ICU length of stay, hospital length of stay, and duration of mechanical ventilation were 7 [5–14] days, 33.5 [19–61.8] days, and 6 [3–11] days, respectively. Among 159 survivors, discharge destination was home, another facility, and nursing home for 71 (44.7%), 87 (54.7%), and 1 (0.7%) patient, respectively. Among non-PICS patients, 22 (62.9%) were discharged home, while among PICS patients, 24 (39.3%) were discharged home from the hospital ($P=0.026$).

Patient-reported outcomes at 6 months

SF-36 score

Fig. 2 shows that SF-36 PCS score and MCS score changed significantly at 6 months compared with the baseline among both PICS and non-PICS patients. Among PICS patients, PCS score at baseline and 6 months was 37.6 [18.9–50.8] and 21.5 [11.4–34.8], respectively ($P<0.001$) and MCS score at baseline and 6 months was 49.7 [42.5–55.0] and 53.4 [44.2–63.4], respectively ($P=0.037$). In contrast, among non-PICS patients, PCS score at baseline and 6 months was 36.7 [11.2–45.4] and 41.0 [30.2–46.0], respectively ($P=0.25$), and MCS score at baseline and 6 months was 53.0 [42.2–58.6] and 56.8 [51.6–64.2], respectively ($P<0.001$). Physical and mental impairment, defined by the worsening of PCS and MCS scores by more than 10 points, occurred in 31 (32.3%) and 14 (14.6%) patients, respectively.

SMQ score

Fig. 2 also shows the change in SMQ score from baseline to 6 months among both PICS and non-PICS patients. Among PICS patients, SMQ score at baseline and 6 months was 43 [33.8–45] and 37 [23.5–43], respectively ($P<0.001$), while it was 43 [40–46] and 43 [41–45], respectively ($P=0.51$), in non-PICS patients. Cognitive impairment, defined by the worsening of SMQ score and SMQ score at 6 months < 40 , occurred in 36 (37.5%) patients.

PICS

In total, 96 patients completed all surveys about physical, mental, and cognitive status. The number of patients whose baseline PCS and MCS scores were < 50 , and SMQ score was < 40 was 74 (77.1%), 46 (47.9%), and 30 (31.3%), respectively. PICS was observed in 61 (63.5%) patients. The proportion of patients with one, two, and three PICS impairments is shown in Figure 3. Table 3 shows the results of multivariate analysis. After adjusting for confounding variables, Population with only mandatory

education (OR: 4.0, 95% CI: 1.1–18.8, P=0.029) was an independent predictor for the occurrence of PICS at 6 months after ICU admission. When education level was analyzed as a continuous variable, the same result was obtained (P=0.009).

Return to work

Among patients who completed the 6-month outcome survey, the data of four patients regarding job were missing. Among the remaining patients, 31 (33.7%) patients (20 in PICS group and 11 in non-PICS group) worked prior to ICU admission. The number of patients who returned to work was 26 (83.9%). All patients of non-PICS group returned to work, while 15 (75%) patients of PICS group returned to work.

Subjective significance of SF-36 score

The subjective significance of change in SF-36 PCS and MCS scores assessed by Global Assessment Rating was not obtained in 3 patients. Additional Table 1 shows the change in SF-36 PCS and MCS scores based on change in Global Assessment Rating. The effect size index between Global Assessment Rating negative change and change in PCS and MCS scores was 0.72 and 0.50, respectively. Area under receiver operating characteristics curve between Global Assessment Rating negative change and change in PCS and MCS scores was 0.69 and 0.64, respectively. These values indicate moderate responsiveness between the change in SF-36 PCS and MCS scores and negative change in Global Assessment Rating. Floor effect of PCS score at baseline and 6 months was 4.3% and 3.2%, respectively. Ceiling effect of PCS score, and floor and ceiling effect of MCS score were not observed. MCID of PCS and MCS scores was 6.5 and 8.0, respectively.

Discussion

Summary of key findings

In total, 192 ICU patients who were expected to receive mechanical ventilation for more than 48 hours were registered in the J-PICS study. The 6-months mortality rate was 30%. The prevalence of PICS at 6 months after ICU admission was 64% among 96 adult ICU survivors. New or worsened physical, mental, and cognitive impairments occurred in 32.3%, 14.6%, and 37.5% patients, respectively. The percentage of patients who had two or more PICS impairments was 17.8%. Population with only mandatory education was associated with the occurrence of PICS. MCID of PCS and MCS scores was 6.5 and 8.0, respectively, as analyzed by anchor-based subjective questions.

Comparison to other studies

This study investigated the co-occurrence of PICS symptoms at 6 months after ICU admission. To the best of our knowledge, only two studies have investigated the co-occurrence of PICS symptoms. One was a small cohort study, wherein, the study population comprised patients with ICU length of stay of at least 2 days [6]. This study reported the occurrence of PICS among the patients and at least one PICS impairment was high in 84% of the patients. The prevalence of two or more PICS impairments was 56%.

Another study [7], which comprised the patients with respiratory failure or shock, reported that the prevalence of PICS was 64% at 3 months and 56% at 12 months after hospital discharge. Two or more PICS impairments occurred in 25% patients at 3 months and 21% patients at 12 months. The results of the latter study were similar to those of our cohort. In contrast, the results of the former study showed a higher prevalence of PICS than the results of our cohort. This difference may be due to the difference in baseline data. In our cohort, baseline SF-36 PCS and MCS scores were lower than NBS. The former study did not evaluate the baseline status; and the latter study excluded patients with preexisting physical, mental, and cognitive impairments. Additionally, the former study reported that PICS occurred in 54% of the survivors and these patients reported worse symptoms after critical illness. The aforementioned observations suggest that the evaluation of baseline status is an important factor for interpreting the prevalence of PICS. The exclusion of patients with preexisting physical, mental, and cognitive impairments was also effective in interpreting the prevalence of PICS. However, that leads to the exclusion of many patients from the study. Therefore, we should evaluate the baseline status of the patients in PICS study.

In the present study, we found significant difference in discharge destination of the patients between the PICS and non-PICS groups. Many patients in the PICS group were transferred to other hospitals where the duration of treatment was associated with the severity of underlying illness. Thus, there is a possibility that hospital length of stay, including that in another hospital, might have affected the PICS symptoms. Therefore, several confounding factors might have affected our results.

In our study, population with only mandatory education was an independent predictor for the occurrence of PICS. This observation is consistent with that of the previous studies, which demonstrated that population with only mandatory education was associated with physical impairment [49], mental impairment [55], and PICS [7]. Socioeconomic position is a broad term which assesses the socioeconomic status, including education level, occupation, and income. Underprivileged is associated with poor health due to behavioral risk factors. The underprivileged persons often have high rates of cardiovascular disease and mortality due to tobacco smoking and less physical activity [55]. Furthermore, those who received less formal education is associated with low income, small social network to use resources for recovery, less coping ability for the rehabilitation process, and less accurate recall [7,49,55,56]. Additionally, financial distress has a direct effect on mental status [57].

In the present study, 84% patients returned to work at 6 months after ICU admission. The proportion of patients returning to work in this study was higher than that reported in the previous study [58,59]. The patients in our study were older and lower proportion of persons worked before ICU admission as compared to previous studies [58,59]. One ARDS longitudinal follow-up study in Canada reported that return to work rates at 1 year and 5 years were 63% (40 of 64 previously employed survivors) and 92% (49 of 53 survivors), respectively [58]. Another ARDS study conducted in the US reported return to work rates at 1 year and 5 years as 49% (28 of 67 previously employed survivors) and 44% (28 of 64 survivors), respectively [59]. These differences were potentially explained by the differences in patient population (e.g., age and diagnosis at admission), and socioeconomic or geographic factors [59]. It has also been

reported that return to work is driven by economic necessity and difficulties in obtaining administrative financial support [60].

We also evaluated the subjective significance of SF-36 score among ICU patients. We evaluated the responsiveness and MCID of only negative change in Global Assessment Rating because the patients with positive change were less in number and our focus was to analyze PICS, i.e., a negative change. This study showed moderate responsiveness between the change in SF-36 PCS and MCS scores and negative change in Global Assessment Rating. A floor effect was observed, but no ceiling effect. Clinically significant differences, rather than statistically significant differences, are important to assess HRQOL. The smallest meaningful change in a score is indicated by MCID. In the present study, we used a 10-point change in PCS or MCS score as the criterion of PICS. However, some studies have used a 5-point change as the criterion of PICS [61]. In the present study, MCID of PCS and MCS scores was 6.5 and 8.0, respectively, as analyzed by anchor-based subjective questions. A 5-point change might have overestimated PICS. MCID can be determined by two approaches: anchor-based method and distribution method. MCID values should be determined by systematic review and evaluation processes such as the modified Delphi method [62]. Our study had a small sample size, and the evaluation was done only by an anchor-based method. Future study with a larger population is required to clarify the MCID of SF-36 scores among ICU patients to standardize the definition of PICS globally.

Strengths and limitations

The present study has several strengths. First, we investigated not only the PICS symptoms but also the co-occurrence of PICS symptoms. Second, we assessed the baseline status of the patients. Third, we evaluated the MCID of SF-36 score change. Fourth, the J-PICS study was the first multicenter observational study of PICS in Japan. To date, a few studies have investigated the occurrence of PICS in Asian populations.

The present study also has some limitations. First, the long-term outcomes were assessed at only one time point. However, a systematic review showed that 60% of the previous studies about PICS have reported a single follow-up assessment. Furthermore, there were 71% articles without baseline assessment of quality of life [63]. We did not do multiple follow-up assessments, but we incorporated baseline assessment in our study. The systematic review about trajectory of recovery after ICU showed that HRQOL, cognitive function, and return to work always rapidly improved during the first 6 months, whereas improved a little after 1 year [64–66]. For this reason, a single follow-up at 6 months is appropriate for assessing the PICS symptoms.

Second, the definition of PICS has not gained a consensus yet. In 425 articles about PICS, 250 different measurement instruments were used [63]. Furthermore, we evaluated the clinimetric property of SF-36. The previous studies assessed all three domains originally defined for PICS using cut-off for each measurement instrument [6,7]. Recently, expert consensus for screening of post discharge impairments was published [67]. However, research agenda about measurement instruments based on follow-up strategies, like in-person interview (in clinic or online), telephone interview, or post questionnaires, were

not decided. Third, our data may have some bias; for example, recall bias because the outcome measures were self-reported by the patients or proxies. Fourth, we could not include a large number of patients in our study, which is also the limitation of majority of the similar studies. This might have affected the mortality rate and prevalence of PICS.

Conclusions

Our data suggest that 64% of ICU survivors had PICS impairments. Approximately 20% of survivors had a co-occurrence of PICS impairments. The less literate was associated with the occurrence of PICS. Future studies are needed to elucidate the modifiable risk factors of PICS to prevent the occurrence of PICS. We attempted to clarify the MCID of SF-36 among the ICU population. MCID of PCS and MCS scores was 6.5 and 8.0, respectively. More research is required for the assessment of MCID of SF-36 among the ICU population. Additionally, generalizability of the definition of PICS is needed.

Abbreviations

APACHE II Acute Physiology and Chronic Health Evaluation II

ARDS Acute respiratory distress syndrome

BMI Body mass index

CI Confidence interval

CCI Charlson comorbidity index

DNAR Do not attempt resuscitation

ECMO Extracorporeal membrane oxygenation

HRQOL Health-related quality of life

IABP Intra-aortic balloon pump

ICU Intensive care unit

IQR Interquartile range

KCGH Kobe City Medical Center General Hospital

MCID Minimal clinically important difference

MCS Mental component scale

NBS Norm-based scoring

OR	Odds ratio
PCS	Physical component scale
PICS	Post-intensive care syndrome
RRT	Renal replacement therapy
SD	Standard deviation
SF-36	36-item Short Form
SMQ	Short-Memory Questionnaire
SOFA	Sequential Organ Failure Assessment

Declarations

Ethics approval and consent to participate

The study protocol was approved by the Kobe City Medical Center General Hospital (KCGH) ethics committee and the ethics committees of all participating hospitals (KCGH approval number: Zn181008).

Consent for publication

Written informed consent was obtained from all patients or authorized surrogates.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by the KCGH research fund. There was no industry involvement in this trial.

Authors' contributions

Study concept and design: DK, SF, and TM

Acquisition of data: All declared authors

Analysis and interpretation of data: DK, SF, and TM

Drafting of the manuscript: DK

Critical revision of the manuscript for important intellectual content: All declared authors.

DK has full access to all the data in the study and takes responsibility for the integrity of the data. All authors have read and approved the final manuscript.

Acknowledgements

The J-PICS study group: Masatoshi Okumura, Aichi Medical University Hospital; Takeshi Morimoto, Hyogo College of Medicine; Takefumi Tsunemitsu, Hyogo Prefectural Amagasaki General Medical Center; Akihiro Takaba, JA Hiroshima General Hospital; Akira Korenaga, Japanese Red Cross Wakayama Medical Center; Tomoya Okazaki, Kagawa University Hospital; Daisuke Kawakami, Kobe City Medical Center General Hospital; Masaaki Hino and Hiromasa Irie, Kurashiki Central Hospital; Michitaka Nakamura, Nara Prefecture General Medical Center; Tomoya Yamashita and Kazuaki Shigemitsu, Osaka City General Hospital; Hisashi Dote, Seirei Hamamatsu General Hospital; Shigeki Fujitani and Mumon Takita, St. Marianna University School of Medicine; Jun Kataoka, Kenji Ishii, Miku Kamada, and Kumi Maruyama, Tokyo Bay Urayasu Ichikawa Medical Center; Tomohiro Adachi and Miki Sorita, Tokyo Women's Medical University Medical Center East; Mami Shibata and Kyohei Miyamoto, Wakayama Medical University Department of Emergency and Critical Care Medicine.

We would like to thank Editage (www.editage.com) for English language editing.

References

1. Needham DM, Davidson J, Cohen H, Hopkins RO, Weinert C, Wunsch H, et al. Improving long-term outcomes after discharge from intensive care unit: Report from a stakeholders' conference. *Crit Care Med*. 2012;40:502–9.
2. Herridge MS, Tansey CM, Matte A, Tomlinson G, Diaz-Granados N, Cooper A, et al. Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med*. 2011;364:1293–304.
3. Pandharipande PP, Girard TD, Jackson JC, Morandi A, Thompson JL, Pun BT, et al. Long-term cognitive impairment after critical illness. *N Engl J Med*. 2013;369:1306–16.
4. Sivanathan L, Wunsch H, Vigod S, Hill A, Pinto R, Scales DC. Mental illness after admission to an intensive care unit. *Intensive Care Med*. 2019;45:1550–8.
5. Righy C, Rosa RG, da Silva RTA, Kochhann R, Migliavaca CB, Robinson CC, et al. Prevalence of post-traumatic stress disorder symptoms in adult critical care survivors: A systematic review and meta-analysis. *Crit Care*. 2019;23:213.
6. Maley JH, Brewster I, Mayoral I, Siruckova R, Adams S, McGraw KA, et al. Resilience in survivors of critical illness in the context of the survivors' experience and recovery. *Ann Am Thorac Soc*. 2016;13:1351–60.

7. Marra A, Pandharipande PP, Girard TD, Patel MB, Hughes CG, Jackson JC, et al. Co-occurrence of post-intensive care syndrome problems among 406 survivors of critical illness. *Crit Care Med*. 2018;46:1393–401.
8. Hung M-C, Yan Y-H, Fan P-S, Lin M-S, Chen C-R, Kuo L-C, et al. Estimation of quality-adjusted life expectancy in patients under prolonged mechanical ventilation. *Value Health*. 2011;14:347–53.
9. Wang Z-Y, Li T, Wang C-T, Xu L, Gao X-J. Assessment of 1-year outcomes in survivors of severe acute respiratory distress syndrome receiving extracorporeal membrane oxygenation or mechanical ventilation. *Chin Med J (Engl)*. 2017;130:1161–8.
10. Su Y-X, Xu L, Gao X-J, Wang Z-Y, Lu X, Yin C-F. Long-term quality of life after sepsis and predictors of quality of life in survivors with sepsis. *Chin J Traumatol*. 2018;21:216–23.
11. Wu KK, Cho VW, Chow FL, Tsang AP, Tse DM. Posttraumatic stress after treatment in an intensive care unit. *East Asian Arch Psychiatry*. 2018;28:39–44.
12. Devlin JW, Skrobik Y, Gélinas C, Needham DM, Slooter AJC, Pandharipande PP, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med*. 2018;46:e825–73.
13. Davidson JE, Aslakson RA, Long AC, Puntillo KA, Kross EK, Hart J, et al. Guidelines for family-centered care in the neonatal, pediatric, and adult ICU. *Crit Care Med*. 2017;45:103–28.
14. Harvey MA, Davidson JE. Postintensive care syndrome: Right care, right now...and later. *Crit Care Med*. 2016;44:381–5.
15. The committee on PICS management and improvement of quality of life, Japanese Society of Intensive Care Medicine. A questionnaire survey regarding recognition and clinical practice of post-intensive care syndrome in Japan [in Japanese]. *J Jpn Soc Intensive Care Med*. 2019;26:467–75.
16. Dowdy DW, Eid MP, Sedrakyan A, Mendez-Tellez PA, Pronovost PJ, Herridge MS, et al. Quality of life in adult survivors of critical illness: A systematic review of the literature. *Intensive Care Med*. 2005;31:611–20.
17. Angus DC, Carlet J, 2002 Brussels Roundtable Participants. Surviving intensive care: A report from the 2002 Brussels Roundtable. *Intensive Care Med*. 2003;29:368–77.
18. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40:373–83.
19. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. 2005;173:489–95.
20. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315:801–10.
21. ARDS Definition Task Force; Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, et al. Acute respiratory distress syndrome: The Berlin definition. *JAMA*. 2012;307:2526–33.

22. Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med.* 1996;22:707–10.
23. Ely EW, Inouye SK, Bernard GR, Gordon S, Francis J, May L, et al. Delirium in mechanically ventilated patients: Validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU). *JAMA.* 2001;286:2703–10.
24. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care.* 1992;30:473–83.
25. McHorney CA, Ware JE Jr, Raczek AE. The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Med Care.* 1993;31:247–63.
26. Chrispin PS, Scotton H, Rogers J, Lloyd D, Ridley SA. Short Form 36 in the intensive care unit: Assessment of acceptability, reliability and validity of the questionnaire. *Anaesthesia.* 1997;52:15–23.
27. Fukuhara S, Bito S, Green J, Hsiao A, Kurokawa K. Translation, adaptation, and validation of the SF-36 Health Survey for use in Japan. *J Clin Epidemiol.* 1998;51:1037–44.
28. Fukuhara S, Ware JE Jr, Kosinski M, Wada S, Gandek B. Psychometric and clinical tests of validity of the Japanese SF-36 Health Survey. *J Clin Epidemiol.* 1998;51:1045–53.
29. Rogers J, Ridley S, Chrispin P, Scotton H, Lloyd D. Reliability of the next of kins' estimates of critically ill patients' quality of life. *Anaesthesia.* 1997;52:1137–43.
30. Diaz-Prieto A, Gorriz MT, Badia X, Torrado H, Farrero E, Amador J, et al. Proxy-perceived prior health status and hospital outcome among the critically ill: Is there any relationship? *Intensive Care Med.* 1998;24:691–8.
31. Capuzzo M, Grasselli C, Carrer S, Gritti G, Alvisi R. Quality of life before intensive care admission: Agreement between patient and relative assessment. *Intensive Care Med.* 2000;26:1288–95.
32. Cuthbertson BH, Scott J, Strachan M, Kilonzo M, Vale L. Quality of life before and after intensive care. *Anaesthesia.* 2005;60:332–9.
33. Nessler N, Defontaine A, Launey Y, Morcet J, Mallédant Y, Seguin P. Long-term mortality and quality of life after septic shock: A follow-up observational study. *Intensive Care Med.* 2013;39:881–8.
34. Gardner AK, Ghita GL, Wang Z, Ozrazgat-Baslanti T, Raymond SL, Mankowski RT, et al. The development of chronic critical illness determines physical function, quality of life, and long-term survival among early survivors of sepsis in surgical ICUs. *Crit Care Med.* 2019;47:566–73.
35. Higgins AM, Peake SL, Bellomo R, Cooper DJ, Delaney A, Harris AH, et al. Quality of life and 1-year survival in patients with early septic shock: Long-term follow-up of the Australasian resuscitation in sepsis evaluation trial. *Crit Care Med.* 2019;47:765–73.
36. Heyland DK, Garland A, Bagshaw SM, Cook D, Rockwood K, Stelfox HT, et al. Recovery after critical illness in patients aged 80 years or older: A multi-center prospective observational cohort study.

- Intensive Care Med. 2015;41:1911–20.
37. Fukuhara S, Suzukamo Y. Manual of SF-36v2 Japanese version: iHope International Inc. Kyoto, 2004, 2015.
 38. Koss E, Patterson MB, Ownby R, Stuckey JC, Whitehouse PJ. Memory evaluation in Alzheimer's disease. Caregivers' appraisals and objective testing. *Arch Neurol.* 1993;50:92–7.
 39. Jorm AF. The Informant Questionnaire on cognitive decline in the elderly (IQCODE): A review. *Int Psychogeriatr.* 2004;16:275–93.
 40. Maki N, Ikeda M, Hokoishi K, Nebu A, Komori K, Hirono N, et al. Japanese version of the Short-Memory Questionnaire: memory evaluation in Alzheimer's disease [in Japanese]. *No To Shinkei.* 1998;50:415–8.
 41. Osoba D, Rodrigues G, Myles J, Zee B, Pater J. Interpreting the significance of changes in health-related quality-of-life scores. *J Clin Oncol.* 1998;16:139–44.
 42. Fischer D, Stewart AL, Bloch DA, Lorig K, Laurent D, Holman H. Capturing the patient's view of change as a clinical outcome measure. *JAMA.* 1999;282:1157–62.
 43. Kamper SJ, Maher CG, Mackay G. Global rating of change scales: A review of strengths and weaknesses and considerations for design. *J Man Manip Ther.* 2009;17:163–70.
 44. Bastian K, Hollinger A, Mebazaa A, Azoulay E, Féliot E, Chevreur K, et al. Association of social deprivation with 1-year outcome of ICU survivors: Results from the FROG-ICU study. *Intensive Care Med.* 2018;44:2025–37.
 45. Wang S, Allen D, Perkins A, Monahan P, Khan S, Lasiter S, et al. Validation of a new clinical tool for post-intensive care syndrome. *Am J Crit Care.* 2019;28:10–8.
 46. Milton A, Schandl A, Soliman IW, Meijers K, van den Boogaard M, Larsson IM, et al. Development of an ICU discharge instrument predicting psychological morbidity: A multinational study. *Intensive Care Med.* 2018;44:2038–47.
 47. Angst F. The new COSMIN guidelines confront traditional concepts of responsiveness. *BMC Med Res Methodol.* 2011;11:152.
 48. Mokkink LB, Terwee CB, Patrick DL, Alonso J, Stratford PW, Knol DL, et al. The COSMIN study reached international consensus on taxonomy, terminology, and definitions of measurement properties for health-related patient-reported outcomes. *J Clin Epidemiol.* 2010;63:737–45.
 49. Puthuchery ZA, Gensichen JS, Cakiroglu AS, Cashmore R, Edbrooke L, Heintze C, et al. Implications for post critical illness trial design: sub-phenotyping trajectories of functional recovery among sepsis survivors. *Crit Care.* 2020;24:577.
 50. Kazis LE, Anderson JJ, Meenan RF. Effect sizes for interpreting changes in health status. *Med Care.* 1989;27:S178–89.
 51. Cohen J. *Statistical power analysis for the behavioral sciences.* 2nd ed. Hillsdale: Lawrence Erlbaum; 1988.

52. Terwee CB, Bot SD, de Boer MR, van der Windt DA, Knol DL, Dekker J, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol*. 2007;60:34–42.
53. Stucki G, Liang MH, Fossel AH, Katz JN. Relative responsiveness of condition-specific and generic health status measures in degenerative lumbar spinal stenosis. *J Clin Epidemiol*. 1995;48:1369–78.
54. Turner D, Schünemann HJ, Griffith LE, Beaton DE, Griffiths AM, Critch JN, et al. The minimal detectable change cannot reliably replace the minimal important difference. *J Clin Epidemiol*. 2010;63:28–36.
55. Jones JRA, Berney S, Connolly B, Waterland JL, Denehy L, Griffith DM, et al. Socioeconomic position and health outcomes following critical illness: A systematic review. *Crit Care Med*. 2019;47:e512–21.
56. Dinglas VD, Gellar J, Colantuoni E, Stan VA, Mendez-Tellez PA, Pronovost PJ, et al. Does intensive care unit severity of illness influence recall of baseline physical function? *J Crit Care*. 2011;26:634.e1–7.
57. Khandelwal N, Hough CL, Downey L, Engelberg RA, Carson SS, White DB, et al. Prevalence, risk factors, and outcomes of financial stress in survivors of critical illness. *Crit Care Med*. 2018;46:e530–9.
58. Herridge MS, Tansey CM, Matté A, Tomlinson G, Diaz-Granados N, Cooper A, et al. Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med*. 2011;364:1293–304.
59. Kamdar BB, Sepulveda KA, Chong A, Lord RK, Dinglas VD, Mendez-Tellez PA, et al. Return to work and lost earnings after acute respiratory distress syndrome: a 5-year prospective, longitudinal study of long-term survivors. *Thorax*. 2018;73:125–33.
60. Griffiths J, Hatch RA, Bishop J, Morgan K, Jenkinson C, Cuthbertson BH, et al. An exploration of social and economic outcome and associated health-related quality of life after critical illness in general intensive care unit survivors: a 12-month follow-up study. *Crit Care*. 2013;17:R100.
61. Orwelius L, Nordlund A, Nordlund P, Simonsson E, Bäckman C, Samuelsson A, et al. Pre-existing disease: The most important factor for health related quality of life long-term after critical illness: A prospective, longitudinal, multicentre trial. *Crit Care*. 2010;14:R67.
62. Revicki D, Hays RD, Cella D, Sloan J. Recommended methods for determining responsiveness and minimally important differences for patient-reported outcomes. *J Clin Epidemiol*. 2008;61:102–9.
63. Turnbull AE, Rabiee A, Davis WE, Nasser MF, Venna VR, Lolitha R, et al. Outcome measurement in ICU survivorship research from 1970 to 2013: A scoping review of 425 publications. *Crit Care Med*. 2016;44:1267–77.
64. Gerth AMJ, Hatch RA, Young JD, Watkinson PJ. Changes in health-related quality of life after discharge from an intensive care unit: a systematic review. *Anaesthesia*. 2019;74:100–8.
65. Wolters AE, Slooter AJ, van der Kooi AW, van Dijk D. Cognitive impairment after intensive care unit admission: a systematic review. *Intensive Care Med*. 2013;39:376–86.
66. Kamdar BB, Suri R, Suchyta MR, Digrande KF, Sherwood KD, Colantuoni E, et al. Return to work after critical illness: A systematic review and meta-analysis. *Thorax*. 2020;75:17–27.

67. Mikkelsen ME, Still M, Anderson BJ, Bienvenu OJ, Brodsky MB, Brummel N, et al. Society of Critical Care Medicine's international consensus conference on Prediction and Identification of Long-Term Impairments After Critical Illness. Crit Care Med. 2020;48:1670–79.

Tables

Table 1. Patients' characteristics: overall and stratified by the presence and absence of PICS

	non-PICS (N=35)	PICS (N=61)	P value	Overall (N=192)
Age, years	75 [64-79]	74 [59-81.5]	0.77	74 [64-81]
Male, N (%)	25 (71.4)	43 (70.5)	0.92	125 (65.1)
BMI, kg/m ²	21.9 [19.5-24.4]	22.8 [19.7-25.3] ^a	0.21	21.9 [19.5-25.0] ^b
APACHE II score	21 [18-25]	20 [14.5-25]	0.14	23 [18-28]
SOFA score	7 [4-10]	8 [5-10]	0.21	8 [6-11]
Charlson comorbidity index	1 [0-2]	1 [0-2]	0.9	1 [0-3]
Clinical frailty scale	3 [2-4] ^a	3 [2-4]	0.67	3 [3-4] ^a
DNAR status at ICU admission, N (%)	1 (2.9)	2 (3.3)	1	15 (7.8)
Educational level, N (%)	N=33*	N=60*	0.098	N=177
≤ 9 years	3 (9.1)	15 (25)		49 (27.7)
> 9 years	30 (91.0)	45 (75)		128 (72.3)
Employment status, N (%)	N=34*	N=61*	0.16	N=188
Student	0 (0)	0 (0)		2 (1.1)
Employed or self-employed	11 (32.4)	23 (37.7)		60 (31.9)
Unemployed	3 (8.8)	11 (18)		41 (21.8)
Housework	2 (5.9)	8 (13.1)		21 (11.2)
Retired	18 (52.9)	19 (31.2)		64 (34.0)
Marital status, N (%)			0.64	
Married	25 (71.4)	39 (63.9)		119 (62.0)
Separated or divorced	1 (2.9)	6 (9.8)		16 (8.3)
Widowed	5 (14.3)	9 (15.8)		37 (19.3)
Unmarried	4 (11.4)	7 (11.5)		20 (10.4)
Patient's residential living status, N (%)			0.25	
Lived alone at home	3 (8.6)	12 (19.7)		33 (17.2)
Lived with someone else	32 (91.4)	48 (78.7)		152 (79.2)
Nursing home	0 (0)	1 (1.6)		7 (3.6)

History of treatment with benzodiazepines, N3 (8.6) (%)	7 (11.5)	0.74	18 (9.4)
History of treatment with steroids, N (%)	2 (5.7)	4 (6.6)	1 35 (18.2)
Source of admission to ICU, N (%)			0.19
Emergency department	20 (57.1)	23 (37.7)	95 (49.5)
Hospital floor	5 (14.3)	16 (26.2)	45 (23.4)
Another hospital	1 (2.9)	0 (0)	3 (1.6)
Operating room (elective)	0 (0)	1 (1.6)	4 (2.1)
Operating room (emergency)	9 (25.7)	21 (34.4)	45 (23.4)
Primary diagnosis at the time of admission in ICU, N (%)			0.075
Cardiogenic	2 (5.7)	13 (21.3)	25 (13.0)
Acute respiratory failure	18 (51.4)	16 (26.2)	67 (34.9)
Infection	8 (22.9)	13 (21.3)	56 (29.2)
Trauma	3 (8.6)	8 (13.1)	16 (8.3)
Others	4 (11.4)	11 (18)	28 (14.6)
ARDS, N (%)	5 (14.3)	12 (19.7)	0.51 35 (18.2)
Sepsis, N (%)	11 (31.4)	20 (32.8)	0.89 79 (41.1)
Management in ICU, N (%)			
Inotrope/vasopressor	27 (77.1)	50 (82.0)	0.6 148 (77.1)
Paralytic agent	3 (8.6)	9 (14.8)	0.53 24 (12.5)
Renal replacement therapy in ICU	6 (17.1)	6 (9.8)	0.34 40 (20.8)
ECMO in ICU	0 (0)	4 (6.6)	0.29 7 (3.6)
IABP in ICU	0 (0)	3 (4.9)	0.3 6 (3.1)
Tracheostomy	4 (11.4)	9 (14.8)	0.76 29 (15.1)

Data are presented as median [interquartile range] or number (percentage).

Abbreviations: PICS: Post-intensive care syndrome, BMI: Body mass index, APACHE II: Acute Physiology and Chronic Health Evaluation, SOFA: Sequential Organ Failure Assessment, DNAR:

Do not attempt resuscitation, ICU: Intensive care unit, ARDS: Acute respiratory distress syndrome, ECMO: Extracorporeal membrane oxygenation, IABP: Intra-aortic balloon pump

^a one missing datum

^b three missing data

* the education status and the employment status of the remaining persons were unknown

Table 2. Patients' hospital and 6-months outcome: overall and stratified by the PICS status

	non-PICS (N=35)	PICS (N=61)	P value	Overall (N=192)
Mortality				
ICU mortality	NA	NA		15 (7.8)
Hospital mortality	NA	NA		33 (17.2)
6-month mortality	NA	NA		48 (29.6) ^a
ICU length of stay (days)	7 [5-12]	7 [5-13]	0.78	7 [5-14]
Hospital length of stay (days)	28 [17-47]	39 [19-71.5]	0.10	33.5 [19-61.8]
Days of mechanical ventilation	5 [3-7]	5 [3-11]	0.96	6 [3-11]
Delirium, N (%)	11 (31.4)	20 (32.8)	0.89	61 (31.8)
Discharged from hospital among survivors, N (%)			0.026	N=159
Home	22 (62.9)	24 (39.3)		71 (44.7)
Another facility	13 (37.1)	37 (60.1)		87 (54.7)
Nursing home	0	0		1 (0.7)

Data are presented as median [interquartile range] or number (percentage).

Abbreviations: PICS: Post-intensive care syndrome, ICU: Intensive care unit, NA: not applicable

^a 30 missing data

Figures




Figure 1

Flowchart depicting the enrolment of subjects in the study Abbreviations: ICU: Intensive care unit, SF-36: 36-item Short Form health survey questionnaire, SMQ: Short-Memory Questionnaire, PICS: Post-intensive care syndrome

A. PCS score before and after ICU admission

B. MCS score before and after ICU admission

C. SMQ score before and after ICU admission




Figure 2

Changes in SF-36 PCS, MCS, and SMQ scores before and after ICU admission Abbreviations: SF-36: 36-item Short Form health survey questionnaire, PCS: Physical component scale, MCS: Mental component scale, SMQ: Short-Memory Questionnaire, ICU: Intensive care unit




Figure 3

Occurrence of PICS problems among patients at 6 months after ICU admission Abbreviations: PICS: Post-intensive care syndrome

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Graphicalabstractimage.jpg](#)
- [AdditionalFigure1.pptx](#)
- [AdditionalTable1R.pdf](#)