Visceral Adipose Tissue Promotes Pressure-Induced Heart Failure Associated with Circulating Fatty Acids

Wenjiao Shi

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Yuanzhi Peng

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Jianer Du (✉ dujianer@xinhumed.com.cn)

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Xiaojian Weng (✉ wengxiaojian@xinhumed.com.cn)

Shanghai Jiaotong University School of Medicine Xinhua Hospital

Research

Keywords: Heart failure, visceral adipose tissue, arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid

Posted Date: November 30th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-76236/v3>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Visceral adipose tissue (VAT) is the main source of circulating fatty acids (FAs) that provides the energy substrate for the heart. Till now, studies have not shown a clear association between individual circulating FAs and heart failure (HF). In this study, we aimed to investigate the changes in circulating FAs in HF mice and their association with VAT by removing epididymal white adipose tissue (eWAT). Here, we found that the serum levels of four fatty acids, namely arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid, were significantly decreased in pressure-induced HF mice via transverse aortic constriction (TAC) surgery accompanied with cardiac enlargement and fibrosis. Most importantly, removal of eWAT in mice led to a significant decrease in the levels of the above-mentioned fatty acids without any significant difference between the HF and sham groups. Accordingly, cardiac enlargement and fibrosis were significantly alleviated. We concluded that VAT excision alleviated TAC-induced cardiac failure by decreasing serum arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels.

1. Introduction

Heart failure (HF) is a major public health issue prevalent in approximately 2% of the population in Western countries and is associated with high morbidity and mortality rates despite pharmacological and non-pharmacological treatment [1]. Evidence suggests that development of HF is associated with abnormalities in cardiac energy metabolism [2]. The failing heart shows an energy deficit with lower levels of adenosine triphosphate (ATP) than does a normal healthy heart, probably because of altered energy substrate availability and impaired mitochondrial oxidative capacity [3]. In the healthy adult heart, the majority of ATP is produced by mitochondrial oxidation of fatty acids. Long-chain fatty acids are crucial oxidizable metabolic substrates for the heart tissue. After esterification into phospholipids, long-chain fatty acids work as building blocks of cellular membranes and participate in associated signal pathways [4]. However, circulating long-chain FAs are heterogeneous, and the specific FAs mainly utilized in cardiomyocyte metabolism during HF development remains unknown.

Changes in the cardiac environment cause cardiac cells to release regulatory peptides which have pathophysiological roles in organs distal to the heart, such as adipose tissue, muscle, and liver, where cell death, growth, fibrosis, and remodeling are regulated [5, 6]. Natriuretic peptides are regulatory peptides released from the heart. In addition to mediating natriuresis, diuresis, and vasodilation, natriuretic peptides enhance lipolysis and lipid mobilization [7, 8]. Moreover, it has also been reported that these natriuretic peptides induce changes in white adipose tissue (WAT) depots promoting the development of brown adipose tissue (BAT) characteristics (such as increased thermogenesis and energy expenditure), a phenomenon termed browning or beiging [9]. These findings provide further evidence of crosstalk between the heart and the peripheral adipose tissues. VAT is not only responsible for lipid storage by lipogenesis, but also for breakdown of fatty acids (FAs) by lipolysis, consequently generating substrates for energy metabolism via β -oxidation [10]. FAs derived from VAT via lipolysis mainly serve as oxidizable

substrates for the heart, and may be responsible for HF. We hypothesized that surgically removing VAT may decrease some kinds of circulating FAs levels, thus preventing HF.

In this study, we aimed to investigate the circulating long-chain fatty acids levels in pressure-induced cardiac failure models to elucidate the potential FAs mainly utilized in cardiomyocyte metabolism. In addition, we examined whether VAT excision could alleviate pressure-induced HF through changes in the levels of some FAs. This study will provide a new idea for the prevention and treatment of heart failure in the future.

2. Materials And Methods

2.1 Animals

Male C57BL/6J wild type mice were purchased from Slac Laboratory Animals (Shanghai, China), and were studied at 10 weeks of age. Animals were housed at 22°C with a 12-h light/12-h dark cycle with free access to water and standard chow. The study conforms to the relevant guidelines and regulations of the Bio-X Institutes of Shanghai Jiao Tong University (Shanghai, China) and was performed after securing approval from the Animal Care and Use Committee of Shanghai Xinhua Hospital affiliated to Shanghai Jiao Tong University School of Medicine (Shanghai, China).

2.2 Experimental models

Ten weeks old C57BL/6J male mice (20-25g) were divided into two groups. In one group, mice were anesthetized with 80-100 mg/kg ketamine and 5-10 mg/kg xylazine intraperitoneally, and then the epididymal white adipose tissue (eWAT) and the main visceral adipose tissue (VAT) were removed. The other group served as normal controls with eWAT reserved. Both groups were then randomly divided into sham and heart failure (HF) groups. The HF mice were established by transverse aortic constriction (TAC). Briefly, mice were anesthetized with sodium pentobarbital (50 mg/kg intraperitoneally), and after opening the chest wall, the thoracic aorta was identified after blunt dissection through the intercostal muscles. Silk suture (7-0) was placed around the transverse aorta and tied around a 26-gauge blunt needle, which was subsequently removed. Sham-operated mice underwent a similar surgical procedure without aortic constriction. After 11 weeks, cardiac structure and function were determined by echocardiography. Mice were then euthanized under isoflurane anesthesia by cervical dislocation, and the hearts were weighed and collected for further analysis.

2.3 Echocardiography

Echocardiography was performed 11 weeks after TAC under the guidance of M-shaped curves of two-dimensional ultrasonic echocardiogram. Every 3 continuous means of the cardiac cycle were selected as the test value. Some cardiac diameters such as fractional shortening (FS), diastolic left ventricular posterior wall (LVPWd), and diastolic left ventricular internal diameter (LVIDd) were quantified as previously described [11]

2.4 Processing of specimens

All mice were weighed before opening the chest wall. The heart was weighed after being taken out, leaving the left ventricle intact (including the inter-ventricular septum) for further weighing. Then, the weight index of the left ventricle (weight of the left ventricle/the body weight) was calculated. Mouse serum was isolated from whole blood, and blood samples were incubated for 60 min at room temperature to induce coagulation. Afterwards, samples were centrifuged at 5000 rpm for 5 min at 4°C and the supernatant (serum) was collected and stored at -80°C until further analysis.

2.5 Hematoxylin and eosin/Masson's staining

Heart tissue samples were formalin-fixed, paraffin-embedded and stained with Hematoxylin/Eosin (HE) and Masson. Cardiac myocyte cross sectional area was assessed using HE-stained and Masson-stained sections of myocardium. After staining, the myocytes were cut into transverse sections and the perimeter of the cell borders of 50 myocytes were measured using CellSens (Olympus). Myocytes were selected in up to ten 400x microscopic fields each separated into 20 equally sized squares. Only one transversally myocyte per square was measured to increase representativeness.

2.6 Quantitative real time PCR (qRT-PCR)

Total RNA was isolated from heart samples with TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.), purified with the RNEasy Kit (Qiagen, Hilden, Germany), and reverse transcribed using the oligo (dT) primers with the Transcriptor First Strand cDNA Synthesis kit (cat. no. 04896866001; Roche diagnostics, Shanghai, china), as previously described [12]. Samples were analyzed for mRNA levels using SYBR-green Master Mix (cat. no. 04887352001; Roche diagnostics), and amplification was monitored using the CF X96 Real-Time PCR system (Bio-Rad, Munich, Germany) with the following thermocycling conditions: 5 s at 95°C followed by 30 s at 60°C for 42 cycles. The reaction mixture was composed of 10 µl SYBR premix Ex Taq II, 0.2 µl primers, 8.8 µl H₂O, and 1 µl DNA. Data were normalized to the expression levels of *GAPDH* (internal control) and were presented as arbitrary units normalized to wild type expression levels. Primer sequences in the present study are presented in **Table 1**.

2.7 Analysis of fatty acids

Mouse serum was isolated from whole blood. For FAs serum profiling, 100 µl of murine serum was hydrolyzed under alkaline-methanolic conditions, neutralized and diluted in methanol (1:10) containing internal FA standards. Analysis of fatty acids was performed using HPLC (Agilent 1200 HPLC system), coupled with an Agilent 6460 triple quad mass was performed, as previously described [13]. Serum concentration of FFA was measured using a HR-NEFA kit (WAKO), as previously described [13]. Triglycerides (TG) in serum were determined using a triglyceride assay kit (DiaSys GmbH), according to the manufacturer's instructions.

2.8 Statistical analysis

Quantitative data were expressed as mean $\pm$ standard error of the mean (SEM). Comparison of mean values between groups was evaluated by the following tests: two-way ANOVA followed by post Bonferroni tests, two-way repeated measures ANOVA followed by post Bonferroni tests, one-way ANOVA followed by Tukey's or Bonferroni multiple comparison test, or unpaired t tests, as appropriate, and data were analyzed with GraphPad Prism software. Exact value of number is provided for each type of experiments. Statistical significance was assumed at $P < 0.05$. Vertical lines in the histograms indicate standard error of the mean (SEM).

3. Results

3.1 Removal of eWAT attenuates TAC-induced cardiac enlargement

To further evaluate the cardiac structure changes, mice were subjected to 11 weeks of pressure overload by transverse aortic constriction (TAC), or sham operations (Sham). The heart weights and cardiac characteristics are presented in Fig 1. Heart weight-to-body weight ratios (HW/BW) and weight index of the left ventricle (LW/BW) were significantly increased in TAC mice compared to sham mice ($P < 0.01$), however, the ratios were significantly decreased in TAC mice with eWAT excision compared to TAC mice with eWAT reserved ($P < 0.05$). Echocardiographic measurements revealed that LVPWd and LVIDd were significantly increased ($P < 0.05$) while FS% was notably decreased ($P < 0.01$) in TAC mice compared to sham mice. However, LVPWd showed a more pronounced decrease in eWAT excised mice compared to eWAT reserved mice following TAC ($P < 0.05$). LVIDd diameters were not significantly altered between eWAT excision and eWAT reserved mice undergoing TAC. In addition, FS% was significantly increased in TAC mice with eWAT reserved compared to eWAT excised ($P < 0.05$). These data showed that there was certain improvement in myocardial hypertrophy and cardiac efficiency after the intervention.

3.2 Removal of eWAT changes cardiac morphology in TAC-induced HF mice

The cardiac morphological changes are presented in Fig 2. The size of the hearts increased markedly in size and weight compared to sham group at 11 weeks after TAC. TAC-induced cardiac enlargement was attenuated after eWAT excision. Accordingly, the myocardial area in cardiac cross section increased significantly in TAC-induced mice whereas this effect was diminished after eWAT excision. Cardiac fibrosis was significantly induced during pressure overload in HF mice as shown by Masson staining. In the TAC group, myocyte hypertrophy and fiber derangement were found with a significant amount of collagen piling in the intercellular space. The above changes were effectively alleviated after eWAT removal.

3.3 Removal of eWAT decreases mRNA expression of *ANP*, *COL1*, and *COL3A1* in pressure-induced HF mice

In keeping with the morphological changes described above, marker genes for cardiac failure such as the *ANP*, *BNP*, *MYH7*, *CTGF*, *COL1*, and *COL3A1* were markedly induced by TAC in WT-hearts (Fig. 3). Cardiac hypertrophy was significantly induced during pressure overload in HF-mice as shown by augmented

mRNA expression of *ANP* ($P<0.01$), *BNP* ($P<0.01$), and *MYH7* ($P<0.01$). Cardiac fibrosis was significantly induced during pressure overload in HF-mice as shown by augmented mRNA expression of the fibrotic marker genes *CTGF* ($P<0.01$), *COL1* ($P<0.01$) and *COL3A1* ($P<0.01$). However, compared to TAC-induced HF mice with eWAT reserve, we observed a significant decrease in mRNA levels of *ANP* ($P<0.01$), *COL1* ($P<0.05$) and *COL3A1* ($P<0.05$) by TAC in eWAT excision mice. In addition, we found moderate decrease in mRNA levels of *BNP*, *MYH7*, and *CTGF* in eWAT excision mice induced by TAC, whereas the result was not statistically significant. Together, these data show that deletion of eWAT protects the heart against pressure-induced cardiac hypertrophy and fibrosis.

3.4 Removal of eWAT attenuates circulating arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels in pressure-induced HF mice

To recognize potential FAs involved in the regulation of cardiac function in pressure-induced HF mice, we examined 24 kinds of FAs, triglycerides (TGs), and free circulating FAs (FFA) in serum. As depicted in TAC and sham groups presented in Fig. 4, in eWAT reserve mice, pressure overload resulted in a significant reduction of selected circulating FAs including arachidic acid ($P<0.001$), behenic acid ($P<0.001$), lignoceric acid ($P<0.01$), and docosapentaenoic acid ($P<0.05$) after TAC (Fig. 4K, R, V, U). The serum TG level was significantly increased ($P<0.05$). In addition, the serum FFA levels were not different between the two groups. Next, we measured whether the lack of eWAT regulated the release of these preferential FAs in a selective manner. We found that in sham and TAC groups, eWAT excision significantly attenuated the circulating arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels compared to eWAT reserve groups ($P<0.05$). But the serum TG and FFA were not different between the two groups. Most importantly, after excision of eWAT, the circulating arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels were not significantly different between the sham and TAC groups. Taken together, pressure-induced cardiac damage was accompanied by changes in circulating FAs among which arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid displayed the most apparent variation, and these changes can be reversed by deletion of eWAT.

4. Discussion

In this study, we aimed to investigate the serum FAs in pressure-induced HF mice with eWAT reserved and eWAT excised. Mice without aortic constriction that underwent a similar surgical procedure served as controls. We demonstrated for the first time that from among multitudinous serum FAs, arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels were significantly decreased in pressure-induced cardiac failure model. In addition, removal of eWAT significantly alleviated cardiac structure and function in HF mice accompanied by a decrease in these four serum FAs levels, indicating non-cardiac tissues such as eWAT participate in the regulation of cardiac failure.

The role of WAT in HF development and cardiac metabolism has been previously studied, with most studies focusing mainly on cardiac lipase function [14, 15]. In this study, we concentrated on VAT and the impact of adipose tissue lipolysis on the development of pressure-induced cardiac failure. This study

revealed that the serum triacylglycerol (TG) level was significantly increased in pressure-induced HF mice compared to controls. Our findings corroborated previous studies that reported a positive association between increased adipose tissue lipolysis and impairment of cardiac function in heart failure models [16, 17]. Circulating FAs are liberated from WAT via hydrolysis of TG which is catalyzed by two major adipose tissue lipases, hormone-sensitive lipase (HSL) and adipose triglyceride lipase (ATGL), under hormonal control [18]. Deletion of ATGL in adipose tissue prevents pressure-induced LV failure by decreasing adipose tissue lipolysis [19]. Consistent with previous studies, our findings showed that removal of eWAT significantly alleviated pressure-induced cardiac hypertrophy and fibrosis, as evidenced by decreased mRNA expression of cardiac hypertrophy and fibrosis genes *ANP*, *COL 1*, and *COL3A1*.

Fatty acids are involved in post-translational modification of proteins through protein acylation and activation of protein kinases. Recent experimental findings have indicated that several cardiac genes are under the control of fatty acids [20, 21]. However, there are multiple FAs present in circulating bloodstream, and it is not clear which of them are mainly involved in cardiac energy metabolism. We analyzed serum FAs using HPLC, which was previously utilized in cardiovascular research for the identification of new molecular co-mediator of exercise-induced cardiac hypertrophy [13]. Our study found that serum arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels were significantly decreased in pressure-induced cardiac injury. Arachidic acid, behenic acid, and lignoceric acid are long-chain saturated FAs that are found in circulation. Circulating long-chain saturated FAs with 20 carbons or more are integrated biomarkers of diet and metabolism [22]. The long-chain saturated FAs are known components of ceramides that are involved in apoptosis and cardiac dysfunction [23, 24]. The heart of genetically engineered mice with reduced ceramides containing long-chain saturated FAs showed increased fibrosis, endoplasmic reticulum stress, and apoptosis [25]. Lemaitre *et al*/ have identified an association of higher levels of plasma phospholipid very-long-chain saturated fatty acids with lower risk of incident HF in a prospective cohort of older American adults [22]. Similar results were obtained in our study, indicating that decreased arachidic acid, behenic acid, and lignoceric acid were probably involved in pressure-induced cardiac failure. In addition, we also found that the serum level of docosapentaenoic acid, a long-chain unsaturated fatty acid, was significantly decreased in pressure-induced HF mice. In addition, we found no increased levels of FAs in serum of pressure-induced heart failure mice. This eliminated the probability that these reduced fatty acids were converted to other fatty acids. Thus, we hypothesized that these four fatty acids are involved in cardiac function impairment during pressure overload. However, this needs to be verified by analyzing lipid class in cardiac tissue through further studies.

The current study found that mice after removal of eWAT showed a marked reduction of these four fatty acids in both normal and pressure-induced HF mice compared to mice with eWAT reserved, which indicated that these four fatty acids may be principally derived from eWAT. In addition, we observed a slightly, but not significantly, decreased TG and FFA levels. eWAT serves as a main VAT, which can more easily produce FFA than subcutaneous adipose tissue (SAT) because VAT has higher expression of adrenergic receptors 1, 2, and 3 on the cell membrane [26, 27]. Surprisingly, no significant differences were found in circulating FFA and TG after eWAT removal in this study. One possible explanation for this

might be that VST is mainly distributed around abdominal organs including epididymal adipose tissue, omental adipose tissue, mesenteric adipose tissue, retroperitoneal adipose tissue, perirenal adipose tissue, etc [28]. Although eWAT was removed in this study, other visceral adipose tissues may have also played a role in producing FFA. Another possible explanation might be because of the difference in the experimental period selected in this study. There are, however, other possible explanations that need to be further studied. The mechanisms underlying the differential impact of adipose tissue on cardiometabolic risk have started to be unraveled; the inability of the subcutaneous adipose tissue to expand in response to positive energy balance serves as an important mechanism that necessitates further studies on the subcutaneous adipose tissue.

5. Conclusion

Taken together, the results of this study demonstrate for the first time that serum arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels are significantly decreased in pressure-induced heart failure. Furthermore, we showed that the decreased levels of these four FAs can be significantly reversed by removing eWAT from mice before inducing HF. However, whether these four fatty acids accumulate in cardiomyocytes remains unknown. Further research should be carried out to explore the accumulation of these fatty acids in cardiomyocytes and the damage to the mitochondrial function of cardiomyocytes, so as to provide new therapeutic targets for clinical treatment of heart failure.

Declarations

Data availability statement

The data used to support the findings of this study are available with the corresponding author upon request.

Funding statement

This work was supported by the National Natural Science Foundation of China (grant number 81901990).

Declaration of Competing Interests

The authors declare that there are no conflicts of interest.

Acknowledgement

We declare the manuscript has been submitted as preprint on Research Square. A draft version of our preprint is now available privately. In addition, the authors would like to thank all the colleagues who contributed to this study. This study was supported by a grant from the National Natural Science Foundation of China (grant number 81901990).

References

1. Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail*, 2016. 18(8): 891-975.
2. Voros G, Ector J, Garweg C, et al. Increased Cardiac Uptake of Ketone Bodies and Free Fatty Acids in Human Heart Failure and Hypertrophic Left Ventricular Remodeling. *Circ Heart Fail*, 2018. 11(12): e004953.
3. Conway MA, Allis J, Ouwerkerket R, et al. Detection of low phosphocreatine to ATP ratio in failing hypertrophied human myocardium by ³¹P magnetic resonance spectroscopy. *Lancet*, 1991. 338(8773): 973-976.
4. Vusse GJ, Bilsen M, Glatz JF. Cardiac Fatty Acid Uptake and Transport in Health and Disease. *Cardiovasc Res*, 2000. 45(2): 279-93.
5. Loncar G, Fülster S, Haehling SV, Popovic V. Metabolism and the heart: An overview of muscle, fat, and bone metabolism in heart failure. *Int J Cardiol*, 2013. 162(2): 77–85.
6. Nakamura, M. and J. Sadoshima, Heart over mind: metabolic control of white adipose tissue and liver. *Embo Molecular Medicine*, 2014. 6(12): 1521-1524.
7. Sengenès C, Berlan M, Gliszinski ID, et al. Natriuretic peptides: a new lipolytic pathway in human adipocytes. *FASEB J*, 2000. 14(10): 1345-51.
8. Moro C, Crampes F, Sengenès C, et al. Atrial natriuretic peptide contributes to physiological control of lipid mobilization in humans. *FASEB J*, 2004. 18(7): 908-10.
9. Bordicchia M, Liu D, Amri EZ, et al. Cardiac natriuretic peptides act via p38 MAPK to induce the brown fat thermogenic program in mouse and human adipocytes. *J Clin Invest*, 2012. 122(3): 1022-36.
10. Bodis, K. and M. Roden, Energy metabolism of white adipose tissue and insulin resistance in humans. *Eur J Clin Invest*, 2018. 48(11): e13017.
11. Kawahara Y, Tanonaka K, Daicho T, et al. Preferable Anesthetic Conditions for Echocardiographic Determination of Murine Cardiac Function. *J Pharmacol Sci*, 2005. 99(1): 95-104.
12. Shi W, Guo ZX, Ji Y, et al. The protective effect of recombinant globular adiponectin on testis by modulating autophagy, endoplasmic reticulum stress and oxidative stress in streptozotocin-induced diabetic mice. *Eur J Pharmacol*, 2020. 879: 173132.
13. Foryst-Ludwig A, Kreissl MC, Benzet V, et al. Adipose Tissue Lipolysis Promotes Exercise-induced Cardiac Hypertrophy Involving the Lipokine C16:1n7-Palmitoleate. *J Biol Chem*, 2015. 290(39): 23603-23615.
14. Zierler KA, Jaeger D, Pollaket NM, et al, Functional cardiac lipolysis in mice critically depends on comparative gene identification-58. *J Biol Chem*, 2013. 288(14): 9892-904.
15. Kienesberger PC, Pulinilkunnit T, Sunget MMY, et al. Myocardial ATGL overexpression decreases the reliance on fatty acid oxidation and protects against pressure overload-induced cardiac dysfunction. *Mol Cell Biol*, 2012. 32(4): 740-50.

16. Polak J, Kotrc M, Wedellova Z, et al. Lipolytic Effects of B-Type Natriuretic Peptide1–32 in Adipose Tissue of Heart Failure Patients Compared With Healthy Controls. *J Am Coll Cardiol*, 2011. 58(11): 1119-1125.
17. Lommi J, Kupari M, Yki-Järvinen H. Free Fatty Acid Kinetics and Oxidation in Congestive Heart Failure. *Am J Cardiol*, 1998. 81(1):45-50.
18. Young, S.G. and R. Zechner, Biochemistry and pathophysiology of intravascular and intracellular lipolysis. *Genes Dev*, 2013. 27(5): 459-484.
19. Salatzki J, Foryst-Ludwig A, Bentele K, et al. Adipose tissue ATGL modifies the cardiac lipidome in pressure-overload-induced left ventricular failure. *PLoS Genet*, 2018. 14(1): e1007171.
20. Bilsen, M.V., J.E.D. Vries, and G.J.V.D. Vusse, Long-term effects of fatty acids on cell viability and gene expression of neonatal cardiac myocytes. *Prostaglandins Leukot Essent Fatty Acids*. 1997. 57(1):39-45.
21. Brandt, J.M., F. Djouadi, and D.P. Kelly, Fatty Acids Activate Transcription of the Muscle Carnitine Palmitoyltransferase I Gene in Cardiac Myocytes via the Peroxisome Proliferator-activated Receptor α . *J Biol Chem*, 1998. 273(37): 23786-23792.
22. Lemaitre RN, McKnight B, Sotoodehnia N, et al. Circulating Very Long-Chain Saturated Fatty Acids and Heart Failure: The Cardiovascular Health Study. *J Am Heart Assoc*, 2018. 7(21): e010019.
23. Ohno Y, Suto S, Yamanaka M, et al. ELOVL1 production of C24 acyl-CoAs is linked to C24 sphingolipid synthesis. *Proc Natl Acad Sci U S A*, 2010. 107(43): 18439-44.
24. Park TS, Hu Y, Noh HL, et al. Ceramide is a cardiotoxin in lipotoxic cardiomyopathy. *J Lipid Res*, 2008. 49(10): 2101-2112.
25. Lee SY, Kim JR, Hu Y, et al. Cardiomyocyte Specific Deficiency of Serine Palmitoyltransferase Subunit 2 Reduces Ceramide but Leads to Cardiac Dysfunction. *J Biol Chem*, 2012. 287(22): 18429-18439.
26. Arner P, Hellström L, Wahrenberg H, Brönnegård M. Beta-adrenoceptor expression in human fat cells from different regions. *J Clin Invest*. 1990. 86(5): 1595-600. .
27. Karpe, F. and K.E. Pinnick, Biology of upper-body and lower-body adipose tissue–link to whole-body phenotypes. *Nat Rev Endocrinol*, 2015. 11(2): 90-100.
28. Wajchenberg BL. Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr Rev*. 2000. 21(6): 697-738..

Table

Table. 1 Primers used in real-time fluorescence quantitative PCR

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
<i>ANP</i>	AGATCTGCCCTCTTGAAAAGCA	TCGAGCAGATTTGGCTGTTATC
<i>BNP</i>	TCCAGAACAATCCACGATGC	GCAGCTTGAACATATGTGCCATC
<i>MYH7</i>	GCGGACATTGCCGAGTCCAG	GCTCCAGGTCTCAGGGCTTCACA
<i>CTGF</i>	GGACACGAACTCATTAGAC	TCTCACTTTGGTGGGATAG
<i>COL1</i>	TTCTCCTGGCAAAGACGGAC	CGGCCACCATCTTGAGACTT
<i>COL3A1</i>	AGCGGCTGAGTTTATGACG	AGCACAGGAGCAGGTGTAGA
<i>GAPDH</i>	CATGACAACCTTTGGCATCGT	GGATGCAGGGATGATGTTCT

ANP: atrial natriuretic peptide; *BNP*: brain natriuretic peptide; *COL1*: collagen 1; *COL3A1*: collagen 3a1; *GAPDH*: glyceraldehyde 3-phosphate dehydrogenase.

Figures

Figure 1

Changes in heart weight and cardiac structure in the four groups. A: Representative M-Mode images of the echocardiographic analysis. B and C: Representative images of the hearts. D: Heart weight (HW)/ body weight (BW) ratio (mean and SEM, n = 8). E: Left ventricle weight (LW)/ body weight (BW) ratio (mean and SEM, n = 8). F: Left ventricular posterior wall in diastol (LVPW; d) (mean and SEM, n = 8). G: Left-ventricular internal diameter in diastole (LVID; d) (mean and SEM, n = 8). H: Fractional shortening [%] (FS) (mean and SEM, n = 8). sham: normal control group. TAC: transverse aortic constriction induced heart failure group. eWAT: epididymal white adipose tissue. *P<0.05 vs. sham + eWAT reserve, **P<0.01 vs. sham + eWAT reserve, #P<0.05 vs. TAC + eWAT reserve, ##P<0.01 vs. TAC + eWAT reserve; two-way ANOVA (Bonferroni post-test).

Figure 1

Changes in heart weight and cardiac structure in the four groups. A: Representative M-Mode images of the echocardiographic analysis. B and C: Representative images of the hearts. D: Heart weight (HW)/body weight (BW) ratio (mean and SEM, n = 8). E: Left ventricle weight (LW)/body weight (BW) ratio (mean and SEM, n = 8). F: Left ventricular posterior wall in diastol (LVPW; d) (mean and SEM, n = 8). G: Left-ventricular internal diameter in diastole (LVID; d) (mean and SEM, n = 8). H: Fractional shortening [%]

(FS) (mean and SEM, n = 8). sham: normal control group. TAC: transverse aortic constriction induced heart failure group. eWAT: epididymal white adipose tissue. *P<0.05 vs. sham + eWAT reserve, **P<0.01 vs. sham + eWAT reserve, #P<0.05 vs. TAC + eWAT reserve, ##P<0.01 vs. TAC + eWAT reserve; two-way ANOVA (Bonferroni post-test).

Figure 2

Cardiac morphological changes in the four groups. A: Representative microscopic cross-sections of the hearts stained with hematoxylin/ eosin (H/E). B: Representative microscopic cross-sections of the hearts stained with Masson. sham: normal control group. TAC: transverse aortic constriction induced heart failure group. eWAT: epididymal white adipose tissue.

Figure 2

Cardiac morphological changes in the four groups. A: Representative microscopic cross-sections of the hearts stained with hematoxylin/ eosin (H/E). B: Representative microscopic cross-sections of the hearts stained with Masson. sham: normal control group. TAC: transverse aortic constriction induced heart failure group. eWAT: epididymal white adipose tissue.

Figure 3

Removal of eWAT decreases mRNA expression of ANP, COL1, and COL3A1 in TAC-induced HF mice. The mRNA expression of ANP, BNP, MYH7, CTGF, COL1, and COL3A1 were detected by real-time fluorescence quantitative polymerase chain reaction (RT-PCR) analysis. Relative gene expression levels were normalized to that of GAPDH. Data are presented as x-fold over sham mice (mean and SEM, n = 8). Statistical significance was determined using two-way analysis of variance (*P < 0.05 vs. sham + eWAT reserve, **P < 0.01 vs. sham + eWAT reserve, #P < 0.05 vs. TAC + eWAT reserve, ##P < 0.01 vs. TAC + eWAT reserve). ANP: Atrial natriuretic peptide; BNP: Brain natriuretic peptide; sham: normal control group. TAC: transverse aortic constriction induced heart failure group. eWAT: epididymal white adipose tissue.

Figure 3

Removal of eWAT decreases mRNA expression of ANP, COL1, and COL3A1 in TAC-induced HF mice. The mRNA expression of ANP, BNP, MYH7, CTGF, COL1, and COL3A1 were detected by real-time fluorescence quantitative polymerase chain reaction (RT-PCR) analysis. Relative gene expression levels were normalized to that of GAPDH. Data are presented as x-fold over sham mice (mean and SEM, n = 8). Statistical significance was determined using two-way analysis of variance (* $P < 0.05$ vs. sham + eWAT reserve, ** $P < 0.01$ vs. sham + eWAT reserve, # $P < 0.05$ vs. TAC + eWAT reserve, ## $P < 0.01$ vs. TAC + eWAT reserve). ANP: Atrial natriuretic peptide; BNP: Brain natriuretic peptide; sham: normal control group. TAC: transverse aortic constriction induced heart failure group. eWAT: epididymal white adipose tissue.

Figure 4

Removal of eWAT attenuates circulating arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels. A-X: Profile of selected serum FAs analyzed by rapid resolution HPLC/Tandem MS. Y: Analysis of serum triglycerides (TG). Z: Analysis of serum free fatty acids (FFA). Data are presented as x-fold over sham mice (mean and SEM, n = 8). Statistical significance was determined using two-way analysis of variance (*P<0.05, **P<0.01, ***P<0.001).

Figure 4

Removal of eWAT attenuates circulating arachidic acid, behenic acid, lignoceric acid, and docosapentaenoic acid levels. A-X: Profile of selected serum FAs analyzed by rapid resolution HPLC/Tandem MS. Y: Analysis of serum triglycerides (TG). Z: Analysis of serum free fatty acids (FFA). Data are presented as x-fold over sham mice (mean and SEM, n = 8). Statistical significance was determined using two-way analysis of variance (*P<0.05, **P<0.01, ***P<0.001).