

Elastography for the Diagnosis of High-Suspicion Thyroid Nodules Based on the 2015 American Thyroid Association Guidelines: A Multicenter Study

Hairu Li

Second Affiliated Hospital of Harbin Medical University

Yulan Peng

Sichuan University West China Hospital

Yan Wang

Sixth People's Hospital Affiliated to Shanghai Communication University

Hong Ai

First Affiliated Hospital of Xi'an Communication University

Xiaodong Zhou

Xijing Hospital Affiliated to The Fourth Military Medical University

Lichun Yang

Third Affiliated Hospital of Kunming Medical University

Kun Yan

Tumor Hospital of Beijing University

Ying Xiao

Xiangya Hospital of Center-south University

Lisha Liu

Tumor Hospital Affiliated to Xinjiang Medical University

Baoming Luo

Sun Yat-sen Memorial Hospital of Sun Yat-sen University

Qiang Yong

Anzhen Hospital Affiliated to Capital Medical University

Shuzhen Cong

People's Hospital of Guangdong Province

Shuangquan Jiang

Second Affiliated Hospital of Harbin Medical University

Xin Fu

Second Affiliated Hospital of Harbin Medical University

Buyun Ma

West China Hospital of Sichuan University

Yi Li

Sixth People's Hospital Affiliated to Shanghai Communication University

Xixi Zhang

First Affiliated Hospital of Xi'an Communication University

Xue Gong

Xijing Hospital Affiliated to The Fourth Military Medical University

Haitao Chen

Third Affiliated Hospital of Kunming Medical University

Wenyong Liu

Tumor Hospital of Beijing University

Ling Tang

Xiangya Hospital of Center-south University

Xiaoyu LV

Xinjiang Medical University Affiliated Tumor Hospital

Xinbao Zhao

Sun Yat-sen Memorial Hospital of Sun Yat-sen University

Liang Li

Beijing Anzhen Hospital Affiliated to Capital Medical University

Kehong Gan

People's Hospital of Guangdong Province

Jiawei Tian (✉ jwtian20041@163.com)

Second Affiliated Hospital of Harbin Medical University

Research article

Keywords: Elastography, thyroid, cancer, ultrasound

Posted Date: January 20th, 2020

DOI: <https://doi.org/10.21203/rs.2.17002/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on April 3rd, 2020. See the published version at

<https://doi.org/10.1186/s12902-020-0520-y>.

Abstract

Background: An accurate diagnosis for high-suspicion nodules based on the 2015 American Thyroid Association (ATA) guidelines would reduce unnecessary invasive examinations. Elastography is a useful tool for discriminating benign and malignant thyroid nodules.

Aims: To investigate the diagnostic efficiency of elastography for high-suspicion thyroid nodules based on the 2015 ATA guidelines in the Chinese population.

Materials and Methods: Thyroid nodules with high-suspicion characteristics based on the 2015 ATA guidelines were subjected to conventional ultrasound (US) and ultrasound strain elastography (USE) examinations at 12 hospitals from 4 geographic regions across China. Cytology/histology of thyroid nodules was used as a reference method. Receiver operating characteristic (ROC) curves were plotted to evaluate the diagnostic performance of the elasticity score (ES) and strain ratio (SR). Logistic regression analysis was used to determine the predictors of malignancy.

Results: Overall, a total of 1445 thyroid nodules (834 malignant, 611 benign) from 12 centers were included in the final analysis. The areas under the curve of the ES and SR were 0.828 and 0.732, respectively. The sensitivity, specificity, accuracy, positive predictive value (PPV) and negative predictive value (NPV) of the ES were 92.4%, 60.7%, 79.0%, 76.3% and 85.5%, respectively, and those of the SR were 81.1%, 50.1%, 68.9%, 65.9% and 67.9%, respectively. The combination of the Thyroid Imaging Reporting and Data System (TI-RADS) and ES led to a significant increase in the sensitivity and NPV (97.1% and 91.9%, respectively) compared with the TI-RADS alone. Logistic regression analysis showed that microcalcifications (OR=5.290), taller than wide (OR=12.710), irregular margins (OR=10.117), extrathyroidal extension (ETE; OR=6.412), the ES (OR=3.741) and the SR (OR=1.083) were independent predictors of malignant thyroid nodules. The sensitivity, specificity, accuracy, PPV and NPV of the ES were all superior in nodules ≥ 1 cm than in those < 1 cm (95.0% vs 90.4%, 68.8% vs 56.8%, 85.9% vs 74.4%, 85.2% vs 69.9%, and 87.8% vs 84.2%, respectively).

Conclusions: Elastography combined with the ES is a valuable tool for the assessment of high-suspicion thyroid nodules based on the 2015 ATA guidelines, especially in nodules ≥ 1 cm.

1. Background

Thyroid nodules are common diseases, and their incidence has increased rapidly during the past decades. Thyroid nodules detected by ultrasound (US) have been detected in up to 50% of the general population[1]. Although most nodules are benign, asymptomatic and do not require treatment, a reliable diagnosis is still necessary to achieve the optimal therapeutic schedule[2]. The early detection and differentiation of malignant thyroid nodules from overall thyroid nodules are particularly important for planning treatment and evaluating prognosis. A diagnostic thyroid US examination is recommended for all patients with thyroid nodules according to the American Thyroid Association (ATA) guidelines[3]. Although certain sonographic features, such as taller than wide, blurred margins, irregular borders, internal microcalcifications, hypoechogenicity and marked hypoechogenicity[4], are conventional indicators of malignancy, the sensitivity and specificity vary largely from 38.8% to 90.9% and 53.0% to 96.6%, respectively[5-7].

Ultrasound strain elastography (USE) has been proposed for differentiating malignant thyroid nodules from benign nodules based on their elasticity and has gradually developed into a widely used US examination method[8]. USE produces an elasticity score (ES) as the qualitative output and the strain ratio (SR, the ratio obtained by dividing the mean strain within the lesion by the mean strain of the surrounding normal tissue) as a semiquantitative output in units of relative strain. USE is based on the principle that softer parts deform more easily than harder parts when body tissues are compressed. Studies have shown that USE can differentiate malignant thyroid nodules from benign nodules with high sensitivity and specificity[9, 10]. However, several studies have found contradictory results in which USE had a lower sensitivity and specificity than conventional US[11, 12]. Thus, the efficacy of USE remains controversial, and additional studies are needed to confirm its efficacy.

According to the 2015 ATA guidelines, the malignant risk of high-suspicion thyroid nodules is >70 - 90% [3]. However, the ATA characteristics of high-suspicion thyroid nodules overlap with degrees 4a to 5 of Kwak's Thyroid Imaging Reporting and Data

System (TI-RADS), the malignancy of which ranges from 3.3% to 87.5%[13]. Depending on the discrepancy between the two methods, we performed a nationwide, multicenter study in China to ascertain the efficacy of USE in differentiating benign and malignant high-suspicion thyroid nodules based on the 2015 ATA guidelines. To our knowledge, this is the first multicenter study in China to investigate the diagnostic performance of strain elastography in differentiating malignant from benign thyroid nodules of high suspicion based on the 2015 ATA guidelines in a large population.

2. Methods

2.1 Patients

Between March 2014 and September 2018, 1819 patients with 1903 suspicious nodules at 12 diagnostic centers from 4 geographic regions across China were evaluated by conventional US and USE (Figure 1). The inclusion criteria were solid hypoechoic nodules with one or more of the following features: irregular margins (specifically defined as infiltrative, microlobulated, or spiculated), microcalcifications, taller than wide, disrupted rim calcifications with a small extrusive hypoechoic soft tissue component, or evidence of extrathyroidal extension (ETE)[3]. The exclusion criteria were as follows: cystic nodules or cystic mixed nodules, nodules with a tumor size >3 cm, nodules with eggshell calcifications, and nodules located in the isthmus. In total, 1445 thyroid nodules were classified as high-suspicion nodules and included in the final analysis. The male/female ratio was 304/1141. The mean age of the patients with benign nodules was 48.30 ± 10.86 years, and the mean age of the patients with malignant nodules was 42.49 ± 11.39 years. The ethics committee of each hospital approved this study. Written or oral consent was obtained from all patients.

2.2 Equipment

All studies were performed using the same type of US machine, i.e., HI VISION Ascendus, HI VISION Avius or HI VISION PREIRUS (Hitachi Medical, Tokyo, Japan), which had both conventional B-mode US and elastography capabilities. A 5-13 MHz linear transducer probe was used for conventional US and USE examinations.

2.3 US and USE examinations

Both conventional US and real-time USE were conducted and recorded by two skilled sonographers at each center with at least 5 years of conventional US experience and 1 year of USE experience. Final results were obtained when the results of the two examiners were identical. If the results of the two examiners were inconsistent, then a superior physician was invited to make a joint decision. All examiners were blinded to the results of the pathological diagnosis. The examination method, procedure, and conditions used by the inspectors remained consistent for all patients. For the elastography examination, the USE/US double-image display mode was selected. Once the default thyroid conditions were selected, they were not changed throughout the process of the USE examination. Next, the region of interest (ROI) of USE was selected as follows: the lesion area was no more than 1/3 of the ROI area to ensure that sufficient surrounding normal tissue was available as a reference. The ROI included a small amount of anterior cervical muscle tissue, the thyroid gland, and the thyroid gland capsule, and cervical blood vessels and tracheal tissue were avoided to the greatest extent possible. The ROI was not set within 2 mm of the skin surface.

According to Itoh et al[14], the elastograms of the thyroid lesions were classified into 5 types. A score of 1 indicated even strain throughout the entire hypoechoic lesion (Figure 2a). A score of 2 indicated strain in most of the hypoechoic lesion, with certain areas of no strain (Figure 2b). A score of 3 indicated strain at the periphery of the hypoechoic lesion and no strain at the center of the lesion (Figure 2c). A score of 4 indicated no strain throughout the entire hypoechoic lesion (Figure 2d). A score of 5 indicated no strain throughout the entire hypoechoic lesion and surrounding area (Figure 2e).

SR measurement: The operator traced the outline of the lesion in the two-dimensional figure or the elastic image on the longitudinal views. Then, the area adjacent to the target lesion, which was at the same depth as the lesion, was selected as a reference. The SR was automatically calculated by the software.

2.4 Reference standard

Fine-needle aspiration (FNA) was performed as recommended by the 2015 ATA guidelines[3]. Patients with suspicious or malignant cytology were referred for surgery. Some of the patients who underwent thyroidectomy did not receive FNA due to certain reasons, such as highly suspicious malignancy or refused FNA. The patients who underwent surgery were subjected to a pathological section examination, and the pathology results were used as the gold standard. Benign nodules confirmed by FNA were followed up for 6-12 months and without lesion growth, i.e., an increase in the diameter less than 20% or an increase in the volume less than 50%. Cytology and histology were performed by experienced local pathologists with at least 5 years of working experience who were blinded to the US and USE results.

2.5 Statistical analysis

The measurement data are expressed as the mean \pm standard deviation. A t test was used for comparisons between the two groups. The counting data are presented as cases (percentages), and the chi-square test was used for comparisons between the two groups. Logistic regression analysis was used for risk factor analysis of both benign and malignant nodules. ROC curves and AUC values were used to examine the diagnostic effect of each index on diseases. All statistical tests were performed on both sides, and $p < 0.05$ was considered statistically significant.

3. Results

3.1 Clinicopathologic data

A total of 1819 patients with 1903 nodules were enrolled from 12 centers in this multicenter study. One hundred and twenty-six patients were excluded due to the absence of adequate data on USE, cytology or histology results or were lost to follow-up, and 332 nodules were not high-suspicion nodules. Therefore, the final analysis comprised 1445 thyroid nodules, including 834 malignant nodules and 611 benign nodules. All 834 malignant nodules and 190 benign nodules were confirmed by surgery, and the remaining 421 benign nodules were confirmed by a cytologic examination. Of the 834 malignant nodules, 690 were papillary carcinomas, 133 were papillary microcarcinomas, 7 were follicular carcinomas, and 4 were medullary carcinomas. The final diagnosis of benign histology was adenoma in 38 nodules, nodular goiter in 121 nodules, follicular nodule in 9 nodules, chronic lymphocytic thyroiditis in 13 nodules, subacute granulomatous thyroiditis in 3 nodules, hyperplasia of the thyroid tissue in 1 nodule, inflammatory lesion in 4 nodules and tuberculosis in 1 nodule.

3.2 Diagnostic performance of conventional US

The results of the diagnostic performance of conventional US and USE are shown in Table 1. All of the parameters were significantly different between the two groups ($p < 0.005$). The sensitivity, specificity, accuracy, positive predictive value (PPV) and negative predictive value (NPV) were calculated for each suspicious criterion. Irregular margins and ETE had a high sensitivity (93.9% and 83.0%, respectively) but very low specificity (14.1% and 23.2%, respectively). Taller than wide and disrupted rim calcifications with a small extrusive hypoechoic soft tissue component had a high specificity (91.0% and 98.9%, respectively) but low sensitivity (37.1% and 4.7%, respectively). The sensitivity and specificity of microcalcifications were moderate (49.9% and 65.6%, respectively). As Kwak's TI-RADS is the mainstream classification of malignant degree, we analyzed the diagnostic performance of Kwak's TI-RADS. Kwak's TI-RADS score showed a sensitivity of 62.2%, a specificity of 70.2%, a PPV of 74.0%, an NPV of 57.7% and an AUC under the ROC curve of 0.69 in the overall assessment (Table 1, Figure 3).

3.3 Diagnostic performance of USE

There were 21 nodules with an ES of 1 (20 benign, 1 malignant), 198 nodules with an ES of 2 (181 benign, 17 malignant), 215 nodules with an ES of 3 (170 benign, 45 malignant), 473 nodules with an ES of 4 (169 benign, 304 malignant), and 538 nodules with an ES of 5 (71 benign, 467 malignant). There were significant differences in the ES and SR values between the malignant and benign groups ($p < 0.001$). The malignant percentage increased with the increase in ES, with values of 4.8%, 8.6%, 20.9%, 64.3% and 86.8%, respectively.

The mean (\pm SD) ES of the malignant and benign nodules was 3.15 ± 1.072 and 4.46 ± 0.703 , respectively ($p < 0.001$). Figure 4 shows the ROC curve of the ES and SR in differentiating malignant and benign nodules. The AUC of the ES was 0.828. The optimal cut-off value of the ES was 3.5 in this study. The sensitivity, specificity, accuracy, PPV and NPV of the ES were 92.4%, 60.7%, 79.0%, 76.3%

and 85.5%, respectively. The mean (\pm SD) SR of the malignant and benign nodules was 3.26 ± 2.06 and 6.15 ± 7.21 , respectively ($p<0.001$). The AUC of the SR was 0.732. The optimal cut-off value of the SR was 2.99. The sensitivity, specificity, accuracy, PPV and NPV of the SR were 81.1%, 50.1%, 67.9%, 68.9% and 65.9%, respectively (Table 1). The combination of the TI-RADS with ES led to a significant increase in the sensitivity, NPV (97.1% and 91.9%, respectively) and the AUC under the ROC curve (0.819) (Figure 3).

3.4 Logistic regression analysis for predicting malignant thyroid nodules

In the univariate and multivariate logistic regression analyses, microcalcification (OR=1.887 and 5.29, $p<0.001$, respectively), taller than wide (OR=5.950 and 12.71, $p<0.001$, respectively), irregular margins (OR=2.515 and 10.117, $p<0.001$, respectively), the ES (OR=4.473 and 3.741, $p<0.01$, respectively) and the SR (OR=1.504 and 1.083, $p<0.01$, respectively) were statistically significant predictors of malignant thyroid nodules (Table 2). ETE was a predictor of malignant nodules in the univariate analysis but not in the multivariate analysis ($p>0.05$).

3.5 Diagnostic performance of the ES and SR in nodules of different sizes

According to the 2015 ATA thyroid nodule guidelines, nodules ≥ 1 cm in the largest dimension with a high-suspicion sonographic pattern are recommended for FNA; thus, we compared the diagnostic performance of the ES and SR in nodules ≥ 1 cm and <1 cm. There were 877 nodules <1 cm and 568 nodules ≥ 1 cm. The male/female ratios were 180/697 and 124/444, respectively, and the mean age of the patients was 45.58 ± 11.001 years for those with nodules <1 cm and 44.01 ± 12.268 years for those with nodules ≥ 1 cm. The results presented in Table 3 demonstrate that the AUC, sensitivity, specificity, accuracy, PPV and NPV of the ES were all superior in nodules ≥ 1 cm than in those <1 cm (95.0% vs 90.4%, 68.8% vs 56.8%, 85.9% vs 74.4%, 85.2% vs 69.9%, and 87.8% vs 84.2%, respectively), with especially high accuracy. The specificity, accuracy and PPV of the SR were also higher in nodules ≥ 1 cm than in those <1 cm (59.3% vs 45.6%, 72.7% vs 64.8%, and 78.8% vs 62.6%, respectively). The above data indicate that elastography with ES has an excellent diagnostic performance in nodules ≥ 1 cm.

4. Discussions

With the development of high-frequency ultrasonic devices, more thyroid nodules have been diagnosed. Although most thyroid nodules are benign, a diagnosis differentiating malignant from benign nodules is still of vital importance for determining the appropriate therapy. A differential diagnosis is still a difficult problem for clinicians. Elastography has emerged as a valuable technology to assist in the diagnosis of thyroid nodules due to its ability to reflect the stiffness of tissue that is related to malignant lesions. A large number of studies have shown the excellent performance of elastography in the differentiation of malignant and benign nodules[15-17]. Nevertheless, several studies have revealed the opposite result[11, 12]. The present study was the first multicenter study in China with a large sample size to determine the value of real-time strain elastography in the discrimination of malignant from benign thyroid nodules with high-suspicion characteristics based on the 2015 ATA guidelines. The results showed that the combination of USE with the ES had good efficacy for the discrimination of malignant from benign high-suspicion thyroid nodules, with a high sensitivity (92.4%), accuracy (79.0%) and NPV (85.5%), especially in nodules ≥ 1 cm (95.0%, 85.9%, and 87.8%, respectively). Additionally, logistic regression analysis showed that the ES and SR were independent predictors of malignancy.

US is the preferred technology for thyroid nodules due to its convenience, maneuverability and low cost. Microcalcifications, nodule hypoechoogenicity compared with the surrounding thyroid or strap muscles, irregular margins (defined as either infiltrative, microlobulated, or spiculated), and taller than wide measured on a transverse view are traditionally recognized characteristics of malignant thyroid nodules[3]. All the conventional US features were significantly different between malignant and benign nodules in this study. Logistic regression analysis showed that microcalcifications, taller than wide, irregular margins and ETE were independent predictors of malignancy. However, previous studies have reported largely varied sensitivities and specificities[5-7]. The diagnostic efficacy of conventional US parameters in this study was also unsatisfactory. Although irregular margins and ETE yielded good sensitivity, their specificity was rather poor. Similarly, despite their high specificity, the sensitivity of taller than wide and disrupted rim calcifications with a small extrusive hypoechoic soft tissue component was unacceptable. The accuracy of conventional US parameters was also limited.

Elastography takes advantage of the change in elasticity of soft tissues resulting from specific pathological or physiological processes, and the low elasticity observed on USE is highly correlated with malignancy. Since the emergence of elastography, it has shown an outstanding sensitivity, specificity, PPV and NPV[16,18- 20], with the specificity and PPV reaching up to 100%. Elasticity, which is absent in conventional US, is the single feature with the best diagnostic performance[7], as well as a potent predictor of malignant thyroid nodules[21, 22]. With an increase in the ES, the percentage of malignancy increased significantly in this research, confirming that the ES is correlated with malignancy. In this research, we used the 5 scoring parameters proposed by Itoh, who defined an ES >3.5 as the cut-off value; the sensitivity, specificity, accuracy, PPV and NPV were 92.4%, 60.7%, 79.0%, 76.3% and 85.5%, respectively, with an AUC of 0.828. However, contrary opinions have also been presented. A multicenter study by Moon et al reported that elastography showed an inferior performance in the differentiation of malignant and benign thyroid nodules relative to grayscale US features[11]. They investigated all solid nodules, including hyperechogenic, isoechogenic, hypoechogenic and marked hypoechogenic nodules, in their research. In this study, we recruited solid hypoechoic nodules according only to the recommendations of the 2015 ATA guidelines. Another study with three scoring methods showed that USE had a limited sensitivity and PPV in detecting malignant thyroid nodules and was not superior to conventional US[12]. A retrospective study involving 197 thyroid nodules utilized the iU22 system for elastography and classified the color mapping as blue versus not blue, showing that the ES and SR had a limited ability in differentiating benign from malignant thyroid nodules[23]. However, Magri et al showed that the strain index was significantly higher in malignant thyroid nodules than in benign thyroid nodules and displayed a good diagnostic performance[24]. It can be speculated that the instruments and scoring method used are among the factors that affect the diagnostic efficacy of elastography.

Elasticity is a qualitative method; thus, it is inevitably affected by the practitioner. The SR, which is an objective and semiquantitative technology, was introduced to address this disadvantage[25]. However, whether the ES or SR is more accurate is a long-standing controversy[20, 26-28]. Furthermore, the cut-off value of SR varied largely in different studies[28, 29]. In this research, using an SR>2.99 as the cut-off value, the sensitivity, specificity, accuracy, PPV and NPV were 81.1%, 50.1%, 68.9%, 65.9% and 67.9%, respectively, with an AUC of 0.738, showing no advantage over the ES. Though the SR is a semiquantitative method, it is defined as the ratio obtained by dividing the mean strain within the lesion by the mean strain of the surrounding normal tissue; thus, thyroid echogenicity surrounding thyroid nodules will inevitably affect the SR. As a result, it might be inferred that no additional information is provided by the SR, which is more time consuming than the ES, in the discrimination of malignant from benign thyroid nodules. Although shear wave elastography (SWE) has drawn more attention in recent years and has been reported to be more accurate than USE in some studies[30-32], Tian et al's and Hu et al's meta-analysis showed that USE had a better diagnostic performance than SWE, with comparable specificity between methods[33, 34]. Even more recently, a prospective study comprising 243 nodules revealed that USE yielded the highest performance compared with the TI-RADS score and SWE[35]. The aforementioned data confirm that USE is still a promising diagnostic tool for discriminating malignant from benign thyroid nodules.

Another controversial topic is whether the combined application of US and ES may provide better results for thyroid nodule characterization. The results of the present research demonstrated that the sensitivity and NPV were dramatically higher with the combination than with the TI-RADS and ES alone, and the accuracy was higher with the combination than with the TI-RADS alone but not with the ES alone, consistent with the study by Cantisani[35]. However, the specificity decreased after the two were combined.

According to the 2015 ATA guidelines, nodules ≥ 1 cm in the largest dimension with a high-suspicion sonographic pattern are recommended for FNA. We compared the diagnostic efficacy of elastography in nodules ≥ 1 cm and < 1 cm, and the results showed that the AUC of the ES for nodules ≥ 1 cm was 0.875, with excellent sensitivity, accuracy, PPV, and NPV and good specificity. The aforementioned parameters were all superior to those of nodules < 1 cm. The SR demonstrated a similar trend, except it had a lower sensitivity and NPV in nodules ≥ 1 cm than in nodules < 1 cm. Wang's research also showed a similar result: elastography yielded higher sensitivity for nodules larger than 1 cm[36]. It might be concluded that elasticity is a helpful tool for discriminating benign and malignant thyroid nodules. FNA is not popular in many developing countries because of the lack of skillful cytologists and the high dependence on the skill and experience of the operator and cytologist[37]. Furthermore, FNA is an invasive technique that is time consuming and costly. Thus, the use of FNA is limited in certain areas.

This study has certain limitations. Cytology without lesion growth was also accepted as a reference method for benign lesions, as suggested by international guidelines, which may have resulted in selection bias. However, including only histology would increase the number of malignant lesions[16]. Furthermore, the number of follicular carcinomas and medullary carcinomas, which was determined by morbidity, was low, consistent with previous studies[16, 38]. No reproducibility tests were performed in this study because all US diagnoses were performed by two physicians.

5. Conclusions

In summary, strain elastography combined with the ES is an effective technique for the discrimination of benign and malignant nodules among high-suspicion thyroid nodules based on the 2015 ATA guidelines, especially in nodules ≥ 1 cm.

Declarations

Ethics approval and consent to participate

The ethics committees of The Second Affiliated Hospital of Harbin Medical University, West China Hospital of Sichuan University, Sixth People's Hospital Affiliated to Shanghai Communication University, First Affiliated Hospital of Xi'an Communication University, Xijing Hospital Affiliated to The Fourth Military Medical University, Third Affiliated Hospital of Kunming Medical University, Tumor Hospital of Beijing University, Xiangya Hospital of Central-south University, Tumor Hospital Affiliated to Xinjiang Medical University, Sun Yat-sen Memorial Hospital of Sun Yat-sen University, Beijing Anzhen Hospital Affiliated to Capital Medical University, and People's Hospital of Guangdong Province approved this study. Written or oral consent was obtained from all patients.

Consent to publish

Consent for publication was obtained from all patients.

Availability of data and materials

The datasets analyzed during the current study are available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable.

Authors' contributions

T JW, P YL, WY, AH, Z XD, Y LC, YK, XY, L LS, L BM, YQ and C SZ were responsible for the conception and design of the study. L HR, J SQ, FX, M BY, LY, Z XX, GX, C HT, L WY, TL, L XY, Z XB, LL, and G KH were responsible for acquisition of the data. LHR analyzed the data and drafted the article. TJW made the final approval of the version to be submitted. All authors have read and approved the manuscript.

Acknowledgments

Not applicable.

Abbreviations

ATA: American Thyroid Association;

ROC: receiver operating characteristic;

ES: elasticity score;

SR: strain ratio;

PPV: positive predictive value;

NPV: negative predictive value;

US: ultrasound;

USE: ultrasound strain elastography;

TI-RADS: Thyroid Imaging Reporting and Data System;

ETE: extrathyroidal extension;

ROI: region of interest;

FNA: fine-needle aspiration;

AUC: area under the curve

References

1. Gharib H, Papini E, Paschke R. Thyroid nodules: a review of current guidelines, practices, and prospects. *Eur J Endocrinol* 2008;159:493-505.
2. Hegedus L, Bonnema SJBennedbaek FN. Management of simple nodular goiter: current status and future perspectives. *Endocrine Reviews* 2013, 24(1):102-132.
3. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid* 2016, 26(1):1-133.
4. Iannuccilli JD, Cronan JJ, Monchik JM. Risk for malignancy of thyroid nodules as assessed by sonographic criteria: the need for biopsy. *Journal of Ultrasound in Medicine* 2004, 23(11):1455-1464.
5. Zhang YF, He Y, Xu HX. Conventional US, US Elasticity Imaging, and Acoustic Radiation Force Impulse Imaging for Prediction of Malignancy in Thyroid Nodules. *Radiology* 2015, 274(1):307-307.
6. Brito JP, Gionfriddo MR, Alaa AN, et al. The accuracy of thyroid nodule ultrasound to predict thyroid cancer: systematic review and meta-analysis. *Journal of Clinical Endocrinology & Metabolism* 2013, 99(4):1253-1263.
7. Remonti LR, Kramer CK, Leitao CB, et al. Thyroid ultrasound features and risk of carcinoma: a systematic review and meta-analysis of observational studies. *Thyroid* 2015, 25(5):538-550.
8. Andrej L, Tatsuya H, Ryo A, et al. Thyroid gland tumor diagnosis at US elastography. *Radiology* 2005, 237(1):202-211.
9. Seyed Amirhossein R, Hadduck TA, Gelareh S, et al. Comparative effectiveness of elastographic and B-mode ultrasound criteria for diagnostic discrimination of thyroid nodules: a meta-analysis. *Ajr American Journal of Roentgenology* 2013, 200(6):1317-1326.
10. Jiasi S, Jingyu C, Xuemei W. Real-time ultrasound elastography for differentiation of benign and malignant thyroid nodules: a meta-analysis. *Journal of Ultrasound in Medicine* 2014, 33(3):495-502.
11. Moon HJ, Sung JM, Kim EK, et al. Diagnostic performance of gray-scale US and elastography in solid thyroid nodules. *Radiology* 2012, 262(3):1002.
12. Unlütürk U, Erdoğan MF, Demir O, et al. Ultrasound elastography is not superior to grayscale ultrasound in predicting malignancy in thyroid nodules. *Thyroid* 2012, 22(10):1031-1038.
13. Shin JH, Baek JH, Chung J, et al. Ultrasonography Diagnosis and Imaging-Based Management of Thyroid Nodules: Revised Korean Society of Thyroid Radiology Consensus Statement and Recommendations. *Korean J Radiol* 2016, 17(3):370-395.

14. Ako I, Ei U, Eriko T, et al. Breast disease: clinical application of US elastography for diagnosis. *Nippon Ishinkin Gakkai Zasshi* 2006, 8(4):305-311.
15. Trimboli P, Guglielmi R, Monti S, et al. Ultrasound sensitivity for thyroid malignancy is increased by real-time elastography: a prospective multicenter study. *The Journal of clinical endocrinology and metabolism* 2012, 97(12):4524-4530.
16. Friedrich-Rust M, Vorlaender C, Dietrich CF, et al. Evaluation of Strain Elastography for Differentiation of Thyroid Nodules: Results of a Prospective DEGUM Multicenter Study. *Ultraschall Med* 2016, 37(3):262-270.
17. Cantisani V, Maceroni P, D'Andrea V, et al. Strain ratio ultrasound elastography increases the accuracy of colour-Doppler ultrasound in the evaluation of Thy-3 nodules. A bi-centre university experience. *European radiology* 2016, 26(5):1441-1449.
18. Rago T, Santini F, Scutari M, et al. Elastography: new developments in ultrasound for predicting malignancy in thyroid nodules. *Journal of Clinical Endocrinology & Metabolism* 2007, 92(8):2917-2922.
19. Tatar IG, Kurt A, Yilmaz KB, et al. The role of elastosonography, gray-scale and colour flow Doppler sonography in prediction of malignancy in thyroid nodules. *Radio Oncol* 2014, 48(4):348-353.
20. Guazzaroni M, Spinelli A, Coco I, et al. Value of strain-ratio on thyroid real-time sonoelastography. *La Radiologia medica* 2014, 119(3):149-155.
21. Wolinski K, Szkudlarek M, Szczepanek-Parulska E, et al. Usefulness of different ultrasound features of malignancy in predicting the type of thyroid lesions: a meta-analysis of prospective studies. *Pol Arch Med Wewn* 2014, 124(3):97-104.
22. Szczepanek-Parulska E, Wolinski K, Stangierski A, et al. Comparison of diagnostic value of conventional ultrasonography and shear wave elastography in the prediction of thyroid lesions malignancy. *PloS one* 2013, 8(11):e81532.
23. Seong M, Shin JH, Hahn SY. Ultrasound Strain Elastography for Circumscribed Solid Thyroid Nodules without Malignant Features Categorized as Indeterminate by B-Mode Ultrasound. *Ultrasound Med Biol* 2016, 42(10):2383-2390.
24. Magri F, Chytiris S, Zerbini F, et al. Maximal stiffness evaluation by real-time ultrasound elastography, an improved tool for the differential diagnosis of thyroid nodules. *Endocr Pract* 2015, 21(5):474-481.
25. Ding J, Cheng HD, Huang J, et al. An improved quantitative measurement for thyroid cancer detection based on elastography. *European journal of radiology* 2012, 81(4):800-805.
26. Cantisani V, Grazhdani H, Ricci P, et al. Q-Elastosonography of Solid Thyroid Nodules: Assessment of Diagnostic Efficacy and Interobserver Variability in a Large Patient Cohort. *European Radiology* 2014, 24(1):143-150.
27. Cantisani V, D'Andrea V, Mancuso E, et al. Prospective evaluation in 123 patients of strain ratio as provided by quantitative elastosonography and multiparametric ultrasound evaluation (ultrasound score) for the characterisation of thyroid nodules. *La Radiologia medica* 2013, 118(6):1011-1021.
28. Chong Y, Shin JH, Ko ES, et al. Ultrasonographic elastography of thyroid nodules: is adding strain ratio to colour mapping better? *Clinical radiology* 2013, 68(12):1241-1246.
29. Ning CP, Jiang SQ, Zhang T, et al. The value of strain ratio in differential diagnosis of thyroid solid nodules. *European journal of radiology* 2012, 81(2):286-291.
30. Kyriakidou G, Friedrich-Rust M, Bon D, et al. Comparison of strain elastography, point shear wave elastography using acoustic radiation force impulse imaging and 2D-shear wave elastography for the differentiation of thyroid nodules. *PloS one* 2018, 13(9):e0204095.
31. Chen BD, Xu HX, Zhang YF, et al. The diagnostic performances of conventional strain elastography (SE), acoustic radiation force impulse (ARFI) imaging and point shear-wave speed (pSWS) measurement for non-calcified thyroid nodules. *Clinical hemorheology and microcirculation* 2017, 65(3):259-273.
32. Lin P, Chen M, Liu B, et al. Diagnostic performance of shear wave elastography in the identification of malignant thyroid nodules: a meta-analysis. *European radiology* 2014, 24(11):2729-2738.
33. Tian W, Hao S, Gao B, et al. Comparison of Diagnostic Accuracy of Real-Time Elastography and Shear Wave Elastography in Differentiation Malignant From Benign Thyroid Nodules. *Medicine* 2015, 94(52):e2312.
34. Hu X, Liu Y, Qian L. Diagnostic potential of real-time elastography (RTE) and shear wave elastography (SWE) to differentiate benign and malignant thyroid nodules: A systematic review and meta-analysis. *Medicine* 2017, 96(43):e8282.

35. Cantisani V, David E, Graždani H, et al. Prospective Evaluation of Semiquantitative Strain Ratio and Quantitative 2D Ultrasound Shear Wave Elastography (SWE) in Association with TIRADS Classification for Thyroid Nodule Characterization. *Ultraschall Med* 2019, 40(4):495-503.
36. Wang F, Chang C, Chen M, et al. Does Lesion Size Affect the Value of Shear Wave Elastography for Differentiating Between Benign and Malignant Thyroid Nodules? *Journal of ultrasound in medicine* 2018, 37(3):601-609.
37. Liu BJ, Li DD, Xu HX, et al. Quantitative Shear Wave Velocity Measurement on Acoustic Radiation Force Impulse Elastography for Differential Diagnosis between Benign and Malignant Thyroid Nodules: A Meta-analysis. *Ultrasound in medicine & biology* 2015, 41(12):3035-3043.
38. Yang J, Song Y, Wei W, et al. Comparison of the effectiveness of ultrasound elastography with that of conventional ultrasound for differential diagnosis of thyroid lesions with suspicious ultrasound features. *Oncology letters* 2017, 14(3):3515-3521.

Tables

Table 1 Diagnostic value of US and USE for the diagnostic of malignant thyroid nodules

Criteria	Benign	Malignant	Sensitivity(%)	Specificity(%)	Accuracy(%)	PPV(%)	NPV(%)	<i>P</i>
Microcalcification								
Yes	211	416	49.9	65.6	56.4	66.3	48.9	0.01
no	400	318						
Taller than wide								
yes	55	309	37.1	91.0	59.8	84.9	51.4	0.01
no	556	525						
Irregular margins								
Yes	525	83	93.9	14.1	60.1	59.9	62.8	0.01
no	86	51						
ETE								
Yes	469	693	83.0	23.2	57.7	59.6	50.0	0.01
No	142	141						
Disrupted rim calcification								
Yes	7	39	4.7	98.9	44.4	84.8	43.2	0.01
No	604	795						
Kwak's TI-RADS								
4a	5	2	62.2	70.2	65.6	74.0	57.7	0.01
4b	424	313						
4c	182	367						
5	0	152						
Elasticity score								
>3	240	771	92.4	60.7	79.0	76.3	85.5	0.01
≤3	371	63						
SR value								
>2.99	305	676	81.1	50.1	68.9	65.9	67.9	0.01
≤2.99	306	158						

PPV: positive predictive value; NPV: negative predictive value; ETE: Extrathyroidal extension; TI-RADS: Thyroid Imaging Reporting and Data System; SR: strain ratio.

Table 2 Logistic regression analysis for predicting malignant thyroid nodules

Criteria	Univariate analysis			Multivariate analysis		
	OR	95%CI	<i>P</i>	OR	95%CI	<i>P</i>
Microcalcification	1.887	1.522-2.339	0.000	5.29	3.472-8.078	0.001
Taller than wide	5.950	4.361-8.118	0.000	12.71	7.625-21.188	0.000
Irregular margins	2.515	1.748-3.618	0.000	10.117	5.88-17.404	0.000
Rim calcification	4.233	1.880-9.529	0.000	2.569	0.791-8.348	0.117
ETE	1.475	1.137	1.915	6.412	3.909-10.519	0.000
Elasticity score	4.473	3.824-5.231	0.000	3.741	3.129-4.474	0.000
SR value	1.504	1.405-1.610	0.000	1.083	1.028-1.142	0.003

SR: strain ratio; OR: odds ratio; CI: confidence interval.

Table 3. Diagnostic performance of Elasticity score and SR in different size of nodules

	AUC	Sensitivity	Specificity	Accuracy	PPV	NPV	<i>P</i>
Elasticity score							
≥1	0.875	95.0	68.8	85.9	85.2	87.8	0.000
≠1	0.800	90.4	56.8	74.4	69.9	84.2	0.000
SR							
≥1	0.779	79.8	59.3	72.7	78.8	60.8	0.000
≠1	0.709	82.1	45.6	64.8	62.6	69.6	0.000

PPV: positive predictive value; NPV: negative predictive value; SR: strain ratio

Figures




Figure 1

Nodule selection among the included patients.




Figure 2

US characteristics of different ES. a. Ultrasound elastography image of a fibroadenoma in a 45-year-old man. On the left side, the nodule shaded in green and red was scored as 1. b. Ultrasound elastography image of a nodular goiter in a 45-year-old woman. On the left side, the nodule colored in green with focal blue spots was scored as 2. c. Ultrasound elastography image of a tiny papillary thyroid microcarcinoma in a 34-year-old woman. On the left side, the nodule half in green and half in blue was scored as 3. d. Ultrasound elastography image of a tiny papillary thyroid microcarcinoma in a 39-year-old man. On the left side, the nodule shaded in blue with focal green spots was scored as 4. e. Ultrasound elastography image of a papillary thyroid microcarcinoma in a 60-year-old woman. On the left side, the entire nodule and its surrounding area in blue were scored as 5.




Figure 3

ROC curve of the TI-RADS, ES and TI-RADS combined with ES in differentiating malignant and benign nodules.




Figure 4

ROC curve of the ES and SR value in differentiating malignant and benign nodules.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Languagecertificate.pdf](#)
- [Titlepage.docx](#)