Citrulline deiminase pathway provides ATP and boosts growth of *Clostridium carboxidivorans* P7

Xiangfei Li

Jiangnan University

Rumeng Han

Jiangnan University

Teng Bao

The Ohio State University

Tolbert Osire

Jiangnan University

Xian Zhang

Jiangnan University

Meijuan Xu

Jiangnan University

Taowei Yang

Jiangnan University

Zhiming Rao (✉ raozhm@jiangnan.edu.cn)

Jiangnan University <https://orcid.org/0000-0003-2495-0660>

Research

Keywords: Clostridium carboxidivorans, citrulline, syngas fermentation, alcohol/acid ratio

Posted Date: August 3rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-764358/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

Clostridium carboxidivorans P7 is capable of producing ethanol and butanol from inexpensive and non-food feedstock such as syngas. Achieving improved ethanol and butanol production in the strain for industrial application depends on the energetics and biomass, especially ATP availability.

Results

This study found that exogenous addition of citrulline promoted accumulation of ATP, increased specific growth rate, and reduced the doubling time of *C. carboxidivorans* P7. In heterotrophic fermentation experiments, the addition of citrulline increased intracellular ATP by 3.39-fold, significantly enhancing the production of total alcohol (ethanol + butanol) by 20%. Moreover, in the syngas fermentation experiments, the addition of citrulline improved the level of intracellular ATP and the biomass by 80.5% and 31.6%, respectively, resulting in a 18.6% and 60.3%, increased in ethanol and the alcohol/acid production ratio, respectively.

Conclusions

This is the first report that citrulline could promote the growth of *C. carboxidivorans* P7 and increased the level of intracellular ATP, which is of great significance for the use of *C. carboxidivorans* P7 to synthesize biofuels.

Background

Globally, the main energy source for human survival is petroleum-based non-renewable fossil fuels [1]. Nowadays, fossil fuel depletion, increasing energy consumption, growing CO₂ emissions, and climate change have increased the demand for renewable energy sources [2, 3]. As renewable biomass fuels, ethanol [4] and butanol [5] have high energy density and are compatible with current petroleum energy infrastructure equipment, which has attracted significant attention [6]. However, the production of bio-butanol and bio-ethanol is presently limited owing to the high substrate cost of conventional feedstock [7]. In recent years, gas fermentation has attracted increasing attention as an emerging method for the production of renewable biofuels, such as bio-butanol and bio-ethanol [8, 9]. In the one hand, the feedstock for gas fermentation are relatively abundant and inexpensive, such as exhaust gas from industrial production, and burning fossil fuels, as well as synthetic gas. On the other hand, compared with traditional acetone-butanol-ethanol (ABE) fermentation by solventogenic *Clostridium*, gas fermentation has no need to occupy food resources and precious land resources for humans [10]. However, low solubility of gaseous substrates in water, low ATP generation and poor biomass of solventogenic *Clostridium* are still the major factors limited the production of bio-butanol and bio-ethanol [11, 12].

Clostridium carboxidivorans P7 is one of the strains capable of producing biofuel from gaseous [13]. It can synthesize ethanol and n-butanol by capturing CO₂, CO and H₂ through WLP (Wood-Ljungdahl pathway) [14, 15]. However, whole-genome sequencing (GenBank: CP011803.1) indicates that there is no a complete TCA cycle in *C. carboxidivorans* P7 [13]. Hence, its own energy supply mainly comes from the glycolytic pathway and the synthetic pathway for acetic acid production. What's more, syngas fermentation also has drawbacks such as low biomass and insufficient energy supply, resulting in low biofuel production and high by-products. Although optimization of the appropriate media composition, metal ion composition [16], pH [17, 18], and reactor design [19] can improve cell growth and product

accumulation, it is not of help in increasing intracellular ATP production. Therefore, there is an urgent need to enhance intracellular ATP supply and biomass of *C. carboxidivorans* P7, so as to promote the biofuels production. Nowadays, genetic engineering is regarded as a useful strategy to manipulate the metabolism and to enhance accumulation of ideal products [20–22]. But the molecular genetic manipulation technology of *C. carboxidivorans* P7 had just been established [7], it required relatively long operation time, and the success rate was not high. So, investigators have noted that amino acids, as the basic building blocks of proteins, play a crucial role in the life of cells [23]. It is known that arginine can be hydrolysed to urea and ornithine at the action of arginase. Then, ornithine is reacted with a molecule of carbamoyl phosphate at the reaction of ornithine carbamoyltransferase to generate citrulline. At last, citrulline can be transformed to carbon dioxide, water, ammonia and two molecules of ATP by carbamoyl phosphate synthetase [24]. For instance, the addition of arginine shortened the *Clostridium autoethanogenum* doubling time, increased the intracellular ATP energy level by 5-fold, and weakened the accumulation of by-product acetic acid [12]. Therefore, amino acid metabolism has an enormous potential to improve the cell density and intracellular energy level of *C. carboxidivorans* P7.

This paper assessed the significance of amino acids which could be helpful to promote the biomass, ATP regeneration and the yield of biofuels. It was found that citrulline added could provide additional ATP, enhance biomass and specific growth rate for *C. carboxidivorans* P7 under both heterotrophic and autotrophic conditions. What's more, the addition of citrulline increased the alcohol/acid production ratio by 60.3% under autotrophic conditions. The results provides a new method of thinking for promoting cell growth and increasing energy levels in *C. carboxidivorans* P7.

Results

Citrulline boosted cell growth under heterotrophic conditions

Amino acids, as the basic units of proteins, play an crucial role in cell life [23]. To explore the importance of amino acid addition for cell growth of *C. carboxidivorans* P7, the effects of 20 conventional amino acids on cell growth were studied by serum bottle fermentation in MM520 medium. Fermentation results showed that histidine and aspartic acid promoted cell growth, but the presence of arginine significantly impaired cell growth (Fig. 1A). It was speculated that arginidase (EC: 3.5.3.1) probably converted arginine to urea and ornithine, which consequently may have contributed to increase in the metabolic load of cells and decreased cell growth. KEGG analysis revealed that *C. carboxidivorans* P7 lost the ability to yield citrulline from arginine due to the absence of arginine deiminase. We hypothesized that if cells could generated citrulline from arginine, they could then break down arginine through the ADI (arginine deiminase) pathway, hence providing cells with additional ATP supply. Although the complete ADI metabolic pathway does not exist in *C. carboxidivorans* P7, it does have a complete citrulline degradation pathway. Furthermore, it was known that citrulline could be transformed to carbon dioxide, water, ammonia and two molecules of ATP by carbamoyl phosphate synthetase [24]. Therefore, citrulline was added to explore the effects on the cell density of *C. carboxidivorans* P7. The results indicated that the addition of citrulline significantly promoted the growth capacity of *C. carboxidivorans* P7, and the maximum OD₆₀₀ increased by 18.6% compared with the wild-type strain (Fig. 1A).

To further determine the optimal concentration of citrulline addition, 0.6 g/L, 1.2 g/L and 2.4 g/L citrulline were added to the fermentation medium and incubated under the same conditions. The results showed that the addition of 0.6 g/L citrulline was most beneficial for cell growth. The stable phase of the cells was obviously prolonged and the sugar consumption was significantly promoted. The glucose was depleted by 24 hours after fermentation. Increasing the concentration of citrulline further resulted in decreased cell growth compared with the addition of 0.6 g/L citrulline

(Fig. 1B and 1C). It may be due to the limited use of citrulline by *C. carboxidivorans* P7. As shown in Fig. 1D, there were still a large amount of citrulline residue in the later stage of fermentation under the addition of 1.2 g/L or 2.4 g/L citrulline.

Heterotrophic growth experiments proved that citrulline metabolism promoted cell glycolysis and delayed cell aging

Based on the results above, the addition of citrulline boosted cell growth, accelerated glucose utilization and prolonged cell stability. It is known that citrulline can generate carbamoyl-P under the action of ornithine carbamoyltransferase, then the carbamoyl-P generates CO₂ and ATP under the action of carbamate kinase (Fig. 2D). Therefore, It was speculated that citrulline catabolism provided additional ATP supply for glycolysis, thereby improving the ability of two key energy-consuming enzymes (*glk* and *pfk*), in turn increaseing glucose utilization. Under the same conditions, *C. carboxidivorans* P7 was fermented by serum bottle in MM520-0.6 Cit and MM520 medium for the same time and samples were taken regularly. RT-qPCR was used to analyze the expression levels of *glk* (encoding glucokinase), *pfk* (encoding phosphofructokinase), *otc* (encoding ornithine carbamoyltransferase) and *ck* (encoding carbamate kinase) at 12 h, 24 h and 36 h, respectively. The results showed that in MM520-0.6 Cit medium, *glk*, *pfk*, *otc* and *ck* were significantly up-regulated at 12 h, 24 h, and 36 h (Fig. 2A, 2B and 2C). However, after 24 hours of fermentation, due to the consumption of citrulline and glucose, the relative expression of *glk*, *pfk*, *otc*, and *ck* decreased accordingly. After the fermentation at 96 h, it was found that the cells cultured in MM520-0.6 Cit medium had smooth surface and no residue of glucose particles with Frozen scanning electron microscopy (FSEM) (Fig. 3B). On the contrary, the surface of cells cultured in MM520 medium had a rough surface with more glucose residues (Fig. 3A). Therefore, FSEM analysis further showed that the addition of citrulline contributed to promote the metabolism of glucose by *C. carboxidivorans* P7 and could maintain cell viability.

Citrulline boosted heterotrophic growth of *C. carboxidivorans* P7 while supplying extra ATP

By analyzing the metabolic pathway of citrulline, it was further speculated that citrulline catabolism could provide cells with additional ATP, thereby promoting cell growth. Therefore, under the same conditions, *C. carboxidivorans* P7 was fermented in serum bottles with MM520-0.6 Cit and MM520 media for the same time. The specific growth rate, doubling time, and intracellular ATP levels of the cells were further analyzed. The results showed that the specific growth rate of *C. carboxidivorans* P7 in MM520-0.6 Cit medium increased significantly when the fermentation was performed within 10 h to 30 h (Fig. 4). When fermentation reached 30 h, the cell doubling time of *C. carboxidivorans* P7 in MM520-0.6 Cit medium was 162 h, which was 81.3% shorter than that of MM520 medium. Through analysis of intracellular ATP level, it was found that the ATP level in MM520-0.6 Cit medium was 3.39-fold higher than that in MM520 medium at 24 h. The results herein showed that the addition of citrulline increased intracellular ATP levels under heterotrophic conditions. Thereafter, when citrulline was depleted, the intracellular ATP level decreased and remained at a lower level.

Under heterotrophic conditions, metabolites of *C. carboxidivorans* P7 in MM520 and MM520-0.6 Cit medium were analyzed (Table 2). Addition of citrulline had no significant effect on product conversion of ethanol and butanol. However, since the addition of citrulline could promote the utilization rate of glucose in cells. Although the product conversion rate did not change significantly, the total alcohol yield of fermentation products (the yield of ethanol and butanol was 1.2 g/L and 0.16 g/L, respectively) increased by about 20% (Fig. 4F).

Table 2

Under heterotrophic conditions, the product yields of *C. carboxidivorans* P7 under different culture conditions.

Medium	Acetate (g/g)	Ethanol (g/g)	Butyrate (g/g)	Butanol (g/g)	Total alcohols (g/g)	Total acid (g/g)	Alcohols/Acids (g/g)	C4/C2 (g/g)	C recovery (%)
MM520	0.30 ± 0.01	0.07 ± 0.01	0.15 ± 0.01	0.025 ± 0.002	0.10 ± 0.01	0.46 ± 0.04	0.21	0.47	57
MM520-0.6 Cit	0.29 ± 0.02	0.08 ± 0.01	0.16 ± 0.01	0.024 ± 0.001	0.10 ± 0.01	0.45 ± 0.02	0.22	0.49	56.4

Citrulline boosted autotrophic growth of *C. carboxidivorans* P7 while supplying extra ATP

To further verify whether citrulline could provide cells with ATP and promote cell growth under autotrophic conditions (CO₂/CO/H₂ [50:35:15], 1 atm), *C. carboxidivorans* P7 was fermented in Syngas-GY free and Syngas-0.6 Cit-GY free medium (pH 6.6, 37°C, 150 rpm) and the effects of citrulline improved growth of *C. carboxidivorans* P7 was investigated. Firstly, as shown in Fig. 5A, when citrulline was added to the N₂-GY free medium, the cells could not grow normally, indicating that citrulline could not be used as the only carbon source to maintain cell growth. However, under syngas conditions, exogenous addition of citrulline could increase the biomass of *C. carboxidivorans* P7 by 31.6%. It was also found that the ATP levels in Syngas-0.6 Cit-GY free medium were significantly higher than those in Syngas-GY free medium after fermentation for 48 h. Moreover, at 96 h of fermentation, although the ATP levels in Syngas-0.6 Cit-GY free medium remained almost constant, intracellular ATP levels increased 80.5% in Syngas-0.6 Cit-GY free medium compared to Syngas-GY free medium (Fig. 5B). It was also found that the specific growth rate of *C. carboxidivorans* P7 in Syngas-0.6 Cit-GY free medium increased significantly, while the doubling time of the bacteria greatly shortened after the fermentation reached 40 h (Fig. 5C and 5D).

Under autotrophic conditions, the metabolites of *C. carboxidivorans* P7 were analyzed in Syngas-0.6 Cit-GY free and Syngas-GY free medium (Table 3). The addition of citrulline increased the yield of ethanol and the alcohol/acid ratio by 18.6% and 60.3%, respectively. Although the yield of ethanol reached to 1.21 g/L, however, there was no significant change in butanol production. A possible explanation for the no significant change in butanol production could be that the low production of butanol in *C. carboxidivorans* P7.

Table 3

Under autotrophic conditions, the product yields of *C. carboxidivorans* P7 under different culture conditions.

Medium	Acetate (g/L)	Ethanol (g/L)	Butyrate (g/L)	Butanol (g/L)	Total alcohols (g/L)	Total acid (g/L)	Alcohols/Acids (g/g)	C4/C2 (g/g)
Syngas-GY	0.45 ± 0.05	1.02 ± 0.11	0.12 ± 0.03	0.03 ± 0.01	1.05 ± 0.12	0.57 ± 0.08	1.84	0.10
Syngas-0.6 Cit-GY	0.26 ± 0.04	1.21 ± 0.09	0.16 ± 0.02	0.03 ± 0.01	1.24 ± 0.10	0.42 ± 0.06	2.95	0.13

Discussion

C. carboxidivorans P7 has abilities common to most acetic bacteria, such as *Acetobacterium woodii* [25], *Clostridium formicoaceticum* [26], *Clostridium aceticum* [27], *Clostridium ljungdahlii* [28], *C. autoethanogenum* [29], etc. all have WLP pathways. Conforming to the current mainstream of clean energy, *C. carboxidivorans* P7 is able to produce

biofuels from syngas via the WLP pathway. However, *C. carboxidivorans* P7 has low cell density and insufficient intracellular energy supply during the synthesis gas fermentation process, resulting in industrially ineffective *C. carboxidivorans* P7. In this study, it was found that the addition of citrulline not only improved cell density, but also enhanced the intracellular ATP level in *C. carboxidivorans* P7 under both heterotrophic and autotrophic conditions. Moreover, the addition of citrulline increased the yield of ethanol and the alcohol/acid ratio by 18.6% and 60.3%, respectively. Therefore, this research revealed that the exogenous addition of citrulline increased the biomass and ATP supply, which had important implications for accumulating biofuels in *C. carboxidivorans* P7.

Recently, cell immobilization technology, optimization of fermentation parameters and metabolic engineering have been used to improve cell density and intracellular ATP level. Cell recovery [30] and cell immobilization [31] techniques promoted cell density during the period of fermentation, resulting in a 5-fold increased in butanol production. The control of fermentation processes, such as pH [17], temperature [32, 33], pressure [17] and inoculation amount [34], also had a great influence on cell density, but was not reported to significantly promote the accumulation of intracellular energy ATP. Amino acids are essential elements for life [35], and amino acid metabolism can produce carbon dioxide and other small molecules for cell life activities [36]. Therefore, by adding 5 g/L arginine, [12] not only increased the cell density of *C. autoethanogenum*, but also increased the intracellular energy ATP level by 5-fold. The engineering modification of strains is also of great significance for improving cell fermentation density and intracellular energy ATP accumulation. Ferredoxin oxidoreductase (AOR) [37] provided extra ATP supply for acetic acid bacteria and promoted the accumulation of ethanol and butanol. *C. autoethanogenum* [13] explored the AOR/ADH pathway to increase the level of intracellular energy ATP and increase the yield of ethanol. Overexpression of two glycerol dehydrogenase (GldH) genes (*dhaD1* and *gldA1*) in *Clostridium pasteurianum* further resulted in a 43% increase in glycerol utilization and a significant increase in cell density (> 50%) [38]. At present, the genetic operation system of *C. carboxidivorans* P7 has been preliminarily established [7]. However, there are some problems, such as complex operation, low transduction efficiency and long time consuming, making it difficult to operate in practice. Therefore, cell immobilization and optimization of fermentation parameters are still the main means to improve cell density and intracellular energy ATP supply.

Although this study achieved a synchronous increase in cell density and intracellular energy ATP of *C. carboxidivorans* P7 by exogenous addition of citrulline, the cell utilization of citrulline was not very high. Furthermore, since the genetic operation system of *C. carboxidivorans* P7 is relatively complex and inefficient, it is possible to overexpress the two key enzymes (ornithine carbamoyltransferase (OTC) and carbamate kinase (CK)) of the citrulline pathway in *C. carboxidivorans* P7, to expand the pathway of citrulline metabolism and further improve citrulline utilization of *C. carboxidivorans* P7.

Conclusions

In general, citrulline could promote the accumulation of intracellular energy ATP, increase the specific growth rate and shorten the doubling time of *C. carboxidivorans* P7 under both autotrophic and heterotrophic conditions. Under heterotrophic conditions, the addition of citrulline increased the level of intracellular ATP and the production of total alcohol by 3.39-fold and 20%, respectively. Under autotrophic conditions, the addition of citrulline increased the level of intracellular ATP by 80.5%, the yield of ethanol by 18.6% and the alcohol/acid ratio by 60.3%. In summary, this is the first report to reveal that citrulline could promote the growth of *C. carboxidivorans* P7 and increase intracellular energy ATP, which is of great significance for improving the production of clean energy such as ethanol and butanol by *C. carboxidivorans* P7.

Methods

Strains and culture medium

The strains and primers in the work are summarized in Table 1. *C. carboxidivorans* P7 (DSM 15243) was preserved at -80°C in our laboratory. It was inoculated into the MM520 medium, containing K₂HPO₄ 2.2 g/L, KH₂PO₄ 1.5 g/L, (NH₄)₂SO₄ 1.3 g/L, yeast extract 2 g/L, tryptone 4 g/L, Na-Resazurin solution 0.1%(w/v), L-cysteine-HCl•H₂O 0.5g/L, FeSO₄•7H₂O 0.00125g/L, CaCl₂•2H₂O 0.075 g/L, MgCl₂•6H₂O 0.2 g/L and trace element solution (FeCl₂•4H₂O 1.5 mg/L, ZnCl₂ 0.07 mg/L, MnCl₂•4H₂O 0.1 mg/L, CoCl₂•6H₂O 0.19 mg/L, CuCl₂•2H₂O 0.002 mg/L, NiCl₂•6H₂O 0.024 mg/L, H₃BO₃ 0.006 mg/L, Na₂MoO₄•2H₂O 0.036 mg/L). Dissolved oxygen in the medium was boiled for half an hour to eliminate under the stream of N₂ gas, then added L-cysteine-HCl•H₂O 0.5 g/L and continued to boil for an hour until the color of the medium from blue to colorless. The boiled medium was divided into a serum bottle (50 mL/100 mL) and nitrogen added for 10 to 20 min, until the color changed from a light blue to yellow. Then serum bottles were sealed with gas impermeable butyl rubber septum-type stoppers and aluminium crimp seals. The serum bottles filled with medium were sterilized at 121°C for 20 min. 500 µL sterilized glucose (500 g/L) and 250 µL sterilized L-cysteine-HCl•H₂O (100 g/L) were added into media. Before inoculation, the culture medium should be adjusted to pH ~ 6.6 with 1 M HCl or 1 M NaOH, which was purged with N₂ for 10 min and sterilized. The gas fermentation medium (Syngas-GY free medium): MM520 medium contains no glucose and yeast extract and filled with the gas mixture (50% CO₂, 35% CO, and 15% H₂ at 1 atm) and it should be purged every 48 hours. The gas fermentation medium (N₂-GY free medium): MM520 medium filled with the N₂ and purged every two days, without any glucose and yeast extract added to it.

Table 1
Strains and primers in this study.

Characteristics		Source
Strains		
<i>C. carboxidivorans</i> P7	autotrophic growth on CO, CO ₂ and H ₂	Our lab
Primers 5' → 3'		
RT-PCR- <i>glk</i> F	aggtagctgataaagcat	This study
RT-PCR- <i>glk</i> R	tgaaactcctcctccaat	This study
RT-PCR- <i>pfk</i> F	tcctgagaaaggctacaa	This study
RT-PCR- <i>pfk</i> R	atctgctccaccaatacc	This study
RT-PCR- <i>otc</i> F	gaagaatgggaagaacg	This study
RT-PCR- <i>otc</i> R	ctggcaagcaatgaaga	This study
RT-PCR- <i>ck</i> F	tgagaaaggcactggat	This study
RT-PCR- <i>ck</i> R	tatgggttagtggtgat	This study

Fermentation

All fermentation processes were preformed in serum bottles filled with 50 mL culture medium, and samples were taken at regular intervals for analysis. In order to study the effects of 20 conventional amino acids and citrulline on

the growth of *C. carboxidivorans* P7. Each amino acid (1 g/L) was added to the MM520 medium, and placed in a 37°C incubator for fermentation. So as to further study the effects of different concentrations of citrulline on the growth of *C. carboxidivorans* P7, different concentrations of citrulline (0.6 g/L, 1.2 g/L, 2.4 g/L) were added to the fermentation medium. Heterotrophic medium supplemented with 0.6 g/L citrulline was named MM520-0.6 Cit. Syngas fermentation was performed in Syngas-GY free medium. After inoculation at 5% (v/v), the serum bottle cells were cultured at 37°C, 150 rpm. Syngas-GY free medium supplemented with 0.6 g/L citrulline named Syngas-0.6 Cit-GY free medium. All fermentation conditions were preformed in duplicate.

Analytical methods

The spectrophotometer (UV-1800) was used to determine the Optical Density (OD₆₀₀). Acetate and butyrate were analyzed on an HPX-87H column (Bio-Rad). Detection conditions: Temperature 50°C, Mobile phase: 5 mm H₂SO₄, Detection wavelength: 210 nm, Flow rate: 1 mL/min. Ethanol and butanol were detected according to Cheng [7]. ATP was measured by ATP Assay Kitae (s0026b) purchased from Beyotime Institute of Biotechnology. All of the assays were performed in triplicate.

Gene expression detection via RT-qPCR

The total RNA extraction of the collected cells were performed by using FastPure Cell/Tissue Total RNA Isolation Kit RC101 (Vazyme Biotech Co., Ltd., Nanjing, China). RT-qPCR reactions were conducted with ChamQ Universal SYBR qPCR Master Mix*Q711-02 (Vazyme Biotech Co., Ltd., Nanjing, China). The StepOnePlus 96 real-time PCR system (Applied biological systems Inc, USA) was used to amplify and quantify the PCR samples. The method was as follows: 30s at 95°C, 40 amplification cycles of 10s at 95°C, 30s at 60°C. Relative levels of transcript abundance of the studied genes were calculated by the $2^{-\Delta\Delta CT}$ method. RT-qPCR of each gene was tested with three reactions in parallel.

Scanning electron microscopy

Frozen scanning electron microscopy (FSEM) was used to observe cell morphology of *C. carboxidivorans* P7 under the medium of MM502 and MM520-0.6 Cit. Bacterial cells were cultured in MM520 and MM520-0.6 Cit medium at 37°C incubator for 96 h. Then the bacteria cells were collected by centrifugation at 6000 rpm, resuspended with sterile water, and placed on carbon film-coated copper Grid (230 mesh; Beijing Zhongjing Science and Technology Co., Ltd., Beijing, China). At last, the bacterial cells liquid on the film was dried at 25°C and observed by FSEM.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

All authors have seen and approved the manuscript before submission to Biotechnology for Biofuels.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare no conflicts of interest.

Funding

This work was supported by the National Key Research and Development Program of China (2021YFC2100900), the Science and Technology Project of Xinjiang Production and Construction Corps (No. 2019AB009), the National Natural Science Foundation of China (31870066, 21778024), National First-Class Discipline Program of Light Industry Technology and Engineering (LITE2018-06), the Program of Introducing Talents of Discipline to Universities (111-2-06).

Authors' contributions

XL performed all the experimental works, analyzed the data and coordinated the manuscript draft and revision. TY designed and revised the manuscript critically. RH, TB and XZ helped with data analysis. TO, MX and ZR were involved in proofreading the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Henrich E, Dahmen N, Dinjus E, Sauer J. The Role of Biomass in a Future World without Fossil Fuels. *Chem-Ing-Tech*. 2015; 87(12):1667-1685.
2. Seroussi H. Fate and future climatic role of polar ice sheets. *Nature*. 2019; 566(7742):48-49.
3. Sommer R, Glazirina M, Yuldashev T, Otarov A, Ibraeva M, Martynova L, Bekenov M, Kholov B, Ibragimov N, Kobilov R *et al*. Impact of climate change on wheat productivity in Central Asia. *Agriculture, Ecosystems & Environment*. 2013; 178:78-99.
4. Shen S, Gu Y, Chai C, Jiang W, Zhuang Y, Wang Y. Enhanced alcohol titre and ratio in carbon monoxide-rich off-gas fermentation of *Clostridium carboxidivorans* through combination of trace metals optimization with variable-temperature cultivation. *Bioresour. Technol*. 2017; 239:236-243.
5. Yu M, Zhang Y, Tang I, Yang S. Metabolic engineering of *Clostridium tyrobutyricum* for n-butanol production. *Metab Eng*. 2011; 13(4):373-382.
6. Lee SY, Park JH, Jang SH, Nielsen LK, Kim J, Jung KS. Fermentative butanol production by Clostridia. *Biotechnol Bioeng*. 2008; 101(2):209-228.
7. Cheng C, Li WM, Lin M, Yang ST. Metabolic engineering of *Clostridium carboxidivorans* for enhanced ethanol and butanol production from syngas and glucose. *Bioresour. Technol*. 2019; 284:415-423.
8. Abubackar HN, Veiga MC, Kennes C. Production of acids and alcohols from syngas in a two-stage continuous fermentation process. *Bioresour. Technol*. 2018; 253:227-234.

9. Fernández-Naveira Á, Veiga MC, Kennes C. Glucose bioconversion profile in the syngas-metabolizing species *Clostridium carboxidivorans*. *Bioresour. Technol.* 2017; 244:552-559.
10. Lee S, Yun EJ, Kim J, Lee SJ, Um Y, Kim KH. Biomass, strain engineering, and fermentation processes for butanol production by *solventogenic clostridia*. *Appl. Microbiol. Biotechnol.* 2016; 100(19):8255-8271.
11. Klasson KT, Ackerson MD, Clausen EC, Gaddy JL. Biological conversion of coal and coal-derived synthesis gas. *Fuel.* 1993; 72(12):1673-1678.
12. Valgepea K, Loi KQ, Behrendorff JB, Lemgruber RDSP, Plan M, Hodson MP, Köpke M, Nielsen LK, Marcellin E. Arginine deiminase pathway provides ATP and boosts growth of the gas-fermenting acetogen *Clostridium autoethanogenum*. *Metab Eng.* 2017; 41:202-211.
13. Li N, Yang J, Chai C, Yang S, Jiang W, Gu Y. Complete genome sequence of *Clostridium carboxidivorans* P7T, a syngas-fermenting bacterium capable of producing long-chain alcohols. *J Biotechnol.* 2015; 211:44-45.
14. Ragsdale SW, Pierce E. Acetogenesis and the Wood-Ljungdahl pathway of CO₂ fixation. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics.* 2008; 1784(12):1873-1898.
15. Wood HG. Life with CO or CO₂ and H₂ as a source of carbon and energy. *Faseb J.* 1991; 5(2):156-163.
16. Li D, Meng CX, Wu GX, Xie BT, Han YF, Guo YQ, Song CH, Gao ZQ, Huang ZY. Effects of zinc on the production of alcohol by *Clostridium carboxidivorans* P7 using model syngas. *J Ind Microbiol Biot.* 2018; 45(1):61-69.
17. Abubackar HN, Veiga MC, Kennes C. Biological conversion of carbon monoxide to ethanol: Effect of pH, gas pressure, reducing agent and yeast extract. *Bioresour. Technol.* 2012; 114:518-522.
18. Kundiyana DK, Wilkins MR, Maddipati P, Huhnke RL. Effect of temperature, pH and buffer presence on ethanol production from synthesis gas by "*Clostridium ragsdalei*". *Bioresour. Technol.* 2011; 102(10):5794-5799.
19. Shen Y, Brown R, Wen Z. Syngas fermentation of *Clostridium carboxidivoran* P7 in a hollow fiber membrane biofilm reactor: Evaluating the mass transfer coefficient and ethanol production performance. *Biochem Eng J.* 2014; 85:21-29.
20. Huang H, Chai C, Li N, Rowe P, Minton NP, Yang S, Jiang W, Gu Y. CRISPR/Cas9-Based Efficient Genome Editing in *Clostridium ljungdahlii*, an Autotrophic Gas-Fermenting Bacterium. *Acs Synth Biol.* 2016; 5(12):1355-1361.
21. Kannuchamy S, Mukund N, Saleena LM. Genetic engineering of *Clostridium thermocellum* DSM1313 for enhanced ethanol production. *Bmc Biotechnol.* 2016; 16(S1).
22. Liew F, Henstra AM, Köpke M, Winzer K, Simpson SD, Minton NP. Metabolic engineering of *Clostridium autoethanogenum* for selective alcohol production. *Metab Eng.* 2017; 40:104-114.
23. Hashimoto SI. Discovery and History of Amino Acid Fermentation. *Adv Biochem Eng Biotechnol.* 2017; 159:15-34.
24. Sawers RG. Amino Acid Degradation. *eLS.* 2009.
25. Bertsch J, Muller V. CO Metabolism in the Acetogen *Acetobacterium woodii*. *Appl. Environ. Microbiol.* 2015; 81(17):5949-5956.

26. Bao T, Cheng C, Xin X, Wang J, Wang M, Yang S. Deciphering mixotrophic *Clostridium formicoaceticum* metabolism and energy conservation: Genomic analysis and experimental studies. *Genomics*. 2019; 111(6):1687-1694.
27. Braun K, Gottschalk G. Effect of molecular hydrogen and carbon dioxide on chemo-organotrophic growth of *Acetobacterium woodii* and *Clostridium aceticum*. *Arch Microbiol*. 1981; 128(3):294-298.
28. Younesi H, Najafpour G, Mohamed AR. Ethanol and acetate production from synthesis gas via fermentation processes using anaerobic bacterium, *Clostridium ljungdahlii*. *Biochem Eng J*. 2005; 27(2):110-119.
29. Liew F, Henstra AM, Winzer K, Köpke M, Simpson SD, Minton NP. Insights into CO₂ Fixation Pathway of *Clostridium autoethanogenum* by Targeted Mutagenesis. *Mbio*. 2016; 7(3):e416-e427.
30. Jiang W, Zhao J, Wang Z, Yang S. Stable high-titer n-butanol production from sucrose and sugarcane juice by *Clostridium acetobutylicum* JB200 in repeated batch fermentations. *Bioresour. Technol*. 2014; 163:172-179.
31. Bao T, Zhao JB, Li J, Liu X, Yang ST. n-Butanol and ethanol production from cellulose by *Clostridium cellulovorans* overexpressing heterologous aldehyde/alcohol dehydrogenases. *Bioresour. Technol*. 2019; 285:121316.
32. Richter H, Molitor B, Wei H, Chen W, Aristilde L, Angenent LT. Ethanol production in syngas-fermenting *Clostridium ljungdahlii* is controlled by thermodynamics rather than by enzyme expression. *Energ Environ Sci*. 2016; 9(7):2392-2399.
33. Ramio-Pujol S, Ganigue R, Baneras L, Colprim J. Incubation at 25°C prevents acid crash and enhances alcohol production in *Clostridium carboxidivorans* P7. *Bioresour. Technol*. 2015; 192:296-303.
34. Sun X, Atiyeh HK, Kumar A, Zhang HL, Tanner RS. Biochar enhanced ethanol and butanol production by *Clostridium carboxidivorans* from syngas. *Bioresour. Technol*. 2018; 265:128-138.
35. Kelly B, Pearce EL. Amino Assets: How Amino Acids Support Immunity. *Cell Metab*. 2020; 32(2):154-175.
36. Zampieri M, Hörl M, Hotz F, Müller NF, Sauer U. Regulatory mechanisms underlying coordination of amino acid and glucose catabolism in *Escherichia coli*. *Nat Commun*. 2019; 10(1).
37. Marcellin E, Behrendorff JB, Nagaraju S, DeTissera S, Segovia S, Palfreyman RW, Daniell J, Licona-Cassani C, Quek L, Speight R *et al*. Low carbon fuels and commodity chemicals from waste gases-systematic approach to understand energy metabolism in a model acetogen. *Green Chem*. 2016; 18(10):3020-3028.
38. Agu CV, Ujor V, Ezeji TC. Metabolic engineering of *Clostridium beijerinckii* to improve glycerol metabolism and furfural tolerance. *Biotechnol. Biofuels*. 2019; 12(1).

Figures

Figure 1

Effects of different of amino acids and different concentration of citrulline on the growth of *C. carboxidivorans* P7. (A) The effects of 21 amino acids on the growth of *C. carboxidivorans* P7. (B) Effects of different concentrations of citrulline on the growth of *C. carboxidivorans* P7. (C) Effects of different concentrations of citrulline on sugar consumption of *C. carboxidivorans* P7.

Figure 2

Expression of key genes related to citrulline metabolism pathway and key genes of EMP pathway were analyzed by RT-qPCR during *C. carboxidivorans* P7 heterotrophic fermentation in different medium. RT-qPCR analysis of *C. carboxidivorans* P7 fermented for 12 h (A), 24 h (B) and 36 h (C). (D) Citrulline metabolism pathway. (P7: *C. carboxidivorans* P7 was fermented in MM520 medium. P7-0.6 g/L Cit: *C. carboxidivorans* P7 was fermented in MM520-0.6 Cit medium.)

Figure 3

FESEM analysis of *C. carboxidivorans* P7 fermented in MM520 medium (A) and MM520-0.6 Cit medium (B) for 96 h.

Figure 4

Under heterotrophic conditions, the effects of citrulline on the fermentation of *C. carboxidivorans* P7. *C. carboxidivorans* P7 was cultivated in MM520 medium (A) and MM520-0.6 Cit medium (B). (C) doubling time, (D) Specific growth rate, (E) The concentration of ATP.

Figure 5

Under autotrophic conditions, the effects of citrulline on the fermentation of *C. carboxidivorans* P7. (A) Cell density, (B) The concentration of ATP, (C) Specific growth rate, (D) Doubling time.