Development of a gRNA-tRNA Array of CRISPR/Cas9 in Combination with Grafting Technique to Improve Gene Editing Efficiency of Sweet Orange

Xiaomei Tang

Huazhong Agricultural University: Huazhong Agriculture University

Shulin Chen

Huazhong Agricultural University: Huazhong Agriculture University

Huiwen Yu

Huazhong Agricultural University: Huazhong Agriculture University

Xiongjie Zheng

Huazhong Agricultural University: Huazhong Agriculture University

Fei Zhang

Huazhong Agricultural University: Huazhong Agriculture University

Xiuxin Deng

Huazhong Agricultural University: Huazhong Agriculture University

Qiang Xu (✉ xuqiang@mail.hzau.edu.cn)

Huazhong Agricultural University: Huazhong Agriculture University <https://orcid.org/0000-0003-1786-9696>

Research Article

Keywords: Sweet orange, Traditional breeding approaches, CRISPR)/CRISPR-associated protein (Cas) system

Posted Date: August 5th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-765771/v1>

License: (cc) (i) This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Plant Cell Reports on September 23rd, 2021. See the published version at <https://doi.org/10.1007/s00299-021-02781-7>.

Abstract

Sweet orange is one of the most popular fruit crops worldwide. Traditional breeding approaches in sweet orange is impracticable due to the apomixis and long juvenility, making it difficult to obtain hybrids and selection of ideal genotypes. The development of targeted genome engineering technologies made it possible for the precise modification of target genes. Recently, a more efficient gene editing tool has been emerged based on the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein (Cas) system (Bhaya et al. 2011). The development of CRISPR/Cas9 technology is promising to accelerate the process of genetic improvement in perennial crops.

Introduction

Sweet orange is one of the most popular fruit crops worldwide. Traditional breeding approaches in sweet orange is impracticable due to the apomixis and long juvenility, making it difficult to obtain hybrids and selection of ideal genotypes. The development of targeted genome engineering technologies made it possible for the precise modification of target genes. Recently, a more efficient gene editing tool has been emerged based on the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein (Cas) system (Bhaya et al. 2011). The development of CRISPR/Cas9 technology is promising to accelerate the process of genetic improvement in perennial crops.

Gene editing in sweet orange are usually faced with difficulties including low editing efficiency and low root formation rate, which severely limits the advancement of gene function studies and application in molecular breeding in sweet orange. Recently, the application of *Arabidopsis* YAO promoter driven CRISPR/Cas9 system in citrus rootstock cultivar Carrizo Citrange significantly improved gene editing efficiency than the 35S promoter driven CRISPR/Cas9 system (Jia and Wang 2014; Zhang et al. 2017; Alvarez et al. 2021). Meanwhile, studies have shown that polycistronic tRNA-sgRNA (PTG) /Cas9 is more efficient for multiplex gene editing than traditional CRISPR/Cas9 system (Xie et al. 2015; Wang et al. 2018), and for Carrizo Citrange (Huang et al. 2020). Thus, it is possible to further improve the editing efficiency by adding the PTG to the YAO-driven Cas9 system in sweet orange. In addition, improving the surviving rate of transgenic sweet orange seedlings by avoiding the high risk of root induction process still requires further investigation (Belide et al. 2011). All those limitations impeding the process of generating healthy sweet orange plants with targeted genome editing using CRISPR/Cas9 system. To overcome those limitations, we developed a reliable protocol for the fast and efficient gene-edited sweet orange plants production. The application of *in vitro* shoot grafting technology significantly reduced the growth cycle of transgenic seedlings, and the survival rate of cleft grafting was more than 90% (Fig. 1l). In addition, the gene editing efficiency was significantly improved by short-term heat stress treatment (Fig. 1o). Thus, our strategies provided a reference for the fast and efficient multiplex genome editing of sweet orange.

To further explore the application of PTG/Cas9 system in sweet orange, three gRNAs targeting phytoene desaturase (*PDS*) in Anliu sweet orange were used (Fig. 1a) (Jia and Wang 2014; Zhang et al. 2017). The

three targets were assembled into pCAMBIA1300-pYAO:Cas9-eGFP vector (Fig. 1b). Statistical analysis revealed that the editing efficiency was about 36.7%-50% by Hi-Tom sequencing (Fig. 1c), and the albino phenotype can be precisely induced using the PTG/Cas9 system after 1 to 2 months (Fig. 1d). Sanger sequencing revealed that most of the mutations induced by PTG/Cas9 in sweet orange were small InDels (Fig. 1e-f, Fig. S1).

Genetic transformation in sweet orange has been vastly studied, and the transformation efficiency varies among sweet orange varieties. In this study, we optimized the transformation conditions by using vacuum negative pressure with the *Agrobacterium*-mediated genetic transformation of epicotyls for 5 min by using Anliu genotype (Fig. S2a). The transformation efficiency was 18.22%-21.15% (Fig. S2b). In order to shorten the time and improve the survival rate of transgenic sweet orange, transgenic seedlings were grafted on the sweet orange rootstocks *in vitro* (Fig. 1g-k). Our data showed that the survival rate of cleft grafting was more than 90% (Fig. 1l). As previous studies illustrated that heat stress could increase targeted mutagenesis induced by CRISPR/Cas9. In order to improve the mutation efficiency of grafted mosaic sweet orange (Fig. 1m), 10-month old grafted mosaic plants were exposed to short-term heat stress treatments at 37°C. We observed new albino tissues grown from mosaic seedlings 15 d after heat stress treatments (Fig. 1n, S3c-d), as well as the increased targeted mutagenesis (Fig. 1o).

The endogenous tRNA processing system exists in almost all species, which can not only process and cleave multiple gRNAs in one sgRNA expression cassette, but also act as transcription enhancers to enhance the expression of gRNAs (Xie et al. 2015). In this study, we evaluated the editing efficiency induced by the three PTG/Cas9 constructs using *Agrobacterium*-mediated genetic transformation of sweet orange epicotyls. For the target sites, the majority of the mutated alleles identified consisted of some deletions and therefore caused a shift of reading frame, while the mutation mediated by PTG/Cas9 system was a little different with YAO promoter-driven CRISPR/Cas9 system (Zhang et al. 2017). Meanwhile, we examined the off-target effects of the three PTG/Cas9 constructs, and no off-target mutations were detected in this study (primers used for amplification was listed in Table S1). Here, the editing efficiency of sweet orange induced by the three PTG/Cas9 vectors was not as efficient as the Carrizo Citrange (Zhang et al. 2017). We inferred that this may be related with the expression specificity of the YAO promoter (Li et al. 2010) and a single AtU6 promoter was used to drive the expression of several gRNAs, which requires further processing. The gene-edited seedlings in this study were directly induced from stem sections in 1-2 months (Fig. S2a), while the transgenic plants of Carrizo Citrange were regenerated from callus, in which the prolonged YAO promoter activity may modulate the expression of SpCas9 (Zhang et al. 2017). In addition to promoters, temperature could also regulate the activity of SpCas9 and the mutation efficiency (LeBlanc et al. 2018). We found that the the mutation efficiency of chimera sweet orange plants was also significantly increased after heat stress treatments (Fig. 1o), our result was consistent with previous report in *Arabidopsis* and Carrizo Citrange (LeBlanc et al. 2018).

Cleft grafting *in vitro* could significant improve the survival rate of transgenic seedlings, and this was a suitable method for those plants which are hardly to root or susceptible to various soil pathogenic bacteria (Belide et al. 2011). The combination of grafting and heat stress treatment may be a suitable

method for the knockout of plant reproduction- and development-related genes, which may affect plant regeneration or rooting if they are mutated at early growth stages. Here, we developed a fast and efficient gene editing method for sweet orange, our approach offers a new way to facilitate the efficient multiplex gene editing of sweet orange, and this approach is especially suitable for the genetic improvement of citrus varieties.

Declarations

Author contribution statement X.T., QX. and X.D. conceived and designed the experiments, X.T., S.C., H.Y. and X.Z. conducted the experiments. X.T. and S.C. analyzed data. X.T. wrote the manuscript. Q.X. and F.Z. directed the study and revised the manuscript. All authors read and approved the manuscript.

Acknowledgments We are grateful to Dr. Qi Xie from Institute of Genetics and Development, Chinese Academy of Sciences for providing the *YAO* promoter driven CRISPR/Cas9 vector. This work was supported by the National Key Research and Development Program of China (NO. 2018YFD1000101), National Natural Science Foundation of China (NOs. 31925034 and 31872052), and the Fundamental Research Funds for the Central Universities.

Conflict of interest The authors declare no conflict of interest.

References

1. Alvarez D, Cerda-Bennasser P, Stowe E, Ramirez-Torres F, Capell T, Dhingra A, Christou P (2021) Fruit crops in the era of genome editing: closing the regulatory gap. *Plant Cell Rep* 40 (6):915-930. doi:10.1007/s00299-021-02664-x
2. Belide S, Hac L, Singh SP, Green AG, Wood CC (2011) Agrobacterium-mediated transformation of safflower and the efficient recovery of transgenic plants via grafting. *Plant Methods* 7 (1):12-12
3. Bhaya D, Davison M, Barrangou R (2011) CRISPR-Cas systems in bacteria and archaea: versatile small RNAs for adaptive defense and regulation. *Annu Rev Genet* 45:273-297. doi:10.1146/annurev-genet-110410-132430
4. Huang X, Wang Y, Xu J, Wang N (2020) Development of multiplex genome editing toolkits for citrus with high efficacy in biallelic and homozygous mutations. *Plant Mol Biol* 104 (3):297-307. doi:10.1007/s11103-020-01043-6
5. Jia H, Wang N (2014) Targeted Genome Editing of Sweet Orange Using Cas9/sgRNA. *Plos One* 9:e93806. doi:10.1371/journal.pone.0093806
6. LeBlanc C, Zhang F, Mendez J, Lozano Y, Chatpar K, Irish VF, Jacob Y (2018) Increased efficiency of targeted mutagenesis by CRISPR/Cas9 in plants using heat stress. *Plant J* 93 (2):377-386. doi:10.1111/tpj.13782
7. Li HJ, Liu NY, Shi DQ, Liu J, Yang WC (2010) *YAO* is a nucleolar WD40-repeat protein critical for embryogenesis and gametogenesis in Arabidopsis. *BMC Plant Biol* 10:169. doi:10.1186/1471-2229-

8. Wang Z, Wang S, Li D, Zhang Q, Li L, Zhong C, Liu Y, Huang H (2018) Optimized paired-sgRNA/Cas9 cloning and expression cassette triggers high-efficiency multiplex genome editing in kiwifruit. *Plant Biotechnol J* 16 (8):1424-1433
9. Xie K, Minkenberg B, Yang Y (2015) Boosting CRISPR/Cas9 multiplex editing capability with the endogenous tRNA-processing system. *Proc Natl Acad Sci U S A* 112 (11):3570-3575. doi:10.1073/pnas.1420294112
10. Zhang F, LeBlanc C, Irish VF, Jacob Y (2017) Rapid and efficient CRISPR/Cas9 gene editing in Citrus using the *YAO* promoter. *Plant Cell Rep* 36 (12):1883-1887. doi:10.1007/s00299-017-2202-4

Figures

Figure 1

Fast and efficient gene editing technology in sweet orange by using YAO promoter-driven PTG/Cas9 system in combination with grafting. (a) Schematic diagram of three gRNAs targeting the CsPDS gene. Blue rectangles represent the location of the three gRNAs. (b) Schematic map of the PTG/Cas9 system. Cas9 gene was driven by the Arabidopsis YAO promoter, purple rounded rectangles represent tRNA, diamonds with different colors indicate gRNAs, and the white rounded rectangles show sgRNA scaffold.

(c) Editing efficiency induced by different PTG/Cas9 vectors. (d) Phenotypes induced by PTG/Cas9 system in Anliu sweet orange. (e-f) Sanger sequencing of site-target mutation in Anliu sweet orange. The nucleotide mutations compared with WT (wild type) are displayed in DNA sequence of CsPDS. The sequences in green represent gRNAs, and the PAM sites are in blue. The deleted nucleotides are shown in black dots. The inserted nucleotides are shown in red. (g) Transgenic lines used as scion for grafting. (h) Rootstock preparation for grafting. (i-j) V-shaped scions were grafted onto prepared sweet orange rootstocks. (k) Grafted seedlings were cultured in vitro with running water at 25°C in the cycle of 16 h of light and 8 h of darkness. (l) Statistical analysis of graft survival rate of Anliu sweet orange. (m) Phenotypes of grafted CsPDS gene-edited chimeric seedlings cultivated at 25°C. (n) New albino tissues formed 15 d after heat stress treatment of 37°C. (o) Analyzing of the mutation efficiency in CsPDS gene-edited citrus plants exposed to heat stress. Old and young leaves formed before heat stress (HS) treatment (continuously grown at 25°C) and after heat stress treatment were used.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryfile.docx](#)