

DNA barcode *trnH-psbA* is a promising candidate for efficient identification of forage legumes and grasses

Miguel Loera-Sánchez

ETH Zurich

Bruno Studer

ETH Zurich

Roland Kölliker (✉ roland.koelliker@usys.ethz.ch)

ETH Zurich <https://orcid.org/0000-0003-1959-0402>

Research note

Keywords: DNA barcoding, forages, grasslands, species richness

Posted Date: November 8th, 2019

DOI: <https://doi.org/10.21203/rs.2.17026/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Research Notes on January 17th, 2020. See the published version at <https://doi.org/10.1186/s13104-020-4897-5>.

Abstract

Objective Grasslands are widespread ecosystems that fulfil many functions. Plant species richness (PSR) is known to have beneficial effects on such functions and monitoring PSR is crucial for tracking the effects of land use and agricultural management on these ecosystems. Unfortunately, traditional morphology-based methods are labor-intensive and cannot be adapted for high-throughput assessments. DNA barcoding could aid increasing the throughput of PSR assessments in grasslands. In this proof-of-concept work, we aimed at determining which of three plant DNA barcodes (*rbcLa* , *matK* and *trnH-psbA*) best discriminates 16 key grass and legume species common in temperate sub-alpine grasslands.

Results Barcode *trnH-psbA* had a 100% correct assignment rate (CAR) in the five analyzed legumes, followed by *rbcLa* (93.3%) and *matK* (55.6%). Barcode *trnH-psbA* had a 100% CAR in the grasses *Cynosurus cristatus* , *Dactylis glomerata* and *Trisetum flavescens* . However, the closely related *Festuca*, *Lolium* and *Poa* species were not always correctly identified, which led to an overall CAR in grasses of 66.7%, 50.0% and 46.4% for *trnH-psbA* , *matK* and *rbcLa* , respectively. Barcode *trnH-psbA* is thus the most promising candidate for PSR assessments in permanent grasslands and could greatly support plant biodiversity monitoring on a larger scale.

Introduction

Grasslands are some of the most widespread ecosystems on Earth, covering two fifths of its land surface [1]. They provide roughage for ruminant livestock production and many other environmental services related to carbon sequestration, water flow regulation and soil stabilization [2, 3]. Plant species richness (PSR) is a component of biodiversity with major effects on the ecosystem functioning of grasslands: in experimental grassland plant communities, high levels of PSR stabilize yields and confer tolerance against environmental stressors [4]. Similar effects have been observed in semi-natural grasslands [5]. Assessing PSR is thus crucial for tracking its changes and effects on ecosystem services. However, such assessments have traditionally relied on morphology-based surveys that are labor-intensive and require trained taxonomists, limiting their use for surveying PSR over large scales and long time periods [3]. Furthermore, grasses and legumes (the two plant families of major economic relevance in temperate grasslands) can be taxonomically assessed with highest precision only when certain distinctive morphological characters are on display (e.g., flowering bodies and leaves). Still, some grass and legume species are difficult to distinguish from closely related species. A standardized, precise, high-throughput solution for PSR surveys in grasslands is therefore desirable for large-scale assessments of changes in PSR.

DNA barcoding is a methodology that has been successfully applied for standardizing and increasing the throughput of PSR surveys in ecological studies [6, 7]. DNA barcodes are organellar or nuclear loci that show a high degree of species-level conservation [8, 9]. By comparing newly sequenced DNA barcodes to reference databases, it is possible to assign an unknown biological sample to its correct taxonomy. An

international effort is currently in place to maintain a well-curated, public reference database of DNA barcodes (The Barcode Of Life Datasystems database, BOLD [10]).

In animals, the DNA barcode of choice is the mitochondrial *COI* gene, which can reproducibly differentiate most of the major animal phyla [8]. In plants, in contrast, there is no single DNA barcode with comparable success [11]. Most plant DNA barcodes are located in the chloroplast genome, either within coding sequences (such as *rbcLa* and *matK*) or in intergenic regions (such as *trnH-psbA*) [11, 12]. Some nuclear loci have also been used as DNA barcodes [13]. More than one barcode per plant individual are typically sequenced and used for taxonomical assignments [11, 12]. However, sequencing more than one DNA barcode per plant may not be technically feasible in higher throughput settings, particularly when analyzing mixed-species samples.

The aim of the present study was to determine the best DNA barcode sequences for forage species by screening the BOLD database for promising candidates and sequencing three DNA barcodes (*rbcLa*, *matK* and *trnH-psbA*) from different cultivars of 16 forage plant species that are common in sub-alpine grasslands.

Methods

Plant material and DNA extraction

Seeds of 2–3 cultivars of 16 forage species (*Alopecurus pratensis* L., *Arrhenaterum elatius* L., *Cynosurus cristatus* L., *Dactylis glomerata* L., *Festuca pratensis* Huds., *F. rubra* L., *Lolium perenne* L., *L. multiflorum* Lam., *Lotus corniculatus* L., *Medicago sativa* L., *Phleum pratense* L., *Poa pratensis* L., *Trifolium pratense* L., *T. repens* L. and *Trisetum flavescens* L.) were germinated and their seedlings were transferred into pot trays (77 wells, 50 cm x 32 cm, with compost as substrate). Plants were grown for 3 weeks after which DNA was extracted from three plants per species. A list of the selected plant specimens and cultivar name is presented in Table 1. For grasses, three leaf fragments of ~1 cm and for legumes three young leaflets were harvested. The plant material was freeze-dried for 48 h and pulverized in a QIAGEN TissueLyser II (QIAGEN, Hilden, Germany). DNA was extracted using the NucleoSpin® II kit (Macherey-Nagel, Düren, Germany) and its integrity visually inspected by agarose gel electrophoresis (1% w/v). DNA purity and concentration were determined with a NanoDrop™ spectrophotometer (ThermoFisher Scientific, Waltham, MA, USA).

DNA barcode amplification and sequencing

The BOLD database was screened for DNA barcode sequences for our selected species and close relatives; barcodes *rbcLa*, *matK* and *trnH-psbA* were selected as candidates because they reported the most available sequences. Primer sequences for the three barcodes were obtained from BOLD [10] and were optimized for amplification in the target plant families (Additional file table S1). Each PCR reaction consisted of 15 ng of template DNA, 1 x flexi buffer (Promega, Madison, WI, USA), 2 mM MgCl₂, 200 μM

dNTPs, each primer at 0.4 μ M, 0.75 units of GoTaq® G2 Flexi DNA Polymerase (Promega, Madison, WI, USA) and water to a final volume of 30 μ L.

For *rbclA*, PCR conditions were 5 min at 94°C followed by 33 cycles of 40 s at 94°C, 1 min at 55°C and 40 s at 72°C, followed by a final extension cycle of 10 min at 72°C. For *matK* and *trnH-psbA*, a 5 min at 94°C followed by 50 cycles of 40 s at 94°C, 1 min at 54°C and 40 s at 72°C followed by a final extension cycle of 10 min at 72°C were used.

Amplicons were purified in a MultiScreen PCR96 filter plate (Merck, Darmstadt, Germany). Sequencing reactions were prepared with 1 x BigDye™ Terminator 3.1 Reaction Mix (ThermoFisher Scientific, Waltham, MA, USA), 1 x BigDye™ 3.1 Sequencing Buffer, forward or reverse primer at 0.16 μ M and 800 ng of purified amplicon to a final volume of 5 μ L. The same primers used for PCR were used for sequencing. Capillary electrophoresis was performed on a 3130 ABI (ThermoFisher Scientific, Waltham, MA, USA). The resulting traces were quality filtered and merged using GAP4 [14] with the default settings. All traces and sequences were uploaded to BOLD v4 (project code: SWFRG; http://boldsystems.org/index.php/MAS_Management_DataConsole?codes = SWFRG).

Taxonomical assignments

Sequences of *matK*, *rbclA* and *trnH-psbA* were downloaded from BOLD v4 on May 23, 2019 [10]. Only sequences from the Poaceae and Fabaceae families with no contaminants and longer than 200 bp were included. In total, 6,232 *rbclA*, 11,971 *matK* and 1,236 *trnH-psbA* sequences were present in the downloaded fasta files, which also include the plants from the BOLD project SWFRG (Supplementary Table S2). The taxonomical identifiers of the BOLD fasta files were reformatted to remove spaces and rearrange their informative fields in a consistent manner (*fasta_name_reformat.py* script from <https://github.com/mloera/forage-barcoding>).

Each barcode-specific fasta file was then used to make a blast database and the SWFRG sequences were queried in their corresponding database with blastn using the flag *outfmt = 6* (i.e., tabular format). The resulting blast output tables were parsed with the *blastn_matcher.R* script from the above-mentioned GitHub repository. The script removes self-hits and corrects some misspellings in the taxonomy of queries and hits. The script then compares the taxonomy of the queries and hits at the species- and genus-level. A “match” was called when the taxonomy of a query sequence is equal to the taxonomy of the highest scoring hit or hits. A “taxonomical assignment rate” for each barcode was then calculated as the ratio between the sum of its correct taxonomical assignments and the total number of query sequences.

Results And Discussion

PCR and sequencing results

The primer sequences of *trnH-psbA* and *matK* were adapted to allow for amplification within the target species, while the primer sequences of *rbcLa* did not need any modification (Supplementary Table 2). From the 48 processed specimens, 115 sequences were obtained (46 for *matK*, 43 for *rbcLa* and 41 for *trnH-psbA*) after repeating and optimizing failed amplifications. The size of the sequences ranged from 470 to 588 bp for *rbcLa*, 185 to 888 bp for *matK* and 268 to 614 bp for *trnH-psbA* (Table 1).

Taxonomical assignments

Barcode *trnH-psbA* had a 100% correct assignment rate (CAR) in legumes, followed by *rbcLa* (93.3%) *matK* (57.1%; Table 2). The highest CAR for grasses was 65.4% with *trnH-psbA*, followed by *matK* (48.4%) and *rbcLa* (46.4%). Overall, genus-level CARs were 69.8%, 73.3% and 90.2% for *rbcLa*, *matK* and *trnH-psbA*, respectively. Legumes had also the highest assignment rate on the genus level (100% correct assignments for all barcodes; Table 2), while correct assignments for grass genera were 53.6%, 61.3% and 84.6% for barcodes *rbcLa*, *matK* and *trnH-psbA*, respectively.

The low CARs for grass DNA barcodes could be due to various factors. Some grass species, such as *Poa* spp., are notoriously hard to discriminate morphologically and their phylogeny is subject to controversy [15, 16]. This could have resulted in misidentified reference sequences. Another factor is the high genetic similarity between some grass taxa. For example, the genetic similarity of some species of the *Festuca-Lolium* complex is reported to be >90%, as calculated from transcriptomic data of orthologous genes [17]. This may result in a higher proportion of incorrect taxonomic assignments for such grass species [18].

Barcode *trnH-psbA* makes for a good candidate for large-scale DNA barcoding of forage legumes and some grasses, such as *C. cristatus*, *D. glomerata* and *T. flavescens* (Table S3), although further work is needed to produce reference sequences in more forage species and cultivars. Overall, our results provide the basic tools to implement DNA barcoding in forage species (i.e., family-specific primer pairs and a standard bioinformatic workflow for taxonomic assignments) and can help in choosing an appropriate DNA barcode for high-throughput applications. Such high-throughput applications could greatly enhance biodiversity-monitoring protocols for the study of grassland ecology.

Limitations

- This is exploratory work focused on the most common forage plant species; further work is needed to address less common forage species.
- As a proof of concept, three specimens per species were analyzed.

List Of Abbreviations

- BOLD: Barcode Of Life Datasystems
- CAR: Correct assignment rate

- PSR: plant species richness

Declarations

- Ethics approval and consent to participate

Not applicable

- Consent for publication

Not applicable

- Availability of data and material

The datasets generated and/or analysed during the current study are available in the following GitHub repository: <https://github.com/mloera/forage-barcoding>. Sequencing trace files are found in BOLD under the project name “SWFRG”.

- Competing interests

The authors declare that they have no competing interests.

- Funding

This work was founded by the Swiss Federal Office of Agriculture (FOAG).

- Authors' contributions

ML obtained and analysed the data and drafted the manuscript. RK and BS conceived the study and assisted drafting the manuscript. All authors read and approved the final version of the manuscript.

- Acknowledgements

Data produced and analyzed in this paper were generated in collaboration with the Molecular Ecology group from Agroscope and the Genetic Diversity Centre (GDC), ETH Zurich.

Tables

Table 1. Size of barcode sequences obtained for 48 plants from 16 different species of forage grasses and legumes

BOLD Process ID	Species	Cultivar	Sequence size in bp [number of n's in sequence]		
			<i>rbcL</i> a	<i>matK</i>	<i>trnH-psbA</i>
SWFRG013-19	<i>Alopecurus pratensis</i>	'Alko' (Saatzucht Steinach, DE)	0 ^a	838[0n]	0 ^a
SWFRG014-19	<i>Alopecurus pratensis</i>	'Alopex' (Agroscope, CH)	0 ^a	663[0n]	0 ^a
SWFRG029-19	<i>Alopecurus pratensis</i>	'Alopex' (Agroscope, CH)	552[0n]	185[0n]	569[0n]
SWFRG015-19	<i>Arrhenatherum elatius</i>	'Arone' (Saatzucht Steinach, DE)	549[0n]	436[0n]	512[0n]
SWFRG016-19	<i>Arrhenatherum elatius</i>	'Median' (DLF Životice, CZ)	550[0n]	610[0n]	0 ^a
SWFRG031-19	<i>Arrhenatherum elatius</i>	'Median' (DLF Životice, CZ)	0 ^a	825[0n]	0 ^a
SWFRG030-19	<i>Cynosurus cristatus</i>	'Cresta' (DSP/INH, CH)	541[0n]	513[0n]	466[0n]
SWFRG045-19	<i>Cynosurus cristatus</i>	'Lena' (HBLF, AT)	585[0n]	531[0n]	564[0n]
SWFRG046-19	<i>Cynosurus cristatus</i>	'Rožnovská' (OSEVA PRO, CZ)	529[0n]	870[0n]	569[0n]
SWFRG001-19	<i>Dactylis glomerata</i>	'Barexcel' (Barenbrug, NL)	577[0n]	640[0n]	519[0n]
SWFRG002-19	<i>Dactylis glomerata</i>	'Brennus' (R2n, FR)	546[0n]	865[13n]	576[0n]
SWFRG017-19	<i>Dactylis glomerata</i>	'Reda' (DSP/INH, CH)	534[0n]	866[13n]	561[0n]
SWFRG003-19	<i>Festuca pratensis</i>	'Cosmolit' (Saatzucht Steinach, DE)	547[0n]	579[0n]	558[0n]
SWFRG004-19	<i>Festuca pratensis</i>	'Paradisla' (DSP/INH, CH)	549[0n]	888[7n]	566[0n]
SWFRG019-19	<i>Festuca pratensis</i>	'Pradel' (DSP/INH, CH)	552[0n]	590[0n]	553[0n]
SWFRG007-19	<i>Festuca rubra</i>	'Echo' (DLF-Trifolium, DK)	559[0n]	886[3n]	274[21n]
SWFRG008-19	<i>Festuca rubra</i>	'Pran Solas' (Schweizer, CH)	588[0n]	869[0n]	570[0n]
SWFRG023-19	<i>Festuca rubra</i>	'Roland' (Saatzucht Steinach, DE)	558[0n]	543[0n]	594[0n]
SWFRG005-19	<i>Lolium multiflorum</i>	'Axis' (DSP/INH, CH)	581[0n]	874[0n]	551[0n]
SWFRG006-19	<i>Lolium multiflorum</i>	'Caribu' (DSP/INH, CH)	577[0n]	884[8n]	567[7n]
SWFRG021-19	<i>Lolium multiflorum</i>	'Zebra' (DSP/INH, CH)	571[0n]	586[0n]	551[0n]
SWFRG009-19	<i>Lolium perenne</i>	'Arara' (DSP/INH, CH)	547[0n]	883[3n]	539[5n]
SWFRG010-19	<i>Lolium perenne</i>	'Arvella' (DSP/INH, CH)	582[0n]	481[0n]	567[15n]
SWFRG025-19	<i>Lolium perenne</i>	'Lipresso' (Euro Grass, DE)	488[0n]	835[0n]	614[0n]
SWFRG024-19	<i>Lotus corniculatus</i>	'Lotar' (OSEVA UNI, SK)	544[0n]	399[0n]	294[0n]
SWFRG039-19	<i>Lotus corniculatus</i>	'Lotar' (OSEVA UNI, SK)	548[0n]	509[0n]	414[0n]
SWFRG040-19	<i>Lotus corniculatus</i>	'Polom' (CVRV, VÚRV, CZ)	502[0n]	702[0n]	412[0n]

SWFRG022-19	<i>Medicago sativa</i>	'Artemis' (Barenbrug, NL)	580[0n]	410[0n]	268[14n]
SWFRG037-19	<i>Medicago sativa</i>	'Catera' (Saatzucht Steinach, DE)	526[0n]	435[0n]	438[3n]
SWFRG038-19	<i>Medicago sativa</i>	'Sanditi' (Barenbrug, NL)	548[0n]	432[0n]	445[2n]
SWFRG028-19	<i>Onobrychis viciifolia</i>	'Perdix' (DSP/INH, CH)	550[0n]	576[0n]	289[0n]
SWFRG043-19	<i>Onobrychis viciifolia</i>	'Perly' (DSP/INH, CH)	582[0n]	627[0n]	284[0n]
SWFRG044-19	<i>Onobrychis viciifolia</i>	'Višňovský' (Agrogen, CZ)	543[0n]	694[10n]	287[0n]
SWFRG032-19	<i>Phleum pratense</i>	'Anjo' (ILVO, BE)	540[0n]	0 ^a	0 ^a
SWFRG047-19	<i>Phleum pratense</i>	'Tiller' (DLF-Trifolium, DK)	576[0n]	527[5n]	584[0n]
SWFRG048-19	<i>Phleum pratense</i>	'Toro' (CRA-FLC, IT)	0 ^a	516[1n]	0 ^a
SWFRG011-19	<i>Poa pratensis</i>	'Likollo' (DSV, DE)	470[0n]	865[0n]	540[0n]
SWFRG012-19	<i>Poa pratensis</i>	'Nixe' (Saatzucht Steinach, DE)	571[0n]	868[0n]	576[0n]
SWFRG027-19	<i>Poa pratensis</i>	'Tommy' (DLF-Trifolium, DK)	0 ^a	489[0n]	0 ^a
SWFRG020-19	<i>Trifolium pratense</i>	'Bonus' (Selgen, CZ)	549[0n]	0 ^a	410[0n]
SWFRG035-19	<i>Trifolium pratense</i>	'Diplomat' (DSV, DE)	564[0n]	556[0n]	485[0n]
SWFRG036-19	<i>Trifolium pratense</i>	'Pavo' (DSP/INH, CH)	514[0n]	430[0n]	496[0n]
SWFRG026-19	<i>Trifolium repens</i>	'Beaumont' (CW 090; Barenbrug, NL)	550[0n]	419[0n]	448[0n]
SWFRG041-19	<i>Trifolium repens</i>	'Bombus' (DSP/INH, CH)	579[0n]	481[0n]	471[0n]
SWFRG042-19	<i>Trifolium repens</i>	'Hebe' (Svalöf-Weibull, SE)	571[0n]	444[0n]	447[0n]
SWFRG018-19	<i>Trisetum flavescens</i>	'Gunther' (HBLFA, AT)	504[0n]	859[6n]	570[0n]
SWFRG033-19	<i>Trisetum flavescens</i>	'Gunther' (HBLFA, AT)	575[0n]	586[4n]	571[0n]
SWFRG034-19	<i>Trisetum flavescens</i>	'Trisett51' (Saatzucht Steinach, DE)	558[0n]	887[4n]	568[0n]
Total sequences			43	46	41
			(89.58%)	(95.83%)	(85.42%)

^a repeatedly unsuccessful PCR

Table 2. Species- and genus-level assignment success by barcode

Barcode	Species-level assignment rate			Genus-level assignment rate		
	Overall	Grasses	Legumes	Overall	Grasses	Legumes
<i>rbcLa</i>	62.8 %	46.4%	93.3 %	69.8 %	53.6 %	100.0%
<i>matK</i>	51.1 %	48.4%	57.1%	73.3%	61.3%	100.00%
<i>trnH-psbA</i>	78.0%	65.4%	100.0%	90.2%	84.6 %	100.0%

Table 3. Species-level taxonomic assignment success by family, query species and barcode sequence

Family	Query species	<i>matK</i>	<i>rbcLa</i>	<i>trnH-psbA</i>
Poaceae	<i>Alopecurus pratensis</i>	0/2	0/1	0/1
Poaceae	<i>Arrhenatherum elatius</i>	1/3	0/2	1/1
Poaceae	<i>Cynosurus cristatus</i>	3/3	2/3	3/3
Poaceae	<i>Dactylis glomerata</i>	3/3	2/3	3/3
Poaceae	<i>Festuca pratensis</i>	2/3	0/3	1/3
Poaceae	<i>Festuca rubra</i>	1/3	3/3	2/3
Poaceae	<i>Lolium multiflorum</i>	0/3	2/3	1/3
Poaceae	<i>Lolium perenne</i>	2/3	0/3	2/3
Poaceae	<i>Phleum pratense</i>	1/2	1/2	1/1
Poaceae	<i>Poa pratensis</i>	0/3	1/2	0/2
Poaceae	<i>Trisetum flavescens</i>	2/3	2/3	3/3
Fabaceae	<i>Lotus corniculatus</i>	1/3	3/3	3/3
Fabaceae	<i>Medicago sativa</i>	2/3	3/3	3/3
Fabaceae	<i>Onobrychis viciifolia</i>	2/3	3/3	3/3
Fabaceae	<i>Trifolium pratense</i>	2/2	2/3	3/3
Fabaceae	<i>Trifolium repens</i>	1/3	3/3	3/3

References

1. Reynolds SG. Chapter 1 Introduction. In: Suttie JM, Reynolds SG, Batello C, editors. Grasslands of the World. Food and Agriculture Organization of the United Nations; 2005. <http://www.fao.org/3/y8344e00.htm>.
2. Lindborg R, Bengtsson J, Berg Å, Cousins SAO, Eriksson O, Gustafsson T, et al. A landscape perspective on conservation of semi-natural grasslands. *Agric Ecosyst Environ.* 2008;125:213–22. doi:10.1016/j.agee.2008.01.006.
3. Bengtsson J, Bullock JM, Egoh B, Everson C, Everson T, O'Connor T, et al. Grasslands-more important for ecosystem services than you might think. *Ecosphere.* 2019;10:e02582. doi:10.1002/ecs2.2582.
4. Wang Y, Cadotte MW, Chen Y, Fraser LH, Zhang Y, Huang F, et al. Global evidence of positive biodiversity effects on spatial ecosystem stability in natural grasslands. *Nat Commun.* 2019;10:3207. doi:10.1038/s41467-019-11191-z.
5. Feßel C, Meier IC, Leuschner C. Relationship between species diversity, biomass and light transmittance in temperate semi-natural grasslands: is productivity enhanced by complementary light capture? *J Veg Sci.* 2016;27:144–55. doi:10.1111/jvs.12326.
6. Kress WJ. Plant DNA barcodes: Applications today and in the future. *J Syst Evol.* 2017;55:291–307. doi:10.1111/jse.12254.
7. Kress WJ, García-Robledo C, Uriarte M, Erickson DL. DNA barcodes for ecology, evolution, and conservation. *Trends Ecol Evol.* 2015;30:25–35. doi:10.1016/j.tree.2014.10.008.

8. Hebert PDN, Ratnasingham S, de Waard JR. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc R Soc London Ser B Biol Sci.* 2003;270 suppl_1. doi:10.1098/rsbl.2003.0025.
9. Hebert PDN, Cywinska A, Ball SL, DeWaard JR. Biological identifications through DNA barcodes. *Proc R Soc London Ser B Biol Sci.* 2003;270:313–21. doi:10.1098/rspb.2002.2218.
10. Ratnasingham S, Hebert PDN. BARCODING: bold: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Mol Ecol Notes.* 2007;7:355–64. doi:10.1111/j.1471–8286.2007.01678.x.
11. CBOL Plant Working Group. A DNA barcode for land plants. *Proc Natl Acad Sci U S A.* 2009;106:12794–7. doi:10.1073/pnas.0905845106.
12. Kress WJ, Erickson DL. A Two-Locus Global DNA Barcode for Land Plants: The Coding *rbcL* Gene Complements the Non-Coding *trnH-psbA* Spacer Region. *PLoS One.* 2007;2:e508. doi:10.1371/journal.pone.0000508.
13. Bolson M, Smidt E de C, Brotto ML, Silva-Pereira V. ITS and *trnH-psbA* as Efficient DNA Barcodes to Identify Threatened Commercial Woody Angiosperms from Southern Brazilian Atlantic Rainforests. *PLoS One.* 2015;10:e0143049. doi:10.1371/journal.pone.0143049.
14. Staden R, Judge DP, Bonfield JK. Managing Sequencing Projects in the GAP4 Environment. In: Krawetz SA, Womble DD, editors. *Introduction to Bioinformatics: A Theoretical and Practical Approach.* Totowa, NJ: Humana Press; 2003. p. 327–44. doi:10.1007/978–1–59259–335–4_20.
15. Nosov NN, Punina EO, Machs EM, Rodionov A V. Interspecies hybridization in the origin of plant species: Cases in the genus *Poa* sensu lato. *Biol Bull Rev.* 2015;5:366–82. doi:10.1134/S2079086415040088.
16. Patterson JT, Larson SR, Johnson PG. Genome relationships in polyploid *Poa pratensis* and other *Poa* species inferred from phylogenetic analysis of nuclear and chloroplast DNA sequences. *Genome.* 2005;48:76–87. doi:10.1139/g04–102.
17. Czaban A, Sharma S, Byrne SL, Spannagl M, Mayer KFX, Asp T. Comparative transcriptome analysis within the *Lolium/Festuca* species complex reveals high sequence conservation. *BMC Genomics.* 2015;16:249. doi:10.1186/s12864–015–1447-y.
18. Meyer CP, Paulay G. DNA Barcoding: Error Rates Based on Comprehensive Sampling. *PLoS Biol.* 2005;3:e422. doi:10.1371/journal.pbio.0030422.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [loeraResearchNoteSupp.docx](#)