Water, sanitation and hygiene practices associated with improved height-for-age, weight-for-height and weight-for-age z-scores in Nepal.

Som Kumar Shrestha (✉ somkr.stha@gmail.com)

London School of Hygiene and Tropical Medicine Department of Population Health <https://orcid.org/0000-0002-1188-2606>

Don Vicendese

La Trobe University School of Psychology and Public Health

Bircan Erbas

La Trobe University School of Psychology and Public Health

Research article

Keywords: Underweight; Stunting; Wasting; Underweight; WASH

Posted Date: February 21st, 2020

DOI: <https://doi.org/10.21203/rs.2.17090/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Evidence of the influence of water, sanitation and hygiene (WASH) behaviors on childhood nutritional status is inconsistent. Few studies have examined their interactive effects. This study aimed to examine associations and interactions between WASH variables and preschool child undernutrition. **Methods:** Data from a nationally representative sample of 2352 children assessed during the 2016 Nepal Demographic and Health Survey were analyzed by multi-variable linear regression to understand the association between height-for-age (HAZ), weight-for-height (WHZ) and weight-for-age (WAZ) z-scores and WASH variables. Interactions between WASH variables, sex and area of residence on childhood nutritional status were also examined. **Results:** The mean z-score [standard deviation] for children's WAZ, HAZ and WHZ scores were -1.33 [1.1], -1.52 [1.3] and -0.65 [1.1], respectively. A unit increase in cluster sanitation coverage was associated with an increase of 0.30 (95%CI: 0.12 to 0.48) for WAZ and 0.28 (95%CI: 0.001 to 0.56) for HAZ scores. Household water purification practice was associated with an increase of 0.24 (95%CI: 0.07 to 0.41) in WHZ score. Handwashing practice with water and soap was associated with an increase of 0.15 (95%CI: 0.04 to 0.25) in WAZ and 0.13 (95%CI: 0.01 to 0.24) in WHZ scores. The effect of water purification practice was higher for rural areas compared to urban settings for HAZ scores (p-value for interaction=0.002). **Conclusions:** Consistent with findings from other countries in the South Asian region, findings of this study highlight the potential importance of good WASH practices, and therefore the potential of WASH interventions, to contribute to improved nutritional status in rural Nepal.

Background

Childhood under-nutrition is a major public health problem in developing countries. In 2018, globally, approximately 21.9% under-five children were estimated to be stunted and 7.3% children wasted, and almost two out of five stunted children belonged to South Asia (1). Nepal is among countries having the highest prevalence of childhood undernutrition in South Asian region (2). A recent Nepal Demographic and Health Survey (NDHS) showed almost 36% of under five children were stunted, 10% were wasted and 27% were underweight in 2016 (3).

A public health programme for tackling the burden of under-nutrition in children has been a high priority for the Nepal Government. In 2011, Nepal joined the global movement Scaling Up Nutrition as its commitment to end malnutrition (4). Realizing the importance of each sector's potential to contribute to the national goal to reduce childhood undernutrition (5), the Multi-Sector Nutrition Plan (2013-2017) was endorsed and envisioned inter-sectoral collaboration between various sectors such as education, agriculture, health and water and sanitation. Several international donors have pledged their support to tackle childhood malnutrition through funding nutrition specific interventions delivered from health facilities and implementing nutrition sensitive interventions addressing underlying determinants of undernutrition (6–8). Over the past decade, Nepal has made notable progress on improving underlying determinants of childhood nutrition such as education attainment, WASH coverage, maternal and child health and poverty reduction. The same period also coincides with a substantial reduction in childhood undernutrition, particularly for childhood stunting (3,9). Despite these successes, Nepal still lagged behind achieving the target of sustainable development goals for reducing the prevalence of stunting and underweight by 2017 (10). Moreover, the recent national survey showed widespread disparities in the burden of childhood undernutrition across different socio-economic status, ethnic

communities and geographical regions (3). This shows that the current progress made in reducing childhood undernutrition has failed to benefit children from all population subgroups especially among the poor and vulnerable communities living in rural areas. Childhood undernutrition still remains one of the major public health challenges in Nepal.

Several underlying factors are associated with poor nutritional outcomes among children (11, 12) and the evidence on the adverse consequences of poor WASH practices on child's nutrition wellbeing is also growing (13–15). The most common hypothesis is that poor WASH facilities and practices mediate transmission of faecal pathogens that causes diarrhea (16–18), and diarrhea in turn exacerbate undernutrition (19, 20). However, other authors have argued that the primary causal pathway for influence of WASH factors on childhood undernutrition is tropical enteropathy and therefore, the estimates modeled entirely through diarrhea could have underestimated the contribution of WASH factors on childhood undernutrition (21). The relative contribution of WASH interventions on childhood undernutrition is still unresolved in the current literature (21, 22) with some studies providing conflicting results depending upon study design, nutritional outcomes and geographical settings where studies were conducted (11, 21–23). Despite this, a majority of cross-sectional studies have consistently demonstrated a crucial role of improved WASH practices on childhood nutrition outcomes (6, 11, 24, 25). However, wide spread variation in methodological approaches and study settings have often posed a considerable challenge for interpretation and synthesis of study findings, limiting the prospect of providing definitive policy recommendations. Studies illustrating how different WASH factors encompassing all important components, namely safe drinking water, improved sanitation and handwashing practices, are associated with different forms of undernutrition is limited in the current literature. The fact that different WASH components are known to be independently associated with child nutrition outcomes, indicates that failure to adjust for all WASH components could potentially bias study findings. On the other hand, WASH related variables could interact among themselves in a complex manner with differential impacts on nutrition outcomes (26). Only a few studies have explored evidence of interaction among different WASH components on child's nutrition outcomes (14, 23).

Therefore, this study aimed to determine the role of different components of WASH facilities on various forms of childhood undernutrition. Specifically, we intend to evaluate how WASH factors are associated with different nutritional outcomes, WAZ, HAZ and WHZ scores among under-five children. In addition, this study also explored whether combined WASH facilities delivered any synergetic benefit on different forms of nutrition outcomes and whether any synergetic benefit evident across urban and rural settings and by child's sex. Within these aims, we endeavored to find evidence to guide public health policy to scale-up WASH interventions to further accelerate the progress on tackling persistent undernutrition in developing countries.

Methods

Data source and sampling:

This study analyzed cross-sectional data from the nationally representative sample of the NDHS (2016). A detailed description of the sampling methods and methodology is reported elsewhere (3). Briefly, the sampling frame of the NDHS 2016 survey was the updated version of National Population and Housing Census 2011 conducted in Nepal. Nepal consists of 77 districts distributed across three ecological zones (Mountain, Hill

and Terai) and 7 provinces. Almost one-third (34%) of the population were under age 15 and under-five children account for almost 11% of total population. The multi-stage sampling was used and the samples were selected in two stages for the rural areas and in three stages for the urban areas. Each province was segregated into urban and rural areas that yielded a total of 14 sampling strata. At the first stage, the wards were selected independently in each stratum using a probability proportional to size strategy. In the rural areas, the selected wards with a relatively small household number (average of 104 households) directly served as the primary sampling units (PSUs). In the urban area, wards were selected as the PSUs, however, due to their large size (average of 800 households), each ward was further stratified as Enumeration Areas (EA) and one EA was randomly selected from each PSU. In the second stage, a complete household listing was carried out from the selected sampling clusters (wards in rural and EAs in urban) that served as the sampling frame for the selection of households. Finally, the predetermined fixed number of 30 households was selected from each cluster with systematic random selection methods.

The total of 11,473 households from 383 sample wards or enumeration areas were selected for the survey. The final study comprised a total of 11040 households with a response rate of 99% that included 2428 children below five years of age (Fig 1). The survey used a household questionnaire and observational checklist for collecting the information on household WASH facilities and practices.

Study Variables:

Outcome variables:

The variables for measuring child nutritional outcomes were based on anthropometric measures of children using an internationally recognized standard practice for assessment of malnutrition at population level (27). The primary outcome variables were the z-scores for WAZ, HAZ and WHZ, calculated based on the median of the World Health Organization (WHO) reference population. The detail calculation for deriving z-scores is provided elsewhere (28). The three nutrition outcomes measure different form of childhood undernutrition. Stunting (low HAZ z-score) is regarded as an indication of long-term nutritional deficiency characterized by recurrent or chronic illness. Wasting (low WHZ z-score) is regarded as an indication of a recent event due to diarrhea or illness. Underweight (low WAZ z-score) represents both acute and chronic shortage of nutrition (29).

Children were dichotomously categorised with reference to the median of the WHO reference population (30). Children with a z-score equal or above -2 SD of the reference population were categorized as “adequately nourished” children and below -2 SD as “undernourished” for all nutritional outcomes. We excluded anthropometric measurements with a z-score below -6 SD and above +6 SD as outliers.

Exposure variables:

WASH related variables such as cluster sanitation coverage, water treatment practices, and hand washing behaviors with soap and water were the major exposure variables. Sanitation facilities could be of different types, constructed using different locally available materials. Their true benefit is only achieved depending on how effective such facilities prevent direct exposure of human faeces to immediate surroundings. We categorized household sanitation facilities as either “improved”, which included facilities with sewer

connections, septic system connections, pour-flush latrines, ventilated improved pit latrines and pit latrines with a slab or covered pit and “unimproved” for pit latrines without slabs or platforms or open pit, hanging latrines, bucket latrines and open defecation (31). We defined community sanitation coverage as the percentage of households with improved sanitation facilities within sampled clusters.

Another important WASH variable is access to safe drinking water but the methods for how the safe drinking water variable is defined and included in existing nutritional studies vary markedly. They mainly differ on either use of a protected water source (24, 32), piped water source (11, 33) or household water treatment methods (34–36). In resource constraint settings where a water pipeline system is unreliable and not well maintained, the risk of water contamination in the supply chain remains high (16, 37) potentially diluting the effect measure and strength of association. In Nepal too, most of the piped water system is not sufficiently well maintained and fully functional to deliver the expected services (38). Therefore, we selected household water purification practice as a relatively reliable proxy indicator for safe drinking water without giving too much reliance on the safety of the water supply system.

Hand-washing practice with soap and water is an important WASH component but measuring actual hand washing practices at different critical points within the household level has always been challenging due to the following reasons. Firstly, hand washing practice is usually measured at different critical points that make it difficult for summarizing the behavior for individuals at household level. Secondly, the information on hand washing behavior is usually based on participants’ verbal responses and so may induce affirmative bias since participants are more likely to report positive behaviors than actual practice (39, 40). Therefore to overcome these limitations, we included availability of soap and water measured through direct household observation as a proxy indicator for hand washing practices that addresses the need for monitoring hand washing behavior of all household participants and is also less subject to bias compared to verbal response.

Data Analysis:

A linear regression model was used to obtain β -coefficients that depict an increase in z-scores for WAZ, HAZ and WHZ associated with a unit increase in sanitation coverage. While for other WASH components (household water purification practices and availability of soap and water), they represented the difference in z-score associated with whether or not household practice included water treatment methods and hand-washing behaviors.

We conducted supplementary analyses using binary nutritional outcome variables to present the Odds Ratios (OR) for risk of childhood underweight, stunting and wasting associated with WASH variables to indicate if interpretation of our findings are robust and independent of the model selected for statistical analysis.

We adjusted for common confounding variables described in existing literature such as child’s birth weight, child’s age, child’s sex, wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women’s age, women marital status, women’s education, women’s smoking status, women’s Body Mass Index (BMI), Antenatal Care Visit (ANC) 4th visits, institutional delivery, frequency of watching TV and childhood diarrhea over past 2 weeks.

The statistical significance level was set at a p -value less than 0.05. We used sampling weights to adjust for the complex survey design and non-proportionate selection probability. The interaction between different WASH variables, sex and area of residence on childhood undernutrition outcomes were also examined. For interaction analysis, we considered a p -value of <0.1 as the level of significance to not miss important associations (41). This study used Stata SE 14.1 (StrataCorp, College Station, Texas) for the analysis.

Results

This study included a total of 2352 children after excluding samples with incomplete WASH information ($n=62$) and either missing or outlier values for anthropometric measurements ($n=14$).

Household WASH and under-five children nutritional status:

Household WASH status and the mean z -score for WAZ, HAZ and WHZ among under-five children are presented in Table 1. Among all households, 75.7% had improved sanitation, while average cluster sanitation coverage was 75.3%. Household water purification practice and availability of soap and water at the time of survey were 18.3% and 37.5% respectively. The mean z -scores for HAZ, WHZ and WAZ were -1.52 [1.3], -0.65 [1.1] and -1.33 [1.1] respectively (Table 1). Among all children nationally, 35.8% were classified as stunted, 9.8% as wasted and 27% as underweight (Table 2).

Association between WASH related variables and childhood under-nutrition status:

Unadjusted and adjusted β -coefficients depicting the linear associations between WASH variables and childhood nutritional status are presented in table 3. All WASH related factors were found to be significantly associated with indicators of nutritional status in the unadjusted models. In the adjusted model, the cluster sanitation coverage was associated with an increase of 0.28 (95%CI: 0.001 to 0.56) in HAZ score and 0.30 (95%CI: 0.12 to 0.48) in WAZ score. The association was not significant for WHZ (Table 3).

Households with water purification practice were associated with an increase of 0.24 (95%CI: 0.07 to 0.41) in WHZ score but no association was found for WAZ and HAZ scores (Table 3).

Handwashing practice was associated with an increase of 0.13 (95%CI: 0.01 to 0.24) in WHZ score and 0.15 (95%CI: 0.04 to 0.25) in WAZ score (Table 3).

Interaction between WASH related variables and childhood undernutrition:

We found no evidence of interaction between WASH variables on child's WAZ, HAZ and WHZ scores (Table 4).

There was no evidence of interaction between WASH variables and child's sex for under-nutrition outcomes (result not shown). However, we observed a statistically significant interaction between improved water purification practice and area of residence on child's HAZ score (p -value for interaction = 0.02) as shown in table 5.

Supplementary analysis

We did the supplementary analyses by modelling nutritional variables as dichotomous outcome variables. Household availability of soap and water was associated with decreased odds of childhood stunting (OR: 0.70, 95%CI 0.54 to 0.92). Sanitation coverage (OR: 0.53, 95%CI 0.28 to 0.99) and household water purification (OR: 0.39, 95%CI: 0.21 to 0.73) were associated with childhood wasting. The odds ratio of childhood underweight decreased significantly with a unit increase in sanitation coverage (OR: 0.59, 95%CI 0.38 to 0.93) and household availability of soap and water (OR: 0.65, 95%CI 0.51 to 0.84).

Discussion

This study found significant associations between WASH related factors and different forms of undernutrition among under-five children, both unadjusted and after controlling for potential confounders, however, the role of individual WASH factors differed substantially for different nutritional outcomes. Child HAZ score had a significant positive linear association with cluster sanitation coverage, while the WHZ score was positively linearly associated with household water purification practice and availability of soap and water. Increase in WAZ score was associated with increased sanitation coverage and household availability of soap and water. We found no evidence of interaction between different WASH variables on childhood nutrition outcomes. The influence of water purification practices on childhood stunting differed in urban and rural settings but child's sex did not have any effect on WASH and undernutrition association.

Cluster sanitation coverage was associated with increased z-score for HAZ and WAZ and the findings are consistent with other studies that have evaluated the role of community sanitation on undernutrition (11, 13, 42). Similar to our findings, previous studies from Nepal have also identified community sanitation coverage as a key determinant of improving HAZ score (11). Other studies that examined the association between household sanitation and child nutrition status also reported similar conclusions (12, 23). However, sanitation coverage is known to have positive externalities providing benefits beyond the household level and failure to adjust for community level sanitation could have underestimated overall benefits of sanitation on child's nutritional outcomes (13, 42) in those studies.

The cluster sanitation coverage also had a significant influence on child's WAZ score. A Mali study also showed a positive association where the benefit of sanitation coverage incremented child's WAZ but leveled off after reaching 60% coverage. We found a positive linear association where the z-score for WAZ increased consistently with increase in sanitation coverage. Our study failed to find an association between sanitation coverage and WHZ z-score, in contrast to earlier studies in Nepal (11) and Bangladesh (43) where positive associations were reported. It should be noted that in the supplementary analysis where we modelled childhood wasting (dichotomized version of WHZ z-score using the WHO reference population), cluster sanitation coverage was significantly associated with decreased likelihood of wasting (Supplementary Table 1). These results suggest methodological differences in assumptions about how the outcome variable is defined and functional fit in statistical modeling could explain at least some discrepancies in study findings across different nutritional researches.

In our adjusted model, household water purification practice was significantly associated with increased z-score for WHZ but no evidence was observed for WAZ and HAZ. Prior nutritional studies from Nepal have mainly included piped water supply as an exposure variable (11, 24). Our findings are consistent with those

studies demonstrating an important association between safe drinking water and child's WHZ z-score but not for HAZ (11, 24). Households not having access to safe drinking water, either measured through household water purification practice or piped water supply, might have increased risk of water borne and infectious disease adversely affecting the child's immediate nutritional wellbeing depicting in the form of wasting. However, water quality was not associated with z-score for HAZ and WAZ which indicates its effect being either neutralized for more severe forms of undernutrition or children experiencing catch-up growth with no major impact on long term nutritional health in a Nepalese context. Nevertheless, other studies have reported a significant influence of water treatment practices on childhood stunting (35, 44). The discrepancies in study findings could be due to various reasons such as cultural factors that determine water handling practices (45) and effectiveness of household water purification techniques (46) that determines the quality of water and thus leading to a varying effect on nutritional outcomes. The availability of sufficient water could be another important factor that affects household choices for adopting healthy WASH behaviors (15) and risk of infectious diseases having adverse consequences on nutritional status of children. These factors could partly explain discrepancies in study findings across different settings and locations.

We found a significant positive linear association between household hand washing practice with soap and water and child's z-scores for WAZ and WHZ. We did not observe a significant association between hand washing behavior and child's HAZ but the association was significant in the supplementary analysis that showed decreased odds of stunting associated with hand-washing behavior. In line with our findings, other studies have also reported significant associations between hand-washing behavior and childhood stunting (23, 47). Despite this, intervention trials have failed to report significant effects of handwashing behavior on childhood stunting (36, 40). Possible explanations include poor compliance and challenges associated with sustaining handwashing behavior (36) that could omit nutritional benefits to children. Similarly, intervention studies are often resource intensive and so planned for a relatively shorter duration that may fail to capture longterm nutritional benefits (40).

No evidence of interaction was observed despite consistent associations between increased z-scores for all nutritional outcomes and the combined effect of WASH practices. In contrast, an Indian study showed a stronger inverse association between sanitation and stunting among households with handwashing practice compared to those without such practice (23). Indian study was based on a very large sample (n=34639) and failure to observe any evidence of interaction in our study could be due to insufficient power. However, we found a significant interaction between water purification practice and area of residence on HAZ score, indicating gain in HAZ score associated with water purification was higher in the rural areas compared to the urban settings (Table 5). In rural locations, households mainly depend on unreliable water sources with varying water flow/supply across different times of the year and increased risk of microbial contamination (48). The water treatment practice in such settings could have offered better benefit for child growth. We observed no interaction between WASH related variables and sex of the child for different nutritional outcomes.

We primarily focused our study on child nutrition outcomes variables on a continuous scale (z-score of various nutrition outcomes). However, modelling dichotomous (binary) nutritional variables has also been done often in other nutritional studies (14, 42). Therefore, we did the supplementary analysis modelling dichotomous nutritional outcome variables to assess if study findings are robust irrespective of choice of the statistical model. A majority of associations retained their statistical significance but some deviations were also

observed. In the linear regression model, the associations between hand-washing practice and stunting and sanitation coverage and wasting were found to be significant [supplementary table 1]. Whereas the association between sanitation coverage and stunting and hand-washing practice and childhood wasting lost their significance in the new model (Table: 3). As dichotomization may result in loss of information (49), so we emphasize that the model based on continuous outcome nutritional variables should be considered as a more effective approach for providing a true measure of association.

Strengths and limitation

Using the nationally representative samples from Nepal, our study demonstrated important roles of various WASH components on different forms of childhood undernutrition. The study findings could be generalized to Nepal and possibly to other countries with similar settings. However, our study has some limitations; the cross-sectional nature of the data is limited to establish causality of the association. The final analysis included fewer samples due to missing values ($n = 135$) but we carried out multiple imputation to check if missingness of variables had any effect on the interpretation of our results but we found no substantial change (results not shown). Though, we controlled for most of the confounders as identified in other nutritional studies, the possibility of residual confounding due to other unknown factors could not be fully ignored.

Complementary feeding has an important role in child nutritional health (50, 51). We did not adjust for complementary feeding practices in the final model since the information was limited only to 6-23 months old children. However, we conducted a sensitivity analysis which included the Minimum Acceptable Diet (MAD) by taking samples of 6-23 months old children ($n=744$). This variable summarizes both minimum feeding frequency and diversity apart from breast-milk (52). The analysis showed MAD itself was not significantly associated with nutritional outcomes, WAZ (β -coefficient: -0.02, 95%CI: -0.15 to 0.19), HAZ (β -coefficient: -0.01, 95%CI: -0.24 to 0.22) and WHZ (β -coefficient: -0.5, 95%CI: -0.24 to 0.14). It did not alter interpretation of the significant associations between WASH variables and childhood nutrition outcomes (result not shown), except for hand washing behaviors which were not significantly associated with WHZ (β -coefficient=0.12, 95%CI: -0.07 to 0.31). Therefore, this indicates that not having adjusted for MAD could have overestimated the association between hand washing behaviors and child's WHZ score.

Conclusions

Our study showed that WASH variables had an important role in nutritional outcomes among under-five year old children and that the findings could have important policy implications. The association of particular WASH factors varied widely across different forms of nutrition outcomes. This calls for careful consideration of evidence at the local level before selecting specific WASH interventions or programs in an effort to improve childhood nutritional outcomes.

The combination of various WASH components led to higher gains in z-scores, but we could not be sure if strategies to combine different WASH components would deliver any synergistic effect on childhood nutrition. Further studies with robust study design and sufficient power could elucidate further the potential impact of combining comprehensive WASH strategies on improving childhood nutrition status. They may also assist policy makers to make a comparative assessment of available evidence to address the pervasive burden of undernutrition in developing countries. However, public health policy should continue to target improving all

domains of WASH components that are known to deliver wider societal benefits beyond health and nutrition outcomes.

List Of Abbreviations

ANC: Antenatal Care Visit; CI: Confidence Interval; BMI: Body Mass Index; EA: Enumeration Areas; HAZ: Height-for-age Z-score; MAD: Minimum Acceptable Diet; NDHS: Nepal Demographic and Health Survey; OR: Odds Ratio; PSUs: Primary Sampling Units; WASH: Water, Sanitation and Hygiene; WAZ: Weight-for-age Z-score; WHZ: Weight-for-height Z-score; WHO: World Health Organization

Declarations

Ethics approval and consent to participant:

The study used a secondary data from Nepal Demographic and Health Survey, 2011. The ethical approval for the study was obtained from Nepal Health Research Council, Kathmandu and ICF Institutional Review Board (IRB). Informed consent was obtained from all participants after respondents were explained about purpose, aim, risks and benefits of the study.

Consent for publication:

Not applicable.

Availability of data and materials:

Dataset used in this study could be assessed from https://dhsprogram.com/data/dataset/Nepal_Standard-DHS_2016.cfm?flag=0

Competing interests:

Authors have no potential conflicts of interest to declare.

Funding:

No specific funding was received from this study.

Authors' contributions:

Som Kumar Shrestha developed concept and design of the research, performed data cleaning, data analysis and interpreted the results, wrote the manuscript and acted as a corresponding author. Don Vincendese supervised the data analysis, supported interpretation of results, supported for editing and critically evaluated the manuscript. Bircan Erbas supported for editing and critically evaluated the manuscript.

Acknowledgements:

We thank Professor Keith West from John Hopkins University for his valuable suggestions and feedback during finalization of the manuscript.

References

1. United Nations Children's Fund, WHO, International Bank for Reconstruction and Development/The World Bank. Levels and trends in child malnutrition: key findings of the 2018 Edition of the Joint Child Malnutrition Estimates. Geneva; 2018. Available from: <https://www.who.int/nutgrowthdb/jme-2019-key-findings.pdf?ua=1>.
2. UNICEF. Stop stunting: South Asia headline results 2018-2021. UNICEF; 2019.. Available from: <http://www.unicefrosa-progressreport.org/stopstunting.html>. Accessed 21 July 2019.
3. Ministry of Health, NewERA, ICF. Nepal Demographic and Health Survey 2016: Key Indicators. Kathmandu, Nepal: Ministry of Health; 2017. Available from: <https://www.healthynewbornnetwork.org/hnn-content/uploads/Nepal-DHS-2016-Key-Indicators.pdf>.
4. SUN Movement. Scaling Up Nutrition: Nepal 2015. Available from: <http://scalingupnutrition.org/sun-countries/nepal/>. Accessed 10 Apr 2018.
5. Government of Nepal. Multi-sector nutritional plan: For accelerating the reduction of maternal and child under-nutrition in Nepal. Kathmandu, Nepal: National Planning Commission; 2012.
6. Cunningham K, Singh A, Pandey Rana P, Brye L, Alayon S, Lapping K, et al. Suaahara in Nepal: An at-scale, multi-sectoral nutrition program influences knowledge and practices while enhancing equity. *Maternal & child nutrition*. 2017;13(4):e12415.
7. Future FT. Nepal: Feed The Future; 2019. Available from: <https://www.feedthefuture.gov/country/nepal/>. Accessed 2 Feb 2019.
8. The World Bank. Nepal: Community Actionfor Nutrition Project (Sunaula Hazar Din). 2019. Available from: <http://projects.worldbank.org/P125359/first-1000-days?lang=en>.
9. Ministry of Health, NewERA, ICF. Nepal Demographic and Health Survey 2006. Kathmandu, Nepal: Ministry of Health; 2006.
10. National Planning Commission. Sustainable Development Goals 2016-2030, National (Preliminary) Report. Kathmandu, Nepal: Government of Nepal; 2015.
11. Cunningham K, Headey D, Singh A, Karmacharya C, Rana PP. Maternal and Child Nutrition in Nepal: Examining drivers of progress from the mid-1990s to 2010s. *Global food security*. 2017;13:30-7.
12. Aguayo VM, Nair R, Badgaiyan N, Krishna V. Determinants of stunting and poor linear growth in children under 2 years of age in India: an in-depth analysis of Maharashtra's comprehensive nutrition survey. *Maternal & child nutrition*. 2016;12(S1):121-40.
13. Harris M, Alzua ML, Osbert N, Pickering AJ. Community-level sanitation coverage is more strongly associated with child growth and household drinking water quality than access to a private toilet in rural Mali. *Environmental Science & Technology*. 2017.
14. Merchant A, Jones C, Kiure A, Kupka R, Fitzmaurice G, Herrera M, et al. Water and sanitation associated with improved child growth. *European Journal of Clinical Nutrition*. 2003;57(12):1562.
15. Cumming O, Cairncross S. Can water, sanitation and hygiene help eliminate stunting? Current evidence and policy implications. *Maternal & child nutrition*. 2016;12:91-105.
16. Fewtrell L, Kaufmann RB, Kay D, Enanoria W, Haller L, Colford JM. Water, sanitation, and hygiene interventions to reduce diarrhoea in less developed countries: a systematic review and meta-analysis. *The*

- Lancet infectious diseases. 2005;5(1):42-52.
17. Cairncross S, Hunt C, Boisson S, Bostoen K, Curtis V, Fung IC, et al. Water, sanitation and hygiene for the prevention of diarrhoea. *International journal of epidemiology*. 2010;39(suppl_1):i193-i205.
 18. Ejemot-Nwadiaro RI, Ehiri JE, Arikpo D, Meremikwu MM, Critchley JA. Hand washing promotion for preventing diarrhoea. *The Cochrane Library*. 2015.
 19. Checkley W, Epstein LD, Gilman RH, Cabrera L, Black RE. Effects of acute diarrhea on linear growth in Peruvian children. *American Journal of Epidemiology*. 2003;157(2):166-75.
 20. Checkley W, Buckley G, Gilman RH, Assis AM, Guerrant RL, Morris SS, et al. Multi-country analysis of the effects of diarrhoea on childhood stunting. *International journal of epidemiology*. 2008;37(4):816-30.
 21. Humphrey JH. Child undernutrition, tropical enteropathy, toilets, and handwashing. *The Lancet*. 2009;374(9694):1032-5.
 22. Dangour AD, Watson L, Cumming O, Boisson S, Che Y, Velleman Y, et al. Interventions to improve water quality and supply, sanitation and hygiene practices, and their effects on the nutritional status of children. *Cochrane Database of Systematic Reviews*. 2013(8).
 23. Rah JH, Cronin AA, Badgaiyan B, Aguayo VM, Coates S, Ahmed S. Household sanitation and personal hygiene practices are associated with child stunting in rural India: a cross-sectional analysis of surveys. *BMJ open*. 2015;5(2):e005180.
 24. Dorsey JL, Manohar S, Neupane S, Shrestha B, Klemm RD, West Jr KP. Individual, household, and community level risk factors of stunting in children younger than 5 years: Findings from a national surveillance system in Nepal. *Maternal & child nutrition*. 2018;14(1):e12434.
 25. Headey DD, Hoddinott J. Understanding the rapid reduction of undernutrition in Nepal, 2001–2011. *PloS one*. 2015;10(12):e0145738.
 26. WHO, UNICEF, USAID. Improving nutrition outcomes with better water, sanitation and hygiene: practical solutions for policies and programmes. 2015. Available from: https://apps.who.int/iris/bitstream/handle/10665/193991/9789241565103_eng.pdf?sequence=1.
 27. Assaf S, Kothari MT, Pullum T. An assessment of the quality of DHS anthropometric data, 2005-2014 . *DHS Methodol Reports No 16* [Internet]. 2015;(September):2005–14. Available from: <http://dhsprogram.com/pubs/pdf/MR16/MR16.pdf>
 28. World Health Organization; WHO Child Growth Standards. Vol. 51, *Developmental Medicine & Child Neurology*. 2009.
 29. Rutstein SO. Factors associated with trends in infant and child mortality in developing countries during the 1990s. *Bulletin of the World Health Organization*. 2000;78(10):1256-70.
 30. Onis M De, Bloessner M. WHO global database on child growth and malnutrition. Geneva: WHO; 1997.
 31. World Health Organization. Water Sanitation Hygiene: Key terms: WHO; 2019. Available from: https://www.who.int/water_sanitation_health/monitoring/jmp2012/key_terms/en/. Accessed 02 Feb 2019.
 32. Dearden KA, Schott W, Crookston BT, Humphries DL, Penny ME, Behrman JR. Children with access to improved sanitation but not improved water are at lower risk of stunting compared to children without access: a cohort study in Ethiopia, India, Peru, and Vietnam. *BMC public health*. 2017;17(1):110.

33. Liu E, Balasubramaniam D, Hunt A. Does access to water matter? A study on the nutritional status of primary-aged children in India. *Journal of Public Health*. 2016;38(4):e419-e24.
34. Hartinger S, Lanata C, Hattendorf J, Verastegui H, Gil A, Wolf J, et al. Improving household air, drinking water and hygiene in rural Peru: a community-randomized–controlled trial of an integrated environmental home-based intervention package to improve child health. *International journal of epidemiology*. 2016;45(6):2089-99.
35. Ambadekar N, Zodpey S. Risk factors for severe acute malnutrition in under-five children: a case-control study in a rural part of India. *Public health*. 2017;142:136-43.
36. Arnold B, Arana B, Mäusezahl D, Hubbard A, Colford Jr JM. Evaluation of a pre-existing, 3-year household water treatment and handwashing intervention in rural Guatemala. *International Journal of Epidemiology*. 2009;38(6):1651-61.
37. Prüss-Ustün A, Bartram J, Clasen T, Colford JM, Cumming O, Curtis V, et al. Burden of disease from inadequate water, sanitation and hygiene in low and middle income settings: a retrospective analysis of data from 145 countries. *Tropical Medicine & International Health*. 2014;19(8):894-905.
38. Rautanen S-L, van Koppen B, Wagle N. Community-driven multiple use water services: Lessons learned by the Rural Village Water Resources Management Project in Nepal. *Water Alternatives*. 2014;7(1).
39. Ram P. Practical guidance for measuring handwashing behavior: 2013 Update. Water and Sanitation Program. 2013. Available from: <https://www.wsp.org/sites/wsp/files/publications/WSP-Practical-Guidance-Measuring-Handwashing-Behavior-2013-Update.pdf>. Accessed 15 Sep 2019.
40. Langford R, Lunn P, Brick CP. Hand-washing, subclinical infections, and growth: A longitudinal evaluation of an intervention in Nepali slums. *American Journal of Human Biology*. 2011;23(5):621-9.
41. Shrestha SK, Katelaris C, Dharmage SC, Burton P, Vicendese D, Tham R, et al. High ambient levels of grass, weed and other pollen are associated with asthma admissions in children and adolescents: A large 5-year case-crossover study. *Clin Exp Allergy*. 2018;48(11):1421–8.
42. Fuller JA, Villamor E, Cevallos W, Trostle J, Eisenberg JN. I get height with a little help from my friends: herd protection from sanitation on child growth in rural Ecuador. *International journal of epidemiology*. 2016;45(2):460-9.
43. Raihan MJ, Farzana FD, Sultana S, Haque MA, Rahman AS, Waid JL, et al. Examining the relationship between socio-economic status, WASH practices and wasting. *PloS one*. 2017;12(3):e0172134.
44. Sinharoy SS, Schmidt WP, Cox K, Clemence Z, Mfura L, Wendt R, et al. Child diarrhoea and nutritional status in rural Rwanda: a cross-sectional study to explore contributing environmental and demographic factors. *Tropical medicine & international health*. 2016;21(8):956-64.
45. Moser G, Navarro O, Ratiu E, Weiss K. Cultural background and environmental context of water perception and use. In: Corral-Verdugo V, Garcia-Cadena CH, Frías-Armenta M. *Psychological approaches to sustainability: Current trends in theory, research and applications*. New York: Nova Science Publishers; 2010.
46. Hunter PR. Household water treatment in developing countries: comparing different intervention types using meta-regression. *Environmental science & technology*. 2009;43(23):8991-7.
47. Saxton J, Rath S, Nair N, Gope R, Mahapatra R, Tripathy P, et al. Handwashing, sanitation and family planning practices are the strongest underlying determinants of child stunting in rural indigenous

- communities of Jharkhand and Odisha, Eastern India: a cross-sectional study. *Maternal & child nutrition*. 2016;12(4):869-84.
48. Merz J, Nakarmi G, Shrestha S, Dahal BM, Dongol BS, Schaffner M, et al. Public water sources in rural watersheds of Nepal's Middle Mountains: Issues and constraints. *Environmental Management*. 2004;34(1):26-37.
49. Altman DG, Royston P. The cost of dichotomising continuous variables. *Bmj*. 2006;332(7549):1080.
50. Onyango AW, Borghi E, de Onis M, del Carmen Casanovas M, Garza C. Complementary feeding and attained linear growth among 6–23-month-old children. *Public Health Nutr*. 2014;17(9):1975–83. *Public Health Nutr*. 2014;17(9):1975–83.
51. Zongrone A, Winskell K, Menon P. Infant and young child feeding practices and child undernutrition in Bangladesh: insights from nationally representative data. *Public Health Nutr*. 2012;15(9):1697–704.
52. World Health Organization. Indicators for assessing infant and young child feeding practices: part 2: measurement. 2010. https://apps.who.int/iris/bitstream/handle/10665/44368/9789241599757_eng.pdf. Accessed 15 Sep 2019.

Tables

Table 1: Household water sanitation and handwashing practices and mean z-score for underweight, stunting and wasting among under-five children (n=2352).

Variables	Number (%)
Proportion of household with improved toilet, water purification practices and availability of soap and water	
Household with improved sanitation	1781 (75.7%)
Cluster sanitation coverage percentage [SD]	75.3 [32]
Household with practice of water purification	430 (18.3%)
Household with Soap and water available	882 (37.5%)
Nutritional Outcome Indicators	Mean [SD]
z-score for height-for-age	-1.52 [1.3]
z-score for weight-for-height	-0.65 [1.1]
z-score for weight-for-age	-1.33 [1.1]

Table 2: Under-five children z-score classification for underweight, stunting and wasting (n=2352)

Outcome indicators	Undernourished Children			Adequately nourished Children		
	N (%)			N (%)		
	z-score	z-score between -3	Total	z-score between	z-score greater	Total
	< -3	to <- 2	(z- score <-2)	-2 to +2	than +2	(z-score >-2)
Weight-for-age	284	559 (23.8)	843	1489 (63.3)	20 (0.85)	1509
Stunting)	(12.1)		(35.8)			(64.2)
Weight-for-height	44 (1.9)	187 (8)	231	2092 (88.9)	29 (1.2)	2121
Wasting)			(9.8)			(90.2)
Weight-for-age	124	511 (21.7)	635	1710 (72.7)	7 (0.3)	1717
Underweight)	(5.3)		(27.0)			(73.0)

Table 3: Linear regression coefficient for association between WASH related variables and nutritional status indicators Z-scores of preschool, Nepalese children (n=2320)

Outcomes	Height-for-age (β ^, 95%CI)		Weight-for-height (β ^, 95%CI)		Weight-for-age (β ^, 95%CI)	
	Unadjusted	Adjusted	Unadjusted	Adjusted	Unadjusted	Adjusted
Infant mortality	0.54 (0.35, 0.74)***	0.28 (0.001, 0.56) ^{a,*}	0.66 (0.50, 0.82)***	0.19 (-0.04, 0.42) ^a	0.78 (0.62, 0.94)***	0.30 (0.12, 0.48) ^{a, **}
Under-five mortality	0.37 (0.16, 0.59)**	-0.02 (-0.24, 0.20) ^b	0.52 (0.38, 0.67)***	0.24 (0.07, 0.41) ^{b, **}	0.56 (0.38, 0.73)***	0.11 (-0.06, 0.28) ^b
Infant mortality (ref)	1	1	1	1	1	1
Under-five mortality (ref)	1	1	1	1	1	1
Soap availability	0.46 (0.32, 0.59)***	0.09 (-0.05, 0.22) ^c	0.29 (0.18, 0.40)***	0.13 (0.01, 0.24) ^{c, *}	0.47 (0.35, 0.58)***	0.15 (0.04, 0.25) ^{c, **}
Soap availability (ref)	1	1	1	1	1	1

^aAdjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery frequency of watching TV, childhood diarrhea over past 2 weeks, household water purification practice and household water and soap availability.

^bAdjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery frequency of watching TV, childhood diarrhea over past 2 week, cluster sanitation coverage and household water and soap availability.

^cAdjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery frequency of watching TV, childhood diarrhea over past 2 weeks, cluster sanitation coverage and household water purification practice.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Table 4: Interaction between WASH related variables for childhood undernutrition (n=2320)

Height-for-age			Weight-for-height			Weight-for-age		
With water purification	Without water purification	<i>p</i> -value for interaction	With water purification	Without water purification	<i>p</i> -value for interaction	With water purification	Without water purification	<i>p</i> -value for interaction
0.44 (-0.55, 1.43)	0.21 (-0.06, 0.48)	0.28	0.31 (-0.53, 1.15)	0.20 (-0.04, 0.44)	0.55	0.67 (-0.11, 1.45)	0.29 (0.11, 0.47)**	0.25
With soap and water	Without soap and water		With soap and water	Without soap and water		With soap and water	Without soap and water	
0.07 (-0.21, 0.34)	-0.10 (-0.46, 0.26)	0.87	0.28 (0.05, 0.51)*	0.15 (-0.09, 0.38)	0.28	0.22 (0.02, 0.44)*	-0.05 (-0.27, 0.18)	0.16
With improved sanitation	Without improved sanitation		With improved sanitation	Without improved sanitation		With improved sanitation	Without improved sanitation	
0.69 (-0.07, 0.21)	0.07 (-0.36, 0.49)	0.39	0.14 (0.01, 0.27)*	0.04 (-0.24, 0.31)	0.36	0.15 (0.04, 0.26)**	0.08 (-0.25, 0.40)	0.97

^aAdjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery, frequency of watching TV, childhood diarrhea over past 2 weeks and household water and soap availability.

^b Adjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery, frequency of watching TV, childhood diarrhea over past 2 weeks and household sanitation availability.

^c Adjusted for child’s birth weight, child’s age, child’s sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women’s age, women marital status, women’s education, women’s smoking status, women’s BMI, ANC 4th visits, institutional delivery, frequency of watching TV, childhood diarrhea over past 2 weeks and household water purification practice.

*p<0.05; **p<0.01; ***p<0.001

Table 5: Interaction between household water purification practices and area of residence on childhood stunting (n=2320)

Water purification practice	Height-for-age z-score		
	Urban	Rural	p-value for interaction
	-0.14 (-0.41, 0.13) ^a	0.19 (-0.1, 0.5) ^a	0.02*

^aAdjusted for child’s birth weight, child’s age, child’s sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area o residence, women’s age, women marital status, women’s education, women’s smoking status, women’s BMI, ANC 4th visits, institutional delivery frequency of watching TV, childhood diarrhea over past 2 week, cluster sanitation coverage and household water and soap availability.

*p<0.05

Supplementary Table

Supplementary table: 1 Odds ratio of underweight, stunting and wasting associated with WASH variables (n=2320)

	Stunting (OR, 95%CI)		Wasting (OR, 95%CI)		Underweight (OR, 95%CI)	
	Unadjusted	Adjusted	Unadjusted	Adjusted	Unadjusted	Adjusted
ion	0.47 (0.34, 0.74) 0.64)***	1.16) ^a	(0.47, 0.38 0.57)***	(0.26, 0.53 0.99) ^{a, *}	(0.28, 0.31 0.44)***	(0.21, 0.59 0.93) ^{a, *}
rence)	1	1	1	1	1	1
p	0.55 (0.39, 0.98 0.78)***	1.48) ^b	(0.65, 0.29 0.50)***	(0.16, 0.39 0.73) ^{b, **}	(0.21, 0.42 0.63)***	(0.28, 1.07 1.75) ^b
ence)	1	1	1	1	1	1
	0.47 (0.38, 0.70 0.58)***	0.92) ^{c, **}	(0.54, 0.66 0.91)***	(0.48, 0.82 1.20) ^c	(0.56, 0.40 0.52)***	(0.32, 0.65 0.84) ^{c, **}
ence)	1	1	1	1	1	1

^aAdjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery frequency of watching TV, childhood diarrhea over past 2 weeks, household water purification practice and household water and soap availability.

^bAdjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery frequency of watching TV, childhood diarrhea over past 2 week, cluster sanitation coverage and household water and soap availability.

^cAdjusted for child's birth weight, child's age, child's sex wealth quintiles, use of clean fuel, sex of household head, ecological region, area of residence, women's age, women marital status, women's education, women's smoking status, women's BMI, ANC 4th visits, institutional delivery frequency of watching TV, childhood diarrhea over past 2 weeks, cluster sanitation coverage and household water purification practice.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Supplementary table 2: Descriptive summary of variables adjusted in the model (n=2352)

Explanatory variables	Number	(%)
Ecological region		
Mountain	165	7.0
Hill	861	36.6
Terai	1326	56.4
Types of place of residence		
Urban	1240	52.7
Rural	1112	47.3
Household Wealth quintiles		
Poorest	487	20.7
Poorer	513	21.8
Middle	532	22.6
Richer	510	21.7
Richest	310	13.2
Sex of household head		
Male	1602	68.1
Female	750	31.9
Clean fuel		
Yes	1804	76.7
No	548	23.3
Child's age (months)		
0-11	463	19.7
12-23	508	21.6
24- 35	447	19.0
36-47	473	20.1
48-59	461	19.6
Child's sex		
Male	1230	52.3
Female	1122	47.7
Number of living child		
1	720	30.6
2	847	36.0
3 or more	785	33.4
Birth weight of child		
>= 2500 gram	1265	53.8
<2500 gram	179	7.6
Not weighted at birth	908	38.6
Ever breastfeed by women		
Yes	2303	97.9
No	49	2.1

Women age group		
15-19	195	8.3
20-24	804	34.2
25-29	767	32.6
30-34	369	15.7
35-39	146	6.2
40 or more	71	3.0
Marital status of women		
Currently married	2338	99.4
Formerly married	14	0.6
Women's education level		
No Education	816	34.7
Primary Education	470	20.0
Some Secondary	565	24.0
SLC and Above	501	21.3
Women's smoking Status		
No	2300	97.8
Yes	52	2.2
Women's Nutrition Status		
Normal	1900	80.8
Undernutrition	452	19.2
Completed 4th ANC visits		
No	729	31.0
Yes	1623	69.0
Institutional delivery		
No	1077	45.8
Yes	1275	54.2
Watching TV over last week		
Not at all	901	38.3
Less than once a week	492	20.9
Once a week or more	960	40.8

Figures

Figure 1

Flow-chart for under-five children sampling procedure for NDHS, 2016.