The Appropriate Acquisition Time Interval Following Injection of ^{99m}Tc -Sestamibi with Water Protocol in Single Photon Emission Computed Tomography Myocardial Perfusion Imaging: First Experience in Indonesia

Nora Anggun Prasetyo (✉ noraprasetyo2004@gmail.com)

Universitas Padjadjaran Bandung

Erwin Affandi Soeriadi

Dr. Hasan Sadikin General Hospital

Hendra Budiawan

Dr. Hasan Sadikin General Hospital

Achmad Hussein Sundawa Kartamihardja

Universitas Padjadjaran Bandung

Research Article

Keywords: Acquisition time, SPECT, stress Myocardial Perfusion Imaging procedure, Water protocol

Posted Date: August 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-774654/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

According to EANM/ESC guideline, image acquisitions in stress test should be begun at 30-60 minutes after tracer administration. Our center is a referral hospital for nuclear medicine imaging with many patients but limited number of gamma camera. The shorter time between injection of radiopharmaceutical and imaging acquisition will lead to more examination can be done.

The aim of this study was to evaluate the appropriate acquisition time interval with water protocol in ^{99m}Tc -Sestamibi SPECT myocardial perfusion imaging.

Methods

Patients who were referred to undergo stress MPI between October 2020 to December 2020 were included in this study. Cardiac stress procedure was performed using treadmill with modified Bruce Protocol. Subjects should drink a total of 330 mL water following ^{99m}Tc -Sestamibi injection. Image acquisitions were performed 10 and 30 minutes afterwards. Quantitative assessment was done by calculated target background ratio (TBR). Statistical analysis was performed using student t-test with Microsoft Excel version 2019. P -value < 0.05 was considered to be statistically significance.

Results

Thirty out of 35 subjects were included in this study. Sixteen of them are male and 14 are female with mean of age was 48.7 years old (28 – 80). Mean target background uptake ratio (TBR) in 10- and 30-minutes images were 0.67 (0.44 – 1.11) and 0.76 (0.43 – 1.18) respectively (p -value = 0.15).

Conclusion

There was no significant difference of target to background ratio between acquisition time 10- and 30-minutes following injection of ^{99m}Tc -Sestamibi with water protocol in myocardial perfusion imaging.

Trial registration

Not applicable.

Background

Coronary artery disease (CAD) remains the world's leading cause of death.¹ Single Photon Emission Computed Tomography (SPECT) Myocardial Perfusion Imaging (MPI) using ^{99m}Tc -Sestamibi can be used to diagnose and evaluate the prognostic value of CAD. According to European Association of Nuclear Medicine/European Society of Cardiology (EANM/ESC) guideline, image acquisitions should be begun at 30–60 minutes after tracer administration in stress test.²

Our center is a referral center hospital for nuclear medicine imaging with many patients but a limited number of gamma camera. The aim of this study was to evaluate the appropriate acquisition time interval between injection of radiopharmaceutical and image acquisition using water protocol in SPECT myocardial perfusion imaging.

Methods

Thirty-five patients who referred to department of Nuclear Medicine and Molecular Imaging Dr. Hasan Sadikin General Hospital for MPI procedure were included in this study, between October 2020 to December 2020. The exclusion criteria were as follow: patient have no history of essential fluid restriction, consume anti-motility or pro-motility drug, no pregnant, or lactating at the time of MI procedure, have contraindications to perform exercise stress test using treadmill. Patients also should not consume caffeine containing products for at least 12 hours prior MPI procedure, stop β -blocker treatment for 24 hours, and stop phosphodiesterase inhibitor for 24–48 hour before MPI procedure. All subjects were asked to fast for at least 4 hours prior to MPI procedure. Informed consent was obtained from all the subjects.

Stress procedure was performed using treadmill with modified Bruce Protocol. ^{99m}Tc -Sestamibi was use as radiopharmaceutical. Subjects who reached minimal 85% of maximal heart rate were asked to drink a total of 330 mL water after radiopharmaceutical injection. Image acquisitions were performed at 10 and 30 minutes after radiopharmaceutical injection.

Region of interest (ROI) were drawn on heart and liver on 10- and 30-minutes images and then the target background ratio (TBR) was calculated, as shown in Fig. 1. Statistical analysis was performed using student t-test with Microsoft Excel version 2019 to evaluate the quantitative assessment of heart to liver background ratio. P -value < 0.05 was considered as statistically significance.

Results

A total of 30 out of 35 subjects (86%) were included in this study. Five of 35 subjects were excluded because they have essential fluid restriction, have contraindications to perform exercise stress test using treadmill, and didn't reach optimal target maximal heart rate. Subjects consist of 16 males (53%) and 14 females (47%) with mean age of 48.7 (28–80 years-old). Mean target background ratio (TBR) in 10- and 30-minutes acquisition time interval following injection of radiopharmaceutical were 0.67 (0.44–1.11) and 0.76 (0.43–1.18) respectively (p -value = 0.15), as shown on Fig. 2.

Discussion

Coronary artery disease (CAD) remains the world's leading cause of death with 16% of the world's total death. According to World Health Organization, there were 2 to 8.9 million deaths in 2019.¹ Prevention and early non-invasive diagnosis or early detection with high sensitivity and high specificity must be the

priority to slow down the increasing death caused by CAD. SPECT-MPI using radiopharmaceutical ^{99m}Tc -Sestamibi can be used to diagnose and evaluate the prognostic value of CAD. It shows more cost effective than any other diagnostic modality, and more sensitive than exercise Electrocardiogram (ECG) in detecting myocardial ischemia. It has sensitivity 85–95%, and specificity 75%. The specificity can up to 94% if combine with ECG gated data.^{3–5} Stress images may help determine the degree of inducible ischemia or viable myocardium that are amenable to revascularization, evaluate myocardial viability before revascularization and following-up after revascularization such as coronary artery bypass graft or angioplasty.^{3–6}

Physiological radioactivity uptake in the liver, bowel or stomach could interfere the interpretation of MPI. This phenomenon can be reduced by using iterative reconstruction and attenuation correction but a high tracer activity around the heart is still influence the evaluation of the myocardial.^{7,8} Several methods to reduce the digestive activity have been described, such as filling the stomach with either solids or fluids^{9–11}, increasing the gastrointestinal activity by using metoclopramide drug¹², keeping the patient standing for several minutes after tracer injection¹³, or even ingesting iodinated contrast medium to absorb the emitted gamma-rays¹⁴. Prolonging the delay imaging between injection and image acquisition can lead to a significant decrease in the hepatic activity but also by increasing the activity in the bowel loops adjacent to the inferior wall of the heart.^{11, 12, 14}

Patients should take fatty meal at time between injection and imaging to increase clearance of tracer from the liver and gallbladder as a standard protocol of MPI in some centers.² Our department usually ask patients to drink milk to reduce physiological infra-cardiac organs uptake. However, some of patients are lactose intolerant, and caused diarrhea and abdominal discomfort. Data on Indonesian population showed that the prevalence of lactose malabsorption and intolerance are quite high, between 21–73%.^{15–17} Those patients will feel abdominal discomfort as well as diarrhea after MPI procedure, and sometimes disturb the MPI procedure.¹³ In this study we use water instead of milk, to minimize the infra-cardiac uptake that can interfere interpretation of SPECT MPI. Our previous study compares subjects in first group (drink 500 mL of milk) and second group (drink 330 mL of water) after doing physical stress MPI procedure with acquisition time 30 minutes after ^{99m}Tc -Sestamibi injection, showed that water was as good as milk in reducing the infra-cardiac tissue tracer uptake. In this study, ^{99m}Tc -Sestamibi was used as radiopharmaceutical for MPI. Since this radiopharmaceutical is a lipophilic agent, patients were advice to take solid fatty meal for breakfast on the day of MPI procedure.^{7,8}

Preparation before MPI procedure is important to get a good quality of images. Subjects should not take any food for at least 4 hours to prevent stress-induced gastric distress and minimize splanchnic blood distribution.¹⁸ Subjects must have no contraindications to perform exercise stress test using treadmill. Caffeine containing products must not be consumed for at least 12 hours prior MPI procedure, to avoid decrease vasodilatation respond from vasodilator stress agents, which will lead to reduce sensitivity in

detecting myocardial ischemia. Subjects also suggested to stop β -blocker treatment for 24 hour and phosphodiesterase inhibitor for 24–48 hour before MPI procedure.¹⁹

Out of 35 subjects, 5 subjects were excluded because they didn't reach minimal target heart rate (85% maximal heart rate) and had contraindications to undergo the exercise test. Based on EANM/ESC guideline, patients are said to achieve maximum exercise stress test when they are able to reach $\geq 85\%$ of their age-predicted maximum heart rate (maximal age-predicted heart rate = $220 - \text{age}$).²

In accordance to conventional guideline, image acquisitions should be performed 15–60 minutes after tracer administration.¹⁹ Strauss, et al. said that image acquisitions with ^{99m}Tc based tracer should be performed after liver radioactivity has sufficiently cleared, this was usually between 15–30 minutes after tracer administration.²⁰ While according to EANM/ESC guideline, image acquisitions should be begun at 30–60 minutes after tracer administration.² In this study, MPI was done on 10 and 30 minutes following ^{99m}Tc -Sestamibi injection intravenously. Ten minutes acquisition time following injection of radiopharmaceutical based on the consideration that the time taken by patients between physical stress MPI, and imaging procedure was about 10 minutes. The 10 minutes image after radiopharmaceutical injection was considered as direct imaging, and 30 minutes image as control, based on EANM/ESC guideline, as an appropriate time for image acquisitions.²

All subjects were scanned in supine position with arms raised above the head and support the knees with pillow under the knees, shoulders and arms comfortably positioned in order to reduce movement and pain, particularly in older subjects, to make feel as comfortable as possible to minimize body motion as suggested by Dorbala, et al..¹⁹ Female subjects were imaged without bra, and used chest band to minimize breast attenuation and minimize motion. Chest band could also be used in male subjects to minimize body motion as suggested by Hesse B. et al..²

From the quantitative assessment results, we found that there was no significant difference TBR of heart to liver ratio between acquisition 10 and 30 minutes after radiopharmaceutical injection as seen in Fig. 2, with p-value = 0.15 ($p > 0.05$).

Conclusion

Our quantitative study showed that there was no significant difference in target to background ratio between acquisition time 10- and 30-minutes following injection of ^{99m}Tc -Sestamibi in water protocol SPECT myocardial perfusion imaging procedure. The appropriate acquisition time in stress exercise test using water protocol SPECT myocardial perfusion imaging is 10 minutes following injection of ^{99m}Tc -Sestamibi.

Abbreviations

1. CAD: Coronary Artery Disease

2. EANM: European Association of Nuclear Medicine
3. ECG: Electrocardiogram
4. ESC: European Society of Cardiology
5. ROI: Region of Interest
6. SPECT: Single Photon Emission Computed Tomography
7. TBR: Target Background Ratio

Declarations

Ethics approval and consent to participate

All procedures performed in this study involving human participants were in accordance with the ethical standards of Dr. Hasan Sadikin General Hospital Bandung and had been approved by the Health Research Ethics Committee of Dr. Hasan Sadikin General Hospital Bandung (West Java – Indonesia) (Ethical Approval Number: LB.02.01/X.6.5/311/2020). Written informed consent was obtained from all the patients those were enrolled in the study.

Consent for publication

Consent for publication was included in written informed consent that was obtained from all the patients those were enrolled in the study.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

There was no source of funding for this study.

Authors' contributions

NAP made the concept of this study and design this study. She also did the literature search, performed the study, did the data acquisition, analyzed the datasets, and analyzed the statistic too. She prepared the manuscript, edited it and the reviewed it.

EAS, HB, and AHSK helped in generating the concept of this study and designing this study. They also helped in editing and reviewing the manuscript.

AHSK acted as the guarantor of this study.

Acknowledgements

Not applicable.

References

1. Organization WH. The top 10 causes of death 2021 [Available from: <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>].
2. Hesse B, Tagil K, Cuocolo A, Anagnostopoulos C, Bardies M, Bax J, et al. EANM/ESC procedural guidelines for myocardial perfusion imaging in nuclear cardiology. *Eur J Nucl Med Mol Imaging*. 2005;32(7):855-97.
3. Loong CY, Anagnostopoulos C. Diagnosis of coronary artery disease by radionuclide myocardial perfusion imaging. *Heart*. 2004;90 Suppl 5:v2-9.
4. Zellweger MJ, Lewin HC, Lai S, Dubois EA, Friedman JD, Germano G, et al. When to stress patients after coronary artery bypass surgery? Risk stratification in patients early and late post-CABG using stress myocardial perfusion SPECT: implications of appropriate clinical strategies. *J Am Coll Cardiol*. 2001;37(1):144-52.
5. Zaret BL, Strauss HW, Martin ND, Wells HP, Jr., Flamm MD, Jr. Noninvasive regional myocardial perfusion with radioactive potassium. Study of patients at rest, with exercise and during angina pectoris. *N Engl J Med*. 1973;288(16):809-12.
6. Hachamovitch R, Hayes SW, Friedman JD, Cohen I, Berman DS. Comparison of the short-term survival benefit associated with revascularization compared with medical therapy in patients with no prior coronary artery disease undergoing stress myocardial perfusion single photon emission computed tomography. *Circulation*. 2003;107(23):2900-7.
7. Hambye AS, Delsarte P, Vervaet AM. Influence of the different biokinetics of sestamibi and tetrofosmin on the interpretation of myocardial perfusion imaging in daily practice. *Nucl Med Commun*. 2007;28(5):383-90.
8. Germano G, Chua T, Kiat H, Areeda JS, Berman DS. A quantitative phantom analysis of artifacts due to hepatic activity in technetium-99m myocardial perfusion SPECT studies. *J Nucl Med*. 1994;35(2):356-9.
9. van Dongen AJ, van Rijk PP. Minimizing liver, bowel, and gastric activity in myocardial perfusion SPECT. *J Nucl Med*. 2000;41(8):1315-7.
10. Boz A, Gungor F, Karayalcin B, Yildiz A. The effects of solid food in prevention of intestinal activity in Tc-99m tetrofosmin myocardial perfusion scintigraphy. *J Nucl Cardiol*. 2003;10(2):161-7.
11. Peace RA, Lloyd JJ. The effect of imaging time, radiopharmaceutical, full fat milk and water on interfering extra-cardiac activity in myocardial perfusion single photon emission computed tomography. *Nucl Med Commun*. 2005;26(1):17-24.

12. Weinmann P, Moretti JL. Metoclopramide has no effect on abdominal activity of sestamibi in myocardial SPET. Nucl Med Commun. 1999;20(7):623-5.
13. Hurwitz GA, Clark EM, Slomka PJ, Siddiq SK. Investigation of measures to reduce interfering abdominal activity on rest myocardial images with Tc-99m sestamibi. Clin Nucl Med. 1993;18(9):735-41.
14. Iqbal SM, Khalil ME, Lone BA, Gorski R, Blum S, Heller EN. Simple techniques to reduce bowel activity in cardiac SPECT imaging. Nucl Med Commun. 2004;25(4):355-9.
15. Lukito W, Malik SG, Surono IS, Wahlqvist ML. From 'lactose intolerance' to 'lactose nutrition'. Asia Pac J Clin Nutr. 2015;24 Suppl 1:S1-8.
16. Swagerty DL, Jr., Walling AD, Klein RM. Lactose intolerance. Am Fam Physician. 2002;65(9):1845-50.
17. Hegar B, Widodo A. Lactose intolerance in Indonesian children. Asia Pac J Clin Nutr. 2015;24 Suppl 1:S31-40.
18. O'Malley HZJ, Thrall J. Nuclear medicine and molecular imaging : the requisites. 5th edition. ed. St. Louis, MO: Elsevier Inc.; 2019. pages cm p.
19. Dorbala S, Di Carli MF, Delbeke D, Abbara S, DePuey EG, Dilsizian V, et al. SNMMI/ASNC/SCCT guideline for cardiac SPECT/CT and PET/CT 1.0. J Nucl Med. 2013;54(8):1485-507.
20. Strauss HW, Miller DD, Wittry MD, Cerqueira MD, Garcia EV, Iskandrian AS, et al. Procedure guideline for myocardial perfusion imaging 3.3. J Nucl Med Technol. 2008;36(3):155-61.

Figures

Figure 1

Comparison of 99mTc-MIBI uptake in the target (heart) and the background (liver) on 58 years old male with atypical chest pain

Figure 2

Mean target to background uptake ratio in 10- and 30-minutes post injection of radiopharmaceutical