Optical solitons in Bragg gratings fibers for the nonlinear (2+1)-dimensional Kundu-Mukherjee-Naskar equation using two integration schemes

Elsayed M. E. Zayed, Reham.M.A.Shohib & Mohamed E. M. Alngar
Mathematics Department, Faculty of Science, Zagazig University,
Zagazig-44519, Egypt

Abstract

The current work handles for the first time, dispersive optical solitons in fiber Bragg gratings for the nonlinear (2+1)-dimensional Kundu-Mukherjee-Naskar equation. Two integration schemes, namely, the modified Kudryashov's approach and the addendum to Kudryashov's methodology are applied. Dark, bright and singular soliton solutions are obtained. Also, combo bright-singular soliton solutions are introduced.

Keywords: Two integration technologies; Optical soliton solutions; Fiber Bragg gratings; Kundu-Mukherjee-Naskar equation.

PACS Numbers: 02.30.Jr; 02.30.Ik; 05.45.Yv

OCIS Codes: 060.2310; 060.4510; 060.5530; 190.3270; 190.4370

1 Introduction

Fiber Bragg gratings (FBGs) are of tremendous interests to physicists and engineers during the past couple of decades or more because of its applications. FBGs are applied in all optical communication systems and thus have been widely studied with several methods[1–19]. Optical nonlinearities, such as Kerr law, quadratic–cubic law, log-law, cubic–quintic–septic law, triple-power law, power-law, parabolic-law, anti-cubic law, dual-power law and parabolic-nonlocal law have been investigated by FBGs using integration technologies. Also, optical solitons in FBGs have been gained with governing models that describe the dynamics of soliton propagation, such as the Biswas–Milovic equation, Lakshmanan–Porsezian–Daniel model, Fokas–Lenells equation, Kundu–Mukherjee–Naskar model, Biswas–Arshed equation, complex Ginzburg–Landau equation, Kudryashov's model, Chen–Lee–Liu equation, Sasa–Satsuma equation, Kaup–Newell equation, Gerdjikov–Ivanov equation, Kundu–Eckhaus equation, Triki–Biswas model and Manakov model. The current paper handles solitons in FBGs with the nonlinear (2+1)-dimensional

Kundu-Mukherjee-Naskar equation that has solved by aid of two integration schemes. The governing model in FBGs is formulated in this the paper for the first time. Optical solitons of the model equation are revealed by the modified Kudryashov's approach [17-19] and the addendum to Kudryashov's methodology [13, 17-19]. These solitons are reported in this paper for the first time. The details are sketched through, after a thorough introduction to the governing model

1.1 Governing model

The dimensionless form of the nonlinear (2+1)-dimensional Kundu-Mukherjee-Naskar model in polarization-preserving fibers is written as [9, 10]:

$$iq_t + aq_{xy} + ib(qq_x^* - q^*q_x)q = 0, \quad (1)$$

where $q(x, y, t)$ is a complex-valued function representing the wave profile, while a and b are real-valued constants. The first term is the linear temporal evolution, the second term represents the dispersion term, while the third term represents the nonlinearity term and $i = \sqrt{-1}$.

In Bragg gratings fibers, Eq.(1) can be written, for the first time as:

$$iu_t + a_1v_{xy} + i[(b_1u^2 + c_1v^2)u_x^* - (d_1|u|^2 + e_1|v|^2)u_x] + i\alpha_1u_x + \beta_1v + \sigma_1u^*v^2 = 0 \quad (2)$$

and

$$iv_t + a_2u_{xy} + i[(b_2v^2 + c_2u^2)v_x^* - (d_2|v|^2 + e_2|u|^2)v_x] + i\alpha_2v_x + \beta_2u + \sigma_2v^*u^2 = 0 \quad (3)$$

where $u(x, y, t)$ and $v(x, y, t)$ are complex-valued functions that represent the wave profiles, while $a_j, b_j, c_j, d_j, e_j, \alpha_j, \beta_j, \sigma_j$ ($j = 1, 2$) are real-valued constants. Here, a_j are the coefficients of dispersion terms. The parameters b_j, c_j, d_j, e_j ($j = 1, 2$) are the coefficients of nonlinearity. Next, α_j, β_j and σ_j give the inter-modal dispersions (IMD), the detuning parameters and the four wave mixing (4WM) parameters, respectively. Yildirim [11] has discussed the birefringent fibers of Eq.(1) using the modified simple equation approach.

The main objective of this article is to apply the modified Kudryashov's approach and the addendum to Kudryashov's method to find the dark, bright and singular soliton solutions of Eqs.(2) and (3).

The organization of this article can be written as: The mathematical preliminaries are introduced in section 2. In sections 3 and 4, we give the solutions of the system (2) and (3). In section 5, the numerical simulations for some solutions are designed. In section 6, conclusions are illustrated.

2 Mathematical preliminaries

In this section, we suppose that Eqs.(2) and (3) have the solutions:

$$\begin{aligned} u(x, y, t) &= P_1(\xi) \exp[i(-\kappa_1x - \kappa_2y + \omega t + \theta_0)], \\ v(x, y, t) &= P_2(\xi) \exp[i(-\kappa_1x - \kappa_2y + \omega t + \theta_0)], \end{aligned} \quad (4)$$

and

$$\xi = B_1 x + B_2 y - \rho t, \quad (5)$$

where B_1 , B_2 , ρ , κ_1 , κ_2 , ω and θ_0 are all non-zero real constants to be determined. The parameters B_1 and B_2 represent direct cosines and are related to the inverse widths of the soliton in the x - and y - directions, respectively, while ρ is the soliton velocity. From the phase component, κ_1 and κ_2 give the frequencies of the solitons along the x - and y - directions, respectively, while ω is the wave number and θ_0 is the phase constant. Here, $P_1(\xi)$ and $P_2(\xi)$ are real valued functions which stand for the pulse shapes. If we substitute (4) and (5) into Eqs.(2) and (3) and separate the real and imaginary parts, we deduce that

$$\begin{aligned} a_1 B_1 B_2 P_2'' + (B_1 - a_1 \kappa_1 \kappa_2) P_2 - \omega P_1 + \kappa_1 (b_1 - d_1) P_1^3 \\ + [\kappa_1 (c_1 - e_1) + \sigma_1] P_1 P_2^2 = 0, \end{aligned} \quad (6)$$

$$\begin{aligned} a_2 B_1 B_2 P_1'' + (B_2 - a_2 \kappa_1 \kappa_2) P_1 - \omega P_2 + \kappa_2 (b_2 - d_2) P_2^3 \\ + [\kappa_2 (c_2 - e_2) + \sigma_2] P_2 P_1^2 = 0, \end{aligned} \quad (7)$$

and

$$(\alpha_1 B_1 - \rho) P_1' - a_1 (\kappa_2 B_1 + B_2 \kappa_1) P_2' + B_1 (b_1 - d_1) P_1^2 P_1' + B_1 (c_1 - e_1) P_2^2 P_1' = 0, \quad (8)$$

$$(\alpha_2 B_2 - \rho) P_2' - a_2 (\kappa_1 B_2 + B_1 \kappa_2) P_1' + B_2 (b_2 - d_2) P_2^2 P_2' + B_2 (c_2 - e_2) P_1^2 P_2' = 0. \quad (9)$$

Setting

$$P_2(\xi) = A P_1(\xi), \quad (10)$$

where A is a non zero constant, such that $A \neq 1$. Now, Eqs.(6) - (9) become

$$a_1 B_1 B_2 A P_1'' + [A (B_1 - a_1 \kappa_1 \kappa_2) - \omega] P_1 + [\kappa_1 (b_1 - d_1) + A^2 \kappa_1 (c_1 - e_1) + A^2 \sigma_1] P_1^3 = 0, \quad (11)$$

$$a_2 B_1 B_2 P_1'' + (B_2 - a_2 \kappa_1 \kappa_2 - A \omega) P_1 + A [A^2 \kappa_2 (b_2 - d_2) + \kappa_2 (c_2 - e_2) + \sigma_2] P_1^3 = 0, \quad (12)$$

and

$$[\alpha_1 B_1 - \rho - a_1 A (\kappa_2 B_1 + B_2 \kappa_1)] P_1' + [B_1 (b_1 - d_1) + A^2 B_1 (c_1 - e_1)] P_1^2 P_1' = 0, \quad (13)$$

$$[A (\alpha_2 B_2 - \rho) - a_2 (\kappa_1 B_2 + B_1 \kappa_2)] P_1' + A B_2 [A^2 (b_2 - d_2) + c_2 - e_2] P_1^2 P_1' = 0. \quad (14)$$

Integrating Eqs.(13) and (14) with zero-integration constants, we have

$$[\alpha_1 B_1 - \rho - a_1 A (\kappa_2 B_1 + B_2 \kappa_1)] P_1 + \frac{1}{3} B_1 [b_1 - d_1 + A^2 (c_1 - e_1)] P_1^3 = 0, \quad (15)$$

$$[A(\alpha_2 B_2 - \rho) - a_2(\kappa_1 B_2 + B_1 \kappa_2)] P_1 + \frac{1}{3} A B_2 [A^2(b_2 - d_2) + c_2 - e_2] P_1^3 = 0. \quad (16)$$

Setting the coefficients of the linearly independent functions of Eqs. (15) and (16) to zero, yields

$$\rho = \alpha_1 B_1 - a_1 A(\kappa_2 B_1 + B_2 \kappa_1), \quad (17)$$

$$\rho = \frac{A\alpha_2 B_2 - a_2(\kappa_1 B_2 + B_1 \kappa_2)}{A}, \quad (18)$$

and the constraints conditions

$$b_1 - d_1 + A^2(c_1 - e_1) = 0, \quad (19)$$

$$A^2(b_2 - d_2) + c_2 - e_2 = 0. \quad (20)$$

Eqs.(11) and (12) are equivalent under the constraint conditions:

$$a_1 A = a_2, \quad (21)$$

$$A(B_1 - a_1 \kappa_1 \kappa_2) - \omega = B_2 - a_2 \kappa_1 \kappa_2 - A\omega, \quad (22)$$

$$\kappa_1(b_1 - d_1) + A^2 \kappa_1(c_1 - e_1) + A^2 \sigma_1 = A[A^2 \kappa_2(b_2 - d_2) + \kappa_2(c_2 - e_2) + \sigma_2]. \quad (23)$$

From (21) and (22), we have the wave number of the soliton:

$$\omega = \frac{AB_1 - B_2}{(1 - A)}. \quad (24)$$

Eq.(11) can be rewritten in the form:

$$P_1''(\xi) + L_1 P_1(\xi) + L_2 P_1^3(\xi) = 0, \quad (25)$$

where

$$L_1 = \frac{A(B_1 - a_1 \kappa_1 \kappa_2) - \omega}{a_1 B_1 B_2 A}, L_2 = \frac{\kappa_1(b_1 - d_1) + A^2 \kappa_1(c_1 - e_1) + A^2 \sigma_1}{a_1 B_1 B_2 A}, \quad (26)$$

provided $a_1 B_1 B_2 A \neq 0$. In the next two sections, we solve Eq. (25) using the following two methods:

3 The modified Kudryashov's method

According to this method, we balance $P_1''(\xi)$ with $P_1^3(\xi)$ in Eq.(25), we get:

$$M + 2p = 3M \implies M = p. \quad (27)$$

Now, the following cases can be considered.

Case1. Setting $p = 1$, then $M = 1$. Thus, Eq.(25) has the formal solution:

$$P_1(\xi) = A_0 + A_1 Q(\xi), \quad (28)$$

where A_0 and A_1 are parameters, provided $A_1 \neq 0$. The function $Q(\xi)$ satisfies the differential equation:

$$Q'(\xi) = Q(\xi)[Q(\xi) - 1] \ln K, \quad 0 < K \neq 1. \quad (29)$$

Substituting (28) along with (29) into (25), setting all the coefficients of $Q(\xi)^l$ ($l = 0, 1, 2, 3$) to zero, we have the results

$$A_0 = -\sqrt{-\frac{\ln^2 K}{2L_2}}, A_1 = 2\sqrt{-\frac{\ln^2 K}{2L_2}}, L_1 = \frac{1}{2} \ln^2 K, \quad (30)$$

provided $L_2 < 0$. Substituting (30) along with the well-known solutions of Eq.(29) obtained in [17-19] into (28), we get the solutions:

$$u(x, y, t) = -\sqrt{-\frac{\ln^2 K}{2L_2}} \left[1 - \frac{2}{1+\epsilon K^{(\xi+\xi_0)}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (31)$$

and

$$v(x, y, t) = -A\sqrt{-\frac{\ln^2 K}{2L_2}} \left[1 - \frac{2}{1+\epsilon K^{(\xi+\xi_0)}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (32)$$

The solutions (31), (32) can be rewritten in the form

$$u(x, y, t) = -\sqrt{-\frac{\ln^2 K}{2L_2}} \left[1 - \frac{2}{1+\epsilon \{\cosh[(\xi+\xi_0) \ln k] + \sinh[(\xi+\xi_0) \ln k]\}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (33)$$

and

$$v(x, y, t) = -A\sqrt{-\frac{\ln^2 K}{2L_2}} \left[1 - \frac{2}{1+\epsilon \{\cosh[(\xi+\xi_0) \ln k] + \sinh[(\xi+\xi_0) \ln k]\}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (34)$$

which represent the combo bright-singular soliton solutions. In particular, if $\epsilon = 1$, we have the dark soliton solution:

$$u(x, y, t) = -\sqrt{-\frac{\ln^2 K}{2L_2}} \tanh \left[\frac{(\xi+\xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (35)$$

and

$$v(x, y, t) = -A\sqrt{-\frac{\ln^2 K}{2L_2}} \tanh \left[\frac{(\xi+\xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (36)$$

while if $\epsilon = -1$, we have the singular soliton solution:

$$u(x, y, t) = -\sqrt{-\frac{\ln^2 K}{2L_2}} \coth \left[\frac{(\xi+\xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (37)$$

and

$$v(x, y, t) = -A\sqrt{-\frac{\ln^2 K}{2L_2}} \coth \left[\frac{(\xi+\xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (38)$$

Case 2. Setting $p = 2$, then $M = 2$. Thus, Eq.(25) has the formal solution:

$$P_1(\xi) = A_0 + A_1 Q(\xi) + A_2 Q^2(\xi), \quad (39)$$

where A_0, A_1 and A_2 are parameters, provided $A_2 \neq 0$. The function $Q(\xi)$ satisfies the differential equation:

$$Q'(\xi) = Q(\xi) [Q^2(\xi) - 1] \ln K, \quad 0 < K \neq 1. \quad (40)$$

Substituting (39) along with (40) into (25), setting all the coefficients of $Q(\xi)^l$ ($l = 0, \dots, 6$) to zero, we have the results

$$A_0 = -\frac{1}{2}\sqrt{-\frac{8\ln^2 K}{L_2}}, A_1 = 0, A_2 = \sqrt{-\frac{8\ln^2 K}{L_2}}, L_1 = 2\ln^2 K, \quad (41)$$

provided $L_2 < 0$. Substituting (41) along with the well-known solutions of Eq.(40) obtained in [17-19] into (39), we have the solutions:

$$u(x, y, t) = -\frac{1}{2}\sqrt{-\frac{8\ln^2 K}{L_2}} \left[1 - \frac{2}{1+\epsilon K^{(2\xi+\xi_0)}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (42)$$

and

$$v(x, y, t) = -\frac{1}{2}A\sqrt{-\frac{8\ln^2 K}{L_2}} \left[1 - \frac{2}{1+\epsilon K^{(2\xi+\xi_0)}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (43)$$

The solutions (42), (43) can be rewritten in the following form of the combo bright-singular soliton solutions:

$$u(x, y, t) = -\frac{1}{2}\sqrt{-\frac{8\ln^2 K}{L_2}} \left[1 - \frac{2}{1 + \epsilon \{ \cosh[(2\xi + \xi_0) \ln K] + \sinh[(2\xi + \xi_0) \ln K] \}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (44)$$

and

$$v(x, y, t) = -\frac{1}{2}A\sqrt{-\frac{8\ln^2 K}{L_2}} \left[1 - \frac{2}{1 + \epsilon \{ \cosh[(2\xi + \xi_0) \ln K] + \sinh[(2\xi + \xi_0) \ln K] \}} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (45)$$

In particular, when $\epsilon = 1$, we have the dark soliton solution:

$$u(x, y, t) = -\frac{1}{2}\sqrt{-\frac{8\ln^2 K}{L_2}} \tanh \left[\frac{(2\xi + \xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (46)$$

and

$$v(x, y, t) = -\frac{1}{2}A\sqrt{-\frac{8\ln^2 K}{L_2}} \tanh \left[\frac{(2\xi + \xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (47)$$

while if $\epsilon = -1$, we have the singular soliton solution:

$$u(x, y, t) = -\frac{1}{2}\sqrt{-\frac{8\ln^2 K}{L_2}} \coth \left[\frac{(2\xi + \xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (48)$$

and

$$v(x, y, t) = -\frac{1}{2}A\sqrt{-\frac{8\ln^2 K}{L_2}} \coth \left[\frac{(2\xi + \xi_0)}{2} \ln K \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (49)$$

Simliarly, we can find many other solutions by choosing other values for p and M .

4 The addendum to Kudryashov's method

According to this method, we balance $P_1''(\xi)$ with $P_1^3(\xi)$ in Eq.(25), we get the relation (27). Let us now discuss the following cases:

Case1. Choose $p = 1$, then $M = 1$. Now, we have the formal solution

$$P_1(\xi) = \chi_0 + \chi_1 R(\xi), \quad (50)$$

where χ_0 and χ_1 are parameters, provided $\chi_1 \neq 0$. Here $R(\xi)$ satisfies the differential equation:

$$R'^2(\xi) = R^2(\xi) [1 - \chi R^2(\xi)] \ln^2 K, \quad 0 < K \neq 1, \quad (51)$$

where χ is a constant. Substituting (50), (51) in (25) and setting all the coefficients of $[R(\xi)]^{q_1} [R'(\xi)]^{q_2}$, ($q_1 = 0, 1, 2, 3$, $q_2 = 0, 1$), to zero, we have the results

$$\chi_0 = 0, \chi_1 = \sqrt{\frac{2\chi \ln^2 K}{L_2}}, L_1 = -\ln^2 K, \quad (52)$$

provided $\chi L_2 > 0$ and $L_1 < 0$. Substituting (52) along with the well-known solutions of Eq.(51) obtained in [17-19] in (50), we have the solutions:

$$u(x, y, t) = \sqrt{\frac{2\chi \ln^2 K}{L_2}} \left(\frac{4S}{4S^2 K^{\xi+\chi} K^{-\xi}} \right) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (53)$$

and

$$v(x, y, t) = A \sqrt{\frac{2\chi \ln^2 K}{L_2}} \left(\frac{4S}{4S^2 K^{\xi+\chi} K^{-\xi}} \right) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (54)$$

The solutions (53), (54) can be rewritten in the form:

$$u(x, y, t) = \sqrt{\frac{2\chi \ln^2 K}{L_2}} \left[\frac{4S}{(4S^2 + \chi) \cosh(\xi \ln K) + (4S^2 - \chi) \sinh(\xi \ln K)} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (55)$$

and

$$v(x, y, t) = A \sqrt{\frac{2\chi \ln^2 K}{L_2}} \left[\frac{4S}{(4S^2 + \chi) \cosh(\xi \ln K) + (4S^2 - \chi) \sinh(\xi \ln K)} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (56)$$

which represent the combo bright-singular soliton solution. In particular, when $\chi = 4S^2$, we have the bright soliton solution:

$$u(x, y, t) = \sqrt{\frac{2\ln^2 K}{L_2}} \operatorname{sech}(\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (57)$$

and

$$v(x, y, t) = A \sqrt{\frac{2\ln^2 K}{L_2}} \operatorname{sech}(\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (58)$$

provided $L_2 > 0$, while if $\chi = -4S^2$, we have the singular soliton solutions:

$$u(x, y, t) = \sqrt{-\frac{2\ln^2 K}{L_2}} \operatorname{csch}(\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (59)$$

and

$$v(x, y, t) = A \sqrt{-\frac{2\ln^2 K}{L_2}} \operatorname{csch}(\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (60)$$

provided $L_2 < 0$.

Case 2. Choose $p = 2$, then $M = 2$. Now, we have the formal solution

$$P_1(\xi) = \chi_0 + \chi_1 R(\xi) + \chi_2 R^2(\xi), \quad (61)$$

where χ_0, χ_1 and χ_2 are parameters, provided $\chi_2 \neq 0$. Here $R(\xi)$ satisfies the differential equation:

$$R'^2(\xi) = R^2(\xi) [1 - \chi R^4(\xi)] \ln^2 K, \quad 0 < K \neq 1, \quad (62)$$

where χ is a constant. Substituting (61), (62) in (25) and setting all the coefficients of $[R(\xi)]^{q_1}$ $[R'(\xi)]^{q_2}$, ($q_1 = 0, \dots, 6$, $q_2 = 0, 1$) to zero, we have the results

$$\chi_0 = 0, \chi_1 = 0, \chi_2 = \sqrt{\frac{8\chi \ln^2 K}{L_2}}, L_1 = -4 \ln^2 K, \quad (63)$$

provided $\chi L_2 > 0$ and $L_1 < 0$. Substituting (63) along with the well-known solutions of Eq.(62) obtained in [17-19] into (61), we have the solutions:

$$u(x, y, t) = \sqrt{\frac{8\chi \ln^2 K}{L_2}} \left(\frac{4S}{4S^2 K^{2\xi} + \chi K^{-2\xi}} \right) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (64)$$

and

$$v(x, y, t) = A \sqrt{\frac{8\chi \ln^2 K}{L_2}} \left(\frac{4S}{4S^2 K^{2\xi} + \chi K^{-2\xi}} \right) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (65)$$

The solutions (64), (65) can be rewritten in the following form of the combo bright-singular soliton solutions:

$$u(x, y, t) = \sqrt{\frac{8\chi \ln^2 K}{L_2}} \left[\frac{4S}{(4S^2 + \chi) \cosh(2\xi \ln K) + (4S^2 - \chi) \sinh(2\xi \ln K)} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (66)$$

and

$$v(x, y, t) = A \sqrt{\frac{8\chi \ln^2 K}{L_2}} \left[\frac{4S}{(4S^2 + \chi) \cosh(2\xi \ln K) + (4S^2 - \chi) \sinh(2\xi \ln K)} \right] e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}. \quad (67)$$

In particular, when $\chi = 4S^2$, we have the bright soliton solution:

$$u(x, y, t) = \sqrt{\frac{8 \ln^2 K}{L_2}} \operatorname{sech}(2\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (68)$$

and

$$v(x, y, t) = A \sqrt{\frac{8 \ln^2 K}{L_2}} \operatorname{sech}(2\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (69)$$

provided $L_2 > 0$, while if $\chi = -4S^2$, we have the singular soliton solution:

$$u(x, y, t) = \sqrt{-\frac{8 \ln^2 K}{L_2}} \operatorname{csch}(2\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (70)$$

and

$$v(x, y, t) = A \sqrt{-\frac{8 \ln^2 K}{L_2}} \operatorname{csch}(2\xi \ln K) e^{i(-\kappa_1 x - \kappa_2 y + \omega t + \theta_0)}, \quad (71)$$

provided $L_2 < 0$.

Similiarly, we can find many other solutions by choosing other values for p and M .

5 Numerical simulations

In this section, we introduce the graphs of the absolute values of some solutions for Eqs.(35) and (36), Eqs.(48) and (49) and Eqs.(57) and (58). Let us now examine Figures, 1-3. as it illustrates some of our solutions obtained in this paper. To this purpose, we choose some special values of the obtained parameters.

Figure-1 shows the profile of the absolute values of the dark soliton solutions (35) and (36).

Figure 1: The numerical simulations of the absolute values of the solutions (35) and (36) with the parameter values $a_1 = 0.02, a_2 = 0.03, \kappa_1 = 0.2, \kappa_2 = 0.1, b_1 = 0.1, d_1 = 0.2, c_1 = 0.01,$

$e_1 = 0.03, \sigma_1 = -0.05, B_1 = 0.2, B_2 = 0.3, y = 0.5, \alpha_1 = 0.2, K = 3, \xi_0 = 0.1, -5 \leq x, t \leq 5.$

Figure-2 shows the profile of the absolute values of the singular soliton solutions (48) and (49).

Figure 2: The numerical simulations of the absolute values of the solutions (48) and (49) with the parameter values $a_1 = 0.02, a_2 = 0.03, \kappa_1 = 0.2, \kappa_2 = 0.1, b_1 = 0.1, d_1 = 0.2, c_1 = 0.01, e_1 = 0.03, \sigma_1 = -0.05, B_1 = 0.2, B_2 = 0.3, x = 0.2, \alpha_1 = 0.2, K = 4, \xi_0 = 0.1, -5 \leq y, t \leq 5$.

Figure-3 shows the profile of the absolute values of the bright soliton solutions (57) and (58).

Figure 3: The numerical simulations of the absolute values of the solutions (57) and (58) with the parameter values $a_1 = 0.02, a_2 = 0.03, \kappa_1 = 0.2, \kappa_2 = 0.1, b_1 = 0.1, d_1 = 0.2, c_1 = 0.01, e_1 = 0.03, \sigma_1 = 0.05, B_1 = 0.2, B_2 = 0.3, y = 0.5, \alpha_1 = 0.2, K = 2, \xi_0 = 0.1, -5 \leq x, t \leq 5$.

From the above Figures, one can see that the obtained solutions possess the dark, singular and bright soliton solutions. Also, these Figures express the behaviour of these solutions which give some perspective readers how the behaviour solutions are produced.

6 Conclusions

This paper studies the nonlinear (2+1)-dimensional Kundu-Mukherjee-Naskar equation in fiber Bragg gratings by the aid of two mathematical approaches which are the modified Kudryashov's approach and the addendum to Kudryashov's technique. Optical solitons to the coupled system (2) and (3) are regained. These optical solitons are emerged from constraint conditions. The coupled system (2) and (3) is studied in this paper for the first time. The results of the current paper have not been determined before. Also, the graphs of Eqs. (35), (36), (48), (49), (57) and (58) are given by choosing special values of constants. Profiles of dark, singular and bright soliton solutions respectively are represented in Figs. 1, 2 and 3 which express the behavior of optical solitons in fiber Bragg gratings.

ACKNOWLEDGMENTS

The authors also declare that there is no conflict of interest.

References

- [1] A. Biswas, J. Vega-Guzman, M.F. Mahmood, S. Khan, Q. Zhou, S.P. Moshokoa, M. Belic, Solitons in optical fiber Bragg gratings with dispersive reflectivity, *Optik*. Volume 182, 119–123. (2019).
- [2] A. Biswas, M. Ekici, A. Sonmezoglu, M.R. Belic, Solitons in optical fiber Bragg gratings with dispersive reflectivity by extended trial function method, *Optik*. Volume 182, 88–94. (2019).
- [3] A. Biswas, M. Ekici, A. Sonmezoglu, M.R. Belic, Optical solitons in fiber Bragg gratings with dispersive reflectivity for parabolic law nonlinearity by extended trial function method, *Optik*. Volume 183, 595–601. (2019).
- [4] A. Biswas, M. Ekici, A. Sonmezoglu, M.R. Belic, Optical solitons in fiber Bragg gratings with dispersive reflectivity for quadratic-cubic nonlinearity by extended trial function method, *Optik*. Volume 185 50–56. (2019).

- [5] A. Biswas, J. Vega-Guzman, M.F. Mahmood, M. Ekici, Q. Zhou, S.P. Moshokoa, M.R. Belic, Optical solitons in fiber Bragg gratings with dispersive reflectivity for parabolic law nonlinearity using undetermined coefficients, *Optik*. Volume 185, 39–44.(2019).
- [6] A. Biswas, Optical soliton perturbation with Radhakrishnan–Kundu–Lakshmanan equation by traveling wave hypothesis, *Optik*, 171, 217-220. (2018).
- [7] Y. Yıldırım, A. Biswas, M. Ekici, H. Triki, O. G. Gaxiola, A. K. Alzahrani, M. R. Belic, Optical solitons in birefringent fibers for Radhakrishnan–Kundu–Lakshmanan equation with five prolific integration norms, *Optik*, 208, 164550. (2020).
- [8] S. Arshed, A. Biswas, P. Guggilla, A.S. Alshomrani, Optical solitons for Radhakrishnan–Kundu–Lakshmanan equation with full nonlinearity, *Phys. Lett. A* 384, 126191. (2020).
- [9] Y. Yıldırım, Optical solitons to Kundu–Mukherjee–Naskar model with modified simple equation approach, *Optik*, 184, 247-252. (2019).
- [10] M. Ekici, A. Sonmezoglu, A. Biswas, M. R. Belic, Optical solitons in (2+1)–Dimensions with Kundu–Mukherjee–Naskar equation by extended trial function scheme, *Chin. J. Phys*, 57, 72–77. (2019).
- [11] Y. Yıldırım, Optical solitons to Kundu–Mukherjee–Naskar model in birefringent fibers with modified simple equation approach, *Optik*, 184, 121-127. (2019).
- [12] N.A. Kudryashov, D.V. Safonova, A. Biswas, Painleve analysis and a solution to the traveling wave reduction of the Radhakrishnan–Kundu–Lakshmanan equation, *Reg. Chaot. Dyn.* 24, 607614. (2019).
- [13] N. A. Kudryashov. “Method for finding highly dispersive optical solitons of nonlinear differential equations". *Optik*. Volume 206, 163550. (2020).
- [14] N. A. Kudryashov. “First Integrals and General Solution of the Complex Ginzburg – Landau Equation". *Appl. Math. Comput.* Volume 386, 125407. (2020).
- [15] N. A. Kudryashov. “First integrals and solutions of the traveling wave reduction for the Triki – Biswas equation". *Optik*. Volume 185, 275–281. (2019).
- [16] N. A. Kudryashov. “General solution of traveling wave reduction for the Kundu – Mukherjee – Naskar equation". *Optik*. Volume 186, 22–27. (2019).
- [17] E. M. E. Zayed, T. A. Nofal, K. A. Gepreel, R. M. A. Shohib, M.E. M. Alngar, “Cubic-quartic optical soliton solutions in fiber Bragg gratings with Lakshmanan–Porsezian–Daniel model by two integration schemes", *Optical and Quantum Electronics*. Volume 53, 249 (2021).
- [18] E. M.E. Zayed, K. A. Gepreel, M. E.M. Alngar, “Addendum to Kudryashov’s method for finding solitons in magneto-optics waveguides to cubic-quartic NLSE with Kudryashov’s sextic power law of refractive index", *Optik*. Volume 230, 166311 (2021).

- [19] E. M.E. Zayed, R. M.A. Shohib, K. A. Gepreel, M. M. El-Horbaty, M. E.M. Alngar, "Cubic–quartic optical soliton perturbation Biswas–Milovic equation with Kudryashov’s law of refractive index using two integration methods" *Optik*. Volume 239, 166871 (2021).