A Geospatial Examination of Specialist Care Accessibility and Impact on Health Outcomes for Patients With Acute Traumatic Spinal Cord Injury in Nsw, Australia: A Whole Population Record Linkage Study

Lisa Nicole Sharwood (✉ lisa.sharwood@sydney.edu.au)

Sydney Medical School Faculty of Medicine and Health University of Sydney <https://orcid.org/0000-0001-8977-6144>

David Whyatt

University of Western Australia Faculty of Health and Medical Sciences

Bharat P Vaikuntam

The University of Sydney Faculty of Medicine and Health

Christiana L Cheng

Praxis Institute

Vanessa K Noonan

Praxis Institute

Anthony P Joseph

RNSH: Royal North Shore Hospital

Jonathon Ball

RNSH: Royal North Shore Hospital

Ralph E Stanford

Prince of Wales Hospital

Mei-Ruu Kok

University of Western Australia Faculty of Health and Medical Sciences

Samuel R Withers

Australian Institute of Orthopedic Robotics

James W Middleton

NSW Agency for Clinical Innovation: New South Wales Agency for Clinical Innovation

Research article

Keywords: acute traumatic spinal cord injury, geographic information science, geospatial methods, SCIU, trauma centre, health services pathways

Posted Date: October 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-78716/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Health Services Research on April 1st, 2021. See the published version at <https://doi.org/10.1186/s12913-021-06235-4>.

Abstract

Background

Timely treatment is essential for achieving optimal outcomes after traumatic spinal cord injury, and expeditious transfer to a specialist spinal cord injury unit is recommended within 24 hours from injury. Previous research in New South Wales found only 57% of TSCI patients were admitted to specialist spinal cord injury unit for acute post-injury care; 73% transferred within 24 hours from injury. We evaluated pre-hospital and inter-hospital transfer practices to identify opportunities for improving care delivery.

Methods

This record linkage study included administrative pre-hospital, admissions and costs data for all patients aged ≥ 16 years with incident traumatic spinal cord injury in New South Wales (2013-2016). We investigated impacts of geographical disparities on pre-hospital and inter-hospital transport decisions from injury location using geospatial methods to better understand the post-injury care pathways impact on patient outcomes.

Results

Inclusion criteria identified 316 patients, geospatial analysis showed that over half (53%, $n=168$) of all patients were injured within 60 minutes road travel of a SCIU, yet only 28.6% ($n=48$) were directly transferred to a specialist spinal cord injury unit. Direct transfers received earlier operative intervention (median (IQR) 12.9(7.9) hours), compared with patients transferred indirectly to specialist spinal cord injury unit (median (IQR) 19.5(18.9) hours), and had lower risk of complications (OR 3.2 v 1.4, $p<0.001$).

Conclusions

Getting patients with acute traumatic spinal cord injury patients to the right place at the right time is dependent on numerous factors; some are still being triaged directly to non-trauma services which delays specialist and surgical care and increases complication risks. More stringent adherence to recommended guidelines would prioritise direct specialist spinal cord injury unit transfer for patients injured within 60 minutes radius, enabling the benefits of specialised care.

Background

Traumatic spinal cord injury (TSCI) is a devastating condition with lifelong physical, psychosocial and economic impacts (1–5). Acute post-injury phase management is time-sensitive, with rapid access to specialist care deemed essential for achieving optimal outcomes. Expeditious transfer to a specialist spinal cord injury unit (SCIU) is recommended within 24 hours following injury (6–8). Previous research in New South Wales (NSW), Australia (9) found only 57% of patients with confirmed TSCI admitted to a SCIU for acute care, and of these, only 73% were transferred within 24 hours from injury. Rurality of

residence was associated with more timely admission to SCIU in this study (9), geospatial information system (GIS) data for injury incidents was unable to support this claim.

Specialist health services are often concentrated in high population density areas, though may often serve large geographical areas with substantial, distributed populations. Time and distance consequently challenge equitable access to these services for the whole population. NSW is the most populous Australian state, covering an area of 809,444 km². NSW specialist health services include the State Spinal Cord Injury Service (SSCIS) network, providing care for approximately 5,500 people living with spinal cord injury (2019)(10). TSCI is a high acuity, resource intensive injury requiring ongoing acute and rehabilitative care; in 2017-18, there were 3,888 re-hospitalisations of people living with spinal cord injury in NSW.

Epidemiological, health services studies are increasingly using geospatial methods (11–13) to better understand and target both injury prevention and post-injury care (14). Triage optimisation using geospatial data and scenario modelling has demonstrated significantly reduced time to SCIU admission in a Canadian study (15). Similar modelling (without geospatial data) showed that optimizing patient-care pathways can achieve significant health system cost reductions from direct in comparison with indirect admissions to SCIU (16). Studying individuals with acute traumatic brain injury (TBI), Brown et al (12) found significant disparity in survival rates between rural and urban areas using geospatial analysis. Specifically, in the most rural areas, TBI fatality rates were 13 deaths per 100,000 persons higher than those in the most urban area (95% confidence interval 12.15–13.86; $P < 0.001$) (12). A similar analysis across the NSW trauma system demonstrated a significantly higher adjusted mortality rate for patients with any traumatic injury treated in regional services compared with metropolitan major trauma services (17). Definitive care at a Major Trauma Service (MTS) was associated with a 41% lower likelihood of death compared to definitive care at a Regional Trauma Service (RTS) (OR 0.59 95%CI 0.35–0.97) (17).

Lack of access to specialist care services, including major trauma management (18), timely and appropriate surgical interventions within a critical time window and other factors such as increased rates of complications are likely contributing to higher mortality rates and poor long term outcomes for survivors. There is an indication internationally that regardless of prescribed, evidence-based pathways for trauma patients and injury locations within acceptable proximity to specialist services, some acute trauma patients are still being triaged to non-trauma services (19). Delivery of health services must account for the fact that injuries will occur in a range of geographic locations, from urban to very remote areas. Triaging systems that direct patient pathways through pre-hospital and acute care settings must therefore offer equitable opportunity for optimal outcomes regardless of the incident location. Geospatial analysis for victims of burn injury, as well as bicyclist crash, has been previously conducted in NSW, targeted towards informing public health interventions and aid policy makers plan for service provision (20, 21). The extent to which geospatial variables impact pathways to direct SCIU admission for patients with TSCI in Australia has not been examined.

The aim of this study was to use geospatial methods to investigate the impact of geographical location on pathways and timing of admission to specialist care services for individuals sustaining acute TSCI across the state of NSW, Australia. Analyses will control for relevant variables and assess associations with patient outcomes, including time to surgery and inpatient complications.

The NSW Population Research Ethics Committee (2012/09/420) approved this project.

Methods

Setting and acute Health Services

The state of NSW, Australia, has a population of just over 8 million persons (22) and covers a geographically diverse area > 800,000 km². Around two-thirds of the population reside in the Greater Sydney (suburban) area; the remainder in rural and very remote areas. The state government funded NSW Ambulance Service is the sole emergency medical service, transporting patients via road, fixed wing or helicopter depending on injury severity and geographic location. Road ambulances are not routinely staffed with emergency physicians, whereas some helicopter services are.

Trauma service hospitals are designated as either MTS (equivalent to Level 1 Trauma Service, accredited) or RTS (equivalent to Level 3) (23); critically ill patients can be taken to one of six strategically located MTS or ten RTS. Approximately 200 additional non-trauma designated hospitals (district/regional/local hospitals) are situated across metropolitan and regional health districts around the state. The SSCIS comprised two specialist SCIUs, both located in metropolitan Sydney. One of these is also an MTS (categorised as SCIU for this study), the other a non-trauma designated hospital; both providing specialist spinal surgical services for TSCI. Hospitals were categorized as trauma hospitals (MTS or RTS), specialist hospitals (SCIU) or non-trauma hospitals (district/local) for the analysis.

Being suspected of having TSCI or evidence of TSCI meets major trauma criteria under the NSW state-wide pre-hospital triage criteria applies; whereby 'patients meeting major trauma criteria should be transported to the highest level Trauma Centre located within a 60-minute driving radius, which may include bypassing closer non-trauma hospitals'(24). Recommendation from the SSCIS was that once medically stable, patients with TSCI should be transferred to an SCIU within 24 hours from injury (25).

Population-level record linkage

The NSW Admitted Patient Data Collection (APDC) was used to identify and extract TSCI patient records from all separations/discharges from NSW public hospitals, based on specific TSCI-related International Classification of Diseases, 10th version, Australian Modification (ICD-10-AM) diagnosis codes (26). Probabilistic data linkage was undertaken by the NSW Centre for Health Record Linkage, linking all patients where a TSCI code was either a principal or additional diagnosis, for any separation within the APDC. Included patients were aged 16 years or more, injured between 1 June 2013-30 June 2016, admitted to a NSW hospital and diagnosed as having TSCI using ICD-10AM diagnostic codes (27).

Excluded were those with missing GIS data for these analyses. The first hospital episode and all contiguous care episodes, including nested/non-nested transfers, was deemed the 'index admission'.

TSCI case records were linked with Emergency Department and NSW Ambulance data. All datasets and variables included are detailed in Appendix 1.

Outcome variables

Primary outcome was triage pattern to SCIU (direct or indirect admission vs no admission) for acute incident TSCI patients. Direct admissions included those first transferred to SCIUs, indirect admissions were those transferred to SCIU hospitals from all non-SCIU hospitals. The definitive hospital was defined as the final hospital where the patient received their acute care, prior to either rehabilitation admission, discharge home or death, indicated by acute care-type and separation/discharge destination.

Secondary outcomes considered included time to surgery (for operated patients) and incidence of inpatient complications, adjusted for relevant predictor variables including triage pattern to SCIU. Inpatient complications included pressure injuries, urinary tract infections, respiratory infections, deep vein thrombosis, pulmonary embolus and others.

Geospatial mapping

Python language was used to conduct geospatial analyses, interacting with Google Distance Matrix Application Programming Interfaces (API) to calculate on-road travel distances and times relative to hospitals/SCIUs using injury location GIS co-ordinates. Travel duration was classified as < 60 minutes or > 60 minutes from SCIU, and/or MTS from injury location, calculated for each patient. Population dispersement data (ABS) (22) enabled risk of disease mapping per unit-head of population over the study time-period according to Statistical Area (SA) boundaries. Population sizes for different SAs for the years 2012–2015 were calculated by linear interpolation using the 2011 and 2016 populations. Crude rates/100,000 person-years were calculated, and due to the sample size, SA3 determined the most stable representation, using person-years between 1st June 2013–30th June 2016 (Fig. 1).

Statistical analysis

Descriptive analysis included continuous variables summarised as means (standard deviations) and medians (interquartile range); categorical variables as percentages. Statistically significant differences (< 0.05) between groups were tested using t-tests, Wilcoxon Rank Sum or chi-square tests. Multinomial logistic regression modelling examined associations between predictor variables and direct/indirect/no admission to SCIU. Age, gender, injury mechanism, transport mode, presence of multi-trauma, co-morbidity (CCI), geographic distance from incident location relative to SCIU hospital, travel time to SCIU, and injury severity (ICISS score) were included in the regression analysis for the entire study population. Admission times to 'definitive hospitals' were used from variables denoting the first admission time to the final (definitive) hospital for the acute care stay. For those patients not transferred, this was also the first hospital to which they were admitted. A multinomial logistic regression model (28) was assessed using

Relative Risk Ratios (RRR), obtained by exponentiating the multinomial logit coefficients, e^{coef} . Modelling used the category 'Direct to SCIU' as the referent group; the RRR for the groups 'Indirect to SCIU' and 'No SCIU' indicating risks relative to this referent group. STATA-IC.v.16(29) was used for all analyses.

Results

Demographic and injury profile

During the study period, 534 patients with acute incident TSCI were identified across NSW; 316 (59%) had GIS data for inclusion in geospatial analysis. Injury mechanisms were commonly falls ($n = 149$, 47.2%) and transport crashes ($n = 99$, 31.3%), with 53% ($n = 168$) sustaining multiple trauma. Over one-third of patients ($n = 126$, 39.9%) had higher injury severity scores (ICISS < 0.83)(30); the median (IQR) ICISS score was 0.839 (0.137) (Table 1); over half of all patients sustained a cervical level injury ($n = 170$, 53.8%).

Table 1 compares baseline demographic and injury epidemiology profile for patients admitted directly, indirectly or not at all to SCIU. Patients were more likely to experience direct transfer to a SCIU without comorbid trauma ($p < 0.01$) but higher ICISS ($p < 0.001$), cervical injury ($p < 0.01$), and transferred by air-ambulance ($p < 0.01$). Indirect transfer to SCIU was more likely with two or more additional traumatic injuries ($p < 0.01$) or incomplete injury ($p < 0.01$). Patients not admitted to SCIU at all were older ($p = 0.05$) with lower levels of injury ($p < 0.01$).

Table 1
Comparing patients with TSCI admitted directly or indirectly to SCIU, or not at all.

Variable	Direct admit to SCIU (n = 85, 26.9%)	Indirect admit to SCIU (n = 108, 34.2%)	No acute admit to SCIU (n = 123, 38.9%)	Total (n = 316)	p-value
	n (%)	n (%)	n (%)	n (%)	
Age years (median (IQR))	57.7 (28.9)	52.7 (37.1)	60.6 (38.1)	56.5 (36.3)	0.05
16–30	17 (20)	27 (25)	18 (14.6)	62 (19.6)	
31–45	12 (14.1)	20 (18.5)	17 (13.8)	49 (15.5)	
46–60	22 (25.9)	23 (21.3)	28 (22.8)	73 (23.1)	
61–75	20 (23.5)	24 (22.2)	22 (17.9)	66 (20.9)	
76+	14 (16.5)	14 (12.9)	38 (30.9)	66 (20.9)	
Sex					0.10
Female	16 (18.8)	22 (20.4)	37 (30.1)	75 (23.7)	
Male	69 (81.2)	86 (79.6)	86 (69.9)	241 (76.3)	
Injury mechanism					0.15
Fall	43 (50.6)	45 (41.7)	61 (49.6)	149 (47.2)	
Transport	32 (37.6)	35 (32.4)	32 (26.0)	99 (31.3)	
Self-harm/Assault	1 (1.2)	8 (7.4)	7 (5.7)	16 (5.1)	
Other [‡]	9 (10.6)	20 (18.5)	23 (18.7)	52 (16.5)	
Incident GPS					0.74

[‡]Other included diving, horse-related incidents or not specified

*SEIFA – Socio-Economic Indexes for Areas (31)

at admission

Variable	Direct admit to SCIU (n = 85, 26.9%)	Indirect admit to SCIU (n = 108, 34.2%)	No acute admit to SCIU (n = 123, 38.9%)	Total (n = 316)	p- value
≤ 60 min to SCIU	48 (56.4)	55 (50.9)	65 (52.8)	168 (53.2)	
> 60 min to SCIU	37 (43.5)	53 (49.1)	58 (47.1)	148 (46.8)	
SEIFA Quintiles					0.002
1 (lowest)	8 (9.4)	21 (19.4)	15 (12.2)	44 (13.2)	
2	14 (16.5)	24 (22.2)	29 (23.6)	67 (21.2)	
3	11 (12.9)	23 (21.3)	37 (30.1)	71 (22.5)	
4	15 (17.6)	11 (10.2)	17 (13.8)	43 (13.6)	
5 (highest)	35 (41.2)	28 (25.9)	20 (16.3)	83 (26.3)	
unknown	2 (2.3)	1 (0.9)	5 (4.1)	8 (2.5)	
Charlson (CCI)					0.84
0	60 (70.6)	77 (71.3)	82 (66.7)	219 (69.3)	
1	14 (16.5)	14 (12.9)	19 (15.4)	47 (14.9)	
2+	11 (12.9)	17 (15.7)	22 (17.9)	50 (15.8)	
Multiple Trauma					0.003
No	33 (38.8)	45 (41.7)	70 (56.9)	148 (46.8)	
1 other injury	29 (34.1)	20 (18.5)	23 (18.7)	72 (22.8)	
≥ 2 other injuries	23 (27.1)	43 (39.8)	30 (24.4)	96 (30.4)	

‡Other included diving, horse-related incidents or not specified

*SEIFA – Socio-Economic Indexes for Areas (31)

at admission

Variable	Direct admit to SCIU (n = 85, 26.9%)	Indirect admit to SCIU (n = 108, 34.2%)	No acute admit to SCIU (n = 123, 38.9%)	Total (n = 316)	p- value
ICISS Score					< 0.001
< 0.7 (Most severe)	9 (10.6)	5 (4.6)	11 (8.9)	25 (7.9)	
0.7 to < 0.83	48 (56.5)	26 (24.1)	27 (21.9)	101 (31.9)	
0.83 to < 0.89	17 (20)	29 (26.8)	23 (18.7)	69 (21.8)	
0.89 to < 0.95	8 (9.4)	21 (19.4)	29 (23.6)	58 (18.4)	
0.95 to 1.00	3 (3.5)	27 (25)	33 (26.8)	63 (19.9)	
Highest Level of Injury					< 0.001
Cervical	54 (63.5)	66 (61.1)	50 (40.6)	170 (53.8)	
Thoracic	22 (25.9)	27 (25)	34 (27.6)	83 (26.3)	
Lumbar	9 (10.6)	15 (13.9)	39 (31.7)	63 (19.9)	
Extent of Injury					< 0.001
Complete	24 (28.2)	23 (21.3)	5 (4.1)	52 (16.5)	
Incomplete	43 (50.6)	57 (52.8)	39 (31.7)	139 (43.9)	
Conus /cauda equina	11 (12.9)	17 (15.7)	38 (30.9)	66 (20.9)	
Unspecified	7 (8.2)	11 (10.2)	41 (33.3)	59 (18.7)	
Mode of Arrival					< 0.001

‡Other included diving, horse-related incidents or not specified

*SEIFA – Socio-Economic Indexes for Areas (31)

at admission

Variable	Direct admit to SCIU (n = 85, 26.9%)	Indirect admit to SCIU (n = 108, 34.2%)	No acute admit to SCIU (n = 123, 38.9%)	Total (n = 316)	p- value
Ambulance	56 (65.9)	99 (91.7)	115 (93.5)	270 (85.4)	
Air ambulance	18 (21.2)	5 (4.6)	3 (2.4)	26 (8.2)	
Others	11 (12.9)	4 (3.7)	5 (4.1)	20 (6.3)	
‡Other included diving, horse-related incidents or not specified					
*SEIFA – Socio-Economic Indexes for Areas (31)					
# at admission					

Case rates across the state of NSW

GIS location of all cases were mapped, and crude rates calculated per 100,000 person-years (Fig. 1). Plotting case distribution per SA boundary SA3 was selected as the most stable representation.

Travel times from incident location

Around one quarter of study patients (n = 85, 26.9%) were directly admitted to SCIU from the injury scene, 170 (53.8%) patients went first to an MTS/RTS; leaving 61 patients (19.3%) who went first to local/district hospitals (Table 2). A further 108 (34.2%) were transferred indirectly to SCIU, from one of these other initial destinations, less than 60% transferred within 24 hours from time of injury (n = 64, 59.2%). Almost 30% (n = 93, 29.4%) of all patients remained in an MTS as their definitive hospital, 69 (74.2%) had been admitted directly from the injury scene. A further 11 (1.8%) were transferred in from an RTS and 13 (13.9%) from local/district hospitals.

Analysing the GIS data of injury incident locations relative to the proximity to hospital types, we found over half (53%, n = 168) of all patients were injured within 60 minutes road travel from an SCIU, yet only 28.6% of these (n = 48) were directly transferred to SCIU. Of the remaining patients injured within the 60-minute radius, less than half (48%, n = 76) were first transferred to an MTS; 22 (14%) first taken to an RTS. For forty percent (n = 125) of all patients the injury incident location was outside the 60 minutes travel radius of any MTS/SCIU, yet 20% went directly to an MTS and 20% to SCIU.

Table 2 shows GIS calculated summary of road travel distances and times calculated from injury incident locations to first hospitals, nearest SCIU and trauma centre by class of first hospital, analysing separately 26 (8.2%) patients who arrived by air-ambulance to their first hospital. For this group, Table 2 shows the actual times and distances ('direct' calculations) using hospital admission times and GIS data, also using GIS data to provide the 'on road' scenarios, had these patients travelled instead by road.

Table 2
Distances and times to first hospital from injury location GIS, by mode of transport

Hospital Class	First Hospital n (%)	Distance (km) to first hospital Median (IQR)	Time (mins) to first hospital Median (IQR)	Definitive Hospital n (%)	Time (hrs) to definitive hospital*** Median (IQR)
Travelled by road ambulance – GIS calculated					
SCIU	67 (23.1)	20.3 (72.7)	27.6 (51.6)	170 (58.6)	12.6 (39.9)
MTS	116 (40.0)	7.7 (9.9)	15.6 (11.4)	90 (31.0)	23.3 (43.6)
RTS	46 (15.8)	13.4 (29.5)	18.0 (23.1)	17 (5.9)	19.9 (44.4)
Local	61 (21.0)	8.7 (27)	12.4 (19.5)	13 (4.5)	2.9 (0)
Total	290	9.7 (20.0)	16 (17.8)	290	14.4 (42.9)
Travelled by air ambulance – direct distances/ recorded times					
SCIU	18 (69.2)	120 (141.9)	176 (279)	23 (88.5)	8.9 (5.2)
MTS	7 (26.9)	78.4 (34.8)	153 (33)	3 (11.5)	N/A
RTS	1 (3.8)	217.3 (0)	261 (0)	N/A	N/A
Travelled by air ambulance – if had travelled by road (not adjusted for 'on scene' time)					
SCIU	18 (69.2)	151.2 (184.7)	111.7 (102.9)	N/A	23 (88.5)
MTS	7 (26.9)	93.4 (58.8)	68.2 (47.5)	N/A	3 (11.5)
RTS	1 (3.8)	268.3 (0)	185.5 (0)	N/A	N/A
Total	26	122.8 (133.2)	168.5 (100.2)	N/A	26
** Ambulance pick up date/time to definitive hospital admission date/time for those transferred from first hospital					

Differences between the first and second last column above indicate movement between hospital classes, for patients transferred from first to definitive hospitals (n = 135, 42.7%). For this transferred group, the median (IQR) time from first to definitive hospital admission was 11.7(38.5) hours; the mean (SD) was 105.7(314.3) hours. Of 193 patients admitted definitively to SCIU, those arriving indirectly, took median (IQR) of 10.5 (38.3) hours, and a mean (SD) of 112.7 (344.4) hours.

For patients *not* taken directly to a SCIU or an MTS but instead taken first to a local/district hospital (n = 61), the additional median (IQR) difference in travel times they would have had to make to reach an MTS would have been 79.9 (218.9) minutes; the mean (SD) 133.4 (130) minutes. Almost half (n = 28, 46%) could have reached their closest MTS hospital by adding less than 30 minutes to their journey (median (IQR) 14.5(12.7) minutes).

Time to surgery

Almost two-thirds (n = 194, 61.4%) had surgery during acute care admissions; around two-thirds of these had surgery performed at SCIU (n = 131, 67.5%). Additionally, 56 (28.9%) received surgery at an MTS, 7 (3.6%) at an RTS.

Times from ambulance call to surgery were available for 98 (51%) patients; 47 (47.9%) of whom were transferred directly from injury scene to SCIU and had surgery within median (IQR) of 12.9 (7.9) hours. This was more rapid than for patients transferred indirectly to SCIU (from MTS/RTS/local hospitals), who had surgery within median (IQR) of 19.5 (18.9) hours. Patients transferred directly from injury location to MTS and who surgery there, were operated within a median (IQR) of 21.8 (115.2) hours.

Comparing times to admission to hospitals where operations took place for the surgery group of 194 patients, Table 3 compares times to surgical hospital admission for the 98 for whom operation times were available with times to surgical hospital admission for the other 96 patients, as well as times to surgery where available. Inter-hospital transfers lengthened times to surgery.

Table 3
First hospital, location of surgery and timing to surgery (median [IQR]) post injury

Hospital Class	As First hospital n (%)	As Surgical Hospital n (%)	Time to surgery median (IQR) (by surgical hospital class)	Time (hrs) to surgical hospital admission* median (IQR)
Time to surgery available (n = 98)				
SCIU	47 (47.9)	90 (91.8)	15.2 (10.9)	6.7 (7.3)
MTS	31 (31.6)	7 (7.1)	21.8 (23.3)	1.4 (0.5)
RTS	8 (8.2)	1 (1.0)	12.7 (0)	1.4 (0)
Local	12 (12.2)	0	NA	NA
Total	98	98	15.2 (11.2)	6.1 (7.2)
Time to surgery not available (n = 96)				
SCIU	18 (18.7)	41 (42.7)	NA	8.1 (9.4)
MTS	45 (46.9)	49 (51.0)	NA	1.4 (8.5)
RTS	20 (20.8)	6 (6.3)	NA	1.0 (0.8)
Local	13 (13.5)	0	NA	0
Total	194	194	NA	3.9 (10.2)
*from ambulance pick up time				

Factors influencing triage pattern to SCIU

The multinomial logistic regression model (Table 4) displays the RRR of various exposure variables to triage patterns; with direct SCIU admission as the base outcome category to no SCIU admission, then indirect SCIU admission. Where the nearest SCIU was within 60 minutes road travel from the injury incident the likelihood of no SCIU admission was significantly reduced (RRR 0.281, $p = 0.003$, 95% CI 0.123–0.646). The relative risk of indirect SCIU transfer was somewhat lower (RRR 0.511) than no SCIU. Patients with two or more concurrent traumatic injuries were 5.2 times more likely to experience indirect (rather than direct) transfer to SCIU (RRR 5.18, $p = 0.002$). Patients with lower injury severity (ICISS 0.95–1.00) were around 10 times more likely to have no admission to SCIU than direct transfer compared to patients with higher injury severity (ICISS < 0.7), (RRR 10.1, $p = 0.03$, 95% CI 1.194–85.885). Patients with complete or incomplete TSCI were significantly less likely to experience no SCIU admission in relation to patients without any completeness of injury coded. Each year older for patients with acute TSCI increased the likelihood of not being admitted to a SCIU at all in the acute admission by 2% per year ($p = 0.032$). Table 4 displays the regression estimates for the entire cohort.

Table 4

Multinomial regression model – relative risk of exposure variables comparing Indirect and No transfer to SCIU, with reference of direct admission to SCIU

Triage Patterns	RRR*	Std. Err	P-value	95% Confidence Interval
Direct SCIU	(as base outcome for No SCIU)			
No SCIU				
age (years)	1.02	0.01	0.034	1.002–1.039
SCIU > 60 min	0.281	0.119	0.003	0.123–0.646
Female sex	0.959	0.44	0.928	0.391–2.355
Injury Mechanism				
Other (reference)				
Falls	0.629	0.343	0.388	0.21–1.83
Transport	0.559	0.343	0.344	0.17–1.86
Self-harm /Assault	11.449	16.289	0.087	0.70–186.13
Injury level				
Cervical (reference)				
Thoracic	1.838	0.983	0.255	0.64–5.24
Lumbar	3.917	5.142	0.298	0.29–51.33
Extent of spinal cord Injury				
Not coded (reference)				
Complete	0.061	0.044	< 0.001	0.015–0.25
Incomplete	0.288	0.159	0.024	0.09–0.85
Conus/cauda-equina	0.108	0.136	0.077	0.01–1.27
Arrival Mode				
Missing (reference)				
Road Ambulance	8.139	6.273	0.007	1.79–36.86
Air ambulance	0.61	0.606	0.619	0.09–4.27
ICISS Score				
< 0.7 (base)				

*Relative Risk Ratios

Triage Patterns	RRR*	Std. Err	P-value	95% Confidence Interval
0.7 to < 0.83	0.696	0.471	0.592	0.18–2.62
0.83 to < 0.89	1.558	1.154	0.55	0.36–6.66
0.89 to < 0.95	2.275	1.929	0.332	0.43–11.99
0.95 to 1.00	10.125	11.045	0.034	1.19–85.88
Multiple trauma				
0 (base)				
One additional injury	0.35	0.158	0.02	0.15–0.85
Two or more	0.942	0.503	0.911	0.33–2.68
Direct SCIU	(as base outcome for Indirect SCIU)			
Indirect SCIU				
Age (years)	0.998	0.01	0.84	0.98–1.02
SCIU > 60 min	0.511	0.22	0.11	0.22–1.17
Female sex	0.96	0.45	0.93	0.38–2.43
Injury Mechanism				
Other (reference)				
Falls	0.51	0.28	0.22	0.17–1.49
Transport	0.56	0.34	0.34	0.17–1.84
Self-harm/Assault	9.02	12.77	0.12	0.56–144.74
Injury level				
Cervical (reference)				
Thoracic	0.61	0.32	0.35	0.22–1.69
Lumbar	0.23	0.27	0.22	0.02–2.48
Extent of Injury				
Missing (reference)				
Complete	1.774	1.21	0.40	0.46–6.77
Incomplete	1.625	1.03	0.44	0.47–5.62
Conus/cauda-equina	0.502	0.59	0.56	0.05–5.01
*Relative Risk Ratios				

Triage Patterns	RRR*	Std. Err	P-value	95% Confidence Interval
Arrival Mode				
Other (reference)				
Road ambulance	20.148	18.13	0.001	3.45–117.52
Air ambulance	1.768	1.84	0.58	0.23–13.55
ICISS Score				
< 0.7 (reference)				
0.7 to < 0.83	2.04	1.52	0.34	0.47–8.79
0.83 to < 0.89	9.98	8.14	0.005	2.02–49.38
0.89 to < 0.95	18.69	17.09	0.001	3.11–112.25
0.95 to 1.00	33.87	39.64	< 0.001	24.14–58.76
Multiple trauma				
0 (reference)				
One additional injury	0.65	0.29	0.35	0.26–1.59
Two or more	5.18	2.77	0.002	1.82–14.75
*Relative Risk Ratios				

Triage patterns influencing complication risk

The potential impact of triage patterns on the risk of experiencing inpatient complications for patients with acute TSCI was assessed and is shown in Table 5. Patients experiencing indirect transfer to SCIU had an increased risk of developing inpatient hospital complications compared with patients transferred directly to SCIU. Complete TSCI, additional comorbidities (CCI) and older age also increased the risks of developing inpatient complications.

Table 5
Triage pattern impact on risk of developing a medical complication as an inpatient

Any complication*	Odds Ratio	Std.Err	P-value	95% Conf. Interval
Triage Patterns				
Direct SCIU	1.392	0.492	0.35	0.696–2.783
Indirect SCIU	3.243	1.113	0.001	1.656–6.353
No SCIU (reference)				
Demographic				
Female sex	1.122	0.357	0.718	0.601–2.095
Age (years)	1.011	0.006	0.050	0.999–1.023
Extent of TSCI				
Complete	6.917	3.672	< 0.001	2.443–19.579
Incomplete	1.447	0.536	0.319	0.7-2.989
Conus/cauda-equina	1.57	0.685	0.301	0.668–3.692
Missing (reference)				
Charlson Index				
No comorbid disease (ref)				
1 additional	1.468	0.57	0.323	0.686–3.143
2 + additional	3.268	1.302	0.003	1.497–7.135
Multiple Trauma				
No additional injuries (ref)				
1 additional injury	1.818	0.609	0.075	0.942–3.506
2 + additional injuries	1.15	0.367	0.661	0.615–2.149
Injury Severity (ICISS)				
0.7 to < 0.83	0.222	0.107	0.002	0.086–0.572
0.83 to < 0.89	0.085	0.043	< 0.001	0.032–0.23
0.89 to < 0.95	0.139	0.07	< 0.001	0.051–0.374
0.95 to 1.00	0.113	0.063	< 0.001	0.038–0.335
*Includes pressure injuries, urinary tract infections/complications, respiratory infections/complications, deep vein thrombosis and pulmonary embolus				

Discussion

This study investigated the impact of geospatial variables on access to specialist care for patients with acute TSCI across NSW, and revealed that despite over half of the patients being injured within a 60-minute road travel time of a SCIU, less than 30% of them experienced a direct SCIU admission. This is poor by international comparison; Cheng et al (15) in Canada demonstrating 77% of patients with acute TSCI and within 40 km of a SCIU admitted directly to a specialist unit. In this study, patients transferred directly to a SCIU underwent earlier surgery (median (IQR) of 7.3 (6; 19) hours) than patients with indirect admission to SCIU, who waited 10 hours longer to have their surgery (median (IQR) 17.5 (11; 31) hours). For many patients, indirect admission places them outside the recommended time window for surgery of ≤ 24 hours after injury, and at greater risk for medical complications and compromised recovery (32).

As specialist health services, the SCIUs in NSW are located in areas of high population density, however, serve large geographical areas. Increased travel time between the incident location and SCIU reduces the likelihood of direct transfer, indicating lack of equitable access to specialist care from all areas of NSW. The fact that 70% of patients who were injured within 60 minutes of road travel to a SCIU did not experience direct transfer, suggests variations in clinical practice that leads to inequity in patient treatment. Further, almost one-fifth of patients were taken first to non-trauma designated hospitals ($n = 61, 19.3\%$), consistent with previous research showing that despite recommended pathways for patients with TSCI in NSW (25) some acute trauma patients are still being triaged to non-trauma services (19).

Aeromedical retrieval offers reduced travel times for the regional or very remote patient, demonstrated by these findings, however, there are resource restraints to this mode of transport. Scenario modelling could demonstrate the impact of changing transport modes to patients for whom distances by road are impediments to timely acute care. The additional resource required to retrieve patients with acute TSCI aeromedically needs to be justified by comprehensive economic evaluation. The cost per mission was estimated in 2011 at between \$9,300 and \$19,000 (33); the cost benefit of such retrieval outlay over the long term costs of a TSCI has not been sufficiently explored in Australia.

Patients were less likely to experience direct transfer to SCIU when they had less severe spinal cord impairment, such as conus and cauda equina injury. This may be appropriate, however, increasing age also reduced the likelihood of direct transfer; and older patients are known to have higher risks of hospital complications and poorer outcomes with delayed intervention (8). As surgical intervention may not have been appropriate in some older patients, we are unable to ascertain if there had been a discussion with the SCIU regarding the appropriateness of transfer to the Specialist Unit. It would seem likely that the combination of an incomplete spinal cord injury and increasing age represents the central cord syndrome group (although this was not specifically identified in this study). This group typically present after minor falls and may not be immediately recognised as TSCI(34), so may have missed being treated according to the transfer guidelines.

Limitations

This study had several limitations. As we required geographic variables, we were unable to include the complete cohort of patients with acute incident TSCI identified across the study in this analysis. Therefore, a selection bias was possible, given omission of certain patients. However, patients not included were found to have similar characteristics across relevant variables (age, injury mechanism and completeness of injury), this was therefore not deemed significantly impactful on our findings.

Conclusion

This study used geospatial methods to analyse pathways of care for patients with acute TSCI in NSW, who received care in a SCIU by either direct or indirect admission and has highlighted specific areas for optimisation on a health system level. Our finding that only 30% of patients who sustained injuries within a 60-minute travel radius to a SCIU were directly admitted, is significantly lower than international comparisons and advocates improvement in early transport pathways to TSCI care.

Regardless of injury incident location, it is evident that there are transport modes and decisions can achieve early and direct SCIU admission with timely surgery. These findings complement previous research showing cost savings and reduced secondary complications achievable by streamlining care pathways(16) for patients with TSCI. Evidence informed policy in health service optimisations can provide timely and definitive surgical treatment for those patients suffering an acute TSCI, leading to improved neurological outcomes and reduced secondary complications and mortality.

List Of Abbreviations

APDC Admitted Patient Data Collection

CCI Charlson comorbidity index

GIS Geographic information system

ICD10-AM International Classification of Diseases, 10th version, Australian

Modification

ICISS International Classification Injury Severity Score

MTS Major trauma service

NSW New South Wales

RRR Relative Risk Ratios

RTS Regional trauma service

SA Statistical area

SCIU Spinal cord injury unit

SSCIS State Spinal Cord Injury Service

TSCI Traumatic spinal cord injury

TBI Traumatic brain injury

Declarations

Ethics approval and consent to participate. The NSW Population Research Ethics Committee (2012/09/420) approved this project.

Consent for publication. Not applicable as this project used de-identified administrative data.

Availability of data and materials. The data that support the findings of this study are available from the Centre for Health Record Linkage, New South Wales, Australia, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available.

Competing interests. No authors declare any competing interests.

Funding. LNS was funded by icare™ for the duration of this research. BPV was funded on a doctoral scholarship for the duration of this research. The funder critically reviewed and approved the study protocol developed for this research.

Authors' contributions. LNS conceived of this project and analysed and interpreted the patient data regarding the health services modelling. SRW analysed the data pertaining to geospatial mapping. DW and MK produced the map. All authors read and approved the final manuscript.

Acknowledgements. The authors acknowledge the statistical advice provided by www.theanalysisfactor.com, and thank Professor Roderick McClure for his recommendations on the study reporting.

References

1. Atkins, D, Noonan, V, Santos, A, Lewis, R, Fehlings, M, Burns, A, et al. Secondary Complications in SCI Across the Continuum: Using Operations Research to Predict the Impact and Optimize Management Strategies. Topics in Spinal Cord Rehabilitation. 2012;18(1):57-66.
2. Craig, A, Guest, R, Tran, Y, Middleton, J. Depressive mood in adults with spinal cord injury as they transition from an inpatient to a community setting: secondary analyses from a clinical trial. Spinal Cord. 2017;55:926-34.

3. Craig, A, Nicholson Perry, K, Guest, R, Tran, Y, Dezarnaulds, A, Hales, A, et al. The occurrence of psychological disorders and co-morbidities following spinal cord injury. *Arch Phys Med and Rehab*. 2015;96:1426-34.
4. Access Economics Pty Ltd. The economic cost of spinal cord injury and traumatic brain injury in Australia. . 2009:1-116.
5. Munce, S, Wodchis, W, Guilcher, S, Couris, C, Verrier, M, Fung, K, et al. Direct costs of adult traumatic spinal cord injury in ontario. *Spinal Cord*. 2013;51:64-9.
6. Consortium for Spinal Cord Medicine. Early acute management in adults with spinal cord injury: a clinical practice guideline for healthcare professionals. *Journal of Spinal Cord Medicine*. 2008;31(4):403-79.
7. Mac-Thiong, J, Feldman, D, Thompson, C, Bourassa-Moreau, E, Parent, S. Does Timing of Surgery Affect Hospitalization Costs and Length of Stay for Acute Care following a Traumatic Spinal Cord Injury? *Journal of Neurotrauma*. 2012;29:2816–22.
8. Parent, S, Barchi, S, LeBreton, M, Casha, S, Fehlings, M. The impact of specialized centers of care for spinal cord injury on length of stay, complications, and mortality: a systematic review of the literature. *Journal of Neurotrauma*. 2011;28:1363–70.
9. Sharwood, L, Boufous, S, Muecke, S, Middleton, J. Health Service Pathways Analysis as Evidence Base for Trauma Policy Change *Emerg Med*. 2017;7(3).
10. NSW Agency for Clinical Innovation. Evidence and utilisation of spinal cord injury services NSW: Evidence Series. 2020;SHPN (ACI) 190018(ISBN 978-1-76081-068-9).
11. Vulliamy, P, Faulkner, M, Kirkwood, G, West, A, O'Neill, B, Griffiths, MP, et al. Temporal and geographic patterns of stab injuries in young people: a retrospective cohort study from a UK major trauma centre. *BMJ Open*. 2018;8(10):e023114.
12. Brown, JB, Kheng, M, Carney, NA, Rubiano, AM, Puyana, JC. Geographical Disparity and Traumatic Brain Injury in America: Rural Areas Suffer Poorer Outcomes. *Journal of neurosciences in rural practice*. 2019;10(1):10-5.
13. Auchincloss, AH, Gebreab, SY, Mair, C, Diez Roux, AV. A review of spatial methods in epidemiology, 2000-2010. *Annual review of public health*. 2012;33:107-22.
14. Smedley, WA, Stone, KL, Brown, A, Farley, P, Griffin, RL, Cox, DB, et al. Use of helicopters for retrieval of trauma patients: A geospatial analysis. *J Trauma Acute Care Surg*. 2019;87(1):168-72.
15. Cheng, CL, Noonan, VK, Shurgold, J, Chen, J, Rivers, CS, Hamedani, HK, et al. Geomapping of Traumatic Spinal Cord Injury in Canada and Factors Related to Triage Pattern. *J Neurotrauma*. 2017.

16. Vaikuntam, B, Middleton, J, McElduff, P, Pearse, J, Walsh, J, Cameron, I, et al. Identifying Predictors of Higher Acute Care Costs for Patients With Traumatic Spinal Cord Injury and Modeling Acute Care Pathway Redesign. . Spine (Phila Pa 1976). 2019;4 Mar 2019.
17. Gomez, D, Sarrami, P, Singh, H, Balogh, Z, Dinh, M, Hsu, J. External benchmarking of trauma services in New South Wales: Risk-adjusted mortality after moderate to severe injury from 2012 to 2016. Injury. 2018.
18. Brown, JB, Stassen, NA, Cheng, JD, Sangosanya, AT, Bankey, PE, Gestring, ML. Trauma center designation correlates with functional independence after severe but not moderate traumatic brain injury. J Trauma. 2010;69(2):263-9.
19. Doumouras, AG, Haas, B, Gomez, D, de Mestral, C, Boyes, DM, Morrison, LJ, et al. The impact of distance on triage to trauma center care in an urban trauma system. Prehosp Emerg Care. 2012;16(4):456-62.
20. Goltsman, D, Li, Z, Bruce, E, Maitz, PKM. Geospatial and epidemiological analysis of severe burns in New South Wales by residential postcodes. Burns. 2014;40(4):670-82.
21. Poulos, RG, Chong, SSS, Olivier, J, Jalaludin, B. Geospatial analyses to prioritize public health interventions: a case study of pedestrian and pedal cycle injuries in New South Wales, Australia. International Journal of Public Health. 2012;57(3):467-75.
22. Australian Bureau of Statistics. Australian Demographic Statistics. 2016:Canberra:Australian Bureau of Statistics
23. Lundy, DW, Harvey, EJ, Jahangir, AA, Leighton, RK. Trauma systems in North America. OTA International. 2019;2(S1).
24. Gomez, D, Larsen, K, Burns, BJ, Dinh, M, Hsu, J. Optimizing access and configuration of trauma centre care in New South Wales. Injury. 2019.
25. NSW Health. NSW Critical Care Tertiary Referral Networks and Transfer of Care (ADULTS). Policy Directive. 2018;PD2018_011(Agency for Clinical Innovation):1-37.
26. Australian Consortium for Classification Development. The international statistical classification of diseases and related health problems, tenth revision, Australian modification (ICD-10-AM/ACHI/ACS) (Tenth ed.). 2017:Darlinghurst, NSW:Independent Hospital Pricing Authority.
27. Vaikuntam, BP, Middleton, JW, McElduff, P, Pearse, J, Walsh, J, Cameron, ID, et al. Assessing the impact of care pathways on potentially preventable complications and costs for spinal trauma patients: a data linkage study BMJ Open. 2018;8(11):e023785.

28. Kwak, C, Clayton-Matthews, A. Multinomial Logistic Regression. *Nursing Research*. 2002;51(6):404-10.
29. Stata Statistical Software. 2017;Version 15.1 IC
30. Gedeberg, R, Warner, M, Chen, LH, Gulliver, P, Cryer, C, Robitaille, Y, et al. Internationally comparable diagnosis-specific survival probabilities for calculation of the ICD-10-based Injury Severity Score. *J Trauma Acute Care Surg*. 2014;76(2):358-65.
31. Australian Bureau of Statistics. Technical Paper - Socio-Economic Indexes for Areas (SEIFA) 2018;2033.0.55.001
32. Dvorak, MF, Noonan, VK, Fallah, N, Fisher, CG, Finkelstein, J, Kwon, BK, et al. The influence of time from injury to surgery on motor recovery and length of hospital stay in acute traumatic spinal cord injury: an observational Canadian cohort study. *J Neurotrauma*. 2015;32(9):645-54.
33. Taylor, CB, Stevenson, M, Jan, S, Liu, B, Tall, G, Middleton, PM, et al. An investigation into the cost, coverage and activities of Helicopter Emergency Medical Services in the state of New South Wales, Australia. *Injury*. 2011;42(10):1088-94.
34. Middleton, P, Davies, S, Anand, S, Marial, O, Reinten-Reynolds, T, Middleton, J. The prehospital epidemiology and management of spinal cord injuries in NSW: 2004-2008. *Injury*. 2012;43:480-5.

Figures

Figure 1

Rate per 100,000 person year 0.2 – 0.8 0.9 – 1.2 1.3 – 1.7 1.8 – 2.4 2.5 – 5.1