3-hydroxykynurenine as a Potential Ligand for Hsp70 Proteins and its Effects on Drosophila Memory After Heat Shock

Aleksandr V. Zhuravlev (✉ beneor@mail.ru)

Pavlov Institute of Physiology RAS: Institut fiziologii imeni I P Pavlova RAN <https://orcid.org/0000-0003-2673-4283>

Boris F. Shchegolev

Pavlov Institute of Physiology RAS: Institut fiziologii imeni I P Pavlova RAN

Gennadii A. Zakharov

Pavlov Institute of Physiology RAS: Institut fiziologii imeni I P Pavlova RAN

Polina N. Ivanova

Pavlov Institute of Physiology RAS: Institut fiziologii imeni I P Pavlova RAN

Ekaterina A. Nikitina

Pavlov Institute of Physiology RAS: Institut fiziologii imeni I P Pavlova RAN

Elena V. Savvateeva-Popova

Pavlov Institute of Physiology RAS: Institut fiziologii imeni I P Pavlova RAN

Research Article

Keywords: Drosophila, cardinal, 3-hydroxykynurenine, heat shock, Hsp70, short-term memory

Posted Date: August 9th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-787746/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Molecular Neurobiology on January 14th, 2022. See the published version at <https://doi.org/10.1007/s12035-021-02704-3>.

Abstract

Kynurenine products of tryptophan metabolism are modifiers of the nervous activity and oxidative processes in mammals and invertebrates. 3-hydroxykynurenine (3HOK) in moderate concentrations is a lipid peroxidation inhibitor. However, its accumulation and oxidative auto dimerization lead to oxidative stress development manifested in age-related neurodegenerative diseases (NDDs) and neurologic disorders provoked by acute stress. Different forms of stress, the mostly studied being heat shock response, rely on functioning of heat shock proteins of Hsp70 superfamily. Since kynurenines are called “kids of stress”, we performed computational estimation of affinity of 3HOK and other kynurenines binding to predicted ATP site of *Drosophila* Hsp70 cognate 71 protein (Dhsp71) using Autodock Vina. The binding energy of 3HOK dimer is -9.4 kcal/mol, its orientation within the active site is close to that of ATP. This might be a new mechanism of producing a competitive inhibitor of Hsp70 chaperones that decreases organism ability to adapt to heat shock. We also showed that the *Drosophila melanogaster cardinal* (*cd¹*) mutant with 3HOK excess, serving as a model for Huntington's disease (HD), manifests severe defects of short-term memory after heat shock applied either in adults, or at the prepupal stage.

Introduction

Kynurenine pathway (KP) is a main route of tryptophan catabolism in different animal species and phyla. KP intermediates, collectively called kynurenines, possess a broad spectrum of neurotropic activities [1, 2]. The pivotal role of kynurenines is widely discussed in scholarly literature in connection to age-dependent neurodegenerative diseases (NDDs) manifested as progressive memory loss and neuroinflammation [3, 4].

The major prerequisite to NDDs development is the concentration ratio of different kynurenines. L-kynurenine (KYN), 3-hydroxykynurenine (3HOK) and quinolinic acid (QUIN) exert either neuroprotective, or neurodegenerative effects. As to 3HOK, its high concentration is a critical factor for development of many NDDs. In moderate concentrations, 3HOK and 3-hydroxyanthranilic acid (3HAA) are inhibitors of lipid peroxidation that can ameliorate the toxic action of reactive oxygen species (ROS) [5, 6]. However, 3HOK accumulation and autodimerization lead to oxidative stress development, as manifested in NDDs and under acute stress [7-9]. 3HOK and QUIN generate hydrogen peroxide, which induces hyperproduction of toxic free radicals within a cell [10]. On the contrary, kynurenic acid (KYNA) is an endogenous protector against neurotoxicity [11, 12].

Contrary to vertebrates, the terminal stage of insect KP is 3HOK dimerization to brown eye pigment xanthommatin (XAN) [13]. The activity of phenoxazinone synthase (PHS) catalyzing transamination of 3HOK to xanthurenic acid is decreased in *Drosophila cardinal* (*cd¹*) mutant [14]. The 2.9-fold rise in the free 3HOK level is pertinent to this strain [15]. In *cd¹*, XAN is formed non-enzymatically due to 3HOK oxidative autodimerization. Total antioxidant capacity in *cd¹* is lower than in the wild-type strain *Canton-S* (CS) [16].

In mammals, KP is induced under stress conditions [2]. The key enzyme of hepatic KP tryptophan-2,3-dioxygenase is activated by corticosterone leading to kynurenines rise and depressive behavior [17]. The brain-specific indole-2,3-dioxygenase is activated by proinflammatory cytokines [18]. Different types of stress induce the universal response originally described in *Drosophila* – massive production of heat shock proteins (HSPs), the stress response proteins having chaperone properties and involved in adaptation [19, 20].

The main diagnostic manifestation of human NDDs is the progressive memory loss. This is the first trait evident in humans not only to physicians, but also to relatives. This presents an ever-growing home and hospital problem. Problem solving lies in dissecting the triangle: learning/memory – HSPs – kynurenines.

Provided that 3HOK is an oxidative stress activator [7], it is reasonable to assess the age-dependent dynamics of learning acquisition and memory retention (learning/memory) in *Drosophila* model, using KP mutants. Conditioned courtship suppression paradigm (CCSP) or a “rejection” paradigm is rather convenient for such an analysis in fly [21-25]. The essence is that naïve *Drosophila* male that has been repeatedly rejected by mated female either during 30 min training for testing STM, or for 5 hrs for testing long term memory (LTM) stop to court another mated female [22].

STM decay, as well as age-dependent memory loss, synaptic pathology and altered brain plasticity were registered beginning from the 13th day of adult life in *cd¹* (high 3HOK/KYNA index), but not in *cn* mutant strain with KYNA excess (low 3HOK/KYNA index) or in the wild-type *Canton-S* [26]. Stress treatment of young flies (1 h exposure to heat shock 37 °C) did not lead to apoptosis only in *cn*, indicating the neuroprotective role of KYNA, but induced it in normal and more intensively in *cd¹* males. Moreover, HS exacerbated the toxic 3HOK action on *Drosophila* causing courtship song defects [27]. Spontaneous locomotor activity is suppressed in aged 40 day-old *cd¹* males [28]. High 3HOK/KYNA index is also a crucial factor for neurodegeneration in *htt*, a *Drosophila* model for Huntington's disease (HD) [29, 30].

The most recent disclosure of HSP and memory link is that of [31]. It proves the assumption that memory processes and the ancient stress–response systems are closely intermingled: any reinforcement used in different learning paradigms, either punishment or reward, utilizes stress reactions. While using *Drosophila* strains with different copy numbers of chaperone Hsp70, in particular strains having a deletion of all six *hsp70* genes, the authors have demonstrated that a low constitutive level of Hsp70 pertinent to *Drosophila* is sufficient for learning and formation of STM and LTM. Moreover, they have revealed major pathways participating in memory formation and consolidation that depend on the presence of *hsp70* in the genome. The roles of heat shock transcription factor 1 (HSF1) and Hsp70 in memory formation and neuroprotection were established [32].

If so, can kynurenines, the “kids of stress” [2], directly affect HSPs functions via binding to any of their important sites? Possibly, the activity of Hsp70 might be suppressed in *cd¹*, making it vulnerable to HS. Therefore, *cd¹* STM not impaired until day 12 under normal conditions might suffer after HS already in

young flies at day 5. To estimate the affinity of kynurenines binding to Hsp70, we performed a computer docking of KYN and its derivatives to the model structure of Dhsp71, a homologue of *Escherichia coli* DNAK and mammalian Hsp70. Also, we used CCSP to reveal putative effects of HS on learning acquisition and memory retention in *cd¹* which might be caused by 3HOK-dependent inactivation of Hsp70.

Material And Methods

Protein modeling and computer docking

Hsp70 structure is characterized by the N-end ATP binding domain where ATP hydrolysis occurs, intermediate linker, and C-end with substrate binding domain [33]. Experimental protein structure of *E. coli* Hsp70 (DNAK) in complex with ATP and Mg^{2+} was taken from RCSB Protein Data Bank (ID: 4jn4_A). *D. melanogaster* ATP-binding domain of Hsp70 cognate 71 protein (Dhsp71) was constructed with the help of MODELLER software [34] using Hsp70 cognate 71 protein sequence (GenBank ID: AAA28628.1) and 4jn4_A in complex with ATP as a 3D template. The model with the highest negative potential energy was selected for docking. The optimal ligands conformations, including that of 3HOK dimer D2', were published in [16].

Flexible ligand and rigid protein structures were prepared for docking using AutoDockTools4 [35]. Gasteiger charges were used; Mg charge in the complex with ATP was set as 0, so that ATP-Mg complex has a total charge of -2 [36]. Docking was performed using Autodock Vina [37] with the following parameters: grid box size: 40 Å, centered on ligand (the experimental ATP position); maximum number of binding modes: 100; exhaustiveness of the global search: 20; other parameters – default. The structures with the highest negative free energies of binding (ΔG) were selected, except for 3HOK where two different ligand orientations were considered (see Results section). The dissociation constants (K_d) were calculated as follows: $K_d = \exp(\Delta G/RT)$, ΔG being expressed in J/mol, $T = 298.15$ K. All figures were prepared using VMD [38].

The structures of Dhsp71 and the docked ligands can be found in Supplementary Data, S1 Docking_coordinates file.

Drosophila strains

Fly strains were taken from the Research Center “Biocollection of Pavlov Institute of Physiology RAS for the study of integrative mechanisms of the nervous and visceral systems”.

The following strains were used:

1. *Canton-S* (*CS*), the wild-type strain.
2. *cardinal* (*cd¹*) with a mutation in phenoxazinone synthase gene leading to 3HOK increase. *cd¹* was out-crossed to *CS* for 20 generations.

Flies were raised on standard yeast–raisin medium with 8 a.m. – 8 p.m. daily illumination cycle at 25±0.5 °C.

Heat shock treatment

Heat shock (HS) treatment was applied at one of the following stages: 1. HS1 – 1st instar larval stage, the mushroom bodies development; 2. HS2 – prepupal stage, the central complex development; 3. HS3 – 5-day-old adult males [39]. Flies were subjected to HS in empty glass vials in a water bath for 30 min at 37 °C. 1 h after HS3 before the experiment flies had a rest in food-containing vials.

Learning and memory

Learning and 3 h STM was assayed in *Drosophila* males in CCSP using retraining test with a fertilized female [22]. Males were kept individually from eclosion until the experiment. Virgin *CS* females were kept in small groups. Before the experiment, *CS* females were fertilized overnight. All experiments were performed in 5 day-old males and females at 25°C in the first half of the day.

For training, a naive male (without courtship experience) was placed together with a fertilized *CS* female in a Plexiglas experimental chamber (15 mm in diameter, 5 mm high) for 30 min. The retraining test was performed immediately after training (for learning acquisition) or 3 h after training (for STM retention). A trained male was tested in a fresh chamber with another fertilized female. After 45 sec of adaptation, the male courtship behavioral elements were recorded for 300 sec using a specially designed program [22]. The following courtship elements were scored: orientation/pursuit, wing vibration, licking and attempted copulation. The non-courtship elements were activity (running), preening and rest. For each group (naive control, learning acquisition, STM), 20 males were tested.

The courtship index (CI), a percentage of time spent in courtship over a 300 sec period [40] was calculated for each male. The learning index (LI) was calculated as follows:

$$LI = ([CI_{na} - CI_{tr}] / CI_{na}) \times 100 = (1 - CI_{tr} / CI_{na}) \times 100 [41],$$

where CI_{na} and CI_{tr} are the mean courtship indices for independent samples of naive and trained males, respectively.

Statistical analysis was performed using a two-sided randomization test [42]. 10,000 permutations were used. The null-hypothesis was rejected at $\alpha R < 0.05$.

Results

Computer docking

ATP docking to D_{NAK} yields a ligand orientation nearly matching to that in the experimental D_{NAK} – ATP complex: the root-mean-square deviation (RMSD) is 0.256 Å for non-hydrogen atoms. Approximately the

same orientation was obtained for Dhsp71 (RMSD 0.502 Å). ATP binding sites are described in Table 1. Mg-triphosphate group is localized above the salt bridge formed by the amino group of Lys 70 and carboxylic group of Glu 171/172. The oxygen atoms of triphosphate group contact Thr 11 – Asn 13 and Ala/Thr 199, while one negatively charged oxygen forms an ionic bond with Lys 70. The ribose group forms hydrogen bonds with Lys 270/266 and Glu 267/263. The adenine group is surrounded by several non-polar groups, such as the hydrophobic part of Arg 345/341 and Ile 271/Cys 267, and also forms a hydrogen bond with Ser 274/270.

Table 1. ATP sites of *E. coli* and *D. melanogaster* HSP70 proteins

DNAK	Dhsp71	Ligand-protein bonds
Thr 11	Thr 11	NH.. O1G(3P): HB
Thr 12	Thr 12	NH.. O2B(3P): HB
Asn 13	Asn 13	NH.. O2B(3P), HD22.. OA1(3P): 2 HB
Lys 70	Lys 70	NH ₃ .. O1G(3P): IB
Ala199	<i>Thr 199</i>	NH.. O2G(3P): HB
Glu 267	Glu 263	COO.. H2'O2'(Rib): HB
Lys 270	Lys 266	NH ₃ .. O2'(Rib): HB, Lys.. Ade: 2 NP
Ile 271	<i>Cys 267</i>	Ile/Cys.. (Ade): NP
Ser 274	Ser 270	OH.. N1(Ade): HB
Gly 342	Gly 338	NH.. O2A(Rib), O5'(Rib): 2 HB
Gln 343	<i>Met 339</i>	Gln/Met.. (Rib): NP
Arg 345	Arg 341	Arg.. (Ade): NP
Mg 1001	Mg 604	Mg.. O1B, O3G (3P): 2 IB
Outside ATP site		
Glu 171	Glu 172	COO.. NH ₃ (3HOK): IB
Asp 194	Asp 194	COO.. NH ₃ (3HOK): IB
Pro 367	Pro 363	CO.. NH ₂ ar (3HOK): HB
Asp 368	Asp 364	COO.. OHar (3HOK): HB

The residues in Dhsp71 that differ from DNAK are marked in italics. Some residues beyond the ATP site important for 3HOK binding are shown. Ade – adenine group, ar – the side chain of 3HOK aromatic group, 3P – triphosphate group, Rib – ribose group. Bond is described as follows: active site residue atom group. ligand atom group (ligand part): the number and type of bonds. HB – hydrogen bond, IB – ionic bond, NP – non-polar (van der Waals or hydrophobic) bond.

The binding energy predicted by Autodock Vina is -12.3 kcal/mol for DNAK and -11.7 kcal/mol for Dhsp71 (Table 2). The reported experimental constant of ATP binding to Hsp70 is 4.4E-06 M [43] or 7.3E-06 M (for DNAK) [44]. This is a significantly lower affinity compared to predicted Kd values, hence docking seems to overestimate the ATP binding energy. The somewhat lower absolute computational ATP binding energy for Dhsp71 can be explained by a lower quality of the model compared to the experimental structure. Some minor differences between the active sites also can affect ATP binding, such as change of polar Gln 343 to hydrophobic Met 339 contacting the adenine group.

Table 2. Free energies and dissociation constants of kynurenines docking to Hsp70 proteins

	DNAK		Dhsp71	
Ligand	ΔG (kcal/mol)	Kd (M)	ΔG (kcal/mol)	Kd (M)
ATP-Mg	-12.3	9.6E-10	-11.7	2.6E-09
3HOK	-7.8	1.9E-06	-7.8/ -7.6*	1.9E-06/ 2.7E-06*
L-3HOKzi-D2'	-7.7	2.3E-06	-9.4	1.3E-07
L-KYN	-7.3	4.4E-06	-7.6	2.7E-06
3HAA	-6.9	8.7E-06	-7.1	6.2E-06
KYNA	-8.1	1.1E-06	-8.7	4.1E-07
QUIN	-7.1	6.2E-06	-7.3	4.4E-07

Ligands description: ATP-Mg – ATP the complex with Mg, 3HOK – 3-hydroxykynurenine, L-3HOKzi_D2' – dimer of 3HOK (D2') in zwitterionic form, 3HAA – 3-hydroxyanthranilic acid, L-KYN – L-kynurenine, KYNA – kynurenic acid, QUIN – quinolinic acid. The values for the second optimal 3HOK – Dhsp71 complex are show by asterisk.

For kynurenines, the absolute binding energy is significantly lower (6.9 – 8.7 kcal/mol). 3HAA has the lowest affinity for DNAK and Dhsp71. KYNA that brings a negatively charged group forming an ionic bond with Lys 70 has the highest affinity for both proteins (see S1 Table for the full description of ligand-receptor interactions). The affinity of L-KYN and 3HOK for Hsp70 active sites are nearly the same,

demonstrating the minor impact of the aromatic OH group. For 3HOK, two different orientations were found with nearly equal binding energies, one of which preferably occurs in complex with DNAK and the other in complex with Dhsp71. Being smaller than ATP and having a zwitterionic or anionic group, kynurenines tend to interact with only one part of the ATP site and the salt pair Lys 70 – Glu 171/172 beyond the site (S1 Fig.). Hence, they do not contact Hsp70 residues that bind ATP ribose group (Ser 274/270) or adenine group (Lys 270/266, Glu 267/263). Some other residues beyond ATP site, such as Pro 367/363 and Asp 368/364, can also be involved in kynurenines' binding.

D2', a non-enzymatically formed 3HOK dimer with the size comparable to ATP, can interact with the both parts of the binding site (Fig. 1). In contrast to 3HOK, D2' forms a hydrogen bond with Glu 267 in DNAK or ionic bond with Lys 267 in Dhsp71. D2' binding can occur in two opposite orientations. In complex with DNAK, it interacts with Lys 70 – Glu 171 by the amino acid moiety of the monomer (2) lacking the aromatic NH₂ group. For Dhsp71, the other D2' monomer (1) contacts the same residues. The second orientation lets D2' to form more bonds with higher affinity to Dhsp71 (K_d 9.4 kcal/mol) which is the median between the respective values for ATP and monomeric kynurenines. However, as Hsp70 structure is rather conservative, we cannot exclude the ability of D2' to bind to DNAK and other Hsp70 proteins in the same orientation. Thus, among the studied compounds D2' is the most potent competitive inhibitor of ATP binding and, therefore, Hsp70 activity.

Learning and short-term memory in *cd¹* after heat shock

Fig. 2 presents the analysis of learning acquisition and STM retention in *cd¹* and the wild-type *CS* males. A strong CCSP expressed as courtship index (CI) and learning index (LI) stays for 3 h (memory retention) at the level reached immediately after training (learning acquisition) in both strains. HS applied at 1st instar larval stage (HS1), prepupal stage (HS2) or before the experiment (HS3) does not affect learning/memory in *CS*. Thus, 5 day-old *CS* males are able to form STM both under normal conditions and after HS applied at different stages of ontogenesis as shown in [45].

As to *cd¹*, its learning/memory performance does not differ from that of *CS* both under normal conditions [26] and following HS1. However, 3 h LI sharply decrease both after HS2 and HS3, being significantly different from the intact control and *CS*. Thus, 5-day-old *cd¹* demonstrates drastic STM disturbances: there is a 5-fold LI reduction after HS applied either at prepupal stage or immediately before the experiment un adults.

Discussion

At organism level, expression and content of stress proteins, particularly of Hsp70, depend on the type of stress and can change both in different organs and over time. HSPs interact with many cell proteins, performing multiple functions – from the regeneration of partly broken protein structure and their protection from proteolysis to defense functions and escort for other proteins.

The most recent and comprehensive reviews on linking the ancient and interconnected systems of stress defense and learning acquisition are those of Zatzepina and coauthors [31, 32]. The mutant flies without *hsp70* genes showed a 4-fold lower 3 h learning and memory scores than control *CS* flies. This clearly pinpointed the involvement of Hsp70 in the learning process, namely in the formation of STM.

Also, *hsp70* males, both intact and treated with heat shock, demonstrated a drastic, 70-fold LTM decrease. The presence of one copy of the *hsp70* gene restored learning, STM and LTM formation. The authors have emphasized the protective role of Hsp70 chaperones in several Parkinson's (PD) and Alzheimer's diseases (AD) models [32]. NDDs, PD, AD and HD are accompanied by imbalance of KP metabolites content [46, 29]. The *Drosophila* AD model demonstrating memory and motor deficits shows the high production of the neurotoxic 3HOK [47]. Therefore, 3HOK is a promising early disease biomarker in human AD.

A rather strong binding of Hsp70 to different proteins under certain conditions can affect their functions. At the same time, issues related to possible binding of HSPs to tryptophan metabolites have never been discussed in literature. If so, what can be the mode of possible interactions between Hsp70 and 3HOK? Hsp70 proteins have a nucleotide binding domain consisting of two lobes with a deep cleft between them, binding ATP and ADP at the bottom [48]. Using Autodock Vina software, we have estimated the affinity of both 3HOK and other kynurenines binding to predicted ATP site of Dshp71. 3HOK is capable of dimer formation [16]. In this study, we demonstrate that 3HOK dimer binds to Dhsp71 protein with a rather high energy (-9.4 kcal/mol), both to the active side part that binds ATP phosphate group, and to the part that binds the adenine group, whereas the monomer binds only to the first part of the site. In liver tissue of healthy people, Hsp70 level is about 180 ng/g [49], in serum its level is about 6 ng/ml [50]. The average level of 3HOK in *cd¹* tissues is about 396 ug/g [15], being significantly higher compared to mammals [51]. This all makes 3HOK a potent competitive inhibitor of Dhsp71 and, possibly, the related proteins of Hsp70 superfamily. Over the period when the ATP binding site remains occupied, the chaperone cannot fully perform its functions, regulating stress response and memory formation.

The protein-bound form of 3HOK (PB-3HOK) is present in *Drosophila* head capsule containing pigment cells where XAN is synthesized from 3HOK. PB-3HOK was also found in the brain, though in less quantity [28]. Hence, 3HOK can penetrate in *Drosophila* central nervous system and affect cognitive functions in a stress-dependent manner.

How to explain that 5-day-old flies are very good learners without HS, but start demonstrating cognitive defects either on day 13 of adult life, or after exposure to HS at day 5? These are the same *cd¹* mutants prone to stress. In the first case, that is a prolonged stress of solitary confinement until the middle of flies' life before testing learning/memory, in the second case it is an acute HS-provoked stress. Interestingly, the later one, namely, the exposure to HS, exerts its effect only starting from prepupal stage, the time-point of 3HOK synthesis. The first one demonstrates how different types of stress experienced until midlife can gradually start resulting in progressive memory loss. This is the *Drosophila* proof [47] statement that 3HOK is a promising early disease biomarker in human AD.

In our study, we have shown that HS applied at prepupal stages or before the experiment impairs 3 h STM in 5-day-old *cd¹*, not influencing the wild type *CS*. There were no behavioral effects of HS applied at the 1st instar larvae stage. 3HOK dimer D2' was predicted to competitively inhibit Dhsp71, a *Drosophila* homologue of HSP70. Possibly, this impacts 3HOK-dependent decrease of in flies' ability to adapt to HS. Also, this sheds a light on newly emerged notion about the role of active forgetting or memory erasure in memory consolidation [58]. Therefore, *cd¹* learning/memory performance dependent on HS strengthens the findings of Zatsepina et al. [31] on the close interconnection of the two ancient systems of HS response and learning/memory.

In summary, a synergistic effect of HS and 3HOK-dependent oxidative stress has been revealed in *cd¹* model of memory disturbances. *Drosophila* Hsp70 homologue is among the putative targets of 3HOK and its dimer was predicted to inhibit ATP binding to protein active site. As both KP and HSP are conservative among different animal phyla, we can expect similar effects in human disorders provoked by 3HOK excess, such as Huntington's disease. Thus, our finding helps to reveal the molecular mechanisms of stress-dependent exacerbation of neurophysiological and cognitive disorders, associated with kynurenines' misbalance in the brain.

Abbreviations

AD – Alzheimer's diseases, CCSP – conditioned courtship suppression paradigm, CI – courtship index, 3HAA – 3-hydroxyanthranilic acid, 3HOK – 3-hydroxykynurenine, HD – Huntington's disease, HS – heat shock, HSPs – heat shock proteins, HSF1 – heat shock transcription factor 1, KP – kynurenine pathway, L-KYN – kynurenine, KYNA – kynurenic acid, LI – learning index, LTM – long-term memory (LTM), NDDs – neurodegenerative disorders, PD – Parkinson's disease, PHS – phenoxazinone synthase, QUIN – quinolinic acid, RMSD – root-mean-square deviation, ROS – reactive oxygen species, STM – short-term memory, XAN – xanthommatin.

Declarations

Acknowledgements

Not applicable

References

1. Moroni F (1999) Tryptophan metabolism and brain function: focus on kynurenine and other indole metabolites. *Eur J Pharmacol* 375:87–100. [https://doi.org/10.1016/s0014-2999\(99\)00196-x](https://doi.org/10.1016/s0014-2999(99)00196-x)
2. Lapin IP (2004) Stress, Anxiety, Depression, Alcoholism, Epilepsy. (Neurokynurenines Mechanisms and New Approaches to Treatment). DEAH, St. Petersburg

3. Schwarcz R, Bruno JP, Muchowski PJ, Wu HQ (2012) Kynurenines in the mammalian brain: when physiology meets pathology. *Nat Rev Neurosci* 13:465-477. <https://doi.org/10.1038/nrn3257>
4. Campbell BM, Charych E, Lee AW, Möller T (2014) Kynurenines in CNS disease: regulation by inflammatory cytokines. *Front Neurosci* 8:12. <https://doi.org/10.3389/fnins.2014.00012>
5. Christen S, Peterhans E, Stocker R (1990) Antioxidant activities of some tryptophan metabolites: possible implication for inflammatory diseases. *Proc Natl Acad Sci USA* 87:2506–2510. <https://doi.org/10.1073/pnas.87.7.2506>
6. Zhuravlev AV, Zakharov GA, Shchegolev BF, Savvateeva-Popova EV (2016) Antioxidant properties of kynurenines: density functional theory calculations. *PLoS Comput Biol* 12:e1005213. <https://doi.org/10.1371/journal.pcbi.1005213>
7. Okuda S, Nishiyama N, Saito H, Katsuki H (1998) 3-hydroxykynurenine, an endogeneous oxidative stress generator, causes neuronal cell death with apoptotic features and region selectivity. *J Neurochem* 70:299–307. <https://doi.org/10.1046/j.1471-4159.1998.70010299.x>
8. Guidetti P, Reddy PH, Tagle DA, Schwarcz R (2000) Early kynurenergic impairment in Huntington's disease and in a transgenic animal model. *Neurosci Lett* 283:233-235. [https://doi.org/10.1016/s0304-3940\(00\)00956-3](https://doi.org/10.1016/s0304-3940(00)00956-3)
9. Kessler M, Terramani T, Lynch G., Baudry M (1989) A glycine site associated with *N*-methyl-*D*-aspartic acid receptors: characterization and identification of a new class of antagonists. *J. Neurochem.* 52:1319–1328. <https://doi.org/10.1111/j.1471-4159.1989.tb01881.x>.
10. Schwarcz R, Pelliccari R (2002) Manipulation of Brain Kynurenines: Glial Targets, Neuronal Effects, and Clinical Opportunities. *J Pharmacol Exp Ther* 303:1-10. <https://doi.org/10.1124/jpet.102.034439>
11. Foster AC, Vezzani A, French ED, Schwarcz R (1984) Kynurenic acid blocks neurotoxicity and seizures induced in rats by the related brain metabolite quinolinic acid. *Neurosci Lett* 48:273–278. [https://doi.org/10.1016/0304-3940\(84\)90050-8](https://doi.org/10.1016/0304-3940(84)90050-8).
12. Urenjak J, Obrenovitch TP (2000) Neuroprotective potency of kynurenic acid against excitotoxicity. *Neuroreport* 11:1341-1344. <https://doi.org/10.1097/00001756-200004270-00038>
13. Linzen B (1974) The tryptophan to ommochrome pathway in insects. *Advances in Insect Physiol* 10:117-246. [https://doi.org/10.1016/S0065-2806\(08\)60130-7](https://doi.org/10.1016/S0065-2806(08)60130-7)
14. Phillips JP, Forrest HS, Kulkarni AD (1973) Terminal synthesis of xanthommatin in *Drosophila melanogaster*. 3. Mutational pleiotropy and pigment granule association of phenoxazinone synthetase. *Genetics* 73:45-56

15. Howells AJ, Summers KM, Ryall RL (1977) Developmental patterns of 3-hydroxykynurenine accumulation in white and various other color mutants of *Drosophila melanogaster*. *Biochem Genet* 15:1049–1059. <https://doi.org/10.1007/BF00484496>
16. Zhuravlev AV, Vetrovoy OV, Savvateeva-Popova EV (2018) Enzymatic and non-enzymatic pathways of kynurenines' dimerization: the molecular factors for oxidative stress development. *PLoS Comput Biol* 12:e1006672. <https://doi.org/10.1371/journal.pcbi.1006672>
17. Gibney SM, Fagan EM, Waldron AM, O'Byrne J, Connor TJ, Harkin A (2014) Inhibition of stress-induced hepatic tryptophan 2,3-dioxygenase exhibits antidepressant activity in an animal model of depressive behaviour. *Int J Neuropsychopharmacol* 6:917-928. <https://doi.org/10.1017/s1461145713001673>
18. Miller AH, Haroon E, Raison CL, Felger JC (2013) Cytokine targets in the brain: impact on neurotransmitters and neurocircuits. *Depress Anxiety* 30: 297-306. <https://doi.org/10.1002/da.22084>
19. Ritossa F (1962) A new puffing pattern induced by heat shock and DNP in *Drosophila*. *Experientia* 18:571-573. <https://doi.org/10.1007/BF02172188>
20. Evgen'ev MB, Garbuz DG, Zatsepina OG (2014) Heat shock proteins and whole body adaptation to extreme environments. Springer, Berlin
21. Siegel RW, Hall JC (1979) Conditioned responses in courtship behavior of normal and mutant *Drosophila*. *Proc Natl Acad Sci USA* 76:3430-3434. <https://doi.org/10.1073/pnas.76.7.3430>
22. Kamyshev NG, Iliadi KG, Bragina JV (1999) *Drosophila* conditioned courtship: two ways of testing memory. *Learn Mem* 6:1–20
23. Savvateeva-Popova EV, Popov AV, Grossman AI, Nikitina EA, Medvedeva AV, Peresleni AI, Korochkin LI, Moe G., Davidowitz E, Pyatkov KI, Myasnyankina E, Zatsepina OG., Schostak N, Zelentsova ES, Evgen'ev MB (2007) Pathogenic chaperone-like RNA induces congophilic aggregates and facilitates neurodegeneration in *Drosophila*. *Cell Stress Chaperones* <https://doi.org/10.1379/csc-222r.1>
24. Savvateeva-Popova EV, Popov AV, Grossman A, Nikitina EA, Medvedeva AV, Peresleni AI, Molotkov DA, Kamyshev NG., Pyatkov KI, Zatsepina OG., Schostak N, Zelentsova ES, Pavlova G., Panteleev D, Riederer P, Evgen'ev MB (2008) Non-coding RNA as a trigger of neuropathologic disorder phenotypes in transgenic *Drosophila*. *J Neural Transm (Vienna)* 115:1629-1642. <https://doi.org/10.1007/s00702-008-0078-8>
25. Kuzin BA, Nikitina EA, Cherezov RO, Vorontsova JE, Slezinger MS, Zatsepina OG., Simonova OB, Enikolopov G.N, Savvateeva-Popova EV (2014) Combination of Hypomorphic Mutations of the *Drosophila* Homologues of Aryl hydrocarbon receptor and Nucleosome assembly protein Family Genes Disrupts Morphogenesis, Memory and Detoxification. *PLOS One* 9: e94975. <https://doi.org/10.1371/journal.pone.0094975>

26. Savvateeva E, Popov A, Kamyshev N, Bragina J, Heisenberg M, Senitz D, Kornhuber J, Riederer P (2000) Age-dependent memory loss, synaptic pathology and altered brain plasticity in the *Drosophila* mutant *cardinal* accumulating 3-hydroxykynurenine. *J Neural Transm* 107:581–601. <https://doi.org/10.1007/s007020070080>
27. Savvateeva-Popova EV, Popov AV, Heinemann T, Riederer P (2003) *Drosophila* mutants of the kynurenine pathway as a model for ageing studies. *Adv Exp Med Biol* 527:713–722. https://doi.org/10.1007/978-1-4615-0135-0_84
28. Zhuravlev AV, Vetrovoy OV, Ivanova PN and Savvateeva-Popova EV (2020) 3-hydroxykynurenine in regulation of *Drosophila* behavior: the novel mechanisms for cardinal phenotype manifestations. *Front Physiol* 11:971. <https://doi.org/10.3389/fphys.2020.00971>
29. Campesan S, Green EW, Breda C, Sathyaikumar K., Muchowski PJ, Schwarcz R, Kyriacou CP, Giorgini F (2011) The kynurenine pathway modulates neurodegeneration in a *Drosophila* model of Huntington's disease. *Curr Biol* 21:961-966. <https://doi.org/10.1016/j.cub.2011.04.028>
30. Green EW, Campesan S, Breda C, Sathyaikumar KV, Muchowski PJ, Schwarcz R, Kyriacou CP, Giorgini F (2012) *Drosophila* eye color mutants as therapeutic tools for Huntington disease. *Fly* 6:117–120. <https://doi.org/10.4161/fly.19999>
31. Zatsepina OG, Nikitina EA, Shilova VY, Chuvakova LN, Sorokina S, Vorontsova JE, Tokmacheva EV, Funikov SY, Rezvykh AP, Evgen'ev MB (2021) Hsp70 affects memory formation and behaviorally relevant gene expression in *Drosophila melanogaster*. *Cell Stress Chaperones* 26:575-594. <https://doi.org/10.1007/s12192-021-01203-7>
32. Zatsepina OG, Evgen'ev MB, Garbuz DG. (2021) Role of a Heat Shock Transcription Factor and the Major Heat Shock Protein Hsp70 in Memory Formation and Neuroprotection. *Cells* 7:1638. <https://doi.org/10.3390/cells10071638>
33. Sharma D, Masison DC (2009) Hsp70 Structure, Function, Regulation and Influence on Yeast Prions. *Protein Pept Lett* 16: 571–581 <https://doi.org/10.2174/092986609788490230>.
34. Sali A, Blundell TL (1993) Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol* 234:779-815. <https://doi.org/10.1006/jmbi.1993.1626>.
35. Morris GM, Huey R, Lindstrom W, Sanner MF, Belew RK, Goodsell DS, Olson AJ (2009) Autodock4 and AutoDockTools4: automated docking with selective receptor flexibility. *J Comput Chem* 16:2785-2791. <https://doi.org/10.1002/jcc.21256>.
36. Storer AC, Cornish-Bowden A (1976) Concentration of MgATP²⁻ and other ions in solution. Calculation of the true concentrations of species present in mixtures of associating ions. *Biochem J* 1:1–5. <https://doi.org/10.1042/bj159000>.

37. Trott O, Olson AJ (2010) AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization and multithreading. J Comput Chem 31:455-461. <https://doi.org/10.1002/jcc.21334>
38. Humphrey W, Dalke A, Schulten K (1996) VMD: visual molecular dynamics. J Mol Graph 14:33-38 [https://doi.org/10.1016/0263-7855\(96\)00018-5](https://doi.org/10.1016/0263-7855(96)00018-5)
39. Nikitina E, Tokmatcheva E, Savateeva-Popova E (2003) Heat shock during the development of central structures of the *Drosophila* brain: memory formation in the I (1) ts403 mutant of *Drosophila melanogaster*. Russian Journal of Genetics. 39:33-40. <https://doi.org/10.1023/A:1022062609102>
40. Koemans TS, Oppitz C, Donders RAT van Bokhoven H, Schenck A., Keleman K, Kramer JM, (2017) *Drosophila* Courtship Conditioning As a Measure of Learning and Memory. J Vis Exp 124: 55808. <https://doi.org/10.3791/55808>
41. Gallagher M, Rapp PR (1997) The use of animal models to study the effects of aging on cognition. Annu Rev Psychol 48:339-370. <https://doi.org/10.1146/annurev.psych.48.1.339>
42. Rohlf FJ, Sokal RR (1981) Biometry: the principles and practice of statistics in biological research. Freeman, New York
43. Nguyen B, Hartich D, Seifert U, Rios PL (2017) Thermodynamic Bounds on the Ultra- and Infra-affinity of Hsp70 for Its Substrates. Biophys J 2:362-370. <https://doi.org/10.1016/j.bpj.2017.06.010>
44. Theyssen H, Schuster HP, Packschies L, Bukau B, Reinstein J (1996) The second step of ATP binding to DnaK induces peptide release. J Mol Biol 5:657-70. <https://doi.org/10.1006/jmbi.1996.0606>
45. Nikitina E, Kaminskaya A, Molotkov D, Popov A, Savvateeva-Popova V (2014) Effect of heat shock on courtship behavior, sound production, and learning in comparison with the brain content of *limk1* in *Drosophila melanogaster* males with altered structure of the *limk1* gene. Zh Evol Biokhim Fiziol 50:154-166
46. Breda C, Sathyaikumar KV, Sograte Idrissi S, Notarangelo FM, Estranero JG, Moore GG, Green EW, Kyriacou CP, Schwarcz R, Giorgini F (2016) Tryptophan-2,3-dioxygenase (TDO) inhibition ameliorates neurodegeneration by modulation of kynurenine pathway metabolites. Proc Natl Acad Sci USA 119:5435-5440. <https://doi.org/10.1073/pnas.1604453113>
47. Thomas J, Smith H, Smith CA, Coward L, Gorman G., De Luca M, Jumbo-Lucioni P (2021) The Angiotensin-Converting Enzyme Inhibitor Lisinopril Mitigates Memory and Motor Deficits in a *Drosophila* Model of Alzheimer's Disease. Pathophysiology 28:307–319. <https://doi.org/10.3390/pathophysiology28020020>
48. Flaherty KM, DeLuca-Flaherty C, McKay DB (1990) Three-dimensional structure of the ATPase fragment of a 70K heat-shock cognate protein. Nature 346:623-628. <https://doi.org/10.1038/346623a0>

49. Merchant S, Korbelik M (2011) Heat shock protein 70 is acute phase reactant: response elicited by tumor treatment with photodynamic therapy. *Cell Stress Chaperones* 2:153-62.
<https://doi.org/10.1007/s12192-010-0227-5>
50. Lechner P, Buck D, Sick L, Hemmer B, Multhoff G. (2018) Serum heat shock protein 70 levels as a biomarker for inflammatory processes in multiple sclerosis. *Mult Scler J Exp Transl Clin*.
<https://doi.org/10.1177/2055217318767192>.
51. Fuertig R, Ceci A, Camus S M et al (2016) LC–MS/MS-based quantification of kynurenine metabolites, tryptophan, monoamines and neopterin in plasma, cerebrospinal fluid and brain. *Bioanalysis* 8:1903–1917. <https://doi.org/10.4155/bio-2016-0111>
52. Welch WJ (1990) The mammalian stress response: Cell physiology and biochemistry of stress proteins. *Stress proteins in biology and medicine*. New York CSH Press pp 223-278
53. Adachi M, Liu Y, Fujii K, Calderwood SK, Nakai A, Imai K, Shinomura Y (2009) Oxidative stress impairs the heat stress response and delays unfolded protein recovery. *PLoS One* 11:e7719.
<https://doi.org/10.1371/journal.pone.0007719>
54. Takayama S, Reed J & Homma S (2003) Heat-shock proteins as regulators of apoptosis. *Oncogene* 22:9041–9047. <https://doi.org/10.1038/sj.onc.1207114>
55. Ravagnan L, Gurbuxani S, Susin SA, Maise C, Daugas E, Zamzami N, Mak T, Jäättelä M, Penninger JM, Garrido C, Kroemer G. (2001) Heat-shock protein 70 antagonizes apoptosis-inducing factor. *Nat Cell Biol* 9:839-43. <https://doi.org/10.1038/ncb0901-839>
56. Zakharov G.A, Zhuravlev AV, Payalina TL, Kamyshov NG., Savvateeva-Popova EV (2012) The effect of mutations of the kynurenine pathway of tryptophan metabolism on locomotor behavior and gene expression in glutamatergic and cholinergic systems of *D melanogaster*. *Russ J Genet Appl Res* 2:197–204. <https://doi.org/10.1134/S2079059712020141>
57. Koniszewski NDB, Kollmann M, Bigham M et al (2016) The insect central complex as model for heterochronic brain development—background, concepts, and tools. *Dev Genes Evol* 226:209–219.
<https://doi.org/10.1007/s00427-016-0542-7>
58. Davis RL, Zhong Y (2017) The biology of forgetting-A perspective. *Neuron* 95:490-503.
<https://doi.org/10.1016/j.neuron.2017.05.039>

Figures

Figure 1

Ligand docking into the ATP site of Dhsp71 model structure a. The complex with ATP; b. the complex with 3HOK dimer D2' (L-3HOKzi_D2'). Hydrogen bonds with $\geq 155^\circ$ angle and $\leq 3.2 \text{ \AA}$ distance between the heavy atoms are shown by dotted lines. See Table 1 and S1 Table for the whole pattern of ligand-protein bonds.

Figure 2

Learning and memory abilities in *Drosophila* strains under normal conditions and heat shock CI: box and whisker plots, the median is shown as an orange line. X axis: N – naïve, 0 h – learning, 3 h – STM; Y axis: percentage. Statistical differences: \$ from CI (naïve), & from LI (0 h), # from CS; two-sided randomization test, $\alpha R < 0.05$, $N = 20$.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [S1Dockingcoordinates.txt](#)
- [S1Figure.tiff](#)
- [S1Table.pdf](#)