Metabolome and Transcriptome Profiling Reveals Anthocyanin Contents and Anthocyanin-Related Genes of Chimeric Leaves in *Ananas comosus* Var. *Bracteatus*

Xuzixin Zhou

Sichuan Agricultural University - Chengdu Campus <https://orcid.org/0000-0001-8048-5287>

Yanbin Xue

Sichuan Agricultural University - Chengdu Campus

Meiqin Mao

Sichuan Agricultural University - Chengdu Campus

Yehua He

South China Agricultural University College of Horticulture

Mark Owusu Adjei

Sichuan Agricultural University - Chengdu Campus

Wei Yang

Sichuan Agricultural University - Chengdu Campus

Hao Hu

Sichuan Agricultural University - Chengdu Campus

Jiawen Liu

Sichuan Agricultural University - Chengdu Campus

Lijun Feng

Sichuan Agricultural University - Chengdu Campus

Huiling Zhang

Sichuan Agricultural University - Chengdu Campus

Jiaheng Luo

Sichuan Agricultural University - Chengdu Campus

Xi Li

Sichuan Agricultural University - Chengdu Campus

Lingxia Sun

Sichuan Agricultural University - Chengdu Campus

Zhuo Huang

Sichuan Agricultural University - Chengdu Campus

Jun Ma (✉ Junma365@hotmail.com)

Sichuan Agricultural University <https://orcid.org/0000-0001-8605-4971>

Research article

Keywords: Ananas comosus var, bracteatus, anthocyanins, CIELAB, Metabolomics, Transcriptomic, MYB transcription factor

Posted Date: October 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-79383/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Genomics on May 7th, 2021. See the published version at <https://doi.org/10.1186/s12864-021-07642-x>.

Abstract

Background: *Ananas comosus* var. *bracteatus* is a colorful plant used as a cut flower or landscape ornamental. The unique foliage color of this plant includes both green and red leaves and, as a trait of interest, deserves investigation. In this study, green and red parts of chimeric leaves of *Ananas comosus* var. *bracteatus* were sampled and anthocyanin accumulation differences were comparatively analyzed at phenotypic, cellular and molecular levels.

Results: A CIELAB results indicated that the a^* values and L^* values samples had significant differences between two parts. Freehand sections showed that anthocyanin presented limited accumulation in the green leaf tissues but obviously accumulation in the epidermal cells of red tissues. Transcriptomic and metabolomic analyses were performed by RNA-seq and LC-ESI-MS/MS. Among the 508 identified metabolites, 10 kinds of anthocyanins were detected, with 6 significantly different between the two samples. The cyanidin-3,5-O-diglucoside content that accounts for nearly 95.6% in red samples was significantly higher than green samples. RNA-Seq analyses showed that 11 out of 40 anthocyanin-related genes were differentially expressed between the green and red samples. Transcriptome and metabolome correlations were determined by nine quadrant analyses, and 9 anthocyanin-related genes, including *MYB5* and *MYB82*, were correlated with 7 anthocyanin-related metabolites in the third quadrant in which genes and metabolites showing consistent change. Particularly, the PCCs between these two MYB genes and cyanidin-3,5-O-diglucoside were above 0.95.

Conclusion: Phenotypic colors are closely related to the tissue structures of different leaf parts of *Ananas comosus* var. *bracteatus*, and two MYB transcription factors might contribute to differences of anthocyanin accumulation in two parts of *Ananas comosus* var. *bracteatus* chimeric leaves. This study lay a foundation for further researches on functions of MYBs in *Ananas comosus* var. *bracteatus* and provides new insights to anthocyanin accumulation in different parts of chimeric leaves.

Background

Anthocyanin is a prominent pigment found in plant organs and has become an indicator of the ornamental or edible value of plants[1–3]. Several enzyme-coding structural genes of anthocyanin biosynthesis have been reported in previous studies[4], and these genes were mainly distributed in three pathways, namely, phenylpropanoid biosynthesis, flavonoid biosynthesis and anthocyanin biosynthesis[5]. At the transcriptional level, anthocyanin biosynthesis is mainly regulated by MYB, basic helix-loop-helix (bHLH) and WD40 transcription factors, which form a complex to directly bind with promoters of anthocyanin biosynthetic genes, such as *chalcone synthesis (CHS)*, *chalcone isomerase (CHI)*, *flavanone 3-hydroxylase (F3H)*, *flavonoid 3'-hydroxylase (F3'H)*, *flavonoid 3'5'-hydroxylase (F3'5'H)*, *dihydroflavonol 4-reductase (DFR)*, *anthocyanidin reductase (ANS)* and *UDP-glucose: flavonoid 3-O-glucosyltransferase (3GT)*[6–8]. After biosynthesis, anthocyanins are transported into vacuoles by a series of transporters, including glutathione S-transferases (GSTs), multidrug resistance-associated protein (MRP), and multidrug and toxic compound extrusion (MATE)[9–11]. According to previous reports

about anthocyanin related transcription factors in *Arabidopsis thaliana*, the MYB transcription factor TRANSPARENT TESTA 2 (TT2) regulates the expression of late biosynthesis genes to pigment the seed coat by combining with factors TT8 (bHLH42) and TTG1 (WD40)[12, 13]. In petunia petal cells, anthocyanin biosynthesis is also regulated by the MYB-bHLH-WD40 complex, but some MYBs, such as AtMYB12 and VvMYBF1, regulate flavonoid biosynthesis without bHLH and WD40 cofactors[14, 15]. In higher plants, anthocyanin biosynthesis is a complicated process that is easily influenced by environmental factors, including light, temperature, moisture, pH, metal ions, and plant hormones[16–20]. However, the molecular mechanism of anthocyanin accumulation in *Ananas comosus* var. *bracteatus* leaves is still largely unknown.

A. comosus var. *bracteatus* is an herbaceous monocot distributed primarily in tropical regions[21]. It has high decorative and ornamental value due to its chimeric leaves, which present a regular distribution of green and red colors. Therefore, *A. comosus* var. *bracteatus* is often cultivated as a commodity for the international market[22]. In addition, carbon assimilation of *A. comosus* var. *bracteatus* occurs via the crassulacean acid metabolism (CAM) pathway, thus allowing for effective utilization of water under drought conditions[23, 24]. Therefore, this plant is mostly used as an indoor potted plant or even as an outdoor landscape plant without much need for irrigation. With the increase in market demands for new varieties of colorful plants, *A. comosus* var. *bracteatus* has successfully attracted attention from researchers[25, 26], and this research is of great significance for further revealing the mechanisms of *A. comosus* var. *bracteatus* leaf color. However, detailed information on the anthocyanins in *A. comosus* var. *bracteatus* leaf tissue has not been formally reported. Moreover, the regular and specific distribution of pigments in different parts of *A. comosus* var. *bracteatus* leaves, especially the specific distribution of anthocyanins, has not been thoroughly investigated in previous research. Therefore, integrated research on the accumulation and distribution of anthocyanin in *A. comosus* var. *bracteatus* leaves is urgently needed.

Plant coloration is often measured by two main methods: Royal Horticulture Society Color Chart (RHSCC) and the CIELAB color coordinate[27]. With the development of technology, the RHSCC has gradually been replaced by advanced colorimetric devices that can digitize the brightness and saturation of color[28]. These devices based on the CIELAB color coordinate principle can accurately measure the color value by a series of indexes. Freehand cross sectioning is the prominent method used to investigate pigments in plant tissue structures and cells, allowing researchers to directly examine how pigments influence phenotypes at the cellular level. Previous studies on the dissection of the colorful leaves of *A. comosus* var. *bracteatus* have been inadequate, although studies on the anatomies of other plants, such as *Pilea cadierei* Gagnep.[29], *Aloe*[30] and *Bruguiera parviflora*[31], can be used as a reference. Additionally, molecular profiling approaches, such as metabolomics and transcriptomics, are effective for exploring the underlying mechanisms responsible for a plant's phenotype. In recent years, LC-ESI-MS/MS and RNA-seq techniques have been rapidly developed and become popular approaches for identifying a considerably large number of plant metabolites and providing broad coverage of transcript abundance, respectively[5, 32, 33]. Transcriptome analysis combined with metabolome analysis enables the comprehensive and detailed exploration of molecular mechanisms.

In this study, *A. comosus* var. *bracteatus* chimeric leaves were selected as experimental materials to carry out phenotypic, cellular and molecular analyses by CIELAB colorimeter, freehand sectioning, LC-ESI-MS/MS and RNA-seq. Comparative analysis indicated the locations of anthocyanins in the leaf tissues, types of anthocyanins and genes involved in regulating anthocyanin biosynthesis in *A. comosus* var. *bracteatus* leaves. The results obtained in this study lay the foundation for molecular breeding and optimized cultivation as well as provide a reference for further research on chimeric leaf color.

Results

Leaf coloration

The distribution of the color values of the GR and RE samples was determined with the CIELAB coordinates: the L^* values ranged from 46.03 to 77.42, the a^* values ranged from -5.68 to 23.58, the b^* values ranged from 14.55 to 17.40, and the C^* values ranged from 15.40 to 28.42. Figure 2 shows that the L^* value of GR was significantly lower than that of RE for both surfaces, indicating that the color of GR was darker than that of RE (Fig. 2A). Moreover, the a^* value of GR was negative for both surfaces and that of RE was significantly higher than that of GR and was red. The b^* value of both surfaces had no significant difference between GR and RE. The C^* value of the RE adaxial surface was approximately 2 times higher than that of GR, but there was no significant difference between the two phenotypes on the abaxial surfaces. The CIE locations of the two samples indicated that GR mainly contained green and yellow colors, while RE mainly contained red and yellow colors.

Anatomical Cross Section

GR and RE samples were separated for anatomical analysis. The GR consisted of the epidermis, mesophyll and vein tissues. In the central green part of the chimeric leaves, there were large water storage cells that are typical structural characteristics of drought-tolerant plants (Fig. 3A). Mesophyll cells of the central green part of the chimeric leaf contained chloroplasts and showed a green color. However, no water storage tissues were observed in the red margins of the chimeric leaves, and the mesophyll cells in this area did not have a green color. According to previous reports, chlorophylls and carotenoids mainly exist in photosystem I (PS I), photosystem II (PS II) and the light harvesting complex (LHC) to adjust light absorption and conversion[34–36]. These two kinds of pigments coexist together in the thylakoids, which increases the likelihood that yellow pigments, such as carotenoids, will be covered by green pigments. On the other hand, chloroplasts were invisible in freehand sections in the RE chimeric leaf samples, and the middle mesophyll mainly consisted of colorless cells (Fig. 3B). Red cells were obviously observed in the epidermis and 1–2 layers of mesophyll under the epidermis in the RE sample.

Comparison of Anthocyanin-related Metabolites in the GR and RE Samples of *A. comosus* var. *bracteatus*

The pigment contents and relative proportions of GR and RE are shown in Fig. 4. The chlorophyll content in GR was approximately 57 times higher than that in RE, while anthocyanins were undetectable in GR

(Fig. 4A). According to Fig. 4B, chlorophyll, accounting for 85.9% of all pigments, was the most prominent pigment in GR. The total pigment content of RE was less than half of the GR pigment content, with chlorophyll and carotenoid contents of no more than 0.05 mg/g. However, total anthocyanins accounted for 93.2% of the total pigments in RE, followed by a small proportion of chlorophyll and carotenoid. To determine the types of anthocyanins in RE, anthocyanin-related metabolites were analyzed by UPLC-ESI-MS/MS. First, a total of 508 metabolites were identified in GR and RE, and the most abundant metabolites were flavonoids, organic acids, alkaloids, lipids, amino acids and derivatives (Table S1). The cluster heatmap, PCA and OPLS-DA plots showed differences in metabolites between two phenotypes (Fig. 5A-5C). Among these metabolites, 153 flavonoids were detected, including 10 kinds of anthocyanins (Table S1a). After selecting differentially produced metabolites (DPMs) based on an absolute $\log_2$ fold-change ≥ 1 among the total metabolites with a VIP value ≥ 1 , 164 DPMs were shown in the filtered results (Table S2). The volcano plots indicated that there were 112 upregulated and 52 downregulated metabolites between the GR and RE samples (Fig. 5B). In addition, 42 differentially produced flavonoids accounted for approximately 25% of the total DPMs, including flavonoids, flavonols, dihydroflavones, dihydroflavonols, flavonoid carbonosides, anthocyanins and other flavonoids. Additionally, 67 DPMs were annotated to the KEGG database, and they were mainly enriched in “Metabolic pathways” and “Biosynthesis of secondary metabolites” (Table S3). A total of 65 metabolic pathways were enriched, including anthocyanin-related pathways such as “phenylpropanoid biosynthesis” (Ko00940), “flavonoid biosynthesis” (Ko00941), and “anthocyanin biosynthesis” (Ko00942) (Fig. 5C, File S1). In the anthocyanin-related pathways, 15 metabolites including 6 kinds of anthocyanins showed significantly different production levels. Among these anthocyanins, cyanidin derivatives accounted for 97% of the total anthocyanin content (Table 1, Fig. 4B). In particular, cyanidin-3,5-O-diglucoside (pme1777) accounted for 98% of cyanidin derivatives, probably being the most prominent pigment responsible for the RE phenotype.

Anthocyanin-related Differentially Expressed Genes (DEGs) in the GR and RE Samples of *A. comosus* var. *bracteatus*

According to the reference transcriptome, 20905 genes were obtained by using RNA-seq technology (Table S4). The clustering heatmap indicated that GR and RE had different gene expression patterns (Fig. 6A). Differentially expressed genes (DEGs) were selected by an absolute $\log_2$ fold-change ≥ 1 with a false discovery rate (FDR) ≤ 0.05 and compared to six public databases (KEGG, NR, SwissProt, Tremble, GO and KOG) to determine their potential functions. In total, 1086 DEGs were selected, and volcano figures revealed that there were 619 downregulated and 467 upregulated DEGs between the GR and RE samples (Fig. 6B, Table S5). KEGG enrichment results indicated that 365 DEGs were enriched in KEGG pathways, including 201 DEGs involved in “Metabolic pathways” and 115 DEGs involved in “Biosynthesis of secondary metabolites” (Fig. 6C, 6D, Table S6). The GO bar plot indicated that the number of DEGs involved in metabolic process, cellular processes, cells, cell parts, catalytic activity, binding, etc. was significantly greater than 100 (Fig. 6E). A total of 121 KEGG pathways were enriched including anthocyanin-related pathways such as “phenylpropanoid biosynthesis”, “flavonoid biosynthesis”,

“anthocyanin biosynthesis”, and “ABC transporters” (File S2). In addition, 110 transcription factors (TFs) that were differently expressed in GR and RE were identified, including 19 MYB and MYB-related factors and 6 bHLH factors (Table S7). Among these TFs, *MYB5* (Aco023263.1.path1) and *MYB82* (Aco023266.1.path1) annotated “transcription factor TT2” and “transcription factor TT2-like”, respectively, and had a probable relationship with anthocyanin biosynthesis. Finally, 40 anthocyanin-related genes were selected for comparison between GR and RE and are shown in Tables 2 and 11 of these genes were DEGs between GR and RE.

qPCR Validation

To validate the RNA-Seq results, eight anthocyanin related genes including *PAL* (Aco020618.1.path1), *CHS* (Aco016200.1.path1), *CHI* (Aco012547.1.path1), *F3'H* (Aco003857.1.path1), *DFR* (Aco006769.1.path1), *3GT* (Aco012126.1.path1), *ABCA3* (Aco006845.1.path1) and *MYB5* (Aco023263.1.path1) were selected, and their relative expression levels in the GR and RE phenotypes were analyzed by using qPCR. The expression patterns of these genes are generally consistent with the RNA-Seq results (Fig. 7).

Correlation Analysis of the Relationship between Genes and Metabolites Involved in Anthocyanin Biosynthesis of *A. comosus* var. *bracteatus*

In this study, a pathway map containing metabolites and structural genes related to anthocyanin biosynthesis was constructed based on the metabolome and transcriptome analyses with the KEGG database (Fig. 8). Metabolites and genes were labeled in different colors, indicating production and expression levels between GR and RE. The expression of 1 *PAL* (Aco020618.1.path1) and 1 *FLS* (Aco011051.1.path1) in GR was significantly higher than that in RE. In contrast, expression of 2 *CHIs* (Aco012547.1.path1 and Aco014232.1.path1), 1 *F3H* (Aco018609.1.path1), 1 *F3'H* (Aco003857.1.path1), 1 *DFR* (Aco006769.1.path1) and 1 *3GT* (Aco012126.1.path1) in RE was significantly higher than that in GR. However, early metabolites such as naringenin chalcone (pme2960), naringenin (pme0376) and dihydroquercetin (DHQ, mws0044) did not show significant differences between GR and RE. In the late stage of anthocyanin biosynthesis, the production level of pelargonidin-3,5-diglucoside (pme1793), cyanidin-3-O-rutinoside (pme1773), cyanidin-3,5-O-diglucoside (pme1777), delphinidin-3,5-O-diglucoside (pmf0116) and delphinidin-3-O-glucoside (pme1398) in RE was significantly higher than that in GR. In particular, cyanidin-3,5-O-diglucoside accounted for 95.6% of all detected anthocyanins in RE, followed by other types of anthocyanins, which were present only at extremely low levels. Additionally, the existence of delphinidin derivatives suggested that purple or blue colors did exist in RE but were probably masked by the high proportion of cyanidin derivatives.

To understand the relationship between anthocyanin-related genes and metabolites, correlation tests were carried out. In the nine-quadrant plot, metabolites and genes in the third and seventh quadrants showed consistent changes (Fig. 9A, File S3). The nine-quadrant results showed that 1734 genes that had Pearson correlation coefficient (PCC) values ≥ 0.8 , and 168 metabolites were identified in the third

and seventh quadrants. Among these genes, 9 anthocyanin-related genes were identified and showed correlations with 7 anthocyanin-related metabolites in the third quadrant (Fig. 9B). *CHI* (Aco012547.1.path1), *F3'H* (Aco003857.1.path1), *F3H* (Aco018609.1.path1), *DFR* (Aco006769.1.path1) and *MYB5* (Aco023263.1.path1) were positively correlated with the content of luteolin-7-O-glucoside (pme2459), delphinidin-3-O-glucoside (pme1398), cyanidin-3-O-rutinoside (pme1773), pelargonidin-3,5-diglucoside (pme1793), cyanidin-3,5-O-diglucoside (pme1777) and delphinidin-3,5-O-diglucoside (pmf0116). *MYB82* (Aco023266.1.path1) were positively correlated with the content of delphinidin-3-O-glucoside, cyanidin-3-O-rutinoside, cyanidin-3-O-rutinoside, cyanidin-3,5-O-diglucoside and delphinidin-3,5-O-diglucoside. Naringenin-7-O-rutinoside (mws1066) was positively correlated with *CHI* (Aco014232.1.path1), *3GT* (Aco012126.1.path1) and *ABCA3* (Aco006845.1.path1).

Discussion

CIELAB Values Are Closely Associated with the Leaf Organization Structures and Pigment Distribution of *A. comosus* var. *bracteatus*

The plant coloration shown in nature is not only attributed to different kinds and proportions of pigments distributed in plant organs but is also attributed to the internal structures of plant organs. As shown in Fig. 2, GR had a lower L^* value than RE, which seemed to be related to a dark phenotype even when the chlorophyll content of GR far exceeded that of RE. Connected with leaf organization structures, this result might be related to the thickness of the GR leaf areas, which are much thicker than the RE areas. Therefore, it was not difficult to infer that the thicker the leaf tissue, the more mesophyll cells exist. Moreover, the cell shape and cell permeability in leaf tissues can be quite different, which easily leads to refraction and reflection of light[37–39]. Additionally, the physical green color itself is a mix of cyan and yellow colors, which are among the three primary colors in the CMYK color model. With the increase in tissue thickness, layers of green color tended to become dark, which also made the L^* value of GR low. In contrast, the RE leaf areas were much thinner than the GR areas, and the mesophyll cells were colorless, with a few cells having anthocyanin in the epidermis. Thus, there was a relatively lower chance of light refraction and reflection in RE. In addition, different from the green color, the red color itself is one of three primary colors. The L^* value of RE was much higher than that of GR even when the thickness of the colorless mesophyll tissue reached a certain level. This result suggested that the RE phenotype was not mainly influenced by the middle transparent cells but was mainly related to the anthocyanin content in the epidermis.

The b^* values of GR and RE were not significantly different, and both were light yellow. The results suggested that these tissues were similar in some respects. According to Fig. 3, we inferred that the yellow color of both GR and RE was probably derived from the keratinized cell walls of epidermal cells, and the specialized cell wall structure is typical for plants under long-term environmental stress. To reduce the evaporation of water and adapt to the high-temperature environment, tropical plants such as *A. comosus* var. *bracteatus* have to develop a series of features to overcome these challenges. Thick wall tissue, such as keratinized cell walls, is a tropical plant feature that protects the water in plants from

excessive evaporation[40]. From the chemical composition perspective, the thick-walled tissue mainly contains fatty acids that have hydrophobic functions. Therefore, we infer that the epidermis of both GR and RE showed slight coloration when certain thresholds were reached for fatty acid levels and tissue thickness.

Anthocyanins are diversified and variable in *A. comosus* var. *bracteatus*

Pelargonidins, cyanidins and delphinidins are three branches of metabolites in anthocyanin biosynthesis and are derived from the catalysis of *F3H*, *F3'H* and *F3'5'H*, respectively. However, not all plants have these three key enzymes; in particular, certain plants lack *F3'5'H*, which contributes to blue/purple anthocyanin biosynthesis. For example, blue/purple varieties of roses, carnations, lilies and chrysanthemums hardly exist in nature due to the lack of delphinidin-based anthocyanins [41]. According to the metabolome analysis of *A. comosus* var. *bracteatus*, the peak areas of pelargonidin derivatives, cyanidin derivatives and delphinidin derivatives were detected in both GR and RE. Moreover, *F3H*, *F3'H* and *F3'5'H* had certain FPKM values in samples of the two phenotypes, which suggested that the three branches of different anthocyanidin biosynthesis exist in *A. comosus* var. *bracteatus*. However, the significant difference in the relative contents among the three derivatives indicated that the predominance of catalysis probably has an important influence on these compounds, and DFR was confirmed to have substrate specificity, according to previous reports. For instance, not dihydromyricetin (DHM) but dihydroquercetin (DHQ) was efficiently catalyzed by DFR in petunia to produce cyanidins and derivatives, but red petunia flowers turned orange with suppression of *F3'H* by RNAi[42, 43]. In addition, overexpression of *viola F3'5'H* and *iris DFR* and downregulation of *rose DFR* resulted in efficient delphinidin accumulation in rose petals that showed blue/purple phenotypes [44]. The results indicated that the substrate specificity of DFR is different in different plants. In *A. comosus* var. *bracteatus*, the cyanidin-type anthocyanin content was significantly higher than the other two types of anthocyanin, which suggested that *A. comosus* var. *bracteatus* DFR predominantly uses DHQ as a substrate. Additionally, DHM derives from the catalysis of DHQ (Fig. 7), so it is possible that competition for DHQ substrates between *F3'5'H* and DFR was responsible for the significant difference in cyanidin-type and delphinidin-type anthocyanin contents. These mechanisms provide a possible approach for regulating the leaf colors of *A. comosus* var. *bracteatus* by biotechnology.

MYB Family is One of Important Roles in Anthocyanin accumulation and Distribution in *Ananas comosus* var. *bracteatus*

Although few anthocyanin-related structural genes have been reported in past studies[45, 46], these genes have been found to be regulated by different kinds of proteins. Among these proteins, the MYB family has been reported to be involved in regulating anthocyanin biosynthesis[4, 47, 48]. MYB factors control anthocyanin biosynthesis genes mainly by two ways: MYB proteins interact with bHLH and WD40 proteins to form a MBW protein complex and then activate structural genes; and early biosynthetic genes (*CHS*, *CHI*, *F3H*, *F3'H*, *FLS*) that produce flavonols are regulated by independent MYB regulators (MYB11, MYB12)[46]. In the transcriptome files of *A. comosus* var. *bracteatus*(Table S6), two MYB genes, *MYB5*

(Aco023263.1.path1) and *MYB82*(Aco023266.1.path1), were annotated as “transcription factor TT2 (A)” and “transcription factor TT2-like (A)”, respectively, which suggested that they probably have similar functions as *TT2*. *TT2* is a determinant gene in *Arabidopsis thaliana* that encodes the R2R3-MYB domain protein, and it is involved in proanthocyanin biosynthesis via interactions with TT8 and TTG1[49]. In addition, Jan and co-workers also showed that *FaMYB9/FaMYB11*, which is the functional homologue of *AtTT2*, interacts with *FabHLH3* and *FaTTG1* to regulate proanthocyanin biosynthesis in strawberry fruits[50]. Moreover, *MYB5* and *MYB82* were shown to have high PCC values (all above 0.9) with anthocyanin metabolites, including delphinidin-3-O-glucoside(pme1398), cyanidin-3-O-rutinoside(pme1773), pelargonidin-3,5-diglucoside (pme1793), cyanidin-3,5-O-diglucoside(pme1777), delphinidin-3,5-O-diglucoside (pmf0116) and luteolin-7-O-glucoside (pme2459). Previous reports and our results indicated a potential relationship between the two MYB factors and anthocyanin accumulation. However, our results suggested that the accumulation and specific distribution of anthocyanin are not only regulated by the MYB family but also influenced by the biosynthesis of other metabolites, such as photosynthetic pigments, flavone, flavonols, etc. These metabolites may share some common substrates with anthocyanin, which probably leads to competition. For example, unstable anthocyanidins are first stabilized *in vivo* with UDP-glucose by 3G7[51] We infer that UDP-glucose is likely derived from glycogen, which is also used for the biosynthesis of other metabolites and organelles, such as chloroplasts, sterols, photosynthetic pigments, etc.[52]. Compared with GR, glycogen probably flows more into the UDP-glucose branch to increase anthocyanin biosynthesis because of incomplete photosynthetic functions in RE. Additionally, anthocyanin biosynthesis is one of several flavonoid biosynthesis branches. From phenylalanine to different types of anthocyanins, different branches occur on the pathway that may compete the common substrates. The flows of substrates in these metabolites in different samples probably results in phenotypic differences. Generally, the mechanisms underlying these factors will lead to the regular and specific pigment distribution and accumulation in the chimeric leaves of *A. comosus* var. *bracteatus*, and these mechanisms are worthy of further study.

Conclusions

Analyses at the phenotypic, cellular and molecular levels showed that (1) CIELAB color values are closely associated with leaf tissue structures and anthocyanins hardly accumulate in green tissues but clearly accumulate in epidermal cells in the RE areas of *A. comosus* var. *bracteatus* leaves. (2) Ten different types of anthocyanins belonging to the three branches of the anthocyanin biosynthesis pathway were detected in both GR and RE samples, and cyanidin-3,5-O-diglucoside accounted for the highest proportion in RE and was present in significantly higher levels than those in GR. (3) *MYB5* and *MYB82* are two probable transcription factors that result in the specific accumulation and distribution of anthocyanin in different parts of *A. comosus* var. *bracteatus* leaves.

Materials And Methods

Plant Materials and Sample Preparation

The chimeric leaves of two-year-old *A. comosus* var. *bracteatus* from an experimental field at Sichuan Agricultural University were used in this study. Two types of samples were collected: the green central parts (GR) and the red edges (RE) from chimeric leaves (Fig. 1). After being washed completely, the two types of samples were collected for color estimation, pigment content determination and anatomical cross sectioning. Additionally, samples were cut into small pieces and immediately frozen with liquid nitrogen. The frozen samples were stored at -80°C for further transcriptome and metabolome analyses.

Leaf Color Estimation

In this assay, the two types of samples were measured by a CM-2600d color analyzer (Konica Minolta CM-2600d, Japan) for color digitization. The measurement of leaf color was based on the Commission Internationale de l'Éclairage (CIE D65/10°) scale[53, 54]. CIE color indexes include the lightness component L^* and two chromatic components a^* and b^* . L^* values from 0 to 100 indicate the range from black to white; values from $-a^*$ to $+a^*$ represent a decrease in green and an increase in red; values from $-b^*$ to $+b^*$ represent a decrease in blue and an increase in yellow. In addition, C^* represents chroma, which is calculated by the following equation: $C^*=(a^{*2}+b^{*2})^{1/2}$ [55]. The color data were finally arranged and visualized by Excel 2019 and Photoshop 5.0.

Anatomical Cross Sectioning

To obtain a better understanding of the anthocyanin distribution in the tissues of GR and RE samples, the samples were cut into blocks of an appropriate size for freehand cross sectioning. Blocks were further cut by a pair of overlapping blades, and a section from the gap between the blades was immediately transferred onto a glass slide sealed with a cover glass. Sections were observed by a microscope (XSP-15CA, Shanghai CSOIF Co., Ltd) and photographed by Cellsens Standard software.

Chlorophyll and Carotenoid Content Determination

The chlorophyll and carotenoid contents of GR and RE samples were determined. A sample of 0.1 g was cut into slices and transferred into a 15 mL centrifuge tube with 5 mL 95% ethanol for extraction. The tube was kept in the dark until the samples became colorless. Chlorophyll and carotenoid absorption was measured at 665, 649 and 470 nm with a UV-Vis spectrophotometer (UV1901 s/UV1901PCS, Shanghai Youke Instrument Co., Ltd.) equipped with 1.0 cm quartz cells, and the concentration was calculated following the method of Arnon [56].

Anthocyanin Content Determination

The total anthocyanin contents of GR and RE samples were measured and calculated by using a pH differential protocol[57, 58]

Sample Extraction for Ultra-Performance Liquid Chromatography (UPLC)/MS Analysis

Freeze-dried leaves stored at -80°C were crushed using a mixer mill (MM 400, Retsch) with a zirconia bead for 1.5 min at 30 Hz. Then, 100 mg of the powder was weighed and extracted overnight at 4°C with 0.6 ml 70% aqueous methanol. Following centrifugation at 10,000 g for 10 min, the extracts were absorbed (CNWBOND Carbon-GCB SPE Cartridge, 250 mg, 3 ml; ANPEL, Shanghai, China, www.anpel.com.cn/cnw) and filtrated (SCAA-104, 0.22 µm pore size; ANPEL, Shanghai, China, <http://www.anpel.com.cn/>) before the UPLC-MS/MS analysis.

UPLC Conditions

The sample extracts were analyzed using a UPLC-ESI-MS/MS system (UPLC, Shim-pack UFLC SHIMADZU CBM30A system, www.shimadzu.com.cn/; MS, Applied Biosystems 4500 Q TRAP, www.appliedbiosystems.com.cn/). The analytical conditions for the UPLC analysis included an Agilent SB-C18 column (1.8 µm, 2.1 mm*100 mm), and the mobile phase for solvent A consisted of pure water with 0.1% formic acid and for solvent B consisted of acetonitrile. Sample measurements were performed with a gradient program that employed the starting conditions of 95% A and 5% B. A linear gradient to 5% A and 95% B over 9 min was programmed, and a composition of 5% A and 95% B was maintained for 1 min. Subsequently, a composition of 95% A and 5.0% B was adjusted within 1.1 min and maintained for 2.9 min. The column oven was set to 40°C, and the injection volume was 4 µl. The effluent was alternatively connected to an ESI-triple quadrupole-linear ion trap (QTRAP)-MS.

ESI-QTRAP-MS/MS

LIT and triple quadrupole (QQQ) scans were acquired on a triple quadrupole-linear ion trap mass spectrometer (Q TRAP) and API 4500 Q TRAP UPLC/MS/MS System equipped with an ESI Turbo Ion-Spray interface. The system was operated in positive and negative ion mode and controlled by Analyst 1.6.3 software (AB Sciex). The ESI source operation parameters were as follows: ion source, turbo spray; source temperature, 550°C; ion spray voltage (IS), 5500 V (positive ion mode)/-4500 V (negative ion mode); ion source gas I (GSI), gas II(GSII), curtain gas (CUR), 50, 60, and 30.0 psi, respectively; collision gas (CAD), high. Instrument tuning and mass calibration were performed with 10 and 100 µmol/L polypropylene glycol solutions in QQQ and LIT modes, respectively. QQQ scans were acquired via MRM experiments with collision gas (nitrogen) set to 5 psi. DP and CE for individual MRM transitions were performed with further DP and CE optimization. A specific set of MRM transitions were monitored for each period according to the metabolites eluted within this period.

RNA Library preparation for transcriptome sequencing

A total amount of 3 µg RNA per sample was used as input material for the RNA sample preparations. Sequencing libraries were generated using NEBNext® Ultra™ RNA Library Prep Kit for Illumina® (NEB, USA) following the manufacturer's recommendations, and index codes were added to attribute sequences to each sample. Briefly, mRNA was purified from total RNA using poly-T oligo-attached magnetic beads. Fragmentation was carried out using divalent cations under elevated temperature in NEBNext First Strand Synthesis Reaction Buffer (5X). First-strand cDNA was synthesized using a random hexamer primer and

M-MuLV Reverse Transcriptase (RNase H⁻). Second-strand cDNA synthesis was subsequently performed using DNA Polymerase I and RNase H. Remaining overhangs were converted into blunt ends via the exonuclease/polymerase activities. After adenylation of the 3' ends of DNA fragments, NEBNext Adaptor with a hairpin loop structure was ligated to prepare for hybridization. To select cDNA fragments with lengths of 150~200 bp (preferentially), the library fragments were purified with AMPure XP system (Beckman Coulter, Beverly, USA). Then, 3 µl USER Enzyme (NEB, USA) was used with size-selected, adaptor-ligated cDNA at 37°C for 15 min followed by 5 min at 95°C before PCR. Then PCR was performed with Phusion High-Fidelity DNA polymerase, Universal PCR primers and Index (X) Primer. Finally, PCR products were purified (AMPure XP system) and library quality was assessed on the Agilent Bioanalyzer 2100 system. The clustering of the index-coded samples was performed on a cBot Cluster Generation System using TruSeq PE Cluster Kit v3-cBot-HS (Illumina) according to the manufacturer's instructions. After cluster generation, the library preparations were sequenced on an Illumina HiSeq platform and 125 bp/150 bp paired-end reads were generated. Quality-filtered Illumina reads of the reference data are available in the NCBI BioProject database (<https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJEB33121>) [59].

Gene Annotation and Differential Gene Expression Analysis

All expressed genes were functionally annotated using the following databases: KEGG, Nr, Swiss-Prot, Tremble, KOG, GO, and Pfam. Gene expression levels were measured using FPKM (fragments per kilobase of transcript per million mapped reads) values based on the number of uniquely mapped reads[60]. For genes with more than one alternative transcript, the longest transcript was selected to calculate the FPKM value. Differential expression analysis of the three groups was performed using the DESeq R package (1.22.2). DESeq provides statistical routines for determining differential expression in digital gene expression data using a model based on the negative binomial distribution. The resulting P values were adjusted using the Benjamini-Hochberg approach for controlling the false discovery rate (FDR). Genes with an adjusted P value < 0.05 found by DESeq were considered differentially expressed.

Validation of RNA-seq Data by qPCR

Total RNA was reverse transcribed to generate first strand cDNA using an Evo M-MLV RT for PCR kit (Accurate Biotechnology, Hunan, Co., Ltd.). The *α-tubulin* was used as an internal control for normalization[61]. qPCR was performed on a qPCR instrument (BIORAD CFX96 real-time PCR detection system), and real-time PCRs were performed using a SYBR[®] Green Premix Pro TaqHS qPCR kit (Accurate Biotechnology, Hunan, Co., Ltd.). The 2^{-ΔΔCT} method[62] was used for quantitative analysis of the relative expression levels. The primers used in the qPCR analysis are shown in Table 3.

Correlation Analysis between Metabolome and Transcriptome Data

Pearson correlation coefficients were calculated for metabolome and transcriptome data integration. In this study, log conversion of data was performed uniformly before analysis. For the joint analysis between the metabolome and transcriptome, the cor program from R was used, and the screening

criterion was a Pearson correlation coefficient (PCC) > 0.8. The relationships between metabolomic and transcriptomic data were visualized by using Cytoscape.

Abbreviations

bHLH: basic helix-loop-helix; CHS: chalcone synthesis; F3H: flavanone 3-hydroxylase; F3'H : flavonoid 3'-hydroxylase; F3'5'H: flavonoid 3'5'-hydroxylase; DFR: dihydroflavonol 4-reductase, ANS: anthocyanidin reductase; 3GT: UDP-glucose: flavonoid 3-O-glucosyltransferase; GST: glutathione S-transferase; MRP multidrug resistance-associated protein; MATE: multidrug and toxic compound extrusion; TT2: TRANSPARENT TESTA 2; CAM: crassulacean acid metabolism; RHSCC: Royal Horticulture Society Color Chart; GR: Green sample; RE: red sample; Rc: red cell; PS I: photosystem I; PS II : photosystem II; LHC: the light harvesting complex; DPMs: differentially produced metabolites; DEGs: differentially expressed genes; PCC: Pearson correlation coefficient; DHQ: dihydroquercetin; DHM: dihydromyricetin; DHK: Dihydrokaempferol

Declarations

Acknowledgements

We sincerely thank Hu Zeng, Li Yang and Wenhan Hu (Wuhan Metware Biotechnology Co., Ltd. Wuhan, China) for the technical assistance.

Availability of data and materials

All data generated or analyzed during this study are included in this published article and additional files.

Funding

This study was supported by grants from National Natural Science Foundation of China (31971704; 31770743).

Authors' contributions

XZXZ conceived the study and designed the experiments. XZXZ, MQM, YBX YHH carried out the experiments. XZXZ, HH, LJW, LJF, JHL, MOA and HLZ analyzed the data. XZXZ wrote the manuscript and draw charts and figures. XL, LXS, ZH, YHH and JM guided and revised the article. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. An X, Tian Y, Chen K, Wang X, Hao Y. The apple WD40 protein MdTTG1 interacts with bHLH but not MYB proteins to regulate anthocyanin accumulation. *J PLANT PHYSIOL*. 2012;169(7):710–7.
2. Zhuang W, Wang H, Liu T, Wang T, Zhang F, Shu X, Zhai H, Wang Z. Integrated physiological and genomic analysis reveals structural variations and expression patterns of candidate genes for colored- and green-leaf poplar. *Sci Rep*. 2019; 9(1).
3. Li H, Yang Z, Zeng Q, Wang S, Luo Y, Huang Y, Xin Y, He N. Abnormal expression of bHLH3 disrupts a flavonoid homeostasis network, causing differences in pigment composition among mulberry fruits. *Hortic Res*. 2020;7:83.
4. Chen L, Hu B, Qin Y, Hu G, Zhao J. Advance of the negative regulation of anthocyanin biosynthesis by MYB transcription factors. *PLANT PHYSIOL BIOCH*. 2019;136:178–87.
5. Sun X, Li L, Pei J, Liu C, Huang L. Metabolome and transcriptome profiling reveals quality variation and underlying regulation of three ecotypes for *Cistanche deserticola*. *PLANT MOL BIOL*. 2020;102(3):253–69.
6. Dubos C, Stracke R, Grotewold E, Weisshaar B, Martin C, Lepiniec L. MYB transcription factors in *Arabidopsis*. *TRENDS PLANT SCI*. 2010;15(10):573–81.
7. Das PK, Shin DH, Choi S, Park Y. Sugar-hormone cross-talk in anthocyanin biosynthesis. *MOL CELLS*. 2012;34(6):501–7.
8. Xu W, Grain D, Bobet S, Gourrierc JL, Thévenin J, Kelemen Z, Lepiniec L, Dubos C. Complexity and robustness of the flavonoid transcriptional regulatory network revealed by comprehensive analyses of MYB-bHLH-WDR complexes and their targets in *Arabidopsis* seed. *NEW PHYTOL*. 2014;202(1):132–44.
9. Zhao J, Dixon RA. MATE Transporters Facilitate Vacuolar Uptake of Epicatechin 3'-O-Glucoside for Proanthocyanidin Biosynthesis in *Medicago truncatula* and *Arabidopsis*. *Plant Cell*. 2009;21(8):2323–40.
10. Marrs KA, Alfenito MR, Lloyd AM, Walbot V. A glutathione S-transferase involved in vacuolar transfer encoded by the maize gene *Bronze-2*. *NATURE*. 1995;375(6530):397–400.
11. Mueller LA, Goodman CD, Silady RA, Walbot V. AN9, a *Petunia* Glutathione S-Transferase Required for Anthocyanin Sequestration, Is a Flavonoid-Binding Protein1. *PLANT PHYSIOL*. 2000;123(4):1561–70.
12. Nesi N, Jond C, Debeaujon I, Caboche M, Lepiniec L. The *Arabidopsis* TT2 Gene Encodes an R2R3 MYB Domain Protein That Acts as a Key Determinant for Proanthocyanidin Accumulation in

- Developing Seed. *Plant Cell*. 2001;13(9):2099.
13. Baudry A, Heim MA, Dubreucq B, Caboche M, Weisshaar B, Lepiniec L. TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and proanthocyanidin biosynthesis in *Arabidopsis thaliana*. *Plant J*. 2004;39(3):366–80.
 14. Mehrtens F, Kranz H, Bednarek P, Weisshaar B. The *Arabidopsis* Transcription Factor MYB12 Is a Flavonol-Specific Regulator of Phenylpropanoid Biosynthesis. *PLANT PHYSIOL*. 2005;138(2):1083–96.
 15. Czempl S, Stracke R, Weisshaar B, Cordon N, Harris NN, Walker AR, Robinson SP, Bogs J. The Grapevine R2R3-MYB Transcription Factor VvMYB11 Regulates Flavonol Synthesis in Developing Grape Berries. *PLANT PHYSIOL*. 2009;151(3):1513–30.
 16. Schrader LEKC. Effects of High Temperature and High Solar Irradiance on Sunburn, Quality, and Skin Pigments of Apple Fruit. *Acta Horticulturae*. 2011(903):1025–1039.
 17. Ekici L, Simsek Z, Ozturk I, Sagdic O, Yetim H. Effects of Temperature, Time, and pH on the Stability of Anthocyanin Extracts: Prediction of Total Anthocyanin Content Using Nonlinear Models. *FOOD ANAL METHOD*. 2014;7(6):1328–36.
 18. Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y, Nishizawa T, Matsuda F, Kojima M, Sakakibara H, Shinozaki K, et al. Enhancement of oxidative and drought tolerance in *Arabidopsis* by overaccumulation of antioxidant flavonoids. *Plant J*. 2014;77(3):367–79.
 19. Shen X, Zhao K, Liu L, Zhang K, Yuan H, Liao X, Wang Q, Guo X, Li F, Li T. A Role for PacMYBA in ABA-Regulated Anthocyanin Biosynthesis in Red-Colored Sweet Cherry cv. Hong Deng (*Prunus avium* L.). *PLANT CELL PHYSIOL*. 2014;55(5):862–80.
 20. Zoratti L, Karppinen K, Luengo Escobar A, Hännigsmann H, Jaakola L. Light-controlled flavonoid biosynthesis in fruits. *FRONT PLANT SCI*. 2014; 5.
 21. Martin CE. Physiological ecology of the Bromeliaceae. *The Botanical Review*. 1994;60(1):1–82.
 22. Xue Y, Ma J, He Y, Yu S, Lin Z, Xiong Y, Rafique F, Jiang F, Sun L, Ma M, et al. Comparative transcriptomic and proteomic analyses of the green and white parts of chimeric leaves in *Ananas comosus* var. *bracteatus*. *Peer J*. 2019;7:e7261.
 23. Smith JAC, Winter K. Taxonomic Distribution of Crassulacean Acid Metabolism. *Ecological Studies*. 1996:427–436.
 24. Borland AM, Barrera Zambrano VA, Ceusters J, Shorrocks K. The photosynthetic plasticity of crassulacean acid metabolism: an evolutionary innovation for sustainable productivity in a changing world. *NEW PHYTOL*. 2011;191(3):619–33.
 25. Li X, Kanakala S, He Y, Zhong X, Yu S, Li R, Sun L, Ma J. Physiological Characterization and Comparative Transcriptome Analysis of White and Green Leaves of *Ananas comosus* var. *bracteatus*. *PLOS ONE*. 2017;12(1):e169838.
 26. Lin Z, Xiong Y, Xue Y, Mao M, Xiang Y, He Y, Rafique F, Hu H, Liu J, Li X, et al. Screening and characterization of long noncoding RNAs involved in the albinism of *Ananas comosus* var. *bracteatus* leaves. *PLOS ONE*. 2019;14(11):e225602.

27. Wang LS, Shiraishi A, Hashimoto F, Aoki N, Shimizu K, Sakata Y. Analysis of Petal Anthocyanins to Investigate Flower Coloration of Zhongyuan (Chinese) and Daikon Island (Japanese) Tree Peony Cultivars. J PLANT RES. 2001;114(1):33–43.
28. Li J, Sakata Y, Wang L, Hashimoto F, Shiraishi A, Aoki N. Chemical taxonomy of the Xibei tree peony from China by floral pigmentation. J PLANT RES. 2004;117(1):47–55.
29. VAUGHN KC. WILSON KG. Development of Air Blisters in *Pilea cadieri* Gagnep. and Guillaumin. ANN BOT-LONDON. 1981; 48(4):467–72.
30. Smith GF, van Wyk AE. Systematic leaf anatomy of selected genera of southern African Alooideae (Asphodelaceae). S AFR J BOT. 1992;58(6):543.
31. Parida AK, Das AB, Mittra B. Effects of salt on growth, ion accumulation, photosynthesis and leaf anatomy of the mangrove, *Bruguiera parviflora*. Trees. 2004;18(2):167–74.
32. Wang F, Chen L, Chen H, Chen S, Liu Y. Analysis of Flavonoid Metabolites in Citrus Peels (*Citrus reticulata* “Dahongpao”) Using UPLC-ESI-MS/MS. MOLECULES. 2019;24(15):2680.
33. Li Q, Song J. Analysis of widely targeted metabolites of the euhalophyte *Suaeda salsa* under saline conditions provides new insights into salt tolerance and nutritional value in halophytic species. BMC PLANT BIOL. 2019; 19(1).
34. Ferreira KN. Architecture of the Photosynthetic Oxygen-Evolving Center. SCIENCE. 2004;303(5665):1831–8.
35. Liu Z, Yan H, Wang K, Kuang T, Zhang JGL. Crystal structure of spinach major light-harvesting complex at 2.72 Å resolution. NATURE. 2004;428(6980):287–92.
36. Amunts A, Drory O, Nelson N. The structure of a plant photosystem I supercomplex at 3.4 Å resolution. NATURE. 2007;447(7140):58–63.
37. Shirley JW. An Early Experimental Determination of Snell’s Law. AM J PHYS. 1951;19(9):507–8.
38. Chiu MH, Lee JY, Su DC. Refractive-index measurement based on the effects of total internal reflection and the uses of heterodyne interferometry. Appl Opt. 1997;36(13):2936–9.
39. Mathur J. Cell shape development in plants. TRENDS PLANT SCI. 2004;9(12):583–90.
40. Müller C, Riederer M. Plant Surface Properties in Chemical Ecology. J CHEM ECOL. 2005;31(11):2621–51.
41. Tanaka Y, Brugliera F, Kalc G, Senior M, Dyson B, Nakamura N, Katsumoto Y, Chandler S. Flower Color Modification by Engineering of the Flavonoid Biosynthetic Pathway: Practical Perspectives. Bioscience, biotechnology, and biochemistry. 2014; 74(9):1760–1769.
42. Tsuda S, Fukui Y, Nakamura N, Katsumoto Y, Yonekura-Sakakibara K, Fukuchi-Mizutani M, Ohira K, Ueyama Y, Ohkawa H, Holton TA, et al. Flower color modification of *Petunia hybrida* commercial varieties by metabolic engineering. Plant biotechnology (Tokyo Japan). 2004;21(5):377–86.
43. Forkmann G, Ruhnau B. Distinct Substrate Specificity of Dihydroflavonol 4-Reductase from Flowers of *Petunia hybrida*. Zeitschrift Für Naturforschung C. 1987;42(9–10):1146–8.

44. Katsumoto Y, Fukuchi-Mizutani M, Fukui Y, Brugliera F, Holton TA, Karan M, Nakamura N, Yonekura-Sakakibara K, Togami J, Pigeaire A, et al. Engineering of the Rose Flavonoid Biosynthetic Pathway Successfully Generated Blue-Hued Flowers Accumulating Delphinidin. *PLANT CELL PHYSIOL.* 2007;48(11):1589–600.
45. Springob K, Nakajima J, Yamazaki M, Satio K. Recent advances in the biosynthesis and accumulation of anthocyanins. *NAT PROD REP.* 2003;20(3):288–303.
46. Petroni K, Tonelli C. Recent advances on the regulation of anthocyanin synthesis in reproductive organs. *PLANT SCI.* 2011;181(3):219–29.
47. Aharoni, Vos. Wein. The strawberry FaMYB1 transcription factor suppresses anthocyanin and favonol accumulation in transgenic tobacco. *Plant J.* 2001;28(3):319–32.
48. Stracke R, Favory J, Gruber H, Bartelniewoehner L, Bartels S, Binkert M, Funk M, Weisshaar B, Ulm R. The Arabidopsis bZIP transcription factor HY5 regulates expression of the PFG1/MYB12 gene in response to light and ultraviolet-B radiation. *Plant, Cell & Environment.* 2010; 33.
49. Nesi N, Jond C, Debeaujon I, Caboche M, Lepiniec L. The Arabidopsis TT2 gene encodes an R2R3 MYB domain protein that acts as a key determinant for proanthocyanidin accumulation in developing seed. *Plant Cell.* 2001;13(9):2099–114.
50. Schaart JG, Dubos C, Irene RDLF, Adèle MMLV, Ric CHDV, Jonker HH, Xu W, Routaboul J, Lepiniec L, Bovy AG. Identification and characterization of MYB-bHLH-WD40 regulatory complexes controlling proanthocyanidin biosynthesis in strawberry (*Fragaria × ananassa*) fruits. *NEW PHYTOL.* 2013;197(2):454–67.
51. Springob INJ. M Y. Recent advances in the biosynthesis and accumulation of anthocyanins. *NAT PROD REP.* 2003;20(3):288–303.
52. Lichtenthaler HK. M R, J S. Two independent biochemical pathways for isopentenyl diphosphate and isoprenoid biosynthesis in higher plants. *PHYSIOL PLANTARUM.* 1997;101(3):643–52.
53. Voss D. Relating Colorimeter Measurement of Plant color to the Royal-Horticultural-Society Color Chart. *HORTSCIENCE.* 1992;27(12):1256–60.
54. McGuire GR. Reporting of Objective Color Measurements. *HORTSCIENCE.* 1992;27(12):1254–5.
55. Jean-François G. Colour effects of co-pigmentation of anthocyanins revisited-1. A colorimetric definition using the CIELAB scale. *FOOD CHEM.* 1998;63(3):409–15.
56. Arnon DI. Copper Enzymes in Isolated Chloroplasts. Polyphenoloxidase in Beta Vulgaris. *PLANT PHYSIOL.* 1949;24(1):1–15.
57. Lee J, Durst RW, Wrolstad RE, Eisele TGMM. Determination of Total Monomeric Anthocyanin Pigment Content of Fruit Juices, Beverages, Natural Colorants, and Wines by the pH Differential Method: Collaborative Study. *J AOAC INT.* 2005;88(5):1269–78.
58. Rafaela G, González M. Antioxidant activity in banana peel extracts: Testing extraction conditions and related bioactive compounds. *FOOD CHEM.* 2010;119(3):1030–9.

59. Chen L, VanBuren R, Paris M, Zhou H, Zhang X, Wai CM, Yan H, Chen S, Alonge M, Ramakrishnan S, et al. The bracteatus pineapple genome and domestication of clonally propagated crops. NAT GENET. 2019;51(10):1549–58.

60. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. NAT BIOTECHNOL. 2010;28(5):511–5.

61. Ruixue L, Screening. Cloning and Functional Identification of POR gene in the Albino of Ananas comosus var. bracteatus Leaves Using VIGS Technique. Sichuan: Sichuan Agricultural University;; Cheng Du; 2018.

62. Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. METHODS. 2001;25(4):402–8.

Tables

Table 1. DPMs in anthocyanin-related pathways between GR and RE.

Name	GR	RE	VIP	Log ₂ FC	Type	Cpd_ID	Related Map
3-O-p-Coumaroyl quinic acid	3.69E +05	1.09E +06	1.19E +00	1.56E +00	up	C12208	ko00940, ko00941
Caffeic acid	1.29E +05	5.25E +06	1.13E +00	5.34E +00	up	C01197	ko00940
Ferulic acid	1.47E +05	4.23E +06	1.18E +00	1.52E +00	up	C01494	ko00940
Caffeic aldehyde	1.66E +05	5.33E +04	1.19E +00	-	down	C10945	ko00940
Chlorogenic acid	1.28E +06	4.04E +06	1.19E +00	1.65E +00	up	C00852	ko00940, ko00941
Coniferin	1.17E +06	2.60E +06	1.18E +00	1.15E +00	up	C00761	ko00940
Hesperetin	8.50E +03	4.14E +03	1.11E +00	-	down	C01709	ko00941
Apigenin-8-C-Glucoside	9.70E +06	4.63E +06	1.13E +00	-	down	C01460	ko00941
Delphinidin-3,5-O-diglucoside	1.74E +05	8.57E +05	1.19E +00	2.30E +00	up	C16312	ko00942
Delphinidin-3-O-glucoside	4.98E +05	2.71E +06	1.18E +00	2.45E +00	up	C12138	ko00942
Cyanidin-3-O-rutinoside	1.82E +05	1.73E +06	1.19E +00	3.25E +00	up	C08620	ko00942
Pelargonidin-3,5-diglucoside	1.93E +05	1.16E +06	1.19E +00	2.59E +00	up	C08725	ko00942
Malvidin-3,5-O-diglucoside	1.26E +04	2.57E +04	1.17E +00	1.03E +00	up	-	-
Cyanidin-3,5-O-diglucoside	3.47E +07	1.56E +08	1.19E +00	2.17E +00	up	C08639	ko00942
Columns 2 and 3 represent metabolite peak area average values of different samples.							

Table 2. Flavonoid biosynthesis-related genes.

Name	ID	GR	RE	Log ₂ FC	P value	Type	KEGG_ID
PAL	Aco020618.1.path1*	12.09	5.42	-1.283	4E-05	down	K10775
	Aco007727.1.path1	0.46	0.85	0.755	0.02258	-	
	Aco010091.1.path1	0.25	0.25	-0.159	0.82465	-	
	Aco013943.1.path1	45.43	87.97	0.845	0.00044	-	
4CL	Aco006521.1.path1	91.38	95.52	-0.058	0.74187	-	K01904
	Aco011626.1.path1	1.43	1.68	0.137	0.77702	-	
	Aco017002.1.path1	1.21	1.51	0.213	0.5899	-	
	Aco017294.1.path1	0.06	0	-1.935	0.6324	-	
	Aco019974.1.path1	1.09	1.79	0.618	0.41004	-	
	Aco019975.1.path1	45.62	51.09	0.043	0.78884	-	
	Aco025129.1.path1	30.5	53.74	0.698	0.28868	-	
CHI	Aco012547.1.path1*	26.16	58.11	1.051	0.00038	up	K01859
	Aco014232.1.path1*	45.47	113.57	1.217	2.9E-06	up	
	Aco022584.1.path1	452.78	448.54	-0.13	0.59715	-	
	Aco025098.1.path1	65.10	62.28	-0.176	0.28868	-	
CHS	Aco016200.1.path1	149.59	293.62	0.865	4.5E-06	-	K00660
F3H	Aco018609.1.path1*	4.15	18.20	2.032	3.5E-11	up	K00475
	Aco027900.1.path1	0.11	0.12	-0.036	0.9743	-	
F3'H	Aco003857.1.path1*	27.99	61.63	1.027	3.7E-13	up	K05280
	Aco003885.1.path1	11.03	23.60	0.982	4.2E-10	-	
	Aco003886.1.path1	0.05	0.28	2.272	0.35361	-	
F3'5'H	Aco017169.1.path1	5.70	7.98	0.371	0.12805	-	K13083
	Aco013254.1.path1	0.23	0.72	1.483	0.19733	-	
DFR	Aco006769.1.path1*	3.51	16.73	2.151	1.8E-07	up	K13082
	Aco014886.1.path1	0.19	0.47	1.113	0.41798	-	
	Aco014887.1.path1	7.43	10.95	0.439	0.07125	-	
FLS	Aco000551.1.path1	0	0.04	2.554	0.25109	-	K05278
Columns 3 and 4 represent average FPKM values of GR and RE* represents DEGs.							

Name	ID	GR	RE	Log ₂ FC	P value	Type	KEGG_ID
	Aco011051.1.path1*	3.34	1.30	-1.475	0.00043	down	
	Aco001560.1.path1	20.24	23.25	0.078	0.73827	-	
	Aco002683.1.path1	15.29	25.77	0.642	0.00261	-	
	Aco007045.1.path1	0.03	0	-2.591	0.51669	-	
	Aco012471.1.path1	0.11	0.08	-0.565	0.73375	-	
ANS	Aco007046.1.path1	0.89	0.20	-2.26	0.00911	-	K05277
3GT	Aco012126.1.path1*	20.75	87.45	1.973	1.6E-12	up	K12930
ABCA3	Aco006845.1.path1*	13.99	38.12	1.324	4.9E-09	up	K05643
	Aco007263.1.path1	42.66	32.95	-0.506	0.22723	-	
	Aco008459.1.path1	23.40	25.61	0.014	0.92605	-	
	Aco014363.1.path1	9.14	11.88	0.259	0.22712	-	
MYB82	Aco023266.1.path1*	0.25	1.69	2.666	0.00044	up	K09422
MYB5	Aco023263.1.path1*	5.02	40.80	2.924	1.5E-15	up	
Columns 3 and 4 represent average FPKM values of GR and RE* represents DEGs.							

Table 3. qPCR primer sequences of eight genes.

Gene Name and ID	F Primer (5'-3')	R Primer (5'-3')
PAL(Aco020618.1.path1)	TGAAGTCCGCCGTCAAGA	GCTCCTCCCTAACGAACCTGTA
CHS(Aco016200.1.path1)	GTGCCGCTGCTGTTATCC	TGTTCCCGAACTCGCTCA
CHI(Aco012547.1.path1)	GATTCCTTCTTTGAGGCACT	GTTTGGAAAACTCGGCAAT
F3'H (Aco003857.1.path1)	GGGGAGGATTACAGAAACG	CCCTTGAGGTCCACATTTT
DFR(Aco006769.1.path1)	AGCATCATCCACCTCTTG	TCGTCGTCATCGGTATCAG
3GT(Aco012126.1.path1)	AGCACGTTTCGGCTTAGGAC	AGAGGCAGTTGCTCGGGTC
ABCA3(Aco006845.1.path1)	AGGCGATGGGTTGGCTATGC	AGAAGAAAGTTGCGGCGAGA
MYB5(Aco023263.1.path1)	AGATGAGACCAAGAACCCAC	CCCGCTATCAAAGACCAC

Figures

Figure 1

Color data of adaxial and abaxial surfaces of *Ananas comosus* var. *bracteatus* chimera leaves. (A) CIE color values and coordinate flats of the adaxial surfaces of GR and RE samples. The GR circle represents $L^*=46$, $a^*=-5$, and $b^*=15$; and the RE circle represents $L^*=68$, $a^*=23$, and $b^*=16$. (B) CIE color values and coordinate flats of abaxial surfaces of the GR and RE samples. The GR circle represents $L^*=56$, $a^*=-6$, and $b^*=16$; and the RE circle represents $L^*=77$, $a^*=9$, and $b^*=17$. Error bars represent $\pm$ SE of biological triplicates. The circle represents the rounded number of average values from triplicates of each phenotypic sample.

Figure 2

Overview of pigment distribution in chimeric leaves. (A) Cross sections of GR central and junction parts showed the chlorophyll distribution in the green mesophyll cells. The dotted line represents the boundary of the two leaf parts. (B) Cross sections of RE central and marginal parts showed red pigments (anthocyanins) located in the epidermal cells and 1-2 layers of cells under the epidermis. The bars=100 μm . Rc, red cell.

Figure 3

Pigment contents and relative proportions in GR and RE. (A) Content of pigments in GR and RE. (B) Relative proportion of pigments in GR and RE. Error bars represent $\pm$ SE of biological triplicates.

Figure 4

Metabolite profiles between GR and RE. (A) Heatmap and hierarchical clustering showing differentially produced metabolites (DPMs). (B) PCA plot. (C) OPLS-DA plot. (D) Volcano plots of upregulated and downregulated metabolites. (E) KEGG enrichment plot of DPMs.

Figure 5

Gene expression and enrichment profiles of GR and RE samples of *A. comosus* var. *bracteatus*. (A) Heatmap and hierarchical clustering showing DEGs. (B) Volcano plots showing the number of DEGs in different regulation models. (C) KEGG pathway enrichment plots. (D) KEGG classification showing enriched pathways. (E) Number of DEGs in the GO classification.

Figure 6

Transcription levels of eight selected genes related to anthocyanin biosynthesis between GR and RE samples. Error bars represent $\pm$ SE of biological triplicates.

Figure 7

Metabolites and genes involved in the biosynthesis of anthocyanin in *A. comosus* var. *bracteatus*. Anthocyanins were synthesized through three pathways: phenylpropanoid biosynthesis, flavonoid biosynthesis and anthocyanin biosynthesis. The color scale of metabolites indicates the log₂FC between the two types of samples. The color scale of genes indicates the log₂ transformation value of FPKM. The

percentage represents the ratio of anthocyanins among all detected anthocyanins in GR and RE. The gray circles and hexagons indicate undetectable metabolites and genes, respectively.

Figure 8

Profiles of the correlation analysis of anthocyanin-related genes and metabolites. (A) Nine-quadrant flat. (B) Correlations between anthocyanin-related genes and metabolites.

Figure 9

Chimeric leaves of *Ananas comosus* var. *bracteatus* used in this study. (A) *A. comosus* var. *bracteatus* chimeric plant. Bar=5 cm. (B) Green central parts (GR) and red edges (RE) from chimeric leaves. Bar = ~1 cm.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfiles.rar](#)