Biomimetic nanoparticles enabled by cascade cell membrane coating for direct cross-priming of T cells

Fangjie Chen

Shanghai Jiao Tong University

Lu Wang

Shanghai Jiao Tong University

Jinyao Liu (✉ jyliu@sjtu.edu.cn)

Shanghai Jiao Tong University <https://orcid.org/0000-0002-6044-2033>

Article

Keywords: biomimetic nanoparticles, dendritic cells, T cells

Posted Date: July 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-79607/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Small on November 27th, 2021. See the published version at <https://doi.org/10.1002/smll.202104402>.

1 **Biomimetic** nanoparticles enabled by cascade cell membrane coating
2 **for direct cross-priming of T cells**

3
4 Fangjie Chen¹, Lu Wang¹, Jinyao Liu^{1,2*}
5

6 ¹Shanghai Key Laboratory for Nucleic Acid Chemistry and Nanomedicine, Institute of
7 Molecular Medicine, State Key Laboratory of Oncogenes and Related Genes, Shanghai
8 Cancer Institute, Renji Hospital, School of Medicine, Shanghai Jiao Tong University,
9 Shanghai 200127, China.

10 ²Shanghai Key Laboratory of Gynecologic Oncology, Renji Hospital, School of
11 Medicine, Shanghai Jiao Tong University, Shanghai 200127, China.

12
13 *Corresponding: J.L. (jyliu@sjtu.edu.cn)

Abstract

Despite the activation of T lymphocytes by antigen-presenting cells is responsible for eliciting antigen-specific immune responses, their crosstalking suffers from temporospatial limitations and endogenous influencing factors, which restrict the generation of a strong antitumor immunity. Here, we describe the manipulation of cross-priming of T cells using biomimetic nanoparticles (BNs) enabled by cascade cell membrane coating. BNs are resulted from coating nanoparticulate substrates with cell membranes extracted from dendritic cells (DCs) that are pre-pulsed with cancer cell membrane-coated nanoparticles. With a DC membrane that presents an array of cancer cell membrane antigen epitopes, BNs inherit intrinsic membrane function of DCs. Strikingly, BNs can directly cross-prime T cells and provoke robust yet antigen-specific antitumor responses in multiple mouse models. Combination with clinical anti-programmed death-1 antibodies demonstrates a practical way of BNs to achieve desirable tumor regression and survival rate. This work spotlights the impact of nanoparticles on direct cross-priming of T cells and supports a unique yet modulate platform for boosting an effective adaptive immunity for immunotherapy.

1 **Introduction**

2 Increasing evidence has shown that immunotherapy is able to successfully fight various
3 classes of cancer by properly regulating the immune system^{1,2}. Training T lymphocytes
4 with appropriate antigen specificities stands for a central route to promote tumor
5 regression³. To elicit an antigen-specific antitumor immunity, the stimulation of
6 cytotoxic T cell populations by professional antigen-presenting cells (APCs) is critical⁴⁻
7 ⁶. Delivering antigen and adjuvant into APCs and reinfusing engineered antigen-
8 presenting APCs are two main strategies to stimulate tumor-specific T cell responses^{7,8}.
9 However, these approaches inevitably depend on the crosstalking between T cells and
10 APCs, as cross-priming of T cells relies on surface markers of APCs^{9,10}. Unfortunately,
11 numerous endogenous factors including susceptibility, restricted migration, immune
12 suppression, and metabolic constraints can influence the inherent function of APCs¹¹⁻
13 ¹³. In addition, the communication between APCs and T cells is often inadequate due
14 to temporal and spatial limitations, resulting in suboptimal antigen presentation and
15 insufficient activation of T cells¹⁴⁻¹⁶.

16 To tackle these difficulties, various artificial APCs that mimic the function of
17 endogenous counterparts have been engineered to express proper biological signals,
18 such as major histocompatibility complex (MHC) proteins and costimulatory
19 markers¹⁷⁻¹⁹. These artificial cells have been successfully exploited for engaging with
20 and activating T cells²⁰. In addition to cell-based artificial APCs, particulate APCs
21 anchoring with requisite ligands including CD28 agonists and epitope-loaded MHC on

1 their surface have emerged as elegant analogues to facilitate antigen presentation²¹⁻²⁴.
2 Despite the potential of artificial APCs for adaptive cell-based therapy and the
3 advantage of synthetic particles to penetrate lymph nodes after subcutaneous
4 administration, the construction of artificial systems is complex and time consuming²⁵⁻
5 ²⁷. Furthermore, epitope-embedded MHC multimer complexes can be engineered only
6 individual epitope of interest has been identified beforehand²⁸. Additionally,
7 introducing certain membrane proteins of APCs can solely confer with partial function
8 of the interaction with T cells, largely limiting their application²⁹. Therefore, alternative
9 methods that are facile and versatile yet capable of directly cross-priming T cells even
10 with an array of unidentified tumor-associated antigen epitopes are highly desirable.

11 Here, we report the use of biomimetic nanoparticles (BNs) enabled by cascade cell
12 membrane coating for directly manipulating cross-priming of T cells. Cancer cell
13 membranes with an array of membrane antigens are extracted and coated onto a
14 nanoparticulate substrate that can be easily devoured by dendritic cells (DCs) for
15 antigen processing and presentation. DC membranes presenting related antigen
16 epitopes are subsequently acquired and fused onto a nanoparticulate core to form BNs.
17 Exceptionally, BNs enabled by the cascade coating retain the entire membrane
18 composition and intrinsic function of DC membranes. Different from endogenous DCs,
19 direct cross-priming of T cells by BNs that bypass the need for regular antigen
20 processing and presentation elicits strong antigen-specific T cell responses. We
21 demonstrate the ability of BNs to boost an extraordinary tumor-specific immunity in

multiple tumor models, including ovalbumin expressing B16 (B16-OVA), HPV E6 and E7-expressing TC-1, and Hepa 1-6 tumor-bearing mice. In combination with a clinical immune checkpoint inhibitor, anti-programmed cell death-1 (α PD-1) antibodies, **BNs** evidence a practical path to enhance tumor growth inhibition and improve survival. Our work suggests a simple yet universal platform for the manipulation of the interaction with T cells and supports the potential of engineered materials to directly cross-priming T cells for immunotherapy.

Results and discussion

Preparation of **BNs by cascade cell membrane coating**

Cell membrane coating, which fuses synthetic cores with natural cell membranes, has emerged as a useful approach to endow coated substrates with foreign functions that are intrinsic unachievable³⁰⁻³². For instance, various nanoparticles coated with erythrocyte membranes exhibit prolonged blood circulation due to the presence of a “don’t eat me” signal of CD47-SIRP α on the surface³³. Furthermore, coated nanoparticles can play as decoys to bind and neutralize toxic molecules since the maintenance of the antigenic characteristics of the source cells, such as macrophage and neutrophils³⁴⁻³⁷. More recently, coating with engineered cancer cell membranes expressing MHC and costimulatory marker CD80 has been explored to mimic DCs for antigen presentation²⁶. As primary APCs, DCs execute a vital role for antigen processing and cross-priming of antigen-specific T cells, which can provoke an

1 adaptive immunity for cancer treatment^{38,39}. However, DC membranes presenting
2 identified antigen epitopes have rarely utilized to coat synthetic cores for stimulating
3 tumor-specific immune responses. To build DC membrane-coated nanoparticles, we
4 developed a technique of cascade cell membrane coating, which could display a set of
5 cancer cell associated peptide epitopes on the surface (Figure 1a). First, cell membranes
6 with multiple antigens were acquired from tumor cells⁴⁰ and subsequently fused onto a
7 polymeric nanoparticulate core to form cancer cell membrane-coated nanoparticles
8 (CCNPs). Then, DCs derived from bone marrow (BMDCs) were incubated with
9 CCNPs for antigen processing and presentation. Lastly, membranes displaying a cluster
10 of exclusive tumor-associated antigen epitopes were extracted from CCNPs-pulsed
11 BMDCs and coated onto a polymeric nanoparticulate substrate to generate DC
12 membrane-coated nanoparticles.

13 Transmission electron microscopy (TEM) images showed a clear core-shell
14 structure for CCNPs, in which each poly (lactic-co-glycolic acid) (PLGA)
15 nanoparticulate core was visualized surrounding with a cell membrane coating (Figure
16 1d and Figure S1a). Compared with bare PLGA nanoparticles (Ns), dynamic light
17 scattering (DLS) measurement indicated that the hydrodynamic diameter and surface
18 zeta potential of CCNPs separately increased by ~35 nm (Figure 1b) and ~6 mV (Figure
19 1c). It was worth noting that the membrane proteins were successfully retained after
20 coating onto PLGA nanoparticles (Figure 1e), which was in good agreement with
21 previous studies¹. After adding into BMDCs, CCNPs could be internalized rapidly, as

1 confirmed by confocal laser scanning microscopy (CLSM) images (Figure 1f and
2 Figure S1b). Fluorescent colocalization analysis indicated that CCNPs were devoured
3 in its entirety by DCs, as verified by a highly overlapping signals from the cores and
4 membrane coatings of the coated nanoparticles. As expected, typical membrane
5 antigens including the overexpressed OVA and main melanoma-associated
6 transmembrane protein glycoprotein 100 (gp100) of B16-OVA cells were processed
7 and presented by BMDCs (Figure 1g-i). To quantify epitopes presented on the cell
8 membrane, CCNPs with cell membrane proteins (CMP) of 40, 60, 80, 120, or 160 μg
9 were added into 1×10^6 DCs. An amount of 40 μg OVA complexed with
10 polyetherimide (PEI-OVA) was cultured as a control⁴¹. Importantly, OVA epitope
11 presented by CCNPs-matured DCs increased with the quantity of membrane proteins
12 (Figure 1h and i). Even pulsed with CCNPs that had a total protein amount of 40 μg ,
13 BMDCs exhibited near 5-times higher OVA epitope level on the surface, in contrast to
14 cells primed with equivalent PEI-complexed OVA. In consideration of the fact that
15 actual quantity of OVA delivered by CCNPs was lower than that of PEI-OVA,
16 enhanced presentation suggested that rather than complexed OVA, antigens embedded
17 in cell membranes could be processed more favorably by DCs. It was noted that
18 incubation with free OVA showed negligible corresponding epitope level on the surface
19 of BMDCs (Figure S1c). These results illustrated that coating nanoparticles with cancer
20 cell membranes represented an efficient way to deliver tumor-associated antigens.
21 Similarly, PLGA nanoparticles coated with DC membranes that were extracted from

1 B16-OVA cell membrane-coated nanoparticles (termed as **BDCNs**) appeared a typical
2 core-shell nanostructure (Figure 1j and Figure S3a). The hydrodynamic diameter and
3 surface zeta potential of **BDCNs** approximated to those of DC membrane vesicles
4 (Figure 1k and l). **Similar to other cell membrane-coated nanoparticles⁴², the colloidal**
5 **stabilities of cancer membrane-coated and dendritic cell membrane-coated**
6 **nanoparticles could maintain up to 14 days in both PBS and pure water (Figure S2).**
7 **The physicochemical properties of the nanoparticles were summarized in Table S1.**
8 Taken together, through translocation of B16-OVA cell membranes onto nanoparticles,
9 a **biomimetic** structure of cancer cells was occurred for interaction with DCs and
10 subsequent antigen processing and presentation. Take advantage of cascade cell
11 membrane coating, nanoparticles displaying **autologous** epitopes that were derived
12 from membrane antigens of cancer cells could be accomplished by coating with mature
13 DC membranes.

15 **BNs mediated cross-priming of T cells**

16 We investigated whether **BDCNs** could execute the function of directly activating T
17 cells *in vitro*. To study the interaction with T cells, the binding of **BNs** with the surface
18 receptors of T cells was analyzed by coculture. CD8⁺ T cells were isolated from the
19 spleen of C57BL/6 mouse and incubated with **BDCNs**. To examine the specific
20 adhesion with T cells, both Ns and Ns coated with red blood cell membranes (RBC-Ns)
21 **or cell membranes extracted from DCs without pre-pulsing with CCNPs** were used as
22 controls. As viewed by confocal imaging, T cells were surrounded with numerous
23 **BDCNs**, as shown by fluorescence overlap between the cell membranes and

nanoparticles (Figure 2a and Figure S3b). In comparison with limited non-specific binding of the control nanoparticles, the enhanced fluorescence intensity reflected the specificity of BDCNs to efficiently bind with T cells, which was further supported by flow cytometric analysis (Figure 2b). The activation of T cells could be explained by tightly regulated T cell receptor recognition of antigenic peptides in complex with MHC glycoproteins presented on DC membranes. Then, carboxy fluorescein succinimidyl ester (CFSE) dilution was performed to detect T cell proliferation. CD8⁺ T cells were incubated individually with Ns, BMDCs and BDCNs. PLGA nanoparticles coated with membranes derived from DCs without pretreatment (DCNs) or with PEI-OVA treatment (ODCNs) were prepared as controls. In striking contrast, BDCNs induced significantly increased proliferation of CD8⁺ T cells than any other groups. As given in Figure 2c, BDCNs showed ~2-3 times higher activation of CD8⁺ T cell proliferation in comparison with those of Ns, BMDCs and DCNs, which confirmed the preferential ability of BDCNs to activate CD8⁺ T cells. To further validate whether BDCNs could initiate activation of antigen-specific T cells, OVA positive CD8⁺ T cells were labelled using OVA (SIINFEKL) tetramer (tet) antibody²⁹. Other than nanoparticles coated with membranes that were extracted from DCs without antigen pulsing, BDCNs could directly prime remarkable proliferation of OVA-specific T cells, which was near 10-fold higher than that of DCNs (Figure 2d and e). We then employed antigen-specific epitopes based on OVA and gp100 to assess *in vivo* T cell production after intravenous injection of BDCNs. It was worth mentioning that comparable biodistribution to other commonly used nanoparticles was observed after intravenous injection of BDCNs⁴³. As expected, these nanoparticles accumulated in the spleen, a major organ for DCs residence. Moreover, the accumulation of BDCNs in tumor increased with time post-injection extending from 0 to 6 hours, as demonstrated by the increase of fluorescence intensity in the sectioned tumor tissue (Figure S4). Mice were intravenously administered with 3 doses of phosphate buffered saline (PBS), Ns, DCNs, ODCNs, or BDCNs on day 0, 5, and 10, respectively (Figure 2f). Spleen tissues were collected and

homogenized on day 15 to analyze the percentage of antigen-specific T cells by flow cytometry. Expectedly, the induction of OVA (Figure 2g and i) and gp100 (Figure 2h and j) positive T cells by BDCNs was far exceeded all control groups. In addition, significantly elevated levels of Ki67 expression and interferon-gamma (IFN- γ) production in CD3⁺ T cells further confirmed that BDCNs promoted the total proliferation and activation of T cells (Figure 2k-m). Interestingly, the similar expression of costimulatory markers CD80 and CD86 on DCs indicated that BDCNs had limited influence on DC maturation (Figure S3e and f).

The comparison of activation efficacy between DCs pulsed with PEI-OVA and ODCNs was also studied. As shown in Figure S3g and h, mice vaccinated with ODCNs showed significantly elevated levels of CD3⁺CD8⁺ and OVA-specific T cells, which separately increased by near 2-times compared with mice administrated with DCs. In order to measure their susceptibility to environmental conditions, BDCNs were incubated with CD8⁺ T cells in presence of 10 mM lactic acid, a tumor-derived metabolite that can influence dendritic cell activation and antigen expression⁴⁴. Interestingly, BDCNs showed 1.7-fold higher proliferation of CD8⁺ T cells than that of DCs after incubation for 3 days, demonstrating their less susceptible to the insult of lactic acid (Figure S3c). The insusceptibility of BDCNs was further validated by near 1.4-fold higher activation efficacy compared to DCs after 24 hours storage at 4 °C (Figure S3d). These data implied the superiorities of BNs over living DCs particularly in activation of T cells, insusceptibility to the tumor microenvironment, and storage condition.

Antigen-specific antitumor responses

We next examined the function of BDCNs in tumor-bearing mice, which were subcutaneously inoculated with B16-OVA in the right hind leg. Mice were daily administered with BDCNs at predetermined time points⁴⁵⁻⁴⁷ (Figure 3a). Experimental

1 endpoint was defined as the tumor size of PBS group exceeded 2000 mm³. Peripheral
2 blood, spleen and tumor were sampled for further analysis. Mice treated with PBS and
3 equivalent nanoparticles of Ns, DCNs, or ODCNs were individually used as controls.
4 Encouragingly, mice administered with BDCNs emerged significantly increased
5 percentages of CD8⁺ and Ki67⁺ in CD3⁺ T cells in the spleen, which disclosed systemic
6 facilitation of T cell proliferation (Figure 3b and c). Moreover, the higher levels of
7 CD80, CD86 and MHC-II in splenic DCs represented a more mature phenotype induced
8 by BDCNs (Figure 3d-f). On the contrary, decreased CD3⁺CD4⁺ T cells and increased
9 CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio in comparison to the control groups of PBS, Ns, DCNs
10 and ODCNs might indicate that immunosuppressive environment was relieved (Figure
11 3g and S5a). Furthermore, BDCNs could elevate IFN- γ production from T cells in the
12 spleen tissue (Figure 3h). Particularly, the increase of OVA tet⁺ T cells demonstrated
13 that BDCNs elicited immune responses in an antigen-specific manner (Figure 3i).
14 Consistently, appropriate priming of tumor-infiltrating lymphocytes mediated by
15 BDCNs was significantly higher than all the controls (Figure 3j-p). Expectedly,
16 negligible CD4⁺CD3⁺ T cells were found in tumor site after vaccination with BDCNs
17 (Figure S5b). The higher concentrations of IFN- γ (Figure 3q) and TNF- α (Figure 3r) in
18 serum sampled from BDCNs-treated mice further revealed the elicitation of a robust
19 systemic immune response. Additionally, in contrast to other groups,
20 immunofluorescence staining images depicted more infiltrated CD8⁺ T cells and
21 upgraded cytokine production (Figure 3u and Figure S5c) in the tumor tissue of

1 BDCNs-treated mice. Moreover, immunohistochemistry staining images implied
2 improved accumulation of CD3⁺ T cells and CD8⁺ T cells in tumor site after treatment
3 with BDCNs (Figure S6). The promotion of potent immune responses was further
4 supported by the most efficient inhibition of tumor growth after treatment with BDCNs
5 (Figure 3s, t and Figure S5d). Briefly, BNs could generate a strong antigen-specific
6 antitumor immunity in an aggressive yet OVA-expressing melanoma model.

7 A common tumor model of Hepa 1-6 hepatocellular carcinoma was further
8 developed to verify the generality of BNs for utilization in other cancers without
9 identified antigen epitopes. BNs were prepared by coating PLGA nanoparticles with
10 cell membranes extracted from mature DCs that were pre-pulsed with Hepa 1-6 cell
11 membrane-coated nanoparticles and defined as HDCNs. The tumor-bearing mice were
12 similarly treated, as illustrated in Figure 4a. Consistent with results obtained in B16-
13 OVA tumor model, the percentages of CD8⁺ and Ki67⁺ in CD3⁺ T cells (Figure 4b and
14 c) and the levels of costimulatory signals CD80, CD86 and MHC-II in DCs were
15 remarkably upgraded in the spleen sampled from mice injected with HDCNs, in
16 comparison with the controls of PBS, Ns, and DCNs (Figure 4d-f). The production of
17 IFN- γ from T cells remained the highest level (Figure 4g), while CD3⁺CD4⁺ T cells
18 were downregulated and CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio was upregulated after
19 treatment with HDCNs (Figure 4h and i). Together with the upgraded secretion of TNF-
20 α (Figure 4j) and IFN- γ (Figure 4k) in the serum, these data claimed that HDCNs could
21 initiate a stronger systemic tumor-specific cytotoxic T lymphocyte (CTL) response in

contrast to the controls. In parallel, the proportions of tumor-infiltrating CTLs (Figure 4l), Ki67⁺ (Figure 4m), DC mature markers (Figure 4n-p), CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio (Figure 4q) and inflammatory cytokines (Figure 4t and Figure S7b) were the highest in the HDCNs treatment group, whereas the population of CD3⁺CD4⁺ T cells maintained the lowest in all experimental groups (Figure S7a). The tumor growth after treatment with HDCNs retarded notably, with part of the tumors eradicated completely, suggesting the stimulation of a potent antitumor response (Figure 4r and s). Further supported by haematoxylin eosin (H&E) staining (Figure S7c), the results found in Hepa 1-6 tumor model reasoned that BNs could be exploited to treat tumors in absence of identified antigens.

Cross-validation of BNs-leveraged antitumor immunity

We further cross-validated the antitumor immunity in B16-OVA and HPV E6 and E7-expressing TC-1 tumor models to identify the specific and autologous characteristic of BNs. Similarly, BNs were fabricated by fusing PLGA nanoparticles with cell membranes sampled from BMDCs that were pre-treated with TC-1 cell membrane-coated nanoparticles and abbreviated as TDCNs. Mice inoculated with B16-OVA cells were treated with TDCNs, while TC-1 tumor-bearing mice were inversely injected with BDCNs (Figure 5a). Both PBS and the corresponding BNs were used as controls. In B16-OVA tumor model, the application of TDCNs displayed negligible tumor growth inhibition, which was comparable to that of PBS control (Figure 5b and c). On the

contrary, BDCNs showed notable tumor regression, which resulted in an extension in median survival time by 12 days (Figure 5d). Rather than TDCNs, the distinct potency of BDCNs to inhibit B16-OVA tumor growth explained that the elicitation of an effective tumor-specific immunity depended on the presence of autologous peptide epitopes on the surface. In TC-1 tumor model, BDCNs lost their ability to suppress tumor growth and provided neglectable survival benefit over PBS control (Figure 5e and f). As expected, TDCNs slowed the growth of TC-1 tumor and prolonged the survival over the controls of PBS and BDCNs (Figure 5g). Meanwhile, the body weight of the treated mice had no evident fluctuations during the experiment, clarifying the limited side effects of these BNs (Figure S8). Treatment with BDCNs and TDCNs in two cross animal models demonstrated the specificities and personalities of BNs in the initiation of antitumor immune responses.

Combination with clinical α PD-1 antibody

Immune checkpoint blockade by preventing PD-1 on T cells from binding with programmed death ligand 1 (PD-L1) can activate CTLs to attack cancer cells⁴⁸⁻⁵⁰. The use of such inhibitors has elicited durable clinical response and long-term remissions in a fraction of cancer patients⁵¹. However, a large fraction of patients has failed to respond to these agents, showing a low objective response rate of anti-PD therapy⁵². We speculated that the largely augmented immune responses by direct cross-priming of T cells could address the problem of anti-PD therapy, particularly in solid tumors.

1 Therefore, we combined **BNs** with clinical α PD-1 and evaluated the antitumor
2 responses in a more clinically relevant therapeutic setting. The combined modality was
3 first tested in B16-OVA tumor model. In contrast to α PD-1 alone, indicators of
4 upgraded immune responses associated with T cells and APCs together with
5 inflammatory cytokine IFN- γ both in the spleen (Figure 6a-h) and tumor (Figure 6i-o)
6 warranted extremely positive function of the combination of α PD-1 with **BDCNs**.
7 Moreover, the splenic and intratumoral **CD3⁺CD4⁺ T** (Figure 6p and Figure S9a)
8 decreased tremendously, **implying the lowered** negative effect in immune stimulation.
9 In addition to the priming assay, the strikingly upgraded expressions of IFN- γ and TNF-
10 α in the serum demonstrated the robust systemic immune responses induced by the
11 combination treatment (Figure 6q and r). Immunofluorescence images including TNF-
12 α together with IFN- γ (Figure 6s) and CD8 (Figure S9b) crowded in the tumor site
13 further validated the promotion of a strong antitumor immunity (Figure S9c). In contrast
14 to the limited beneficial effect from α PD-1 therapy alone, the combination therapy
15 delayed the tumor growth, which accompanied with an extension of median survival
16 period from 25 to 40 days (Figure 6t-v). Hepa 1-6 carcinoma model was further used
17 to evaluate the broad applicability of the combination of **BNs** with α PD-1 for
18 implementation in other cancers in absence of identified antigen epitopes. As shown in
19 Figure 7, the stimulation of T cells (Figure 7a-d), DC maturation (Figure 7e-g), the
20 secretion of cytokines IFN- γ and TNF- α (Figure 7i and j) were greatly improved by the
21 combination of **HDCNs** and α PD-1, which reversely suppressed the **CD3⁺CD4⁺ T** cells

(Figure 7h). Different from continuous tumor growth in mice treated with α PD-1, the combination treatment was able to abolish all the tumors (Figure 7k and l), benefiting from the ability of HDCNs to elicit a powerful antitumor immunity.

In summary, we have described the use of **BNs** enabled by cascade cell membrane coating to manipulate the cross-priming of T cells. **BNs** have been fabricated through coating PLGA nanoparticles with cell membranes extracted from DCs that have been matured with cancer cell membrane-coated nanoparticles. Due to the maintenance of an intact composition of DC membranes, **BNs** presenting an array of cancer cell membrane antigen epitopes on the surface can inherit intrinsic membrane function of DCs to cross-prime T cells. Significantly, **BNs** can directly interact with T cells and have elicited a robust antigen-specific antitumor immunity in multiple mouse models including B16-OVA, HPV E6 and E7-expressing TC-1, and Hepa 1-6 tumors. In combination with clinical α PD-1, **BNs** have demonstrated a practical way to accomplish desirable tumor regression and survival rate. To this end, treatment efficacy and injection frequency of BNs need to be rationally optimized by using immunoadjuvants or adjusting dosage, particle size, and the loading of antigens for future translation. Beyond the specific use of **BNs** for cross-priming of T cells, the direct interaction with other immune cells may inspire new methods that can manipulate immune responses for treating other diseases, such as autoimmune disease, infections and inflammations, studies that are currently ongoing.

Data availability

The datasets generated and/or analyzed during the study are available from the corresponding authors upon reasonable request.

Online Content Methods, along with any additional Extended Data display items and SourceData, are available in the online version of the paper; references unique to these sections appear only in the online paper.

References

1. Kroll AV, *et al.* Nanoparticulate delivery of cancer cell membrane elicits multiantigenic antitumor immunity. *Advanced Materials* **29**, 1018-1026 (2017).
2. Zhang Q, *et al.* Neutrophil membrane-coated nanoparticles inhibit synovial inflammation and alleviate joint damage in inflammatory arthritis. *Nature Nanotechnology* **13**, 1182-1190 (2018).
3. Riley RS, June CH, Langer R, Mitchell MJ. Delivery technologies for cancer immunotherapy. *Nature Reviews Drug Discovery* **18**, 175-196 (2019).
4. Hu CM, Zhang L, Aryal S, Cheung C, Fang RH, Zhang L. Erythrocyte membrane-camouflaged polymeric nanoparticles as a biomimetic delivery platform. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 10980-10985 (2011).

- 1 5. Rao L, *et al.* Cancer cell membrane-coated upconversion nanoprobe for highly
2 specific tumor imaging. *Advanced Materials* **28**, 3460-3466 (2016).
- 3 6. Chen Z, *et al.* Cancer cell membrane-biomimetic nanoparticles for homologous-
4 targeting dual-modal imaging and photothermal therapy. *ACS Nano* **10**, 10049-10057
5 (2016).
- 6 7. Vijayan V, Uthaman S, Park IK. Cell membrane-camouflaged nanoparticles: a
7 promising biomimetic strategy for cancer theragnostics. *Polymers* **10**, 983 (2018).
- 8 8. Fang RH, *et al.* Cancer cell membrane-coated nanoparticles for anticancer
9 vaccination and drug delivery. *Nano Lett* **14**, 2181-2188 (2014).
- 10 9. Schetters STT, *et al.* Outer membrane vesicles engineered to express membrane-
11 bound antigen program dendritic cells for cross-presentation to CD8 (+) T cells. *Acta*
12 *Biomaterialia* **91**, 248-257 (2019).
- 13 10. Balyan R, *et al.* Correlation of cell-surface CD8 levels with function, phenotype
14 and transcriptome of naive CD8 T cells. *Immunology* **156**, 384-401 (2019).
- 15 11. Verdijk P, *et al.* Limited amounts of dendritic cells migrate into the T-cell area of
16 lymph nodes but have high immune activating potential in melanoma patients. *Clinical*
17 *Cancer Research* **15**, 2531-2540 (2009).
- 18 12. Chang CC, *et al.* Defective human leukocyte antigen class I-associated antigen
19 presentation caused by a novel beta2-microglobulin loss-of-function in melanoma cells.
20 *Journal of Biological Chemistry* **281**, 18763-18773 (2006).

- 1 13. Cubillos-Ruiz JR, *et al.* ER stress sensor XBP1 controls anti-tumor immunity by
2 disrupting dendritic cell homeostasis. *Cell* **161**, 1527-1538 (2015).
- 3 14. Gottfried E, *et al.* Tumor-derived lactic acid modulates dendritic cell activation and
4 antigen expression. *Blood* **107**, 2013-2021 (2006).
- 5 15. O'Neill LA, Pearce EJ. Immunometabolism governs dendritic cell and macrophage
6 function. *Journal of Experimental Medicine* **213**, 15-23 (2016).
- 7 16. Awate S, Babiuk LA, Mutwiri G. Mechanisms of action of adjuvants. *Frontiers in*
8 *Immunology* **4**, 114 (2013).
- 9 17. Turtle CJ, Riddell SR. Artificial antigen-presenting cells for use in adoptive
10 immunotherapy. *Cancer Journal* **16**, 374-381 (2010).
- 11 18. Butler MO, Hirano N. Human cell-based artificial antigen-presenting cells for
12 cancer immunotherapy. *Immunological Reviews* **257**, 191-209 (2014).
- 13 19. Eggermont LJ, Paulis LE, Tel J, Figdor CG. Towards efficient cancer
14 immunotherapy: advances in developing artificial antigen-presenting cells. *Trends*
15 *Biotechnology* **32**, 456-465 (2014).
- 16 20. Sunshine JC, Green JJ. Nanoengineering approaches to the design of artificial
17 antigen-presenting cells. *Nanomedicine* **8**, 1173-1189 (2013).
- 18 21. Rhodes KR, Green JJ. Nanoscale artificial antigen presenting cells for cancer
19 immunotherapy. *Molecular Immunology* **98**, 13-18 (2018).
- 20 22. Fang RH, Zhang L. Nanoparticle-based modulation of the immune system. *Annual*
21 *Review of Chemical and Biomolecular Engineering* **7**, 305-326 (2016).

- 1 23. Sherman SE, Xiao Q, Percec V. Mimicking complex biological membranes and
2 their programmable glycan ligands with dendrimersomes and glycodendrimersomes.
3 *Chemical Reviews* **117**, 6538-6631 (2017).
- 4 24. Jiang Y, *et al.* Engineered cell-membrane-coated nanoparticles directly present
5 tumor antigens to promote anticancer immunity. *Advanced Materials*, e2001808 (2020).
- 6 25. Zhang S, *et al.* Mimicking biological membranes with programmable glycan
7 ligands self-assembled from amphiphilic janus glycodendrimers. *Angewandte Chemie*
8 *International Edition* **53**, 10899-10903 (2014).
- 9 26. Liao PC, Boldogh IR, Siegmund SE, Freyberg Z, Pon LA. Isolation of
10 mitochondria from *Saccharomyces cerevisiae* using magnetic bead affinity purification.
11 *PloS One* **13**, e0196632 (2018).
- 12 27. Radauer-Preiml I, *et al.* Nanoparticle-allergen interactions mediate human allergic
13 responses: protein corona characterization and cellular responses. *Particle and Fibre*
14 *Toxicology* **13**, 3 (2016).
- 15 28. Yaman S, *et al.* Melanoma peptide MHC specific TCR expressing T-cell membrane
16 camouflaged PLGA nanoparticles for treatment of melanoma skin cancer. *Frontiers in*
17 *Bioengineering and Biotechnology* **8**, 943 (2020).
- 18 29. Luo M, *et al.* A STING-activating nanovaccine for cancer immunotherapy. *Nature*
19 *Nanotechnology* **12**, 648-654 (2017).
- 20 30. Hu CM, *et al.* Nanoparticle biointerfacing by platelet membrane cloaking. *Nature*
21 **526**, 118-121 (2015).

- 1 31. Hu Q, Sun W, Qian C, Wang C, Bomba HN, Gu Z. Anticancer platelet-mimicking
2 nanovehicles. *Advanced Materials* **27**, 7043-7050 (2015).
- 3 32. Hu CM, Fang RH, Copp J, Luk BT, Zhang L. A biomimetic nanosponge that
4 absorbs pore-forming toxins. *Nature Nanotechnology* **8**, 336-340 (2013).
- 5 33. Li H, *et al.* Size dependency of circulation and biodistribution of biomimetic
6 nanoparticles: red blood cell membrane-coated nanoparticles. *Cells* **8**, 881 (2019).
- 7 34. Ou Z, *et al.* Macrophage membrane-coated nanoparticles alleviate hepatic
8 ischemia-reperfusion injury caused by orthotopic liver transplantation by neutralizing
9 endotoxin. *International Journal of Nanomedicine* **15**, 4125-4138 (2020).
- 10 35. Zhang Y, *et al.* Macrophage-membrane-coated nanoparticles for tumor-targeted
11 chemotherapy. *Nano Letters* **18**, 1908-1915 (2018).
- 12 36. Meng QF, *et al.* Macrophage membrane-coated iron oxide nanoparticles for
13 enhanced photothermal tumor therapy. *Nanotechnology* **29**, 134004 (2018).
- 14 37. Bhattacharyya S, Ghosh SS. Transmembrane TNF α -expressed macrophage
15 membrane-coated chitosan nanoparticles as cancer therapeutics. *ACS Omega* **5**, 1572-
16 1580 (2020).
- 17 38. Hlavackova E, *et al.* Dendritic cell-based immunotherapy in advanced sarcoma and
18 neuroblastoma pediatric patients: anti-cancer treatment preceding monocyte harvest
19 impairs the immunostimulatory and antigen-presenting behavior of DCs and
20 manufacturing process outcome. *Frontiers in Oncology* **9**, 1034 (2019).

- 1 39. Tan G, Wang Z, Zhang X, Cai Z, Zhang J. Induction of CTLs by DCs pulsed with
2 k-ras mutant peptide on the surface of nanoparticles in the treatment of pancreatic
3 cancer. *Oncology Reports* **26**, 215-221 (2011).
- 4 40. Ronnie H Fang et al. Cancer cell membrane-coated nanoparticles for anticancer
5 vaccination and drug delivery. *Nano Lett* **14**, 2181-2188 (2014).
- 6 41. Luo M et al. A STING-activating nanovaccine for cancer immunotherapy. *Nat*
7 *Nanotechnol* **12**, 648-654 (2017).
- 8 42. Qiangzhe Zhang et al. Cellular Nanosponges Inhibit SARS-CoV-2 Infectivity. *Nano*
9 *Lett* **20**, 5570-5574 (2020).
- 10 43. Kranz LM et al. Systemic RNA delivery to dendritic cells exploits antiviral defence
11 for cancer immunotherapy. *Nature* **534**, 396-401 (2016).
- 12 44. Gottfried E et al. Tumor-derived lactic acid modulates dendritic cell activation and
13 antigen expression. *Blood* **107**, 2013-2021 (2006)
- 14 45. Yunching Chen , Surendar R Bathula, Qi Yang, Leaf Huang. Targeted nanoparticles
15 deliver siRNA to melanoma. *J Invest Dermatol.* **12**, 2790-2798 (2010).
- 16 46. Diana Anissian et al. Piperine-loaded chitosan-STPP nanoparticles reduce neuronal
17 loss and astrocytes activation in chemical kindling model of epilepsy. *Int J Biol*
18 *Macromol.* **107**, 973-983 (2018).

47. Lakeisha A Lewter et al. Slow-sustained delivery of naloxone reduces typical
naloxone-induced precipitated opioid withdrawal effects in male morphine-dependent
mice. *J Neurosci Res.* **8** (2020).

48. Jiao P, et al. Small molecules as PD-1/PD-L1 pathway modulators for cancer
immunotherapy. *Current Pharmaceutical Design* **24**, 4911-4920 (2018).

49. Topalian SL, Drake CG, Pardoll DM. Immune checkpoint blockade: a common
denominator approach to cancer therapy. *Cancer Cell* **27**, 450-461 (2015).

50. Wu X, et al. Application of PD-1 blockade in cancer immunotherapy.
Computational and Structural Biotechnology Journal **17**, 661-674 (2019).

51. Martin-Romano P, et al. Evidence of pseudoprogression in patients treated with
PD1/PDL1 antibodies across tumor types. *Cancer Medicine* **9**, 2643-2652 (2020).

52. Cai J, et al. The role of PD-1/PD-L1 axis and macrophage in the progression and
treatment of cancer. *Journal of Cancer Research and Clinical Oncology* **145**, 1377-
1385 (2019).

53. Qiangzhe Zhang et al. Cellular nanosponges inhibit SARS-CoV-2 infectivity. *Nano
Lett* **20**, 5570-5574 (2020).

54. Audrey Arrighi et al. Development of PLGA microparticles with high
immunoglobulin G-loaded levels and sustained-release properties obtained by spray-
drying a water-in-oil emulsion. *Int J Pharm* **566**, 291-298 (2019).

Acknowledgments

1 This work was financially supported by the National Natural Science Foundation of
2 China (21875135), the Recruitment Program of Global Youth Experts of China
3 (D1410022), the Shanghai Municipal Education Commission-Gaofeng Clinical
4 Medicine Grant Support (20181704, 20191820), and the Innovative research team of
5 high-level local universities in Shanghai (SSMU-ZLCX20180701).

6

7 **Author contributions**

8 J.L. conceived and designed the experiments. F.C. and L.W. performed all experiments.

9 All authors analyzed and discussed the data. F.C. and J.L. wrote the paper.

10

11 **Competing interests**

12 The authors declare no competing interests.

Figure 1. Preparation and characterization of BNs. (a) Schematic illustration of BNs enabled by cascade cell membrane coating for direct cross-priming of T cells. (b) Sizes and (c) Zeta potentials of Ns, B16-OVA cell membrane vesicles (CMV), and CCNPs, respectively. CCNPs were prepared by extruding PLGA nanoparticles with cell membranes at a cell membrane protein/nanoparticle weight ratio of 1:1. (d) Typical TEM images of Ns (left) and CCNPs (right) negatively stained with uranyl acetate. Scale bar: 100 nm. (e) Western blot analysis of membrane-specific protein marker OVA on B16-OVA cells and CCNPs, respectively. (f) Colocalization of PLGA cores (DiD, red channel) and cancer cell membranes (Neuro-DiO, green channel) upon BMDC uptake by confocal imaging. Cell nucleus was stained with DAPI (blue channel). Scale bar: 10 μ m. (g) Processing and presentation of antigen gp100 associated with B16-OVA cell membranes by BMDCs. (h) Flow cytometric profiles and (i) mean fluorescence intensities of BMDCs pulsed by CCNPs with different amounts of membrane proteins. BMDCs were stained with OVAp-H-2k^b-APC. BMDCs, isotype, and PEI-complexed OVA were used as controls. (j) Representative TEM images of BDCNs and DC cell membrane vesicles (DCCMV) negatively stained with uranyl acetate. Scale bar: 100 nm. (k) Sizes and (l) zeta potentials of DC cell membrane vesicles and BDCNs, respectively. BDCNs were prepared by extruding through a polycarbonate membrane with a pore size of 200 nm. Error bars represent the standard deviation ($n = 3$).

Figure 2. Direct cross-priming of T cells by BNs. (a) Representative confocal images and (b) flow cytometric analysis of primary CD8⁺ T lymphocytes after incubation with Ns, RBC-Ns or BDCNs

1 at 4 °C for 1-hour. Cell nucleus, cell membrane, and BDCNs were separately stained with Hoechst
2 (blue channel), Neuro-DiO (green channel), and DiD (red channel). Scale bar: 10 μm. (c) Flow
3 cytometric analysis of CFSE-labeled primary CD8⁺ T cells cultured with Ns, BMDCs, DCNPs,
4 ODCNs or BDCNs. CFSE dilution was used to quantify the proliferation of CD8⁺ T cells. (d)
5 Representative FACS scatter plots and (e) frequency of SIINFEKL-tetramer CD8⁺ T cells in
6 primary splenocytes after incubation with DCNs or BDCNs. (f) Experimental design for *in vivo*
7 evaluation of T cell activation by injection with BNs in 6-8 weeks old male C57BL/6 mice.
8 Representative FACS scatter plots and frequency of (g, i) H-2K^b OVA Tetramer-SIINFEKL and (h,
9 j) H-2Db gp100 Tetramer-EGSRNQDWL CD8⁺ T cells in splenocytes collected from mice treated
10 with PBS, Ns, DCNs, ODCNs, and BDCNs, respectively. (k) Ki67⁺, (l) CD8⁺ and (m) IFN-γ⁺ in
11 CD3⁺ of splenocytes from the treated mice. Error bars represent the standard deviation (*n* = 3).
12 Significance was assessed using One-way ANOVA with Dunnett's post-hoc test. *****p* < 0.0001,
13 ****p* < 0.001, ***p* < 0.01, and **p* < 0.05.

Figure 3. Antitumor immune responses induced by BNs in B16-OVA tumor-bearing mice. (a) Experimental design for *in vivo* evaluation of antitumor immunity. B16-OVA tumor-bearing mice were intravenously administrated with PBS, Ns, DCNs, ODCNs or BDCNs at predetermined time points and sacrificed once tumor volume in control groups exceeded 2000 mm³. Percentages of (b) CD3⁺ T cells as effector T cells and (c) Ki67⁺, mean fluorescence intensities of DC indicator (d) CD80, (e) CD86 and (f) MHC-II, (g) CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio, (h) IFN- γ ⁺ and (i) OVA

1 tetramer in the spleen. Percentages of (j) CD3⁺ T cells as effector T cells and (k) Ki67⁺, mean
2 fluorescence intensities of DC indicator (l) CD80, (m) CD86 and (n) MHC-II, (o)
3 CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio, and (p) OVA tet⁺ in the tumor. Levels of (q) IFN- γ and (r) TNF- α
4 in the serum. (s) Average and (t) individual tumor growth curves. (u) Immunofluorescence staining
5 images of tumor tissue sectioned from the treated mice. CD3⁺ T cells (green channel), CD8⁺ T cells
6 (red channel), and TNF- α (red channel) were separately labeled. Scale bar: 50 μ m. Error bars
7 represent the standard deviation ($n = 5$). Significance was assessed using One-way ANOVA with
8 Dunnett's post-hoc test. **** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$.

Figure 4. Antitumor immune responses elicited by BNs in Hepa 1-6 tumor-bearing mice. (a) Experimental design for *in vivo* assessment of antitumor immune responses. Hepa 1-6 tumor-bearing mice were intravenously injected with PBS, Ns, DCNs, or HDCNs at the indicated time points and sacrificed once tumor volume in control groups exceeded 2000 mm³. Percentages of (b) CD3⁺CD8⁺ and (c) Ki67⁺ T cells, mean fluorescence intensities of DC indicator (d) CD80, (e) CD86, and (f) MHC-II, (g) IFN- γ , (h) CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio and (i) CD3⁺CD4⁺ in the spleen. Expression of (j) TNF- α and (k) IFN- γ in the serum. Percentages of (l) CD3⁺ CD8⁺ and (m) Ki67⁺ T cells, mean fluorescence intensities of DC indicator (n) CD80, (o) CD86, and (p) MHC-II, and (q) CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio in the tumor tissue. (r) Average and (s) individual tumor growth profiles. (t) Immunofluorescence staining images of tumor sampled from the treated mice. CD3⁺ T cells (green channel), CD8⁺ T cells (red channel), and TNF- α (red channel) were labeled individually. Scale bar: 50 μ m. Error bars represent the standard deviation ($n = 5$). Significance was assessed using One-way ANOVA with Dunnett's post-hoc test. **** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$.

Figure 5. Cross-validation of BNs-leveraged antitumor immunity. (a) Experimental design for *in vivo* validation of antigen-specific antitumor immunity. After inoculation with 3×10^6 cells of B16-OVA or TC-1 on day 0, mice were daily treated with PBS, TDCNs or BDCNs for 5 days after the tumor size reached 50~100 mm³. (b) Average and (c) individual tumor growth profiles of B16-OVA tumor-bearing mice. Survivals of (d) B16-OVA and (e) TC-1 tumor-bearing mice after treatment. (f) Average and (g) individual tumor growth curves of TC-1 tumor-bearing mice. Error bars represent the standard deviation ($n = 8-10$). Significance was assessed using One-way ANOVA with Dunnett's post-hoc test. Survival rates were analyzed using the log-rank test. **** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, and * $p < 0.05$.

Figure 6. Combination of BDCNs with α PD-1 in B16-OVA tumor-bearing mice. After inoculation with B16-OVA on day 0, the mice were treated with intraperitoneal administration of 100 μ g α PD-1 on day 8, 9, 11, 13 and 15 along with daily tail vein injection of BDCNs through day

1 8 to 14. Equivalent free α PD-1 was used as a control. Mice were sacrificed for sampling once tumor
 2 volume in the control group exceeded 2000 mm³ ($n = 5$). Percentages of (a) CD3⁺CD8⁺ and (b)
 3 Ki67⁺ T cells, (c) IFN- γ ⁺, (d) OVA tetramer, mean fluorescence intensities of DC indicator (e) CD80,
 4 (f) CD86 and (g) MHC-II, and (h) CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio in the spleen tissue. Percentages
 5 of (i) CD3⁺ CD8⁺ and (j) Ki67⁺ T cells, (k) OVA tetramer, mean fluorescence intensities of DC
 6 indicator (l) CD80, (m) CD86 and (n) MHC-II, (o) CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio and (p)
 7 CD3⁺CD4⁺ in tumor site. Concentrations of (q) IFN- γ and (r) TNF- α in the serum. (s)
 8 Immunofluorescence staining images of tumor tissue sampled from the treated mice. Both TNF- α
 9 and IFN- γ were labeled as red. Scale bar: 50 μ m. (t) Survival as well as (u) average and (v) individual
 10 tumor growth curves of mice treated with the combination of BDCNs with α PD-1 ($n = 8-10$).
 11 Significance was assessed using One-way ANOVA with Dunnett's post-hoc test. Survival rates
 12 were analyzed using the log-rank test. **** $p < 0.0001$ and *** $p < 0.001$.

Figure 7. Combination of HDCNs with α PD-1 in Hepa 1-6 tumor-bearing mice. After inoculation with Hepa 1-6 on day 0, the mice were treated with intraperitoneal administration of 100 μ g α PD-1 on day 8, 9, 11, 13 and 15 along with daily tail vein injection of HDCNs through day 8 to 14. Equivalent free α PD-1 was used as a control. Mice were sacrificed for sampling once tumor volume in the control group exceeded 2000 mm³. Percentages of (a) CD3⁺ T cells as effector T cells, (b) Ki67⁺, (c) IFN-γ⁺, mean fluorescence intensities of DC indicator (d) CD8⁺/Foxp3⁺CD25⁺CD4⁺ ratio, (e) CD80 and (f) CD86, and (g) MHC-II in the spleen. Levels of (h) CD3⁺CD4⁺, (i) IFN-γ and (j) TNF-α in the serum. (k) Average and (l) individual tumor growth curves of the treated mice. Error bars represent the standard deviation ($n = 5$). Significance was assessed using One-way ANOVA with Dunnett's post-hoc test. **** $p < 0.0001$, *** $p < 0.001$ and ** $p < 0.01$.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryInformationRevised.pdf](#)