

Paraganglioma Presenting Atypical Initial Symptoms: a Case Report

Shu Eguchi

St. Luke's International Hospital

Rintaro Ono (✉ rin.r34@gmail.com)

St. Luke's International Hospital <https://orcid.org/0000-0001-8847-2528>

Takeshi Sato

Keio University School of Medicine

Keigo Yada

St. Luke's International Hospital

Naoki Umehara

St. Luke's International Hospital

Satoshi Narumi

National Research Institute for Child Health and Development

Yosuke Ichihashi

Keio University School of Medicine

Taiki Nozaki

St. Luke's International Hospital

Naoki Kanomata

St. Luke's International Hospital

Tomonobu Hasegawa

Keio University School of Medicine

Miwa Ozawa

St. Luke's International Hospital

Daisuke Hasegawa

St. Luke's International Hospital

Case report

Keywords: Paraganglioma, Polyuria, Weight loss, Proteinuria, Posterior reversible encephalopathy syndrome, Case report

Posted Date: August 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-801163/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Paraganglioma (PGL), an extra-adrenal pheochromocytoma, is a rare tumor, especially in children. Here, we present a case of PGL with atypical initial symptoms.

Case presentation: For seven months, an eight-year-old boy complained of polyuria and weight loss, followed by proteinuria and headache for one month prior to admission. He was admitted to our hospital due to an afebrile seizure. His blood pressure remained markedly elevated even after cessation of the convulsion. Magnetic resonance imaging of the brain revealed posterior reversible encephalopathy syndrome. Abdominal computed tomography showed a mass lesion encasing the left renal artery, measuring 41 mm in length along its major axis. The plasma and urine levels of normetanephrine were elevated. Additionally, iodine-123-metaiodobenzylguanidine scintigraphy showed an abnormal uptake in the abdominal mass with no evidence of metastasis. We tentatively diagnosed him with PGL. Substantial alpha- and beta-blocking procedures were performed, followed by a tumor resection and an extended left nephrectomy on day 31 of hospitalization. Pathological findings confirmed the diagnosis of PGL. The postoperative course was uneventful, and his blood pressure normalized without the use of antihypertensive agents. Genetic testing revealed a known *SDHB* germline mutation. The patient remained asymptomatic for eleven months after surgery without relapse.

Conclusion: It remains challenging to diagnose pheochromocytoma/PGL (PPGL) promptly because PPGL can present with a variety of symptoms. Although PPGL is a rare disease, especially in children, it should be considered in differential diagnosis when various unexplained symptoms persist.

Background

Paraganglioma (PGL), an extra-adrenal pheochromocytoma, is a rare tumor. The incidence of pheochromocytoma/PGL (PPGL) in adults is two to five per million, and the incidence in children is even lower [1]. While the hypersecretion of catecholamines causes the classic triad of headaches, palpitations, and excessive sweating, prompt diagnosis is still challenging.

Case Presentation

An eight-year-old boy, who required developmental follow-up at our hospital for developmental disorders, showed polyuria and weight loss seven months before his admission. One month prior to admission, severe headache and proteinuria were also observed (Fig. 1A). On the day of admission, the patient suffered generalized convulsion and was transported to our hospital. Although his convulsions stopped spontaneously within two minutes, his impaired consciousness persisted, and his blood pressure was elevated at 180/140 mmHg. A physical examination revealed no abnormal findings, and brain magnetic resonance imaging showed high signal intensity in the bilateral cerebellar hemispheres on fluid-attenuated inversion recovery, suggestive of posterior reversible encephalopathy syndrome (Fig. 1B, upper left panel). Abdominal contrast-enhanced computed tomography revealed a partially ill-defined

hypervascular mass with heterogeneous enhancement, measuring 35 × 30 × 41 mm, at the left retroperitoneal space below the renal artery bifurcation. A necrotic area was seen in the mass, and the left kidney was atrophic and poorly enhanced (Fig. 1B, lower left panel). The plasma and urine catecholamine levels were both elevated: plasma noradrenaline: 2.1 ng/mL (reference value: 0.10–0.50 ng/mL); urine noradrenaline: 330 µg/day (reference value: 31–160 µg/day); plasma free normetanephrine: 912 pg/mL (reference value: ≤506 pg/mL); and urine normetanephrine: 1.4 mg/day (reference value: 0.09–0.33 mg/day). Levels of adrenaline, dopamine, and metanephrine were within normal limits in both the plasma and the urine. Iodine-123-metaiodobenzylguanidine scintigraphy showed abnormal uptake in the abdominal mass, and there was no evidence of metastasis (Fig. 1B, right panel). With the clinical diagnosis of noradrenaline-producing PGL, we initiated antihypertensive therapy using a calcium-channel blocker and an alpha-blocker (doxazosin). Although a hypertensive attack with convulsion recurred on the seventh day of admission, the blood pressure was well-controlled, and the dose of doxazosin was gradually increased from the initial dose of 1 mg thereafter. After the dose of doxazosin was increased to 16 mg per day, a beta-blocker (carvedilol) was added for heart rate control. The patient underwent resection of the tumor on the 31st day of hospitalization. A left total nephrectomy was also performed because the tumor was tightly adhered to the left renal vessel, and the left kidney was highly atrophic. During the perioperative and postoperative periods, no adverse events, such as hypertensive or hypotensive attacks, occurred, and the patient was discharged on the 10th postoperative day. Immunohistochemical findings of the tumor cells revealed positivity for chromogranin A and synaptophysin, and the diagnosis of PGL was confirmed (Fig. 1C). The Grading of Adrenal Pheochromocytoma and Paraganglioma (GAPP) score was 4 (moderately differentiated type), and Ki67 was positive in 16% of the cells. The levels of noradrenaline and normetanephrine in both the plasma and the urine normalized after the surgery. All symptoms, including the hypertension, headache, polyuria, and proteinuria, disappeared, and the body weight started to increase. The patient remained asymptomatic without relapse, and his blood pressure remained stable without hypotensive agents for eleven months after the surgery. The genetic testing revealed a known heterozygous *SDHB* germline mutation (c.201-2A > C), which was also detected on his father and paternal grandfather without any history of hypertension or malignant tumor.

Discussion And Conclusion

It remains challenging to diagnose PPGL promptly because PPGL can present with various symptoms. Although hypertension is often the first trigger to suspect PPGL, some cases show normal blood pressure in early stages of the disease [2]. In addition, blood pressure is not routinely measured in a pediatric outpatient clinic. In this case, uncommon symptoms of PPGL, such as polyuria, proteinuria, and weight loss, were observed prior to classic symptoms. Although PPGL is a rare disease, especially in children, it should be considered in differential diagnosis when various unexplained symptoms persist.

According to previous reports from the United States and Europe, approximately 30–40% of PPGL cases are hereditary [3–6]. These reports indicated that pediatric cases are more likely to have germline mutations, multiple site lesions, and higher risks of recurrence than adult cases. Genetic testing should be

considered for PPGL patients, especially in younger cases, and more intensive follow-up is needed regarding hereditary cases. Because there are so few reports of hereditary PPGL from Asia [7], the proportion of germline mutations in Asian patients with PPGL still needs to be clarified.

List Of Abbreviations

GAPP: Grading of Adrenal Pheochromocytoma and Paraganglioma; MRI: Magnetic resonance imaging; PGL: Paraganglioma; PPGL: Pheochromocytoma/Paraganglioma

Declarations

Ethics approval and consent to participate

This study was reviewed and approved by the Institutional Review Board of St Luke's International Hospital and Keio University. Written informed consent was obtained from the parents.

Consent for publication

Written informed consent was obtained from the patient's legal guardian for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was partly supported by a Grant-in-Aid from the Ministry of Health, Labour, and Welfare, Japan (Nanjiseisikkanseisakukenkyujigyo [20FC1020 to Tomonobu Hasegawa.]).

Authors' contributions

SE, RO, and DH conceived and drafted the manuscript; TS, SN, YI, and TH conducted genetic analyses; SE, RO, KY, NU, MO, and DH treated the patient; TN performed the evaluation of radiological imaging and drafted the manuscript; NK performed the pathological evaluation. All authors reviewed and approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Pamporaki C, Hamplova B, Peitzsch M, Prejbisz A, Beuschlein F, Timmers HJLM, et al. Characteristics of Pediatric vs Adult Pheochromocytomas and Paragangliomas. *J Clin Endocrinol Metab.* 2017;102(4):1122-32. <https://doi.org/10.1210/jc.2016-3829>.
2. Olson KR, Nimkin K, Carroll RW, Stathatos N, Ordulu Z. Case 9-2021: A 16-Year-Old Boy with Headache, Abdominal Pain, and Hypertension. *N Engl J Med.* 2021;384(12):1145-55. <https://doi.org/10.1056/NEJMcp2027094>.
3. Jochmanova I, Abcede AM, Guerrero RJS, Malong CLP, Wesley R, Huynh T, et al. Clinical characteristics and outcomes of *SDHB*-related pheochromocytoma and paraganglioma in children and adolescents. *J Cancer Res Clin Oncol.* 2020;146(4):1051-63. <https://doi.org/10.1007/s00432-020-03138-5>.
4. Bausch B, Wellner U, Bausch D, Schiavi F, Barontini M, Sanso G, et al. Long-term prognosis of patients with pediatric pheochromocytoma. *Endocr Relat Cancer.* 2013;21(1):17-25. <https://doi.org/10.1530/ERC-13-0415>.
5. Amar L, Bertherat J, Baudin E, Ajzenberg C, Paillerets BB, Chabre O, et al. Genetic testing in pheochromocytoma or functional paraganglioma. *J Clin Oncol.* 2005;23(34):8812-8. <https://doi.org/10.1200/JCO.2005.03.1484>.
6. King KS, Prodanov T, Kantorovich V, Fojo T, Hewitt JK, Zacharin M, et al. Metastatic pheochromocytoma/paraganglioma related to primary tumor development in childhood or adolescence: significant link to *SDHB* mutations. *J Clin Oncol.* 2011;29(31):4137-42. <https://doi.org/10.1200/JCO.2011.34.6353>.
7. Kimura N, Takayanagi R, Takizawa N, Itagaki E, Katabami T, Kakoi N, et al. Pathological grading for predicting metastasis in phaeochromocytoma and paraganglioma. *Endocr Relat Cancer.* 2014;21(3):405-14. <https://doi.org/10.1530/ERC-13-0494>.

Figures

Fig. 1





Figure 1

A. The trend of the patient's body weight and the course of each symptom are shown with the average body weight of a Japanese boy. B. Upper left panel: The magnetic resonance imaging (MRI) of the patient's brain on admission is shown. Axial fluid-attenuated inversion recovery MRI shows edema in the bilateral cerebellar hemispheres (yellow arrows). Lower left panel: The abdominal contrast-enhanced computed tomography on admission shows a partially ill-defined hypervascular mass, measuring 35 × 30 × 41 mm, at the left retroperitoneal space below the renal artery bifurcation (circle). Noted is a left kidney

with atrophy and poor enhancement (white arrows). Right panel: Iodine-123-metaiodobenzylguanidine scintigraphy shows abnormal uptake in the abdominal mass (red arrow). C. The results of hematoxylin and eosin staining, as well as the immunohistochemical results for chromogranin A and synaptophysin, are shown.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CAREchecklist.pdf](#)