Weakened seasonality of the African rainforest precipitation in boreal winter and spring driven by tropical SST variabilities

Xin-Yue Wang

Institute of Atmospheric Physics Chinese Academy of Sciences

Jiang Zhu

Institute of Atmospheric Physics Chinese Academy of Sciences

Meijiao Xin

Institute of Atmospheric Physics Chinese Academy of Sciences

Chentao Song

Institute of Atmospheric Physics Chinese Academy of Sciences

Yadi Li

Institute of Atmospheric Physics Chinese Academy of Sciences

Yi Zhou

Institute of Atmospheric Physics Chinese Academy of Sciences

Xichen Li (✉ lixichen@mail.iap.ac.cn)

Institute of Atmospheric Physics Chinese Academy of Sciences <https://orcid.org/0000-0001-6325-6626>

Research Letter

Keywords: seasonality of precipitation, African rainforest, Walker Circulation, convergence of moisture

Posted Date: September 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-80118/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Precipitation in the equatorial African rainforest plays an important role in both the regional hydrological cycle and the global climate variability. Previous studies mostly focus on the trends of drought in recent decades or long-time scales. Using two observational datasets, we reveal a remarkable weakening of the seasonal precipitation cycle over this region from 1979 to 2015, with precipitation significantly increased in the boreal winter dry season (+ 0.45 mm/day) and decreased in the boreal spring wet season (-0.78 mm/day), which account for ~ 14% of their respective climatological means. We further use a state-of-the-art atmospheric model to isolate the impact of sea surface temperature change from different ocean basins on the precipitation changes in the dry and wet seasons. Results show that the strengthening precipitation in the dry season is mainly driven by the Atlantic warming, whereas the weakening precipitation in the wet season can be primarily attributed to the Indian Ocean. Warming Atlantic intensifies the zonal circulation over the African rainforest, strengthening moisture convergence and intensifying precipitation in the boreal winter dry season. Warming Indian Ocean contributes more to reducing the zonal circulation and suppressing the convection in the boreal spring wet season, leading to an opposite effect on precipitation. This result has important implication on local ecology as well as global climate system.

Introduction

African rainforest, located in the Congo Basin, is the world's second-largest tropical forest region (Malhi et al. 2013; Mayaux et al. 2013; Fig. 1a). Rainforests play important roles in hydrology and carbon storage due to their rich biodiversity and high ecological function (Baccini et al. 2012; Saatchi et al. 2011). Deep convection over the African rainforest region makes a great contribution to the global climate system (Creese et al. 2019), and even dominates the global tropical rainfall distribution in the transition seasons (Washington et al. 2013). However, long-term drying trend occurred over the equatorial western and central Africa (Asefi-Najafabady and Saatchi 2013; Diem et al. 2014; Hua et al. 2016; Malhi and Wright 2004; Yin and Gruber 2010), resulting in a declining of Congo rainforest greenness and altering the forest structure (Chambers and Roberts 2014; Zhou et al. 2014). Therefore, it is crucial to understand the change of climate in African rainforest regions, in particular precipitation in different seasons.

Rainfall in equatorial Africa is affected by the shift of the intertropical convergence zone (ITCZ, Nicholson 2018; Sultan and Janicot 2000) and is related to West African monsoon (Nicholson 2009; Redelsperger et al. 2006; Thorncroft et al. 2011) and local zonal circulation (Cook and Vizzy 2016; Dezfuli et al. 2015; Leduc-Leballeur et al. 2013). Some regional factors, such as topography, mid-level jets, and mesoscale convective systems all influence precipitation (Jackson et al. 2009; Nicholson 2009; Pokam et al. 2012). Under the impact of these complex factors, the seasonal cycle of equatorial African precipitation is characterized by a bimodal feature, with two rainy seasons (March, April, May, MAM; September, October, November, SON) and two dry seasons (June, July, August, JJA; December, January, February, DJF) (Creese and Washington 2018; Washington et al. 2013; Fig. 1b).

There is a seasonal difference in driving the variation of regional precipitation by sea surface temperature (SST) forcing (Balas et al. 2007; Farnsworth et al. 2011). On intraseasonal timescales, the equatorial SST cooling plays a role in the precipitation along the African coast during boreal spring (Leduc-Leballeur et al. 2013). On interannual timescales, Atlantic SST governs the movement of the ITCZ and further influence boreal summer precipitation in equatorial Africa (Nicholson 2009). The Indian Ocean is more related to rainfall in late summer/early fall, whereas Pacific is important for central Africa in MAM (Balas et al. 2007; DiNezio et al. 2013; Farnsworth et al. 2011). However, ENSO is less directly connected to African rainforest precipitation (Malhi and Wright 2004). Long-term drought in West and Central Africa is mainly linked to the change of Atlantic circulation (Shanahan et al. 2009) and SST (Camberlin et al. 2001; Diem et al. 2014), and warming Indian Ocean also makes an important contribution (Diem et al. 2014; Hoerling et al. 2006; Hua et al. 2016; Richard et al. 2001). In boreal winter/spring season, tropical North Atlantic may force anomalous circulation and increased precipitation in central equatorial Africa (Todd and Washington 2004). Additionally, Pacific SST can also modulate precipitation over eastern and southern Africa by atmospheric bridge (Dezfuli and Nicholson 2013; Hua et al. 2016; Maidment et al. 2015).

Previous studies indicated that long-term dry trend caused the shrinking of Congo rainforest (Zhou et al. 2014). It is also reported that the dry season (JJA) length has been increased by 6.4–10.4 days per decade in the period of 1988–2013 over the Congo rainforest (Jiang et al. 2019). In East Africa, long rains (MAM) precipitation decreased due to abrupt change of tropical Pacific SST (Lyon and DeWitt 2012). Many efforts have been devoted to study the precipitation in dry and wet seasons under global warming and found models show large uncertainties (Creese and Washington 2018; James et al. 2013). Up to now, most of these studies concentrate on either one season or the projection of models in African rainforest precipitation. However, the change in seasonalities between dry and wet seasons in recent decades are not very well investigated because of extremely limited observation stations. Meanwhile, the mechanism causing the change of precipitation seasonality remains unclear.

In the rest of this paper, we focus on the recent decadal trend in dry season of boreal winter (DJF) and boreal spring wet season (AM) over the equatorial African rainforest region, by combining observed datasets and atmospheric model simulations. We describe the datasets, the statistical methods, and the model experiments in “Methods” section, and examine the observed recent long-term trends in the dry and wet seasons in “Results” section. We found that the seasonality of precipitation over the African rain forest is weakening over the past ~ four decades, i.e. precipitation has significantly increased in the boreal winter dry season and decreased in the boreal spring wet season. Further, we investigate the impacts of tropical SSTs on the observed precipitation changes using atmospheric model simulations in “Results” section. The change of specific humidity, moisture flux and Walker Circulation forced by SST over the African rainforest region are presented in “Results” section. Finally, conclusions and discussions are given in last section.

Methods

In this study, we use MODIS land cover map dataset (MCD12C1, Friedl et al. 2010) to describe the distribution of the African rainforest. To examine the precipitation change, we used two observed precipitation datasets, the Global Precipitation Climatology Project (GPCP, Huffman et al. 2009) and the CPC Merged Analysis of Precipitation (CMAP, Xie and Arkin 1997). These datasets are based on gauge stations and satellite observations, and are widely used as reliable precipitation measurements (Yin et al. 2004). The pattern of tropical SST in recent decades has an important impact on precipitation change. To investigate observed SST changes in different basins, we use the Hadley Centre Sea Ice and Sea Surface Temperature dataset (HadISST, Rayner et al. 2003). The same dataset was also used to drive the atmospheric models.

In addition, as the African rainforest is mainly distributed in the equatorial Congo Basin and is characterized by seasonal wetting-drying cycles, we calculate the changes of precipitation over the rainforest (7°E-32°E, 4°S-5°N) in the boreal winter dry season (DJF) and boreal spring wet season (AM) from 1979 to 2015. The Sen's slope method (Sen 1968) are adopted to calculate observed and simulated trends, and their confidence intervals was estimated using a Mann-Kendall test (Gilbert 1987). These two methods are non-parametric and robust in the existence of outliers. For the following model simulations, we also use the same method to analyze.

One of the focus of this study is how SST change in tropical basin in the recent decades influences African rainforest precipitation. However, different basins interact with each other on interannual and decadal timescales. Results in atmospheric models do not reflect inter-basin interaction, and are thus typically used to isolate the role of different basins. As a result, the NCAR atmospheric model, Community Atmosphere Model version 4 (CAM4) is used to investigate the atmospheric responses to the observed SST trend in different ocean basins. We designed three SST forcing experiments by the tropical Atlantic, Indian Ocean and Pacific, respectively. The model is forced by observed SST variability from 1979 to 2015 in the target basin, while the model is driven by climatological SST in the other basins. We conducted 12-member ensemble simulations in the experiments for each ocean basin. By examining the change of the ensemble-mean, we could understand the contribution to precipitation change by the individual ocean basin.

Results

Observed Changes in African rainforest Precipitation

Given regional distribution of African rainforest, we focus on the domain marked by black box in Fig. 1a (7°E-32°E, 4°S-5°N). We further calculate the climatology of area-averaged precipitation in each month (Fig.1b). As suggested in previous studies, a typically bimodal seasonal precipitation feature is apparent in this region. The boreal winter (DJF) is a clear dry season, with climatological precipitation of 3.15~3.31mm/day. In the next few months, climatological precipitation starts to exceed 5mm/day, which means wet season is coming. In particular, in April and May, rainforest receives the highest precipitation in the first half of the year. Accordingly, we only choose the boreal winter dry season (DJF) and boreal

spring wet season (AM) to examine the seasonality of precipitation over the African rainforest regions in past decades.

Fig.2 shows the observed precipitation trend in the boreal winter dry season (DJF; Fig.2a, 2b) and boreal spring wet season (AM; Fig. 2c, 2d) from 1979 to 2015. Remarkably, two observational datasets show similar spatial patterns, with a significant increasing (deep green color) trend in DJF and a decreasing trend (deep brown color) in AM. In the dry season (DJF), the largest positive precipitation anomalies are mainly distributed in the west and east of the domain (Fig.2a, 2b). However, in the wet season (AM), stronger negative precipitation anomalies are not confined to the eastern rainforest, but widely extended to East African Plateau (Fig.2c, 2d). Observations exhibit an opposite sign in two seasons, whereas the drought in the boreal spring appears to be more extensive.

Observed patterns indicate different amounts of precipitation change during the boreal winter dry season (DJF) and boreal spring wet season (AM) (Fig.2). Therefore, we show the quantitative features of precipitation in Fig.3. The results of GPCP and CMAP are nearly identical. Time series of precipitation show an intensification in DJF (Fig. 3a) and a declination in AM (Fig. 3b) since 1979. Spatially-averaged precipitation trends over the rainforest region (Fig. 3c) suggest that the observed precipitation anomalies are statistically significant at the 5% level, with magnitudes of ~ 0.4472 mm/day/37years in DJF and ~ -0.7787 mm/day/37years in AM.

All observational datasets show a dramatic weakening of seasonality of precipitation over Africa rainforest region, i.e. an increased trend in the boreal winter dry season (DJF) and a decreased trend in the boreal spring wet season (AM). The climatological precipitation over this region is ~ 3.2278 mm/day in DJF and ~ 5.6510 mm/day in AM, and increased by ~ 0.4472 mm/day in DJF and decreased by ~ 0.7787 mm/day in past 37 years, accounting for $\sim 14\%$ of their respective climatological means.

To a large extent, the change of convective precipitation is modulated by tropical SST. In past decades, three tropical ocean basins experience different SST trends, namely a warming in the Atlantic, Indian Ocean and western Pacific, and a cooling in central-eastern Pacific. SST changes can be further attributed to the positive phase of the Atlantic Multidecadal Oscillation (AMO) and the cooling phase of the Pacific Decadal Oscillation (PDO). These SST trends play key roles in changing regional or global climate. In particular, the warming Atlantic and Indian Ocean may be more relevant to the change of precipitation in Africa. Therefore, we use atmospheric models to conduct experiments in order to quantify contribution of individual ocean basins to tropical African precipitation.

Impacts of Tropical SST Variabilities on the Precipitation Changes

We first examine observed DJF and AM decadal SST trends (Fig.4). As revealed in many studies, Atlantic and Indo-western Pacific show an apparent warming, but central eastern Pacific shows an evident cooling. However, the amplitude of SST trends varies with season. The magnitude of warming in Atlantic and cooling in central-Pacific in DJF is larger than that in AM, which may exert different effects in the

corresponding season. Next, we add these different SST variabilities to climatological mean SST in atmospheric models respectively to isolate the role of different ocean basins.

In order to test and to verify the simulation of atmospheric models, we compare the observed and simulated precipitation climatology (Fig.5). In the boreal winter dry season (DJF) and boreal spring wet season (AM), the climatological pattern is reasonably reproduced by models, such as spatial distribution and transition of dry and wet season, although models usually slightly overestimate precipitation. As a result, we continue to simulate the change of precipitation by running these models.

Fig.6 displays the precipitation trends by forcing three ocean basins: the tropical Atlantic, Indian Ocean, and Pacific, respectively. During the boreal winter dry season (DJF), the model responses to the tropical Atlantic show the closest resemblance to the observation (Fig.6a), while the responses to tropical Indian Ocean and Pacific are opposite to those in observations (Fig.6b-c). Forced by tropical Atlantic Forcing, the model reproduces the strengthening pattern in central and western rainforest (Fig.6a), implying that Atlantic plays a leading role in the wetting in DJF while Indian Ocean and Pacific have completely opposite effect.

During the boreal spring wet season (AM), significant negative anomalies appear over the African rainforest in three ocean forcing experiments. Compared to observations, simulated trend distribution is slightly different (Fig. 6d-f). Reproduced decreased trends are mainly located in northwestern rainforest, albeit with enhanced trends in the southwest part. However, the precipitation change driven by Indian Ocean and Pacific better resembles that in observation (Fig.5e), indicating that Indian Ocean and Pacific have more important impacts on the drying in the AM.

The simulations in atmospheric models emphasize that the individual ocean basins has different impacts on African rainforest rainfall in the boreal winter dry season (DJF) and boreal spring wet season (AM). The enhanced precipitation in DJF is mainly attributed to the warming Atlantic forcing, whereas the Indian and Pacific Oceans produce an opposite effect. This positive anomalies in DJF cannot be fully explained by atmospheric models. The weakening precipitation in AM is driven by all three oceans, with Indian Ocean and the Pacific making more contributions.

Mechanisms of the Precipitation Changes

Because African precipitation can be affected by the strength of the Walker circulation (Cook and Vizy 2016; Hua et al. 2016), we examine changes of the latter to clarify the processes that link precipitation over the African rainforest and tropical SST. Fig.6 shows the zonal mass stream function (Walker Circulation) trend (shading) and climatology (contour) over the equatorial Africa in atmospheric models. During the boreal winter dry season (DJF), significant positive trends appear in the core region of climatology simulated by Atlantic forcing (Fig. 7a), whereas negative trends appear under Indian Ocean forcing (Fig.7b). Enhanced circulation associated with deep convection is closely linked to higher rainfall in DJF. During the AM, only Indian Ocean generates significant negative trends (Fig.7e), with corresponding stronger divergence and less precipitation. Pacific forcing has no clear sign in either

season (Fig.7c, 7f). The strengthening (weakening) circulation is coherent with increased (decreased) precipitation in simulations.

Changes of specific humidity and moisture flux convergence reflect evolutions in moisture, energy and convection in the air, which are intimately connected with change of precipitation (Chadwick et al. 2016; Gaffen and Ross 1999; Trenberth et al. 2011). Fig.8 shows the simulated changes of specific humidity (moisture) and wind at 825hPa. During the boreal winter dry season (DJF), strengthening moisture anomalies accompanied by typical convergence over the equatorial western African are driven by the tropical Atlantic forcing (Fig.8a). However, forced by the Indian Ocean and Pacific, simulations show the weakening moisture trend and divergence (Fig.8b-c). The comparison of model responses suggests that warming Atlantic increases moisture and further leads to wetting trend in DJF (Fig.6a). During the boreal spring wet season (AM), the model responses all show decreased moisture and enhanced divergence (Fig.8d-f), in particular forced by Indian Ocean and Pacific, causing corresponding dry trend in AM (Fig.6d-f).

Simulated changes of vertically integrated moisture flux convergence (Fig.9) resemble the changes of moisture at 825hPa (Fig.8). In the boreal winter dry season (DJF), the tropical Atlantic forcing triggers strong convergence, implying that more moisture is transported to the equatorial western Africa (Fig.9a). Indian Ocean forcing induces the divergence (Fig. 9b) and Pacific forcing has no obvious signs (Fig.9c). In the boreal spring wet season (AM), the simulation results are consistent regarding more divergence (Fig.9d-f), with a more widespread area driven by Indian Ocean forcing, which have a substantial impact on decreasing precipitation.

Overall, changes of Walker Circulation and moisture flux convergence modulated by SST support the wetting trend in the boreal winter dry season (DJF) and drying trend in the boreal spring wet season (AM). The variation of Atlantic and Indian Ocean forcing with season are more evident. Warming Atlantic drivers intensified Walker circulation over the Africa and thus lead to the increase of moisture convergence associated with deep convection and further strengthens precipitation in the boreal winter dry season. However, warming Indian Ocean weakens Walker circulation and the corresponding convergences in the boreal spring wet season, causing reductions in precipitation.

Conclusions And Discussions

In this study, we examine the changes of precipitation over the African rainforest during the boreal winter dry season (DJF) and boreal spring wet season (AM) from 1979 to 2015 by using observational precipitation datasets. We reveal that the seasonality of precipitation is dramatically weakened in observations, i.e. the wetting trend in the boreal winter dry seasons and the drying trend in the boreal spring wet season. The increased precipitation anomaly is ~ 0.4472 mm/day in DJF and decreased anomaly is ~ 0.7787 mm/day in AM in past 37 years, accounting for 14% of the climatology. As revealed in atmospheric models (CAM4), these changes are driven by tropical seasonal SST forcing, in particular the Atlantic and the Indian Ocean. Simulations show that the warming of the tropical Atlantic plays a key

role in intensifying zonal circulation and moisture convergence over the equatorial Africa in boreal winter dry season, resulting in positive precipitation anomalies. However, warming Indian Ocean makes more contribution to weakening circulation and convection, leading to negative precipitation anomalies in boreal spring wet season.

However, the mechanism of seasonal precipitation change is mainly based on numerical simulations of a state-of-the-art atmospheric model. The relationship between SST and precipitation is extraordinarily complex in real world. In our study, observed intensified precipitation in the boreal winter dry season is only partly explained by Atlantic forcing. The model responses to Indian Ocean and Pacific forcing show negative precipitation anomalies in this dry season. Warming Atlantic plays an important role in strengthening moisture flux and warming Indian Ocean suppresses the convergence in DJF. The whole process tends to be nonlinear, and therefore the specific relationship between different SST and precipitation needs further study. In addition, for the other dry season (JJA) and wet season (SON), it is worthwhile to investigate whether the seasonality of precipitation weakens or strengthens and the roles of SST in the future.

The precipitation is of great importance to African rainforest and even the entire tropics. In recent decades, observed precipitation increased during the boreal winter dry season (DJF) and decreased during the boreal spring wet season (AM) over the African rainforest. The opposite change of precipitation could cause a wetter dry season and a drier wet season, which may dramatically impact on the hydrological and ecosystem over the entire continent and around. Meanwhile, the deep convection associated with precipitation in different seasons, may have a further effect on the global circulation and potentially impact on the global climate system.

Declarations

Availability of data and materials

MODIS land cover map dataset (MCD12C1) is available at https://lpdaac.usgs.gov/get_data. Observational precipitation (GPCP and CMAP) is download from The NOAA Earth System Research Laboratory's Physical Sciences Division (<https://www.esrl.noaa.gov/psd/data/gridded/tables/precipitation.html>). Model datasets and materials analyzed in this study are accessible from the corresponding author under reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by the National Natural Science Foundation of China (Grants 41976193).

Authors' contributions

Xichen Li performed the model experiments. Xin-Yue Wang analyzed the datasets and wrote the manuscript. All authors contributed to the manuscript revision.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grants 41976193).

References

- Asefi-Najafabady S, Saatchi S (2013) Response of African humid tropical forests to recent rainfall anomalies. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368(1625):20120306
- Baccini A, Goetz S, Walker W, Laporte N, Sun M, Sulla-Menashe D, Hackler J, Beck P, Dubayah R, Friedl M (2012) Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature climate change* 2(3):182-185
- Balas N, Nicholson S, Klotter D (2007) The relationship of rainfall variability in West Central Africa to sea-surface temperature fluctuations. *International Journal of Climatology: A Journal of the Royal Meteorological Society* 27(10):1335-1349
- Camberlin P, Janicot S, Pocard I (2001) Seasonality and atmospheric dynamics of the teleconnection between African rainfall and tropical sea-surface temperature: Atlantic vs. ENSO. *International Journal of Climatology: A Journal of the Royal Meteorological Society* 21(8):973-1005
- Chadwick R, Good P, Willett K (2016) A simple moisture advection model of specific humidity change over land in response to SST warming. *Journal of Climate* 29(21):7613-7632
- Chambers JQ, Roberts DA (2014) Drought in the Congo Basin. *Nature* 509(7498):36-38
- Cook KH, Vizzy EK (2016) The Congo Basin Walker circulation: dynamics and connections to precipitation. *Climate Dynamics* 47(3-4):697-717
- Creese A, Washington R (2018) A process-based assessment of CMIP5 rainfall in the Congo Basin: the September–November rainy season. *Journal of Climate* 31(18):7417-7439
- Creese A, Washington R, Jones R (2019) Climate change in the Congo Basin: processes related to wetting in the December–February dry season. *Climate Dynamics* 53(5-6):3583-3602
- Dezfuli AK, Nicholson SE (2013) The relationship of rainfall variability in western equatorial Africa to the tropical oceans and atmospheric circulation. Part II: The boreal autumn. *Journal of Climate* 26(1):66-84
- Dezfuli AK, Zaitchik BF, Gnanadesikan A (2015) Regional atmospheric circulation and rainfall variability in south equatorial Africa. *Journal of Climate* 28(2):809-818

- Diem JE, Ryan SJ, Hartter J, Palace MW (2014) Satellite-based rainfall data reveal a recent drying trend in central equatorial Africa. *Climatic Change* 126(1-2):263-272
- DiNezio PN, Vecchi GA, Clement AC (2013) Detectability of Changes in the Walker Circulation in Response to Global Warming. *Journal of Climate* 26(12):4038-4048
- Farnsworth A, White E, Williams CJ, Black E, Kniveton DR (2011) Understanding the large scale driving mechanisms of rainfall variability over Central Africa. In: *African climate and climate change*. Springer, pp 101-122
- Gaffen DJ, Ross RJ (1999) Climatology and trends of US surface humidity and temperature. *Journal of Climate* 12(3):811-828
- Hoerling M, Hurrell J, Eischeid J, Phillips A (2006) Detection and attribution of twentieth-century northern and southern African rainfall change. *Journal of climate* 19(16):3989-4008
- Hua W, Zhou L, Chen H, Nicholson SE, Raghavendra A, Jiang Y (2016) Possible causes of the Central Equatorial African long-term drought. *Environmental Research Letters* 11(12):124002
- Huffman GJ, Adler RF, Bolvin DT, Gu G (2009) Improving the global precipitation record: GPCP Version 2.1. *Geophysical Research Letters* 36
- Jackson B, Nicholson SE, Klotter D (2009) Mesoscale convective systems over western equatorial Africa and their relationship to large-scale circulation. *Monthly Weather Review* 137(4):1272-1294
- James R, Washington R, Rowell DP (2013) Implications of global warming for the climate of African rainforests. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368(1625):20120298
- Jiang Y, Zhou L, Tucker CJ, Raghavendra A, Hua W, Liu YY, Joiner J (2019) Widespread increase of boreal summer dry season length over the Congo rainforest. *Nature Climate Change* 9(8):617-622
- Leduc-Leballeur M, De Coëtlogon G, Eymard L (2013) Air–sea interaction in the Gulf of Guinea at intraseasonal time-scales: wind bursts and coastal precipitation in boreal spring. *Quarterly Journal of the Royal Meteorological Society* 139(671):387-400
- Lyon B, DeWitt DG (2012) A recent and abrupt decline in the East African long rains. *Geophysical Research Letters* 39(2)
- Maidment RI, Allan RP, Black E (2015) Recent observed and simulated changes in precipitation over Africa. *Geophysical Research Letters* 42(19):8155-8164
- Malhi Y, Adu-Bredu S, Asare RA, Lewis SL, Mayaux P (2013) African rainforests: past, present and future. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368(1625):20120312

- Malhi Y, Wright J (2004) Spatial patterns and recent trends in the climate of tropical rainforest regions. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 359(1443):311-329
- Mayaux P, Pekel J-F, Desclée B, Donnay F, Lupi A, Achard F, Clerici M, Bodart C, Brink A, Nasi R (2013) State and evolution of the African rainforests between 1990 and 2010. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368(1625):20120300
- Nicholson SE (2009) A revised picture of the structure of the “monsoon” and land ITCZ over West Africa. *Climate Dynamics* 32(7-8):1155-1171
- Nicholson SE (2018) The ITCZ and the seasonal cycle over equatorial Africa. *Bulletin of the American Meteorological Society* 99(2):337-348
- Pokam WM, Djotang LAT, Mkankam FK (2012) Atmospheric water vapor transport and recycling in Equatorial Central Africa through NCEP/NCAR reanalysis data. *Climate Dynamics* 38(9-10):1715-1729
- Rayner N, Parker DE, Horton E, Folland CK, Alexander LV, Rowell D, Kent E, Kaplan A (2003) Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research: Atmospheres* 108(D14)
- Redelsperger J-L, Thorncroft CD, Diedhiou A, Lebel T, Parker DJ, Polcher J (2006) African Monsoon Multidisciplinary Analysis: An international research project and field campaign. *Bulletin of the American Meteorological Society* 87(12):1739-1746
- Richard Y, Fauchereau N, Pocard I, Rouault M, Trzaska S (2001) 20th century droughts in southern Africa: spatial and temporal variability, teleconnections with oceanic and atmospheric conditions. *International Journal of Climatology: A Journal of the Royal Meteorological Society* 21(7):873-885
- Saatchi SS, Harris NL, Brown S, Lefsky M, Mitchard ET, Salas W, Zutta BR, Buermann W, Lewis SL, Hagen S (2011) Benchmark map of forest carbon stocks in tropical regions across three continents. *Proceedings of the national academy of sciences* 108(24):9899-9904
- Sen PK (1968) Estimates of the regression coefficient based on Kendall's tau. *Journal of the American Statistical Association* 63(324):1379-8
- Shanahan TM, Overpeck JT, Anchukaitis K, Beck JW, Cole JE, Dettman DL, Peck JA, Scholz CA, King JW (2009) Atlantic forcing of persistent drought in West Africa. *science* 324(5925):377-380
- Sultan B, Janicot S (2000) Abrupt shift of the ITCZ over West Africa and intra-seasonal variability. *Geophysical Research Letters* 27(20):3353-3356
- Thorncroft CD, Nguyen H, Zhang C, Peyrillé P (2011) Annual cycle of the West African monsoon: regional circulations and associated water vapour transport. *Quarterly Journal of the Royal Meteorological Society*

Todd MC, Washington R (2004) Climate variability in central equatorial Africa: Influence from the Atlantic sector. *Geophysical Research Letters* 31(23)

Trenberth KE, Fasullo JT, Mackaro J (2011) Atmospheric moisture transports from ocean to land and global energy flows in reanalyses. *Journal of climate* 24(18):4907-4924

Washington R, James R, Pearce H, Pokam WM, Moufouma-Okia W (2013) Congo Basin rainfall climatology: can we believe the climate models? *Philosophical Transactions of the Royal Society B: Biological Sciences* 368(1625):20120296

Xie PP, Arkin PA (1997) Global precipitation: A 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs. *Bulletin of the American Meteorological Society* 78(11):2539-2558

Yin X, Gruber A (2010) Validation of the abrupt change in GPCP precipitation in the Congo River Basin. *International Journal of Climatology: A Journal of the Royal Meteorological Society* 30(1):110-119

Yin XG, Gruber A, Arkin P (2004) Comparison of the GPCP and CMAP merged gauge-satellite monthly precipitation products for the period 1979-2001. *Journal of Hydrometeorology* 5(6):1207-1222

Zhou L, Tian Y, Myneni RB, Ciais P, Saatchi S, Liu YY, Piao S, Chen H, Vermote EF, Song C (2014) Widespread decline of Congo rainforest greenness in the past decade. *Nature* 509(7498):86-90

Figures

Figure 1

African rainforests and monthly mean precipitation. a The distribution of tropical African rainforest, based on 2011-2015 MCD12C1 datasets. In this study, we focus on the domain (7°E-32°E, 4°S-5°N, the box), covering the most of African rainforest. b The spatial averaged (the box in fig.1a) climatology of monthly precipitation, based on 1979-2015 GPCP and CMAP datasets. Rainfall features bimodal distribution over the African rainforests, with two dry seasons (DJF and JJA) and two wet seasons (MAM and SON). We concentrate on precipitation change during the boreal winter dry season (DJF) and boreal spring wet season (AM).

Figure 2

Seasonal Precipitation changes in observational datasets from 1979 to 2015. Color mapping represents the precipitation change (increase, dark green; decrease, dark brown). Dark dots indicate that precipitation trends are significant at the 5% level. Upper Panels (a, b) show the precipitation change during the boreal winter dry season (DJF) from GPCP and CMAP. Lower Panels (c, d) same as (a, b) but for the boreal spring wet season (AM). The increasing trend in DJF and decreasing trend in AM are significant, implying that the seasonality of precipitation is weakening over the rainforest region.

Figure 3

The statistics of spatial averaged precipitation in observations. a, b Time series of observed precipitation anomalies over the African rainforest (dark box in Fig.2) show an intensified tendency during the boreal winter dry season and a weakened tendency during the boreal spring wet season. c The black lines and thick bars represent precipitation change and 95% confidence intervals. These results show that precipitation changes are statistically significant. From 1979 to 2015, precipitation increased ~ 0.4472 mm/day * 37 years in DJF and ~ -0.7787 mm/day * 37 years in AM over the African rainforest domain.

Figure 4

Observed seasonal HadISST trend from 1979 to 2015. a, b Color mapping represents decadal SST change in DJF and AM. SST exhibits apparent warming in the Atlantic and Indo-western Pacific, and cooling in the central eastern Pacific. However, the magnitude of warming over the Atlantic and cooling over the Pacific are relatively larger in DJF than that in AM. The amplitude of seasonal SST in different seasons may have a different effect on precipitation over the African rainforest.

Figure 5

Climatological precipitation pattern in observation and simulated. Upper Panels (a, b) show observed (the mean of GPCP and CMAP) climatology in DJF and AM. Lower Panels (c, d) same as (a, b) but for the AM. These results suggest that climatological patterns are reasonably reproduced by simulation.

Figure 6

Simulated seasonal precipitation change from 1979 to 2015 in atmospheric models. Color mapping represents the precipitation change (increase, dark green; decrease, dark brown). Dark dots indicate that precipitation trends are significant at the 5% level. a, b and c show the precipitation change during the boreal winter dry season (DJF) forced by the tropical Atlantic, Indian Ocean and Pacific SST. e, f and g same as a, b and c but for the boreal spring wet season (AM). In the dry season (DJF), only the simulated forced by the tropical Atlantic show closest resemblance to the observation. In the wet season (AM), three ocean all play roles in decreasing precipitation.

Figure 7

Simulated seasonal change of Walker Circulation in Atmospheric Models from 1979 to 2015. Color mapping illustrates the circulation change and contour show the climatology. a, b and c show the pattern forced by the tropical Atlantic, Indian Ocean and Pacific during the boreal winter dry season (DJF). d, e and f same as a, b and c but for the boreal spring wet season (AM). Green lines are the African rainforest region. Forced by Atlantic, Walker Circulation and deep convection over the equatorial African rainforest strengthened, causing the increased precipitation in the dry season. Forced by three ocean forcing, especially Indian Ocean forcing, the weakening of Walker Circulation is evident in the wet season, with corresponding stronger divergence and less precipitation.

Figure 8

Simulated seasonal change of moisture and wind at 825hPa in Atmospheric Models. Color mapping illustrates the special humidity change (increase, blue color; decreased, yellow color) and the vector illustrates the wind change. a, b and c show the change forced by the tropical Atlantic, Indian Ocean and Pacific during the boreal winter dry season (DJF). d, e and f same as a, b and c but for the boreal spring wet season (AM). In the dry season (DJF), the model response to Atlantic forcing show the strengthening specific humidity. In the wet season (AM), three models all appear negative specific humidity anomalies but with relative larger magnitudes in Indian Ocean and Pacific forcing.

Figure 9

Simulated seasonal change of vertically integrated moisture flux convergence in Atmospheric Models. Color mapping in a, b and c illustrates the convergence change (increase, blue color; decreased, yellow color) forced by the tropical Atlantic, Indian Ocean and Pacific during the boreal winter dry season (DJF). d, e and f same as a, b and c but for the boreal spring wet season (AM). The changes of convergence agree with specific humidity (Fig.8). In the dry season (DJF), enhanced convergence was driven by Atlantic SST change. However, in the wet season (AM), strengthening divergence occurred in three ocean forcing, with a more widespread area in Indian Ocean forcing.