The Interplay Between Infection Risk Factors of SARS-CoV-2 and Prognosis: A Cross-Sectional Study from a Cohort of Long-Term Care Nursing Home Residents

Laura Soldevila

Institut Català de la Salut

Núria Prat

Institut Català de la Salut

Miquel À. Mas

Institut Català de la Salut

Mireia Massot

Institut Català de la Salut

Ramon Miralles

Institut Català de la Salut

Josep Maria Bonet-Simó

Institut Català de la Salut

Mar Isnard

Institut Català de la Salut

Marta Expósito-Izquierdo

Institut Català de la Salut

Irene García-Sánchez

Institut Català de la Salut

Sara Rodoreda-Noguerola

Institut Català de la Salut

Nemesio Moreno

Institut Català de la Salut

Esther Badia

Institut Català de la Salut

Genís López

Institut Català de la Salut

Javier Sevilla

Institut Català de la Salut

Oriol Estrada

Institut Català de la Salut

Research Article

Keywords: long-term care nursing homes, SARS-CoV-2, epidemiology, mortality, Covid-19

Posted Date: September 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-806540/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Covid-19 pandemic has particularly affected older people living in Long-term Care settings.

Methods: We carried out a cross-sectional analysis of a cohort of Long-term care nursing home residents between March first and June thirty, 2020, who were ≥ 65 years old and on whom at least one PCR test was performed. Socio-demographic, comorbidities, and clinical data were recorded. Facility size and community incidence of SARS-CoV-2 were also considered.

Results: A total of 8021 participants were included from 168 facilities. Mean age was 86.4 years (SD = 7.4). Women represented 74.1%. SARS-CoV-2 infection was detected in 27.7% of participants, and the overall case fatality rate was 11.3% (24.9% among those with a positive PCR test). Epidemiological factors related to risk of infection were larger facility size (pooled aOR 1.73; $P < .001$), higher community incidence (pooled aOR 1.67, $P = .04$), leading to a higher risk than the clinical factor of low level of functional dependence (aOR 1.22, $P = 0.03$). Epidemiological risk factors associated with mortality were male gender (aOR 1.75; $P < .001$), age (pooled aOR 1.16; $P < .001$), and higher community incidence (pooled aOR 1.19, $P = < .001$). There was evidence of clustering for facility and health area when considering the risk of infection and mortality ($P < .001$).

Conclusions: Our results suggest a complex interplay between structural and individual factors regarding Covid-19 infection and its impact on mortality in nursing-home residents.

Introduction

It is estimated that in spite of representing less than five per cent of the elderly population (> 65 years old), 47% of all Covid-19 deaths occurred among residents from Long-Term Care (LTC) nursing homes during the first wave of the SARS-CoV-2 pandemic [1]. This appalling data could be explained by the fact that LTC show the ideal epidemiological conditions for transmission of an airborne infectious agent like SARS-CoV-2 since they are densely populated spaces with large numbers of staff who have extensive contact with highly vulnerable residents. The result has rightly been regarded as a perfect storm [2]. However, more detailed examinations of Covid-19 in LTC have shown that particular structural factors mediated the impact of the disease, including the size of the facility, the quality of care, and understaffing [3-8]. The lack of preparedness as well as short and long-term policy failures also contributed [9,10]. Indeed, besides the direct impact of Covid-19, there have been other secondary consequences due to the lockdowns and other measures intended to contain the infection, among the more important being the exacerbation of comorbidities and delayed medical treatment. Also, psychological, and emotional distress due to social isolation [11-14] and the lack of adequate end-of-life care [15], frequently reported during the first wave of the pandemic, added to the impact.

Most of the previous articles on the impact of Covid-19 in LTC premises have explored risk factors, both individual [16] and structural [6], associated with mortality in LTC residents. In previous articles, we analyzed contextual and clinical factors related to mortality of a local cohort during the first wave of

Covid-19 in Europe [6,17]. In this article we aim to expand previous analysis by focusing on epidemiological factors related to SARS-CoV-2 infection in LTC and the interaction between contextual and societal factors in the evolution and results of SARS-CoV2 outbreaks and clinical prognosis.

Methods

Study population

The study region comprises an area immediately to the north of the city of Barcelona in Catalonia, Spain. Comprising a mixture of urban, semi-rural, and rural municipalities, the area includes a total of 1 986 032 inhabitants, with 190 LTC nursing homes, which as of March first, 2020, housed a total of 10737 residents. Of these LTC, 168 (88.4%) were institutions devoted to elderly adult patients, with a total of 9553 users registered. Lockdown came into force at these institutions on March 15, 2020, and by the end of that month an emergency task force of health professionals had been mobilized to provide back-up support to LTC staff.

Study design

A cross-sectional study was carried out among a cohort study of LTC residents living in the study area between first of March and June 30, 2020, who were ≥ 65 years old and on whom at least one PCR test was performed during the study period. Socio-demographic data (age and gender), and underlying chronic conditions as such as comorbidities, level of functional impairment and advanced conditions were recorded by their primary care teams according to local clinical guidelines [18], laboratory test results (specific Polymerase Chain Reaction henceforth PCR, to detect SARS-CoV-2) and clinical outcome (recovery/death), as well as the size of the LTC facility (number of residents) and cumulative incidence of Covid-19 in the Primary Healthcare Catchment Area where the facility was located (this is the smallest administrative area of the Spanish public healthcare system, typically covering between 15 000 and 25 000 inhabitants attended by a primary health team). Following implemented guidelines, PCR testing was carried out when Covid-19 infection was suspected or when LTC residents had come in close contact with an infected individual (i.e. when three or more confirmed or suspected cases of Covid-19 within the LTC facility including staff members were detected). Regular testing was implemented starting in mid-April 2020 irrespective of infection suspicion or exposure.

Data collection and statistical analysis

Data was entered in the database either by health staff using an in-house app developed *ad hoc*[19] or using the regional health service's digital patient records (socio-demographic and clinical data), alongside PCR test results sent from the reference laboratory serving the region. Data regarding incidence from the catchment area was obtained from the official open-source database [20]. Community incidence excluded all notified cases from LTC located in the specific catchment area. LTC facility capacity (number of residents) and community cumulative incidence were categorized according to interquartile ranges (IQR) of <25, 25-50, 50-75 and >75. PCR tests with inconclusive results were considered negative.

The data was analyzed using Stata v14.0 software (StataCorp, Texas) and R. For the descriptive analysis, we used means, medians, and Standard Deviation (SD) for continuous variables and proportions and a 95% Confidence Interval (95%CI) for categorical variables. Proportions were calculated for all participants with the given data available. For univariate analysis, we used the Chi-Square test to compare categorical variables, and the Student's T test for continuous variables after testing for normality (skewness and kurtosis tests), or their non-parametric counterparts (Fisher test or Wilcoxon test), when necessary. Multilevel mixed-effects logistic regression was used for multivariate analysis to explore for the presence of clustering, and factors (LTC facility and catchment area) and tested by means of regress post-estimation (likelihood ratio test, henceforth LR), against the usual logistic regression model. Crude and adjusted odds ratios (OR and aOR), respective 95%CI and *P* values were estimated.

Laboratory methods (PCR test)

Biological specimens were collected through nasopharyngeal swabs. Specific PCR tests were carried out using the Aptima® SARS-CoV-2 assay following manufacturer instructions [21].

Ethical approval

The study was approved by the Ethics Board of the reference hospital of the study region (Hospital Universitari Germans Trias i Pujol), registered under reference number PI-20-349, and was conducted in accordance with the principles of the Declaration of Helsinki.

Results

Participant characteristics

A total of 9158 residents ≥ 65 years residing in 168 LTC facilities were included in the study, with a median occupancy of 41 (range 12 to 229, IQR 26-41) and distributed over 52 local catchment areas. Of these residents, 8021 (87.6%) underwent at least one PCR test during the study period and were included in the analysis (see Figure 1). Women accounted for 5939 (74.1%) and men for 2079 (25.9%). The mean age was 86.4 years (SD = 7.4, range 65-107), with significant difference between males (83.4 yrs., SD = 7.9) and females (87.4, SD = 6.9; $P < .001$). Overall, 1115 (14.0%) had a cardiovascular comorbidity. The most frequently recorded individual comorbidities were hypertension ($n = 3855$, 48.3%) and dementia ($n = 3666$, 45.9%). A total of 6013 patients presented functional impairment (80.3%) and 4171 (52.0%) an advanced condition. In the four-month study period, SARS-CoV-2 infection was detected in 2225 participants (27.7%), and 909 deaths were reported, of which 554 corresponded to residents with proven SARS-CoV-2 infection (case facility rate, henceforth CFR, 24.9%, see Figure 1). At the LTC facility level, 97 institutions (57.7%) reported at least one confirmed case of SARS-CoV-2 infection, with attack rates ranging from 1% to 95%. The median cumulative incidence of Covid-19 in the 52 catchment areas of the study region during the period of interest was 95 cases/10 000 inhabitants (range 34.7 to 158.2). Table 1 offers a description of the main variables in the total population, and in the subgroup of infected people

and those that dye in the study period. Figure 2 shows the cumulative incidence of SARS-CoV-2 among residents and in catchment areas.

Risk factors associated to SARS-CoV-2 infection and death

Table 2 shows epidemiological risk factors associated with SARS-CoV-2 infection in our analysis. In brief, the risk of infection by SARS-CoV-2 was independent of age, sex, and chronic conditions, except respiratory diseases and cardiovascular disease (P values of .01 and .005, respectively) and low level of functional dependence on daily living (low level of functional dependence score, aOR 1.22; 95%CI 1.0-1.5; $P = .03$). There was a linear correlation between the risk of infection and size of LTC facility (pooled aOR 1.73; 95%CI 1.6-1.9; $P < .001$) and cumulative incidence in the catchment area (pooled aOR 1.77; 95%CI 1.0-3.0; $P = .04$). We observed evidence of clustering by LTC facility and catchment area regarding risk of infection (P value of LR $< .001$). After adjusting for study variables, epidemiological risk factors related to infection that remain associated with mortality included SARS-CoV-2 infection (positive PCR test, aOR 4.26; 95%CI 3.6-5.1; $P < .001$), followed by low functional dependence (aOR 2.42; 95%CI 1.8-3.2; $P < .001$), male gender (aOR 1.75; 95%CI 1.5-2.1; $P < .001$) and older age (pooled aOR 1.16; 95%CI 1.1-1.2; $P < .001$). We observed evidence of clustering regarding mortality with LTC facility and catchment area (P value of LR $< .001$). Furthermore, residents living in nursing homes located in catchment areas with a high incidence of SARS-CoV-2 ($\geq 84/10\ 000$ inhabitants) were at substantially higher risk of mortality (pooled aOR 1.19; 95%CI 1.1-1.3; $P < .001$).

Discussion

Our results indicate that there is a complex interplay between the risk of SARS-CoV-2 infection and subsequent clinical outcome, mediated by individual level factors (age, gender, and chronic conditions) and contextual factors. Particularly, size of residence and community incidence showed the highest influence on infection acquisition, as has been observed previously [6,22], with high OR (3.18 and 2.28, respectively, see Table 2). This may be explained because large nursing homes were more susceptible to a SARS-CoV-2 carrier entering the premises due to the higher number of visiting relatives and working staff. Consequently, larger facilities had a greater likelihood of having one or more cases of Covid-19 compared to smaller ones; (88.1% vs. 37.0%, $P < .001$). However, our data showed highly heterogeneous attack rates once the infection was inside an institution (from 1% to 95%). This may be related to structural factors of each LTC, which showed disparate features (including quality of the service provided and residents' clinical profile), as observed in a previous study involving this study population [6] and a comprehensive study carried out in England [8]. This is consistent with the clustering effect seen here. The correlation between community incidence and risk of infection in LTC has been described previously [23]. At first glance, the straight correlation may reflect higher chances of irruption in LTC of infected individuals from the general population which include LTC staff and visiting relatives before the lockdown. However, it cannot be ruled out that during the first wave and before the lockdown, these institutions could have become, to a certain extent, hubs of SARS-CoV-2 infection in the community. This would have been similar to the role that community health centers played during the 2014-2016 Ebola

crisis in West Africa [24,25]. LTC staff are known to have been heavily infected whilst at work [26,27], especially before the implementation of effective contention measures, and most probably also visiting relatives before the lockdown, and to have spread the infection within the local communities specially in crowded and densely populated areas. If this hypothesis turns out to be true, the rapid intervention in LTC carried out by the health authorities' right at the outset of the Covid-10 pandemic was more relevant than supposed. It is worth noting that we do not consider this to be currently the case due to the strict Covid-19 prevention measures that have been applied in LTC. Moreover, the correlation between mortality and community incidence has been observed elsewhere and by our group [6] in a previous analysis and deserves further assessment using the LTC setting as unit of analysis.

Focusing on gender, we found that being male is not correlated to SARs-CoV-2 infection but it is an independent risk factor for severe Covid-19 disease, as previously observed among elderly populations [16,28]. The underlying reasons have not yet been disentangled, but this strongly suggests some genetic-based susceptibility [29]. In spite of this, considering the disproportion between males/females (25.9% vs. 74.1%), females showed a much higher absolute number of fatalities (600 vs. 309). This cannot simply be explained by the longer life expectancy of women but must also reflect societal gender-based inequalities that increase the likelihood of females ending up in a LTC facility compared to males. Given the disproportionate contribution of LTC residents to the global rate of Covid-19 deaths [1], the observed overrepresentation of female mortality in Spain [30] may be explained by, first, an increased proportion of females in Spanish LTC compared to other countries and, second, the greater impact of the first wave of Covid-19 in LTC observed in this country [30].

Besides these factors, we observed an intriguing interplay between risk of infection and mortality. More autonomous residents (low level of functional dependence) showed a notably higher risk of SARS-CoV-2 infection, but a disproportionate mortality rate as well, as we published previously [17]. This may be explained by the fact that residents with greater autonomy may have had a higher rate of social contacts inside the LTC facility and therefore a higher probability of exposure to infection. The higher risk of mortality once infected may be a consequence of more efficient transmission and/or multiple infections (closeness to and increased frequency of risky contacts) with associated to higher viral loads, which has been correlated with mortality [31]. This association between risk of infection and autonomy as measured by Activities of Daily Living scores has been previously observed but not discussed [3]. The increased risk of infection associated with cardiovascular and respiratory disease could easily be biased in a population with a high prevalence of such conditions. According other reports, besides chronic renal failure [17], we observed a lack of association between negative outcome and other underlying comorbidities described among general populations [28,32,33], and patients with Covid-19 in geriatric care [34]. We hypothesized that in the context of extremely high attack rates in LTC, once SARS-CoV-2 infection occurs tends to behave independently of underlying risk factors, like a primary pathogen. Furthermore, existing medications may have had a substantial effect against infection or severe disease in a population that is usually polymedicated [35].

Our study has some limitations. A substantial number of residents did not undergo PCR testing ($n = 1154$; 12.6%) because it was not possible to mobilize enough material and human resources in time amidst the extremely demanding circumstances of the early stages of the epidemic wave, and only later on was general screening implemented. As in other reports, our data indicate that many deaths occurred in people who were infected with Covid-19 but not tested [36]. The CFR of this sample subset (those that did not undergo a PCR test) was 69.2% ($n = 798$, see Figure 1), which increases the death toll rate of our study population to 18.6%. This toll is close to the expected total mortality rate among our cohort for an entire year [37]. Considering that most fatalities were Covid-19 patients, the specific CFR among SARS-CoV-2-infected residents was around 50%. This estimate lies in the upper bounds of previous estimates [1,27,38]. Therefore, because it included only patients for whom a PCR test was available and thus possibly excludes the most vulnerable patients, our analysis could be to a certain extent biased. Non-tested residents tended to be older (87.4 years vs. 86.4, $p < .001$), although the gender distribution was similar ($p = .2$).

Overall, our results may explain the much higher CFR (around 40%) of older adults living in LTC infected by SARS-CoV-2 compared to the general CFR observed in people older than 80 years old, estimated to be 14.5% [39]. due to the interaction between of structural (institutional), as the clustering of our data suggests, which were in turn consequence of societal and implemented policies, and individual factors (related to chronic/advanced conditions, age, gender and autonomy level, as previously discussed [17], which may be more or less prevalent in each facility. However, it is worth noting that these results correspond to the conditions of the first wave of a pandemic, which tends to be characterized by a lack of preparedness against an unknown pathogen. Therefore, other risk factors correlated to infection and negative outcome in LTC may emerge in later stages of the pandemic.

In spite of the overwhelmingly positive impact of the vaccination of LTC residents [40,41], given the current uncertainty about the role of vaccines in preventing transmission in nursing homes at mid and long term, as well as the impact of new variants of the virus [42], research on Covid-19 in LTC facilities should still be prioritized. The scope of research Covid-19 in nursing homes should be expanded, as well, to ascertain the indirect consequences of not only lockdown measures but also measures intended to mitigate these effects (i.e. to safely allow family visits) [43-45] as well as the ways by which such measures might be avoided in the future [46] Furthermore, long-term policies should address structural factors underlined by our results and previous publications, to create a more adequate elderly care system.

Conclusions

There was a complex interplay between structural and individual factors regarding Covid-19 infection and its impact on mortality in nursing-home residents during the first epidemic wave. The overwhelming impact on LTC residents could be partially explained by the intersect of lack of preparedness and structural factors. Risk of SARS-CoV-2 infection was mainly associated to contextual level factors (i.e. LTC structure), whereas risk of mortality to individual level factors (i.e. frailty). Long-term policies should

be implemented to develop a residential system more resilient to epidemic surges. Secondary effects due to lockdowns and social restrictions among elderly residents should be investigated.

Declarations

Funding: This research did not receive any funding from agencies in the public, commercial, or not-for-profit sectors.

Competing interests: None to declare.

Availability of data and materials: The data underlying this article cannot be shared publicly due to it is subject to Ethical approval before accessing to it. The data will be shared on reasonable request to the corresponding author and with previous Ethical Approval by the Ethics Board of reference.

Authors contribution: LS, NP, MAM, RM, MM, OE and XV conceptualized and designed the study. JB, MI, ME, IG, SR, NM, EB, GL and JS contributed to the data collection, literature revisions and systematization. JS, GL and XV carried out the statistical analysis and performed the tables and figures. LS, MAM, RM, MM and XV drafted the manuscript and made final revisions. All authors critically revised, read and approved the final manuscript

Ethics approval: The study was approved by the Ethics Board of the reference hospital of the study region (Hospital Universitari Germans Trias i Pujol), registered under reference number PI-20-349, and was conducted in accordance with the principles of the Declaration of Helsinki.

Consent to participate: The consent form from participants was waived since all study data was retrospectively obtained from the routine clinical data registration system with previous anonymization.

Consent for publication: All authors consent with the publication in the BMC Geriatrics

References

1. Comas-Herrera A, Zalakaín J, Lemmon E, et al. In: Mortality associated with Covid-19 outbreaks in care homes Mortality associated with Covid-19 outbreaks in care homes: early international evidence, 2020. <https://ltcCovid.org/2020/04/12/mortality-associated-with-Covid-19-outbreaks-in-care-homes-early-international-evidence>. Accessed June 15, 2021.
2. Ouslander JG, Grabowski DC. Covid-19 in Nursing Homes: Calming the Perfect Storm. *J Am Geriatr Soc* 2020;<https://doi.org/10.1111/jgs.16995>
3. Unruh MA, Yun H, Zhang Y et al. Nursing Home Characteristics Associated With Covid-19 Deaths in Connecticut, New Jersey, and New York. *J Am Med Dir Assoc* 2020;<https://doi.org/10.1016/j.jamda.2020.06.019>
4. Abrams HR, Loomer L, Gandhi A, Grabowski DC. Characteristics of U.S. Nursing Homes with Covid-19 Cases. *J Am Geriatr Soc* 2020; <https://doi.org/10.1016/10.1111/jgs.16661>.

5. Chapman S, Harrington C. Policies Matter! Factors Contributing to Nursing Home Outbreaks During the Covid-19 Pandemic Policy. *Polit Nurs Pract* 2020; <https://doi.org/10.1177/1527154420959021>
6. Suñer C, Ouchi D, Mas MA et al. Risk factors for mortality of residents in nursing homes with Covid-19: A retrospective cohort study. *Nat Aging* 2021; <https://doi.org/10.1038/s43587-021-00079-7>
7. Belmin J, Georges S, Franke F et al. Coronavirus Disease 2019 in French Residential Care Facilities: A Nationwide Study. *J Am Med Dir Assoc* 2021; <https://doi.org/10.1016/j.jamda.2021.03.013>
8. Shallcross L, Burke D, Abbott O et al. Factors associated with SARS-CoV-2 infection and outbreaks in long-term care facilities in England: a national cross-sectional survey. *Lancet Healthy Longev* 2021; [https://doi.org/10.1016/S2666-7568\(20\)30065-9](https://doi.org/10.1016/S2666-7568(20)30065-9)
9. Werner RM, Hoffman AK, Coe NB. Long-Term Care Policy after Covid-19 – Solving the Nursing Home Crisis. *N Engl J Med* 2020; <https://doi.org/10.1056/NEJMp2014811>
10. O'Neill D, Briggs R, Holmerová I, Samuelsson O et al. Covid-19 highlights the need for universal adoption of standards of medical care for physicians in nursing homes in Europe. *Eur Geriatr Med* 2020; <https://doi.org/10.1007/s41999-020-00347-6>
11. Miller EA. Protecting and Improving the Lives of Older Adults in the Covid-19 Era. *J Aging Soc Policy* 2020; <https://doi.org/10.1080/08959420.2020.1780104>
12. Simonetti A, Pais C, Jones M et al. Neuropsychiatric Symptoms in Elderly With Dementia During Covid-19 Pandemic: Definition, Treatment, and Future Directions. *Front Psychiatry* 2020; <https://doi.org/10.3389/fpsy.2020.579842>
13. Lombardo FL, Salvi E, Lacorte E et al. Adverse Events in Italian Nursing Homes During the Covid-19 Epidemic: A National Survey. *Front Psychiatry* 2020; <https://doi.org/10.3389/fpsy.2020>.
14. Ayalon L, Zisberg A, Cohn-Schwartz E et al. Long-term care settings in the times of Covid-19: challenges and future directions. *Int Psychogeriatr* 2020; <https://doi.org/10.1017/S1041610220001416>
15. Strang P, Bergström J, Martinsson L et al. Dying From Covid-19: Loneliness, End-of-Life Discussions, and Support for Patients and Their Families in Nursing Homes and Hospitals. A National Register Study. *J Pain Symptom Manage* 2020; <https://doi.org/10.1016/j.jpainsymman.2020.07.020>
16. Panagiotou OA, Kosar CM, White EM et al. Risk Factors Associated With All-Cause 30-Day Mortality in Nursing Home Residents With Covid-19. *JAMA Intern Med* 2021; <https://doi.org/10.1001/jamainternmed.2020.7968>
17. Mas MA, Massot M, Miralles R et al. Clinical factors related to Covid-19 outcomes in institutionalized older adults: cross-sectional analysis from a cohort in Catalonia. *J Am Med Dir Assoc* 2021; <https://doi.org/10.1016/j.jamda.2021.07.004>
18. Contel JC, Ledesma A, Blay C et al. Chronic and integrated care in Catalonia. *Int J Integr Care*. 2015; <https://doi.org/10.5334/ijic.2205>
19. Echeverría P, Mas Bergas MA, Puig J et al. CovidApp as an Innovative Strategy for the Management and Follow-Up of Covid-19 Cases in Long-Term Care Facilities in Catalonia: Implementation Study. *JMIR Public Health Surveill* 2020; <https://doi.org/10.2196/21163>

20. AQUAS. In: Map of Coronavirus Incidence from Catalonia. 2020.
<https://aquas.gencat.cat/ca/actualitat/ultimes-dades-coronavirus/mapa-per-municipis/>. Accessed February 2021
21. Pham J, Meyer S, Nguyen C et al. Performance Characteristics of a High-Throughput Automated Transcription-Mediated Amplification Test for SARS-CoV-2 Detection. *J Clin Microbiol* 2020;<https://doi.org/10.1128/JCM.01669-20>
22. Costa AP, Manis DR, Jones A et al. Risk factors for outbreaks of SARS-CoV-2 infection at retirement homes in Ontario, Canada: a population-level cohort study. *CMAJ* 2021;<https://doi.org/10.1503/cmaj.202756>
23. Chatterjee P, Kelly S, Qi M, Werner RM. Characteristics and Quality of US Nursing Homes Reporting Cases of Coronavirus Disease 2019 (Covid-19). *JAMA Netw Open* 2020;
<https://doi.org/10.1001/jamanetworkopen.2020.16930>.
24. Fischer WA, Hynes NA, Perl TM. Protecting Health Care Workers From Ebola: Personal Protective Equipment Is Critical but Is Not Enough. *Ann Intern Med* 2014;<https://doi.org/10.7326/M14-1953>.
25. Kilmarx PH, Clarke KR, Dietz PM et al. Ebola virus disease in health care workers—Sierra Leone, 2014. *MMWR Morb Mortal Wkly Rep* 2014; 12;63:1168-71.
26. Ladhani SN, Chow JY, Janarthanan R et al. Increased risk of SARS-CoV-2 infection in staff working across different care homes: enhanced Covid-19 outbreak investigations in London care Homes. *J Infect* 2020;<https://doi.org/10.1016/j.jinf.2020.07.027>.
27. Sacco G, Foucault G, Briere O, Annweiler C. Covid-19 in seniors: Findings and lessons from mass screening in a nursing home. *Maturitas* 2020; <https://doi.org/10.1016/j.maturitas.2020.06.023>.
28. Chang TS, Ding Y, Freund MK et al. Prior diagnoses and medications as risk factors for Covid-19 in a Los Angeles Health System. *medRxiv [Preprint]* 2020; <https://doi.org/10.1101/2020.07.03.20145581>.
29. Cao Y, Li L, Feng Z, Wan S et al. Comparative genetic analysis of the novel coronavirus (2019-nCoV/SARS-CoV-2) receptor ACE2 in different populations. *Cell Discov* 2020;<https://doi.org/10.1038/s41421-020-0147-1>
30. Kontis V, Bennett JE, Rashid T et al. Magnitude, demographics and dynamics of the effect of the first wave of the Covid-19 pandemic on all-cause mortality in 21 industrialized countries. *Nat Med* 2020;
<https://doi.org/10.1038/s41591-020-1112-02014;63:1168-71>.
31. Pujadas E, Chaudhry F, McBride R et al. SARS-CoV-2 viral load predicts Covid-19 mortality. *Lancet Respir Med* 2020; [https://doi.org/doi:10.1016/S2213-2600\(20\)30354-4](https://doi.org/doi:10.1016/S2213-2600(20)30354-4).
32. Zheng Z, Peng F, Xu B et al. Risk factors of critical & mortal Covid-19 cases: A systematic literature review and meta-analysis. *J Infect* 2020; <https://doi.org/10.1016/j.jinf.2020.04.021>.
33. Fang X, Li S, Yu H et al. Epidemiological, comorbidity factors with severity and prognosis of Covid-19: a systematic review and meta-analysis. *Aging (Albany NY)* 2020;
<https://doi.org/10.18632/aging.103579>.
34. Hägg S, Jylhävä J, Wang Y et al. Age, Frailty, and Comorbidity as Prognostic Factors for Short-Term Outcomes in Patients With Coronavirus Disease 2019 in Geriatric Care. *J Am Med Dir Assoc* 2020;

- <https://doi.org/10.1016/j.jamda.2020.08.014>.
35. Soldevila L, Valerio-Sallent L, Roure S et al. Drug exposure may have a substantial influence on Covid-19 prognosis among residents of long-term care facilities: an exploratory analysis. *Int J Infect Dis.* 2021;<https://doi.org/10.1016/j.ijid.2021.07.007>.
 36. Dutey-Magni PF, Williams H, Jhass A et al. Covid-19 infection and attributable mortality in UK care homes: Cohort study using active surveillance and electronic records (March-June 2020). *Age Ageing* 2021; <https://doi.org/10.1093/ageing/afab060>.
 37. Amblàs-Novellas J, Santaegència SJ, Vela E et al. What lies beneath: A retrospective, population-based cohort study investigating clinical and resource-use characteristics of institutionalized older people in Catalonia. *BMC Geriatr* 2020;<https://doi.org/10.1186/s12877-020-01587-8>
 38. Graham NSN, Junghans C, Downes R et al. SARS-CoV-2 infection, clinical features and outcome of Covid-19 in United Kingdom nursing homes. *J Infect* 2020; <https://doi.org/10.1016/j.jinf.2020.05.073>.
 39. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (Covid-19) Outbreak in China: Summary of a Report of 72314 Cases From Chinese Center for Disease Control and Prevention. *JAMA* 2020;<https://doi.org/10.1001/jama.2020.2648>
 40. Mor V, Gutman R, Yang X et al. Short-term impact of nursing home SARS- CoV-2 vaccinations on new infections, hospitalizations, and deaths. *J Am Geriatr Soc.* 2021;<https://doi.org/10.1111/jgs.17176>
 41. White EM, Yang X, Blackman C, Feifer RA, Gravenstein S, Mor V. Incident SARS-CoV-2 Infection among mRNA-Vaccinated and Unvaccinated Nursing Home Residents. *N Engl J Med* 2021; <https://doi.org/10.1056/NEJMc2104849>.
 42. National Collaborating Centre for Methods and Tools. In:What strategies mitigate risk of Covid-19 outbreaks and mortality in long-term care facilities? Update 2. <https://www.nccmt.ca/knowledge-repositories/Covid-19-rapid-evidence-service>. Accessed April 2021
 43. Verbeek H, Gerritsen DL, Backhaus R et al. Allowing Visitors Back in the Nursing Home During the Covid-19 Crisis: A Dutch National Study Into First Experiences and Impact on Well-Being. *J Am Me Di Assoc* 2020;<https://doi.org/10.1016/j.jamda.2020.06.020>
 44. Hado E, FrissFeinberg L. Amid the Covid-19 Pandemic, Meaningful Communication between Family Caregivers and Residents of Long-Term Care Facilities is Imperative. *J Aging Soc Policy* 2020; <https://doi.org/10.1080/08959420.2020.1765684>
 45. van Dyck LI, Wilkins KM, Ouellet J et al. Combating Heightened Social Isolation of Nursing Home Elders: The Telephone Outreach in the Covid-19 Outbreak Program. *Am J Geriatr Psychiatry* 2020;<https://doi.org/10.1016/j.jagp.2020.05.026>
 46. Anderson DC, Grey T, Kennelly S, O'Neill D. Nursing Home Design and Covid-19: Balancing Infection Control, Quality of Life, and Resilience. *J Am Med Dir Assoc* 2020;<https://doi.org/10.1016/j.jamda.2020.09.005>

Tables

Table 1: Prevalence of study variables stratified by infection

	Total ^a		PCR+			Death ^b		
	(N = 8021)		(N = 2225; 27.7%)			(N = 909; 11.3%)		
Variable	n	%	n	%	<i>P</i>	n	%	<i>P</i>
Gender								
Male	2079	25.9	597	28.7	.2	309	14.7	< .001
Female	5939	74.1	1626	27.4		600	10.1	
Age in years (m, SD)	86.4	(7.4)	86.5	(7.3)	.2	87.4	(7.1)	< .001
65-74	638	8.0	166	26.0	.2	47	5.2	< .001†
75-79	752	9.4	210	27.9		83	9.1	
80-84	1362	17.0	346	25.4		133	14.6	
85-89	2289	28.6	663	29.0		260	28.6	
≥90	2977	37.1	838	28.2		386	42.5	
Level of functional dependence (low/high) ^c								
High level of functional dependence	6013	80.3	1497	24.9	< .001	357	5.9	< .001
Low level of functional dependence	1475	19.7	447	30.3		178	12.1	
Comorbidities								
Hypertension	3855	48.3	1164	30.2	< .001	488	12.7	.001
Diabetes Mellitus-II	1593	20.0	471	29.6	.08	208	13.1	.02
Chronic renal insufficiency	1564	19.6	451	28.8	.3	230	14.7	< .001
Dementia	3666	45.9	1002	27.3	.4	472	12.9	< .001
Cardiovascular disease ^d	1115	14.0	363	32.6	< .001	170	15.3	< .001
Respiratory disease ^e	824	10.3	262	31.8	.007	124	15.1	< .001
Cerebrovascular disease	211	2.6	46	21.8	.05	28	13.3	.4
Clinical complexity								

Complex Chronic Condition	723	17.3	202	27.9	.9	533	12.8	< .001
Number of residents								
≤ 40 (N = 84)	2060	25.7	1464	17.6	< .001	194	9.4	< .001 ^f
41-72 (N = 44)	1986	24.8	1419	22.3		196	9.9	
73-108 (N = 25)	2049	25.6	1512	31.0		239	11.7	
> 108 (N = 16)	1926	24.0	1407	40.7		280	14.5	
Community incidence								
≤ 54.7	1994	25.4	1811	16.9	< .001	137	6.9	< .001 ^f
54.8-72.9	1903	24.2	1620	28.6		221	11.6	
73.0-83.9	1840	23.4	1515	30.0		200	10.9	
≥ 84.0	2115	26.9	1769	36.8		340	16.1	

a. Includes participants with data available.

b. Includes mortality among participants with PCR test result available.

c. Levels of functional dependence have been calculated using the Barthel score, which provides a range from 0 to 100. The cut-off between high and low functional dependence has been set up at 50.

d. Includes ischemic heart disease and heart insufficiency.

e. Include Chronic Obstructive Pulmonary disease, asthma, and emphysema.

f. *P* test for trend.

Table 2: Risk factors associated with SARS-CoV-2 infection (PCR+)

Variable	Crude ^a			Adjusted		
	OR	95%CI	<i>P</i>	OR	95%CI	<i>P</i>
RT-PCR positive	—	—	—	—	—	—
Gender						
Male	1.11	(0.9-1.3)	.2	1.09	(0.9-1.3)	.3
Female	1	—	—	—	—	—
Age (years)						
65-74	1	—	—	1	—	—
75-79	1.08	(0.8-1.5)	.6	0.98	(0.7-1.4)	0.9
80-84	1.02	(0.8-1.4)	.9	1.04	(0.8-1.4)	0.8
85-89	1.40	(1.1-1.8)	.02	1.34	(1.0-1.8)	.06
≥ 90	1.28	(0.98-1.66)	.07	1.29	(0.96-1.73)	.09
Age (pooled)	1.08	(1.0-1.1)	.008	1.08	(1.0-1.2)	.005
Level of functional dependence (low/high)						
Low level	1.25	(1.1-1.5)	.01	1.22	(1.0-1.5)	.03
Comorbidities						
Respiratory disease	1.37	(1.1-1.7)	.007	1.36	(1.1-1.7)	.01
Cardiovascular disease	1.40	(1.2-1.7)	.001	1.38	(1.1-1.7)	.005
Cerebrovascular disease	0.70	(0.4-1.1)	.1	0.74	(0.5-1.2)	.2
Hypertension	1.05	(0.9-1.2)	.5	1.02	(0.9-1.2)	.8
Dementia	0.99	(0.9-1.1)	.8	0.95	(0.8-1.1)	.6
Chronic renal insufficiency	1.10	(0.9-1.3)	.3	1.03	(0.9-1.3)	.8
Diabetes Mellitus-II	1.09	(0.9-1.3)	.3	1.00	(0.8-1.0)	.9
Clinical complexity						
Complex Chronic condition	0.94	(0.8-1.1)	.4	0.89	(0.8-1.0)	.1
Number of residents ^{b,c}						
≤ 40 (N = 84)	1	—	—	1	—	—
41-72 (N = 44)	1.95	(1.6-2.4)	<	1.47	(1.2-1.8)	<

			.001			.001
73-108 (N = 25)	2.54	(2.1-3.1)	< .001	1.96	(1.6-2.4)	< .001
> 108 (N = 16)	5.31	(4.2-6.7)	< .001	3.18	(2.6-3.9)	< .001
Pooled OR	1.66	(1.6-1.8)	< .001	1.67	(1.6-1.8)	< .001
Community incidence ^{d,e}						
≤ 54.7	1	—	—	1	—	—
54.8-72.9	1.99	(0.4-9.9)	.4	1.92	(1.5-2.4)	< .001
73.0-83.9	1.90	(0.2-8.6)	.4	1.92	(1.5-2.4)	< .001
≥ 84.0	5.70	(1.2-26.2)	.03	2.28	(1.9-2.8)	< .001
Pooled OR	1.68	(1.0-2.7)	.04	1.67	(1.0-2.7)	.04

a.Adjusted by clustering (LTC facility and catchment area).

b.N indicates the number of LTC institutions included in each strata.

c.Adjustments for clustering included only the catchment area.

d.Cumulative cases per 10,000 inhabitants during the study period.

e.Adjustment for clustering included only LTC facility.

Figures

Figure 1

Flow-chart of study population and participants with PCR test available

Figure 2

Cumulative incidence of SARS-CoV-2 infection of LTC residents and corresponding catchment area