Resting-State Functional Connectivity of Salience Network in Schizophrenia and Depression

Huan Huang

Renmin Hospital of Wuhan University

Cheng Chen

Renmin Hospital of Wuhan University

Bei Rong

Renmin Hospital of Wuhan University

Qirong Wan

Renmin Hospital of Wuhan University

Jingang Chen

Zhongxiang Hospital of Renmin Hospital of Wuhan University

Zhongchun Liu

Renmin Hospital of Wuhan University

Yuan Zhou

Chinese Academy of Sciences

Gaohua Wang

Renmin Hospital of Wuhan University

Huiling Wang (✉ hlwang@whu.edu.cn)

Renmin Hospital of Wuhan University

Research Article

Keywords: Resting-state functional connectivity, salience network, schizophrenia, depression, resting-state functional magnetic resonance imaging (rs-fMRI), pattern

Posted Date: August 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-808232/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

To explore the salience network(SN) functional alterations in schizophrenia and depression, resting-state functional magnetic resonance imaging (rs-fMRI) data from 29 patients with schizophrenia (SCH), 28 patients with depression (DEP) and 30 healthy controls(HC) were obtained. The SN was derived from data-driven group independent component analysis (gICA). ANCOVA and post hoc tests were performed to discover the FC differences of SN between groups. The ANCOVA demonstrated a significant group effect in FC with right inferior and middle temporal gyrus (ITG and MTG), left caudate, and right precentral gyrus. Post-hoc analyses revealed an opposite altered FC pattern between SN and right ITG and MTG for both patient groups. The DEP group showed a reduced FC between SN and right ITG and MTG compared with HC whereas the SCH group showed an increased FC. In addition, the SCH group showed decreased FC between SN and left caudate, and enhanced FC between SN and right precentral gyrus compared to the other two groups. Our findings suggest distinct FC of SN in schizophrenia and depression, supporting that the resting-state FC pattern of SN may be a transdiagnostic difference between depression and schizophrenia and may play a critical role in the pathogenesis of these two disorders.

Introduction

Schizophrenia and depression are two of the three most common serious psychiatric disorders. Considerable overlap in symptomatology and clinical features such as emotional and cognitive dysregulation could be seen in both diseases¹, and genetic studies have also identified some common polymorphisms². All these findings suggest that they may share neurobiological commonalities, as well as disease-specific mechanisms.

In recent years, with rapid development of neuroimaging techniques, researchers have widely explored the brain in people with mental disorders, and various structural and functional changes have been associated with both disorders. Among these, resting state functional magnetic resonance imaging (rs-fMRI) has been widely used to explore the biological markers in psychiatry, as they are relatively simple and easy to acquire, particularly in patients with mental disorders^{3,4}. Functional connectivity (FC) is the main research method in this field and represents brain regional co-activation, which is a statistical relationship showing that neural activation levels in spatially diverse brain regions are associated in time dimension⁵. Previous evidence has illustrated that both SCH and DEP have been generally identified in relation to the aberrant functioning and organization of large-scale resting state networks (RSNs)^{3,6-8}. RSNs are defined by the specific FC of ongoing, slowly fluctuating brain activity, including typical default mode network (DMN), the central executive network (CEN), and the salience network (SN), etc⁹.

Studies that direct compare between SCH and DEP are rather less and the results are inconsistent. Schilbach et al assessed transdiagnostic aberrations of resting state FC of DMN in schizophrenia and depression patients from multi-sites using seed-based resting state analyses and their results demonstrated a common decrease of FC between precuneus and bilateral superior parietal lobe in

both disorders, and further identified the reduced FC between DMN and parietal operculum as the diagnosis-specific feature in schizophrenia relative to depression¹⁰. Resting-state FC pattern in the prefrontal cortex has been regarded as a potential transdiagnostic difference between depression and schizophrenia patients¹¹. Jiang et al common/distinct structural deficits and aberrant causal connectivity patterns in the prefrontal-thalamic-cerebellar circuit in SCH and DEP¹². In addition, recent studies support that SN abnormality may play a critical role in the pathogenesis of schizophrenia and depression^{13,14}. The SN includes prominent nodes in the anterior insula (AI) and dorsal anterior cingulate cortex (dACC), as well as the rostral prefrontal cortex (RPFC) and the supramarginal gyrus (SMG). The SN also includes subcortical limbic areas including the striatum, thalamus and amygdala, which have been identified in fMRI studies¹⁵⁻²⁰. The function of SN might be to identify the most homeostatically relevant among multiple internal and external stimuli that constantly accost the brain, and is commonly coactivated across a wide range of cognitive and affective tasks^{17,21}. Moreover, researchers have found that SN manifested the highest levels of time-varying flexibility in connectivity with other networks and suggested the SN is a unique hub for driving inter-network interactions and acts as a “dynamic switch”¹⁹.

However, the FC abnormalities in SN between the two disorders are still unclear.

In this study, we used rs-fMRI to investigate FC patterns of SN in patients with schizophrenia, depression, and a matched group of health controls. We defined the SN from the data-driven group independent component analysis (gICA), and further explored the FC between the SN and the whole brain.

Material And Methods

Participants

This study recruited 30 individuals with schizophrenia (SCH group), 30 individuals with depression (DEP group), and 30 healthy controls (HC group). The participants from three groups were matched for age, gender and years of education. All patients were recruited from the inpatients and outpatients at the department of psychiatry, Renmin Hospital of Wuhan University. The diagnose of schizophrenia or major depression disorder was made by two clinical psychiatrists independently according to the DSM-VI Disorder-Clinical Version (SCID-CV). Symptom severity was assessed using the Positive and Negative Syndrome Scale (PANSS) for schizophrenia patients and the 17-items Hamilton Depression Rating Scale (HAMD-17) for depression patients. All of the patients were at their acute episodes. For the first-episode patients, they have never taken any antidepressants or antipsychotics before being recruited in this study; for the multiple-episode patients, they discontinued their psychotropic drug treatment for more than 6 months. However, all patients were medicated following established medication regimes by their psychiatrist for less than one week. The healthy participants were recruited from the hospital faculty and local community via advertisement. They had neither lifetime psychiatric disorder nor family history of psychosis in their first-degree relatives.

Participants in the three groups were excluded if they had any neurologic abnormalities, organic mental disorders, other serious physical illnesses, dementia, brain injuries, substance abuse or addiction, or contraindications to MRI. Three participants were excluded from data analyses due to bad image quality, leaving 30 healthy controls, 28 patients with depression and 29 patients with schizophrenia to be included in the analyses.

This study was approved by the Ethics Committee of Renmin Hospital of Wuhan University, and all participants provided their written informed consent individually after the study was completely described to each participant. In addition, we confirm that all methods were carried out in accordance with relevant guidelines and regulations.

Imaging Data Acquisition and Preprocessing

Whole-brain imaging data were performed on a 3.0T General Electric (GE) Signa HDxt MR scanner at the department of radiology, Renmin Hospital of Wuhan University. The subjects were instructed to keep their eyes closed without falling asleep, relax, and not to focus their thoughts on anything in particular.

Functional images were collected using a gradient echo planar imaging (EPI) sequence with following main scan parameters: TR = 2000 ms; TE = 30 ms; flip angle = 90°; FOV = 220 mm × 220 mm; matrix = 64 × 64; slice = 32; slice thickness = 4 mm; gap = 0.6 mm. Each functional scan session lasted 8, resulting in 240 volumes to be obtained. In addition, high-resolution T1-weighted structural images were obtained using a magnetization-prepared rapid acquisition gradient echo sequence (TE = 7.8 ms; TR = 3.0 ms; flip angle = 7°; inversion time = 1100ms; FOV = 256mm × 256 mm; matrix= 256 × 256; slice = 188; voxel size = 1mm × 1mm× 1mm).

The functional data preprocessing was performed in MATLAB using the CONN toolbox²² and SPM12(Welcome Trust Centre for Neuroimaging; London, UK). All functional images underwent the default preprocessing pipeline for volume-based analyses, which consisted of (1) removing the first five volumes to allow the subjects' adaptation to the scanning environment and make signal equilibrium, (2) functional realignment and unwrap, (3) functional slice-timing correction,(4) functional outliers detection(ART-based identification of outliers scans for scrubbing),(5)functional direct segmentation and normalization to Montreal Neurological Institute (MNI) space, (6) structural segmentation and normalization to MNI space, (7) functional smoothing using a Gaussian kernel with full-width at half-maximum of 6 mm. For the scrubbing, outliers were defined as composite movement greater than 2 mm or more than 3 standard deviations away from the mean image intensity, individuals were excluded if > 20% of time points were identified as outliers; however, no participants met this exclusion criterion. Then in the denoising step, all data were default band-pass filtered (0.008–0.09 Hz), and the white matter and cerebrospinal fluid were regressed out and head motion related artifacts were reduced using the anatomical component-based noise correction (CompCor) strategy²³.

Independent Component Analysis

Using default settings for CONN 19c, the data-driven ICA was run with a G1 FastICA for estimation of independent spatial components and GICA3 back-projection for individual subject level spatial map estimation. Dimensionality reduction was set to 64 and the number of components was set to 40 by default.

The correlational spatial match-to-template approach was used in CONN to identify the component representing salience network, which showed the best match to the SN template from HCP ($r=0.320$). This component includes SN major cortical nodes including bilateral AI and dACC, as well as bilateral rostral prefrontal cortex, and subcortical nodes can also be detected albeit at a weaker level. Spatial distribution of SN component is highly similar to previous studies^{17,24,25} (Figure 1). Then this component was defined as the network of interest for the following analyses.

Functional connectivity analysis

Pearson's correlation coefficients were calculated between the time series of SN component and that of all voxels in the whole brain, the Fisher's r-to-z transform was applied for increasing normality. The FC strength between SN and brain regions was represented by Pearson's correlation coefficients. The FC maps of SN in each of the three groups were obtained by one-tailed one-sample t-test (voxel-level $p < 0.05$, FWE-corrected), as shown in Figure 2. To investigate the effect of group and differences between groups, one-way analysis of covariance (ANCOVA) and post-hoc LSD tests were performed on the FC maps respectively, controlling for age, gender, years of education, and head-motion covariable (FD value). The results were displayed at a voxel-wise $p < 0.001$ uncorrected and a cluster-wise threshold of $p < 0.05$ FWE-corrected.

Correlation analyses

We further explored the relationships between altered FC of SN and clinical variables. The correlation analyses were performed separately for each patient group. Partial correlations between FC values and illness severity measures (HDMA or PANSS) and DOI were calculated, controlling for age, gender, years of education, and head-motion covariable (FD value).

Results

Demographic and Clinical data

Demographic variables across the three groups were compared with the one-way analysis of variance (ANOVA) or independent sample t tests for continuous variables and Chi-square tests for categorical variables. The three groups had no difference in gender, age and education (all $p \geq 0.05$). There was also no significant difference in duration of illness between the two patient groups (Detailed information can be seen in Table 1).

Functional Connectivity of SN

The ANCOVA demonstrated a significant group effect in FC with right inferior and middle temporal gyrus (ITG and MTG), left caudate, and right precentral gyrus. Subsequently post-hoc LSD analyses revealed an opposite altered FC pattern between SN and right ITG and MTG for both patient groups (Figure 3). The DEP group showed a reduced FC between SN and right ITG and MTG compared with HC whereas the SCH group showed an increased FC. In addition, the SCH group showed decreased FC between SN and left caudate, and enhanced FC between SN and right precentral gyrus compared to the other two groups. The results from the ANCOVA and post hoc analyses are presented in Table 2 and Table 3 respectively.

Correlations between abnormal FC and clinical variable

There was no significant correlation between FC strength with significant group effects and clinical variables in the analyses ($p \geq 0.05$).

Discussion

In the current study, we used a data-driven gICA to identify the SN and further investigated the FC of SN among patients with schizophrenia and depression by the resting state fMRI. We found that the SCH group and the DEP group showed completely opposing FC between SN and three regions, including right ITG and MTG, left caudate and right precentral gyrus. In detail, compared with the HC group, the FC between the SN and the right ITG and MTG was enhanced in the SCH group but decreased in the DEP group; the patients with SCH showed decreased FC between the SN and left caudate whereas increased FC between SN and right precentral gyrus. These findings suggest aberrant resting-state FC pattern of SN may be a transdiagnostic difference between depression and schizophrenia patients and may play a critical role in the pathogenesis of these two disorders.

The role of the SN in the neural mechanism of psychiatric disorders has been the subject of interest for many researchers, given that the SN is crucial to sustain human emotion and cognition, especially during the detection and processing of salient information¹⁵. Both schizophrenia and depression are characterized by impairments of cognitive and emotion function. Many SN-related studies have been conducted in schizophrenia, including patients at first episode and chronic phase, as well as with high-risk²⁴⁻²⁶. The Dong et al conducted a meta-analysis of resting-state FC studies in schizophrenia, and they found SN plays a prenominal role in large-scale brain networks, the dysconnectivity of SN may lead to the failure of schizophrenia patients to differentiate self-representation (inner world) and environmental salience processing (outside world)²⁷. For the individual with depression, the studies focusing on the SN is relatively less, a meta-analysis study reviewed researches using either seed-based correlation or independent component analysis, and found that increased connectivity between the SN and the anterior DMN²⁸.

In this study, we found that compared to the HC group, the FC between the SN and right ITG and MTG was enhanced in SCH group but decreased in the DEP group, and displayed in the opposite direction in two groups. The patients with schizophrenia showed positive FC whereas the patients with depression

showed negative FC. The temporal gyrus plays an important role in the processing of emotion, memory, and mental activities, and the ITG and MTG with a significant group effect identified here are mainly located in the DMN, which reflects an introspective and spontaneous activity in a non-tasking state and is involved in internal psychological processing and external environmental monitoring^{29,30}. Therefore, we assume that SN-DMN dysconnectivity is significantly associated with both schizophrenia and depression, but in a completely different way. In schizophrenia, the positive FC between the SN and temporal gyrus indicated that the ITG and MTG were coactive with the SN at resting-state, which means an impaired segregation of task positive regions (SN) and task negative (the ITG and MTG) brain networks in schizophrenia. In contrast, negative FC was demonstrated between the SN seed and the temporal gyrus, and is even stronger in depression. The negative FC suggests an anticorrelation, which presumably reflects an active downregulation or inhibition in depression.

Another notable SN FC abnormality observed in the present study suggests a disconnection between SN and subcortical caudate during the resting state in schizophrenia, which is not affected in depression. We found that the SCH group exhibited a negative FC between the SN and left caudate, while the DEP and HC groups exhibited a positive FC. The caudate is located in the rostral striatum and commonly regarded as the subcortical region of SN³¹, which means it should have positive FC with SN, as shown in the healthy controls and patients with depression in our study. However, the SN-caudate in the SCH group shifted from positive to negative FC, suggesting a significant functional disconnection within the SN in schizophrenia, and this disconnectivity was consistent with our previous study with first episode schizophrenia, which demonstrated that the reduced FC existed within cortico-striatal-thalamic-cortical (CSTC) sub-circuit of SN²⁵. For depression patients, the SN-caudate FC was not different from the healthy. The relatively reserved FC within SN have also been reported by a recent research do used rsfMRI and ICA to investigate intrinsic FC within and between resting-state networks (RSNs) in drug-free patients with depression, they did not find any abnormal intra-network connectivity related to SN⁶. Therefore, we suppose that the disorganized connectivity within the SN may be only presented in schizophrenia but not depression, since reduced the cortico-striatal connectivity was linked with psychotic symptoms, such as hallucinations and delusions³². Our findings are in line with previous rs-fMRI evidence, which indicates that SN is increasingly considered as having a fundamental role in the pathophysiology of schizophrenia^{26,33-36}.

In addition, this study also found that compared to normal negative FC between SN and precentral gyrus in the DEP and HC groups, the FC was significantly increased and became positive in the SCH group. The precentral gyrus is a key component of sensory and motor movement³⁷, and it is believed to play an important role in social cognition, including perception of facial expressions, emotions, and individual desires. A meta-analysis revealed that the engagement level of precentral gyrus in schizophrenia patients was associated with their performance in facial emotion perception³⁸. Moreover, both reduced gray matter volume³⁹ and local activity(regional homogeneity)⁴⁰ have been reported in schizophrenia. The hyperconnectivity between SN and precentral gyrus in schizophrenia means that the information

interaction was overactive, which may cause emotional and motional processing dysfunction in schizophrenia.

Taking these results together, the current study supports that the SN FC alterations are more pronounced in the SCH group than in the DEP group. Specifically, we found that the patients with schizophrenia showed opposite coupling between the right temporal gyrus, left caudate, and right precentral gyrus and the SN, while the patients with depression exhibited original negative connectivity mode between the temporal gyrus and the SN. In addition, the findings of current study are in agreement with the previous, believing that FC features of SN may play an important role in distinguishing schizophrenia from depression⁷. However, we failed to confirm any relationships between the symptom's severity and disease's duration in the patients as previous researches⁴¹⁻⁴³, which may be due to the relatively small sample size in each group and the high heterogeneity in both diseases.

Conclusions

In summary, our study provides evidence for different FC alterations of the SN in schizophrenia and depression. The findings suggest that the resting-state FC of SN may be a transdiagnostic difference between depression and schizophrenia patients and may play a critical role in the pathogenesis of these two disorders.

Limitation

Several limitations should be considered in this study. First, the sample size was relatively small in each group, which may have yielded unreliable results, a larger number of participant samples is needed to increase the reliability and sensitivity. Second, most patients were taking medication, so the systematically different medications for schizophrenia and depression may have different effects on functional connectivity. This source of variance may have acted as a major confound in transdiagnostic analyses. Third, the study was a cross-sectional design and all the patients were at their acute phase, so it remains unclear when the FC abnormalities of FC emerged and how they change as the disease progresses and longitudinal follow-up studies are needed.

Declarations

Acknowledgments

We thank the all the patients and healthy volunteers for their participation. The work is supported by the Fundamental Research Funds for the Central Universities (2042020kf0128), Health Commission of Hubei Province scientific research project(WJ2021M142) and the Medical Science Advancement Program of Wuhan University (TFLC2018001).

Author contributions

WHL and ZY designed the study. HH, CC, RB and WQR conducted subjects' recruitment and performed the study. HH undertook the imaging data analysis and wrote the first draft of the manuscript. WHL, LZC, CJG, WGH and ZY revised the manuscript. All authors contributed to and have approved the final manuscript.

References

- 1 Anticevic, A., Schleifer, C. & Youngsun, T. C. Emotional and cognitive dysregulation in schizophrenia and depression: understanding common and distinct behavioral and neural mechanisms. *Dialogues Clin Neurosci***17**, 421-434 (2015).
- 2 Cross-Disorder Group of the Psychiatric Genomics, C. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet***381**, 1371-1379, doi:10.1016/S0140-6736(12)62129-1 (2013).
- 3 Parkes, L., Satterthwaite, T. D. & Bassett, D. S. Towards precise resting-state fMRI biomarkers in psychiatry: synthesizing developments in transdiagnostic research, dimensional models of psychopathology, and normative neurodevelopment. *Curr Opin Neurobiol***65**, 120-128, doi:10.1016/j.conb.2020.10.016 (2020).
- 4 Pearlson, G. D. Applications of Resting State Functional MR Imaging to Neuropsychiatric Diseases. *Neuroimaging Clin N Am***27**, 709-723, doi:10.1016/j.nic.2017.06.005 (2017).
- 5 Friston, K. J. Functional and effective connectivity: a review. *Brain Connect***1**, 13-36, doi:10.1089/brain.2011.0008 (2011).
- 6 Luo, L. *et al.* Abnormal large-scale resting-state functional networks in drug-free major depressive disorder. *Brain Imaging Behav*, doi:10.1007/s11682-019-00236-y (2020).
- 7 Han, W. *et al.* Low-rank network signatures in the triple network separate schizophrenia and major depressive disorder. *Neuroimage Clin***22**, 101725, doi:10.1016/j.nicl.2019.101725 (2019).
- 8 Shao, J. *et al.* Common and distinct changes of default mode and salience network in schizophrenia and major depression. *Brain Imaging Behav***12**, 1708-1719, doi:10.1007/s11682-018-9838-8 (2018).
- 9 Damoiseaux, J. S. *et al.* Consistent resting-state networks across healthy subjects. *Proc Natl Acad Sci U S A***103**, 13848-13853, doi:10.1073/pnas.0601417103 (2006).
- 10 Schilbach, L. *et al.* Transdiagnostic commonalities and differences in resting state functional connectivity of the default mode network in schizophrenia and major depression. *Neuroimage Clin***10**, 326-335, doi:10.1016/j.nicl.2015.11.021 (2016).
- 11 Chen, X. *et al.* Transdiagnostic differences in the resting-state functional connectivity of the prefrontal cortex in depression and schizophrenia. *J Affect Disord***217**, 118-124, doi:10.1016/j.jad.2017.04.001

(2017).

- 12 Jiang, Y. *et al.* Aberrant Prefrontal-Thalamic-Cerebellar Circuit in Schizophrenia and Depression: Evidence From a Possible Causal Connectivity. *Int J Neural Syst***29**, 1850032, doi:10.1142/S0129065718500326 (2019).
- 13 Yang, Y. *et al.* Common and Specific Functional Activity Features in Schizophrenia, Major Depressive Disorder, and Bipolar Disorder. *Front Psychiatry***10**, 52, doi:10.3389/fpsy.2019.00052 (2019).
- 14 Jiang, Y. *et al.* Common and distinct dysfunctional patterns contribute to triple network model in schizophrenia and depression: A preliminary study. *Progress in Neuro-Psychopharmacology and Biological Psychiatry***79**, 302-310, doi:10.1016/j.pnpbp.2017.07.007 (2017).
- 15 Seeley, W. W. *et al.* Dissociable intrinsic connectivity networks for salience processing and executive control. *The Journal of neuroscience : the official journal of the Society for Neuroscience***27**, 2349-2356, doi:10.1523/JNEUROSCI.5587-06.2007 (2007).
- 16 Dosenbach, N. U. *et al.* A core system for the implementation of task sets. *Neuron***50**, 799-812, doi:10.1016/j.neuron.2006.04.031 (2006).
- 17 Sridharan, D., Levitin, D. J. & Menon, V. A critical role for the right fronto-insular cortex in switching between central-executive and default-mode networks. *Proceedings of the National Academy of Sciences of the United States of America***105**, 12569-12574, doi:10.1073/pnas.0800005105 (2008).
- 18 Menon, V. & Uddin, L. Q. Saliency, switching, attention and control: a network model of insula function. *Brain structure & function***214**, 655-667, doi:10.1007/s00429-010-0262-0 (2010).
- 19 Dosenbach, N. U. *et al.* Distinct brain networks for adaptive and stable task control in humans. *Proceedings of the National Academy of Sciences of the United States of America***104**, 11073-11078, doi:10.1073/pnas.0704320104 (2007).
- 20 Menon, V. Salience Network. 597-611, doi:10.1016/b978-0-12-397025-1.00052-x (2015).
- 21 Uddin, L. Q. *et al.* Dissociable connectivity within human angular gyrus and intraparietal sulcus: evidence from functional and structural connectivity. *Cereb Cortex***20**, 2636-2646, doi:10.1093/cercor/bhq011 (2010).
- 22 Whitfield-Gabrieli, S. & Nieto-Castanon, A. Conn: a functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connect***2**, 125-141, doi:10.1089/brain.2012.0073 (2012).
- 23 Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage***37**, 90-101, doi:10.1016/j.neuroimage.2007.04.042 (2007).

- 24 Tu, P. C., Hsieh, J. C., Li, C. T., Bai, Y. M. & Su, T. P. Cortico-striatal disconnection within the cingulo-opercular network in schizophrenia revealed by intrinsic functional connectivity analysis: a resting fMRI study. *NeuroImage***59**, 238-247, doi:10.1016/j.neuroimage.2011.07.086 (2012).
- 25 Huang, H. *et al.* Aberrant resting-state functional connectivity of salience network in first-episode schizophrenia. *Brain Imaging Behav***14**, 1350-1360, doi:10.1007/s11682-019-00040-8 (2020).
- 26 Wang, C. *et al.* Disrupted salience network functional connectivity and white-matter microstructure in persons at risk for psychosis: findings from the LYRIKS study. *Psychological medicine***46**, 2771-2783, doi:10.1017/S0033291716001410 (2016).
- 27 Dong, D., Wang, Y., Chang, X., Luo, C. & Yao, D. Dysfunction of Large-Scale Brain Networks in Schizophrenia: A Meta-analysis of Resting-State Functional Connectivity. *Schizophrenia bulletin*, doi:10.1093/schbul/sbx034 (2017).
- 28 Mulders, P. C., van Eijndhoven, P. F., Schene, A. H., Beckmann, C. F. & Tendolkar, I. Resting-state functional connectivity in major depressive disorder: A review. *Neurosci Biobehav Rev***56**, 330-344, doi:10.1016/j.neubiorev.2015.07.014 (2015).
- 29 Raichle, M. E. The brain's default mode network. *Annu Rev Neurosci***38**, 433-447, doi:10.1146/annurev-neuro-071013-014030 (2015).
- 30 Yin, Y. *et al.* Structural and Functional Connectivity of Default Mode Network underlying the Cognitive Impairment in Late-onset Depression. *Sci Rep***6**, 37617, doi:10.1038/srep37617 (2016).
- 31 Grahm, J. A., Parkinson, J. A. & Owen, A. M. The cognitive functions of the caudate nucleus. *Prog Neurobiol***86**, 141-155, doi:10.1016/j.pneurobio.2008.09.004 (2008).
- 32 Orliac, F. *et al.* Links among resting-state default-mode network, salience network, and symptomatology in schizophrenia. *Schizophrenia research***148**, 74-80, doi:10.1016/j.schres.2013.05.007 (2013).
- 33 Peters, S. K., Dunlop, K. & Downar, J. Cortico-Striatal-Thalamic Loop Circuits of the Salience Network: A Central Pathway in Psychiatric Disease and Treatment. *Frontiers in systems neuroscience***10**, 104, doi:10.3389/fnsys.2016.00104 (2016).
- 34 Palaniyappan, L. & Liddle, P. F. Does the salience network play a cardinal role in psychosis? An emerging hypothesis of insular dysfunction. *Journal of psychiatry & neuroscience : JPN***37**, 17-27, doi:10.1503/jpn.100176 (2012).
- 35 Palaniyappan, L., White, T. P. & Liddle, P. F. The concept of salience network dysfunction in schizophrenia: from neuroimaging observations to therapeutic opportunities. *Current topics in medicinal chemistry***12**, 2324-2338 (2012).

- 36 Kapur, S. Psychosis as a state of aberrant salience: a framework linking biology, phenomenology, and pharmacology in schizophrenia. *The American journal of psychiatry***160**, 13-23, doi:10.1176/appi.ajp.160.1.13 (2003).
- 37 Yu, Y. *et al.* Altered Amplitude of Low-Frequency Fluctuations in Inactive Patients with Nonneuropsychiatric Systemic Lupus Erythematosus. *Neural Plast***2019**, 9408612, doi:10.1155/2019/9408612 (2019).
- 38 Jani, M. & Kasperek, T. Emotion recognition and theory of mind in schizophrenia: A meta-analysis of neuroimaging studies. *World J Biol Psychiatry***19**, S86-S96, doi:10.1080/15622975.2017.1324176 (2018).
- 39 Zhou, S. Y. *et al.* Volumetric analysis of sulci/gyri-defined in vivo frontal lobe regions in schizophrenia: Precentral gyrus, cingulate gyrus, and prefrontal region. *Psychiatry Res***139**, 127-139, doi:10.1016/j.psychres.2005.05.005 (2005).
- 40 Zhao, X. *et al.* Abnormalities of regional homogeneity and its correlation with clinical symptoms in Naive patients with first-episode schizophrenia. *Brain Imaging Behav***13**, 503-513, doi:10.1007/s11682-018-9882-4 (2019).
- 41 Hare, S. M. *et al.* Salience-Default Mode Functional Network Connectivity Linked to Positive and Negative Symptoms of Schizophrenia. *Schizophr Bull***45**, 892-901, doi:10.1093/schbul/sby112 (2019).
- 42 Brakowski, J. *et al.* Resting state brain network function in major depression - Depression symptomatology, antidepressant treatment effects, future research. *J Psychiatr Res***92**, 147-159, doi:10.1016/j.jpsychires.2017.04.007 (2017).
- 43 Manoliu, A. *et al.* Insular dysfunction within the salience network is associated with severity of symptoms and aberrant inter-network connectivity in major depressive disorder. *Front Hum Neurosci***7**, 930, doi:10.3389/fnhum.2013.00930 (2013).

Tables

Due to technical limitations, table 1-3 is only available as a download in the Supplemental Files section.

Figures

Figure 1

The salience network component identified from group independent component analysis(gICA) of the resting-state fMRI data of all subjects (n=87)

Figure 2

T maps of functional connectivity with the SN and whole brain in the three groups. One-sample t-test was performed for each group (voxel-wise $p < 0.001$ uncorrected and a cluster-wise threshold of $p < 0.05$, FWE-corrected).

Figure 3

Part(A) Differences of SN functional connectivity in the HC, SCH, and DEP by the ANCOVA(voxel-wise $p < 0.001$ uncorrected and a cluster-wise threshold of $p < 0.05$, FWE-corrected), and the colorbar represents F values; part(B)Post-hoc comparisons found that the schizophrenia group showed increased SN-right ITG and MTG FC and SN- right precentral gyrus, as well as increased SN-left caudate in SCH group compared to the other two groups($p \leq 0.01$), the Depression group showed decreased FC SN-right ITG and MTG FC compared to the other two groups($p \leq 0.05$).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Tables.pdf](#)