MicroRNA-20a mediates the cytotoxicity of natural killer cells in endometriosis via ERG/HLX/STAT4/perforin axis

Li-Juan Chen

Institute of Hematology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology

Bin Hu

The Central Hospital of Wuhan, Tongji Medical College, Huazhong University of Science and Technology

Zhi-Qiang Han

Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology

Jian Ni

Brigham and Women's Hospital, Harvard Medical School

Yong-Ming Zhou

The Affiliated Tianyou Hospital, Wuhan University of Science and Technology

Xue-Xing Chen

Institute of Hematology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology

Hao Zhou (✉ sue_drHAO@126.com)

Institute of Hematology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology

Research

Keywords: Endometriosis, microRNA-20a, ETS transcription factor, H2.0 like homeobox, signal transducer and activator of transcription 4, Perforin, NK cells, Cytotoxicity

Posted Date: December 23rd, 2019

DOI: <https://doi.org/10.21203/rs.2.17459/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

The authors have withdrawn this preprint from Research Square