Single-Cell RNA-seq Reveals LGALS1 and LAG3 as Novel Drivers of Ibrutinib Resistance in Chronic Lymphocytic Leukemia

Hui Jin

the first affiliated hospital of Nanjing medical university

Bin Huang

Nanjing Medical University

Zijuan Wu

First Affiliated Hospital of Nanjing Medical University

Huayuan Zhu

the first affiliated hospital of Nanjing medical university

Hanning Tang

the first affiliated hospital of Nanjing medical university

Jiazhu Wu

the first affiliated hospital of Nanjing medical university

Danling Gu

the first affiliated hospital of Nanjing medical university

Yun Cai

Nanjing medical university

Luqiao Wang

the first affiliated hospital of Nanjing medical university

Xueying Lu

The First Affiliated Hospital of Nanjing Medical University

Wei Xu

The First Affiliated Hospital of Nanjing Medical University

Kening Li

Nanjing Medical University

Qianghu Wang

Nanjing Medical University

Jian-Yong Li (✉ lijianyonglm@126.com)

the first affiliated hospital of Nanjing medical university <https://orcid.org/0000-0002-5878-945X>

Keywords: Chronic lymphocytic leukemia, Ibrutinib resistance, Single-cell RNA sequencing, Galectin-1 inhibitor

Posted Date: August 31st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-810283/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Background

Chronic lymphocytic leukemia (CLL) is a highly heterogeneous malignant lymphoproliferative B-cell disorder that can be treated using ibrutinib, a Bruton's tyrosine kinase inhibitor. However, the ibrutinib resistance of CLL patients has caused widespread concerns, necessitating the development of novel treatment strategies.

Methods

Here, we identified lectin galactoside-binding soluble 1 (LGALS1) and lymphocyte-activating gene 3 (LAG3) as potential markers for ibrutinib-resistant CLL using single-cell RNA sequencing (scRNA-seq), and the results were validated in an ibrutinib-resistant CLL cell line (MEC1-IR) and primary cells from CLL patients. Marker-gene expression was detected while functional analyses were conducted with or without OTX008, a selective Galectin-1 inhibitor. ScRNA-seq revealed that the biological features, gene expression profiles, and clonal signatures of peripheral blood mononuclear cells (PBMCs) from patients with ibrutinib-resistant CLL were distinct from those displayed by PBMCs from ibrutinib-sensitive patients.

Results

A close correlation between LGALS1 and LAG3 expression was observed and these factors were found to be highly expressed in ibrutinib-resistant CLL, with diagnostic and prognostic stratification, indicating that they may serve as drivers of ibrutinib-resistant CLL. Concordantly, LGALS1 and LAG3 expression was higher in ibrutinib-resistant CLL cells and primary cells, and OTX008 suppressed proliferation and induced apoptosis in both cells.

Conclusion

LGALS1 and LAG3 gene panel is promising indicator of ibrutinib-sensitivity and prognosis marker of CLL. LGALS1 inhibitor OTX008 is effective in CLL patients, both those naïve to and those resistant to ibrutinib.

Background

Chronic lymphocytic leukemia (CLL) involves leukemic cell infiltration and accumulation in lymphoid organs and is the most common leukemia in Western populations. Although its incidence is lower in China, the frequency of new cases is increasing and the clinical presentation and rate of CLL progression vary highly [1, 2]. The approval of Bruton's tyrosine kinase (BTK) inhibitors such as ibrutinib and acalabrutinib, and the B cell lymphoma-2 (Bcl-2) inhibitor venetoclax have revolutionized the treatment of CLL [3–7]. While these novel agents alone or in combination induce long lasting and deep remissions in most patients with CLL, their use may be associated with the development of clinical resistance. BTK is a key B cell receptor (BCR) signaling kinase that is amplified in CLL and activates proliferative and anti-apoptotic signals [8]. Ibrutinib is an orally bioavailable BTK inhibitor that eliminates pro-survival pathway

activation and microenvironmental survival signals [9]. As the first FDA-approved BTK inhibitor for CLL, ibrutinib has shown significant clinical benefits in patients [10]. However, current limitations include off-target toxicities and the development of resistance remain an unmet clinical need [11–13]. BTK and PLCG2 are the most frequently described mutations that confer acquired resistance to ibrutinib [14–16], while clonal evolution driven by the selective pressure of drug therapy is central to recurrence and resistance in leukemia. Consequently, a comprehensive and accurate genetic profiling of CLL clonal evolution is vital for precise, individualized treatment.

Next-generation sequencing (NGS) [17, 18] and whole exome sequencing (WES) [16, 19] have helped to clarify the complete clonal structure of CLL patients and analyze chromosomal translocations or key gene mutations in subclones. However, most studies on the molecular mechanisms of ibrutinib-resistance in CLL are based on bulk-sample data. Single-cell RNA sequencing (scRNA-seq) can systematically characterize heterogeneity at a higher resolution to assess gene expression in individual cells [20, 21]. Moreover, this technology can be used to identify pathogenic or drug-resistant cell subtypes and relevant transcriptional modules [22, 23], and thus could be used to discover reliable biomarkers for clinical efficacy and drug-resistance.

Here, we examined the heterogeneity of peripheral blood mononuclear cells (PBMCs) from patients with ibrutinib-sensitive (IBS) and -resistant (IBR) CLL by analyzing bulk and single-cell level gene expression profiles, clinical features, biological properties, and phenotypes. Seven distinct ibrutinib-resistant subpopulations were identified and two candidate genes were screened that contribute toward ibrutinib resistance and poor survival in CLL, lectin galactoside-binding soluble 1 (LGALS1) and lymphocyte-activating gene 3 (LAG3). LGALS1 is located on chromosome 22 and encodes galectin-1 (Gal-1), a protein involved in regulating physiological and pathological processes like cell transformation, proliferation, and adhesion, blood vessel formation, and immunosuppression [24]. LAG3 is an immune checkpoint molecule that was recently suggested as a novel therapeutic target in CLL [25–27]. Together, we hope that these findings may offer a new therapeutic strategy for the treatment of ibrutinib-resistant CLL.

Materials And Methods

ScRNA-seq data processing

ScRNA-seq libraries were sequenced from 150 bp 3' paired cDNAs using HiSeq Xten instruments. Raw data (BCL files) were converted to fastq files using bcl2fastq software (Illumina). Reads were aligned to a human genome reference (GRCh38) and a digital gene expression matrix built using the STAR algorithm in CellRanger (10x Genomics; v3.0.2). Quality control and further analyses were carried out using Seurat [28] (v3.1.5). To eliminate the effect of abnormal cells and technical noise on downstream analysis, we removed cells with abnormal gene counts or a high percentage of unique molecular identifiers (UMIs) mapped to mitochondria for each sample (Table S1). Detailed clinical characteristics were showed in Fig. S1 and Table S2.

ScRNA-seq sample integration and cell type recognition

After log-normalization and variable feature identification based on variance stabilizing transformation, we integrated filtered cells from three IBR, four IBS CLL patients, and two healthy donors using “FindIntegration Anchors” and “IntegrateData” in Seurat with 30 dimensionalities (dims) [28], followed by scaling using “ScaleData”. The statistical significance of principal component analysis scores was determined using “JackStraw” and dimension reduction was performed using the first 20 principal components. Data was visualized using T-distributed stochastic neighbor embedding (t-SNE) with “RunTSNE”, while cell clusters were identified using “FindNeighbors” and “FindClusters” (resolution = 2). Differentially expressed genes (DEGs) were identified using the Wilcoxon Rank Sum test in “FindAllMarkers”, with significance indicated by an average natural logarithm (fold change) ³ 0.25, Bonferroni-adjusted *p*-value < 0.05, and minimum percentage ³ 0.25. Candidate markers were considered with CD14, CD1C, CD79A, CD3E, and KLRD1, and were used to annotate cell clusters. Single cell copy number variation (CNV) was estimated using inferCNV [29] (v0.8.2), with non-B cells as a reference.

B-cell re-clustering and IBR cluster definition

First, we ran the “FindNeighbors” function (30 dims and “k.param” = 15), k-nearest neighbors (SNN) clustering (resolution = 1 into 21 clusters), and t-SNE to visualize integrated B cells. Based on the proportion of B cells from IBR samples, B-cell clusters were categorized into three main subgroups, i.e., IBR (> 50 % of B cells from IBR samples), IBS (> 70 % of B cells from IBS or NC samples). Other clusters are defined as shared clusters. Clusters with < 200 cells were filtered.

Detailed materials and methods can be found in the supplemental data.

Results

Single-cell landscape of PBMCs in IBR and IBS patients with CLL

To study PBMC heterogeneity in CLL patients treated with ibrutinib and the mechanism underlying the development of ibrutinib resistance, we collected PBMCs from three IBR, four IBS CLL patients and two healthy donors, and performed scRNA-seq using a droplet-based 10× Genomics platform. After quality control, we obtained 42,288 cells with a mean number of cells per sample of 4,760. The median number of UMIs and genes per cell was 3,570 and 1,108 respectively. t-SNE visualization of filtered cells following integration and unsupervised clustering, revealed that cells from IBR and IBS samples were distributed in different clusters and suggested that IBR cells display a unique transcriptional pattern (Fig. 1A). To differentiate between cell types, we sorted cells into 36 clusters (Fig. 1B) and annotated them as five major categories, i.e., CD79A⁺ B-cells, CD3E⁺ T-cells, KLRD1⁺ NK-cells, CD14⁺ monocytes, and CD1C⁺ dendritic cell (DC) cells (Fig. 1C, D; Fig. S2).

To elucidate the role of PBMC components in CLL pathology, we calculated the proportion of cell types in each sample (Table S3). B cells exhibited significant proliferation in CLL patients (compared to healthy donors), accounting for > 60 % and < 15 % of cells in each group, respectively, suggesting massive B-cell expansion as a major feature of CLL (Fig. 1E). As chromosomal alteration is a common phenomenon in

CLL [30], we predicted CNV in IBR and IBS patients at the cellular level using InferCNV (Fig. 1F). Notably, one IBR patient carried a deletion of the chromosome 8 short arm, consistent with the findings of Burger [31]. Two IBR patients carried a deletion of the chromosome 9 long arm and one IBR patient carried a deletion of the chromosome 4 short arm, implying that del(9q) and del(4p) may contribute toward IBR. Furthermore, we found that chromosome 6p was often amplified, consistent with the results of Brown [32].

IBR B cells exhibit a unique transcriptional pattern

Next, we investigated B-cell heterogeneity in IBR and IBS patients by re-clustering the 30,417 identified B cells into 21 clusters (Fig. 2A). The cells from IBR patients were distributed in clusters 4, 5, 6, 7, 9, 10, and 19, suggesting transcriptomic differences between B cells in IBR and IBS patients. CytoTRACE analysis revealed that B cells in IBR CLL patients had higher stem index scores than those in IBS patients, implying that B cell stemness may contribute toward IBR ($p < 2.2e-16$, Fig. 2B and 2C).

According to the proportion of B cells from IBR samples, we classified each B-cell cluster into three main subgroups, i.e., IBR, IBS, and shared cluster (Fig. 2D, Table S4). To elucidate the functional differences between IBR and IBS/shared clusters, we performed metabolic enrichment analysis. IBR clusters were significantly enriched in glycolysis and gluconeogenesis, which supply energy and support cancer cell growth (Fig. 2E, Table S5). Previously, Qian et al. [33] reported that glycolysis and TP53-induced glycolysis and apoptosis regulator (TIGAR) inhibit leukemia cell apoptosis and are associated with poor prognosis, suggesting the importance of glycolysis in leukemia cells. We also found that many glycometabolism pathways, such as the pentose phosphate pathway, pentose and glucuronate interconversion, and fructose and mannose metabolism, were enriched in IBR clusters. Adokola showed that extracellular glucose reduction could induce the development of resistance against DNA-damaging therapeutics in normal cells and enhance cancer cell sensitivity to mainstream CLL drugs [34]. Further, we found that glutathione metabolism was enriched in IBR clusters, consistent with Zhang who reported that elevated glutathione levels can increase leukemia cell survival and protect them against drug-induced cytotoxicity [35]. These findings suggest that these energy-related metabolic pathways are important for drug resistance in cancer cells. Together, IBR B cells have higher stemness scores and are enriched in some energy metabolism pathways.

Specific intercellular interactions in the microenvironment of IBR CLL Intercellular communication mediated by ligand-receptor complexes is essential for coordinating biological processes, such as differentiation, development, and inflammation. Therefore, we studied the interaction between B cells and other immune microenvironment cells in IBR and IBS patients by performing cell-cell communication analysis using CellphoneDB. IBR B cells displayed more interactions with monocytes, NK, T, and dendritic cells than IBS B cells, suggesting that IBR B cells may actively build connections with other immune cells to reshape the protective niche (Fig. 2F, Table S6). Previous studies have shown that co-culturing CLL cells and stromal NK cells can increase oxidative phosphorylation in CLL cells, with ATP synthesis promoting their proliferation [36]. Kurtova et al. [37] also found that bone marrow stromal cells (MSC)

provide survival and drug resistance signals for CLL cells against spontaneous and drug-induced apoptosis. Therefore, the interaction between CLL cells and other microenvironment cells is important for the development of drug resistance.

Notably, the LGALS9-CD47 interaction only exists between T cells and IBR, not IBS, B cells (Fig. 2G, Table S7); however, this interaction has not previously been reported in CLL. We also found that other B cell-T cell interactions such as MIF-TNFRSF14 and LILRA4-BST2 were specifically enriched in IBR clusters and may be important for IBS-IBR transition. These findings indicated that IBR B cells might reprogram intercellular interaction patterns in CLL patients.

The expression of LGALS1 and LAG3 is upregulated in IBR CLL

Although genome-level features have been extensively studied in IBR patients with CLL [15, 38, 39], their transcriptional characteristics are rarely reported. To improve our understanding of the mechanism underlying IBR and accurately identify the key regulatory molecules in patients with CLL for targeted treatments, we collected PBMCs from 6 IBR and 40 IBS CLL patients and performed bulk RNA sequencing. Ninety genes were significantly upregulated in the IBR group compared to that in the IBS group, among which LGALS1, LAG3, and PTMS were the top three upregulated genes based on the delta mean and *p* values (Fig. 3A, Table S8). To verify these genes, we identified highly expressed genes in each IBR cluster at the scRNA level (719 genes; Table S9), including LGALS1, LAG3, and PTMS. Moreover, as hematopoietic differentiation is a continuous process, we investigated the developmental trajectory of B cells in IBR and IBS patients using pseudo-time analysis. B cells from IBR samples were distributed at the end of one state track, indicating that IBR cells may evolve from IBS cells (Fig. 3B). We found that the expression of some genes gradually increased along the trajectory from IBS to IBR, suggesting that some signatures may be key factors in the transition of B cells from IBS to IBR (Fig. 3C). Notably, LGALS1 was highly expressed in five single-cell IBR clusters (Fig. 3D), while LGALS1 and LAG3 expression increased along the trajectory from IBS to IBR (Fig. 3E). Interestingly, PTMS and LAG3 were found to be located adjacent to each other on chromosome 12p13.31 and bulk-RNA data indicated co-expression of these genes, implying an association phenomenon.

LGALS1 and LAG3 typically act as immune checkpoints that repress innate and adaptive immune programs [40, 41] by inducing T-cell failure, limiting T-cell survival, promoting regulatory T-cell expansion, inactivating NK cells, and polarizing myeloid cells. To elucidate the function of LGALS1 and LAG3 in CLL patients, we identified genes co-expressed with LGALS1—based on bulk-RNA seq data—including S100A4, S100A6, and COX5B, which are associated with cancer cell invasion[42] and proliferation [43] (Fig. 3F, Table S10). We also identified genes co-expressed with LAG3, such as HSPG2, which is associated with poor prognosis in acute myeloid leukemia[44] (Fig. S3, Table S11).

GO functional enrichment analysis revealed that genes co-expressed with LGALS1 were significantly enriched in mitochondrial ATP synthesis-coupled electron transfer and oxidative phosphorylation (Fig. 3G, Table S12). Mitochondria regulate cell death and reactive oxygen species production [45], and studies have shown that CLL-cell proliferation is accompanied by mitochondrial energy and reactive oxygen

species production and enhanced adaptability to oxidative stress [46, 47]. Thus, LGALS1 may play an important role in mitochondrial energy generation and CLL-cell proliferation. Conversely, genes co-expressed with LAG3 were enriched for pathways related to glucose metabolism (Fig. S4, Table S13), which provides energy for cancer cell growth [45], indicating that LAG3 may be associated with cancer cell proliferation. Together, these findings suggest that energy-related metabolic pathways are related to IBR.

LGALS1 and LAG3 expression correlates with poor outcomes in CLL

Next, we detected and analyzed the correlation between LGALS1/LAG3 mRNA expression and overall survival (OS) in CLL patients ($n = 59$) from our hospital, finding that patients with higher LGALS1 and LAG3 expression showed poor OS (Fig. 4A). After ranking both LGALS1 and LAG3 expression, patients were divided into low and high expression groups based on median expression. Kaplan-Meier survival analyses revealed that patients with high expression of LGALS1 or LAG3 displayed a significantly shorter OS than those with low expression. Moreover, log-rank tests demonstrated that patients with concurrent high LGALS1 and LAG3 expression exhibited a worse OS than those with either high LGALS1 or LAG3 expression. To further validate the survival analysis results, we also analyzed the microarray and survival data in CLL patients ($n = 107$) from the GEO database (GSE22762) [48] (Fig. 4B). These results indicating that LGALS1 and LAG3 gene panel is associated with IBR and poor prognosis in CLL.

To evaluate and confirm the clinical relevance of LAG3 and LGALS1 in CLL, we analyzed the expression pattern of these genes and associated clinical features in CLL patients ($n = 59$; Fig. 4C). Patients with higher level of LAG3 and LGALS1 were found to have higher mutation rate of SF3B1 and NOTCH1. Immunoblots were then performed to compare protein levels in IBS and IBR patients, revealing that Gal-1 and LAG3 protein expression varied in CLL, with higher levels being observed in IBR cells (Fig. 4D, Fig. S5A). As Gal-1 is a secretory protein, we determined its plasma concentration as well as that of LAG3 in 16 treatment-naïve, relapsed, and refractory CLL patients using ELISA. Gal-1 and LAG3 levels were significantly higher in relapsed and refractory CLL patients than those in treatment-naïve patients (Fig. S5B). Together, these results indicate that elevated expression of LGALS1 and LAG3 is associated with poor prognosis in CLL.

Acquired resistance following chronic exposure to ibrutinib leads to upregulation of LGALS1 and LAG3

As LGALS1 and LAG3 may indicate poor prognosis in CLL patients, we sought to confirm these results in a CLL cell line with acquired ibrutinib-resistance (MEC1-IR). MEC1 cells initially exposed to 1 μM ibrutinib returned to a normal growth rate after the recovery period and were then treated continuously with increasing doses of ibrutinib (3, 5, 7, 10, 15, and 20 μM) over a course of approximately 6 months (Fig. 5A). CCK8 assays for cell viability indicated that MEC1 cells with an IC_{50} of 4.56 μM exhibited greater sensitivity to increasing ibrutinib concentrations for 48 h, whereas MEC1-IR cells were resistant to ibrutinib at an IC_{50} 7-fold higher than that of the parental cells (Fig. 5B). Annexin-V/PI staining revealed a pronounced increase in cell death in MEC1 cells treated with ibrutinib for 48 h, but not in MEC1-IR cells

(Fig. 5C, D). Next, we examined LGALS1 and LAG3 expression in parental and resistant cells, with qRT-PCR revealing that LGALS1 and LAG3 expression was markedly higher in MEC1-IR cells than that in the parental cells (Fig. 5E). Expectedly, immunoblotting and ELISA showed that Gal-1 and LAG3 proteins were upregulated in MEC1-IR cells (Fig. 5F-H). These results demonstrate that ibrutinib-resistance acquired in response to chronic ibrutinib exposure leads to LGALS1 and LAG3 upregulation.

LGALS1 inhibitor OTX008 effectively inhibits the growth of ibrutinib-resistant CLL cells

Recent studies have demonstrated the efficacy of OTX008, a small molecule that serves as a Gal-1 inhibitor, in preclinical models of multiple tumors [49]; therefore, we studied the activity of OTX008 against IBS and IBR CLL cells. First, we examined the effect of OTX008 on Gal-1 and LAG3 expression, and found that at low concentrations, OTX008 decreased the expression of Gal-1 and LAG3 at the protein level in a dose-dependent manner in both CLL cell lines and primary cells (CLL-1), but did not exhibit a significant effect on the corresponding mRNA levels (Fig. 6A, B). This observation was validated by immunofluorescence (Fig. 6C) before examining the role of Gal-1 inhibition in cell proliferation and apoptosis *in vitro*. While MEC1-IR displayed higher proliferation than MEC1 cells, the proliferative capacity of both groups was almost completely inhibited by OTX008 (Fig. 6D). To assess the toxicity of OTX008 in CLL cells, we measured viability by flow cytometry using Annexin-V and PI staining. CLL-cell death increased markedly following exposure to 30 μ M OTX008 but did not significantly change with 10 μ M OTX008 (Fig. 6E); however, OTX008 was effective against both IBS and IBR cells the inhibitory effect of which is more obvious in IBR. To determine whether OTX008 was effective against primary CLL cells, we treated PBMCs isolated from 15 CLL patients with OTX008. Trypan blue staining revealed cell vitality variation between CLL patients after *in vitro* culture for 48 h (53–88 %), whilst a significant increase in CLL-cell death was observed in response to 30 μ M OTX008 (Fig. 6F). For subsequent comparative studies, IBS and IBR CLL samples were arbitrarily defined as those in which incubation with 10 μ M ibrutinib resulted in more or less than 20 % cell death. Notably, OTX008 was more effective against CLL cells collected from IBR patients. Overall, our findings demonstrate the potential clinical utility of OTX008 in CLL, particularly for IBR patients.

Discussion

CLL is a highly heterogeneous disease that can be treated using ibrutinib, a BTK inhibitor that has profound activity in CLL; however, increasing clinical use has resulted in the development of ibrutinib-resistance during monotherapy, necessitating the development of novel treatment strategies. Second-generation BTKi include zanubrutinib (BGB-3111) and Orelabrutinib (ICP-022) which offer greater BTK selectivity [50, 51]. While these agents may limit off-target toxicity, they do not overcome common mechanisms of ibrutinib resistance. This study elucidated the transcriptional landscape and cellular composition of PBMCs from patients with IBS and IBR CLL using scRNA-seq. We identified IBR subpopulations and revealed their transcriptional regulation, metabolism, and intercellular communication, finding that LGALS1 and LAG3 were upregulated in IBR primary cells and cell lines and

also distinguished IBR subclones. Furthermore, the LGALS1 inhibitor, OTX008, induced apoptosis and displayed particular efficacy against IBR CLL cells, indicating potential clinical utility in CLL.

Considerable evidence has shown that Gal-1 expression is often increased in tumor tissues [52, 53] and secreted into the tumor extracellular space and tumor-associated stromal cells, and can correlate with tumor aggressiveness and treatment resistance [54, 55]. Croci [56] indicated that Gal-1 contributes toward modulation of BCR signaling and is associated with poor outcomes in CLL, whereas Shokri et al. [57] demonstrated that there is no significant difference between Gal-1 levels in CLL patients and healthy individuals. Thus, the biological function and molecular mechanism of Gal-1 in CLL require further investigation. Here, we revealed that Gal-1 plays a critical role in patients with IBR CLL at the scRNA level, helps to distinguish IBR and IBS clusters, and serves as a potential prognostic indicator and therapeutic target. Moreover, OTX008 directly targets Gal-1 to inhibit proliferation and promote apoptosis in CLL cells, especially in IBR-CLL cells. This may provide new treatment strategies for improving the clinical efficacy of ibrutinib-resistant CLL patients.

Sc-RNA seq also identified pronounced LAG3 expression in IBR clusters. LAG3 is a co-suppressor that is usually expressed by activated T, NK, B, and plasma cell-like dendritic cells, as well as depleted T cells and tumor infiltrating lymphocytes. Many studies have illustrated the role of LAG3 in tumor invasiveness and immunotherapeutic regulation [58], and multiple clinical trials are currently being conducted to assess the efficacy of anti-LAG3 monoclonal antibodies in solid tumors and hematological diseases [26, 59]. Dual PD1 and LAG3 immune checkpoint targeting has also been shown to successfully control CLL development in preclinical mouse models [60]; however, LAG3 expression and its roles in CLL cells and the tumor microenvironment remain unclear. Jana et al. detected increased LAG3 expression in CLL cells, and found it to be correlated with unmutated-IGHV and patient outcomes [27, 61], consistent with our observations in a small cohort. As OS was the shortest in CLL patients with high Gal-1 and LAG3 expression, we propose that targeting Gal-1 and LAG3 with a combination of OTX008 and anti-LAG3 monoclonal antibodies may achieve better efficacy in CLL. Further *in vitro* and *in vivo* studies are required to determine the synergistic effect and toxicities, as well as individual and synergistic efficacies against resistance to other BTKis.

We conducted in-depth analyses of the microenvironment characteristics of PBMC from ibrutinib-resistant CLL patients. Our study presents, for the first time, evidence showing that LGALS1 and LAG3 gene panel as an indicator of ibrutinib sensitivity as well as prognosis marker and potential therapeutic targets of CLL patients at single cell level. We demonstrated that LGALS1 inhibitor OTX008 treatment could restore the sensitivity of CLL cells to ibrutinib. However, the major limitation of this study is the small sample size of patients, and the lack of mechanism study. We will expand the clinical sample collection in our future work, including patients with resistance of ibrutinib and other BTK inhibitors. Also, the molecular mechanism by which LGALS1 regulates ibrutinib-resistance of CLL cells requires further elaboration. Together, our findings expand the current knowledge regarding the development of ibrutinib-resistance of CLL patients, identify improved biomarkers for patient selection, and offer a promising combinatorial therapeutic strategy for IBR CLL patients.

Collectively, our study conducts in-depth analyses of the heterogeneity and microenvironment characteristics of PBMC from ibrutinib-sensitive and -resistant CLL. We present, for the first time, evidence showing that LGALS1 and LAG3 gene panel as the prognosis markers and potential therapeutic targets for CLL patients at single cell level. We demonstrate that LGALS1 inhibitor OTX008 treatment could help to overcome ibrutinib-resistance of CLL cells.

Conclusion

Together, our findings demonstrate that ibrutinib-resistant CLL cells exhibit a unique transcriptional pattern and indicate a hope for future combination therapy. We propose that ibrutinib-resistance is mediated by LGALS1 and LAG3, two promising prognostic biomarkers and potential therapeutic targets in CLL, and that OTX008 may help to overcome ibrutinib-resistance of CLL patients.

Abbreviations

BCR B cell receptor

BTK Bruton tyrosine kinase

CLL Chronic lymphocytic leukemia

DEGs Differentially expressed genes

dims dimensionalities

FCR fludarabine, cyclophosphamide, and rituximab

Gal-1 galectin-1

IGHV immunoglobulin heavy-chain variable region

IBR ibrutinib-resistant

IBS ibrutinib-sensitive

LGALS1 lectin galactoside-binding soluble 1

LAG3 lymphocyte-activating gene 3

MSC bone marrow stromal cell

NGS Next-generation sequencing

OS overall survival

PBMC peripheral blood mononuclear cell

PI3K phosphatidylinositol3-kinase

scRNA-seq single-cell RNA sequencing

TIGAR TP53-induced glycolysis and apoptosis regulator

t-SNE T-distributed stochastic neighbor embedding

UMIs unique molecular identifiers

WES whole exome sequencing

Declarations

Ethics approval and consent to participate

Human sample study was approved by the Medical Ethics Committee of First Affiliated Hospital of Nanjing Medical University. Written informed consent was obtained from all participants.

Consent for publication

Written informed consent was obtained from all patients.

Availability of data and materials

The datasets generated and/or analysed during the current study are available in the GSA under accession number HRA000773.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by grants from National Natural Science Foundation of China (Grant #81720108002, 81970146, 81770166, 81972358 and 91959113), Key Research & Development Program of Jiangsu Province (Grant #BE2017733), Basic Research Program of Jiangsu Province (Grant #BK20180036), Jiangsu Province's Medical Elite Programme (Grant #ZDRCA2016022), Project of National Key Clinical Specialty, Jiangsu Provincial Special Program of Medical Science (Grant #BE2017751), Six Top Talent Project of Jiangsu Province (Grant #WSN-001).

Authors' contributions

H.J., K.N.L., Z.J.W. and H.Y.Z. designed and performed most of the experiments; B.H., K.N.L. and Y.C. performed bioinformatics analysis of scRNA-seq and bulk RNA-seq data; H.Y.Z., H.N.T. and D.L.G. collected the clinical samples; Z.J.W., J.Z.W., L.Q.W. and X.Y.L collected and analyzed the clinical data; H.J., B.H. and Z.J.W. drafted the manuscript; H.J., K.N.L., Q.H.W. and J.Y.L. critically revised the manuscript; J.Y.L., K.N.L., Q.H.W. and W.X. supervised the project. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Editage (<http://www.editage.cn>) for English language editing.

References

1. Yang S, Gale RP, Shi H, Liu Y, Lai Y, Lu J, et al. Is there an epidemic of chronic lymphocytic leukaemia (CLL) in China? *Leukemia research*. 2018;73:16-20.
2. Yang SM, Li JY, Gale RP, Huang XJ. The mystery of chronic lymphocytic leukemia (CLL): Why is it absent in Asians and what does this tell us about etiology, pathogenesis and biology? *Blood reviews*. 2015;29:205-13.
3. Flinn IW, Hillmen P, Montillo M, Nagy Z, Illés Á, Etienne G, et al. The phase 3 DUO trial: duvelisib vs ofatumumab in relapsed and refractory CLL/SLL. *Blood*. 2018;132:2446-55.
4. Sun C, Nierman P, Kendall EK, Cheung J, Gulrajani M, Herman SEM, et al. Clinical and biological implications of target occupancy in CLL treated with the BTK inhibitor acalabrutinib. *Blood*. 2020;136:93-105.
5. Chanan-Khan A, Cramer P, Demirkan F, Fraser G, Silva RS, Grosicki S, et al. Ibrutinib combined with bendamustine and rituximab compared with placebo, bendamustine, and rituximab for previously treated chronic lymphocytic leukaemia or small lymphocytic lymphoma (HELIOS): a randomised, double-blind, phase 3 study. *The Lancet Oncology*. 2016;17:200-11.
6. Byrd JC, Harrington B, O'Brien S, Jones JA, Schuh A, Devereux S, et al. Acalabrutinib (ACP-196) in Relapsed Chronic Lymphocytic Leukemia. *The New England journal of medicine*. 2016;374:323-32.
7. Fischer K, Al-Sawaf O, Bahlo J, Fink AM, Tandon M, Dixon M, et al. Venetoclax and Obinutuzumab in Patients with CLL and Coexisting Conditions. *The New England journal of medicine*. 2019;380:2225-36.
8. Burger JA, Chiorazzi N. B cell receptor signaling in chronic lymphocytic leukemia. *Trends in immunology*. 2013;34:592-601.
9. Byrd JC, Furman RR, Coutre SE, Flinn IW, Burger JA, Blum KA, et al. Targeting BTK with ibrutinib in relapsed chronic lymphocytic leukemia. *The New England journal of medicine*. 2013;369:32-42.
10. de Claro RA, McGinn KM, Verdun N, Lee SL, Chiu HJ, Saber H, et al. FDA Approval: Ibrutinib for Patients with Previously Treated Mantle Cell Lymphoma and Previously Treated Chronic Lymphocytic

- Leukemia. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2015;21:3586-90.
11. Barr PM, Brown JR, Hillmen P, O'Brien S, Barrientos JC, Reddy NM, et al. Impact of ibrutinib dose adherence on therapeutic efficacy in patients with previously treated CLL/SLL. *Blood*. 2017;129:2612-5.
 12. Burger JA, Tedeschi A, Barr PM, Robak T, Owen C, Ghia P, et al. Ibrutinib as Initial Therapy for Patients with Chronic Lymphocytic Leukemia. *The New England journal of medicine*. 2015;373:2425-37.
 13. Mato AR, Nabhan C, Thompson MC, Lamanna N, Brander DM, Hill B, et al. Toxicities and outcomes of 616 ibrutinib-treated patients in the United States: a real-world analysis. *Haematologica*. 2018;103:874-9.
 14. Woyach JA, Ruppert AS, Guinn D, Lehman A, Blachly JS, Lozanski A, et al. BTK(C481S)-Mediated Resistance to Ibrutinib in Chronic Lymphocytic Leukemia. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2017;35:1437-43.
 15. Woyach JA, Furman RR, Liu TM, Ozer HG, Zapatka M, Ruppert AS, et al. Resistance mechanisms for the Bruton's tyrosine kinase inhibitor ibrutinib. *The New England journal of medicine*. 2014;370:2286-94.
 16. Ahn IE, Underbayev C, Albitar A, Herman SE, Tian X, Maric I, et al. Clonal evolution leading to ibrutinib resistance in chronic lymphocytic leukemia. *Blood*. 2017;129:1469-79.
 17. Landau DA, Sun C, Rosebrock D, Herman SEM, Fein J, Sivina M, et al. The evolutionary landscape of chronic lymphocytic leukemia treated with ibrutinib targeted therapy. *Nature communications*. 2017;8:2185.
 18. Raponi S, Del Giudice I, Marinelli M, Wang J, Cafforio L, Ilari C, et al. Genetic landscape of ultra-stable chronic lymphocytic leukemia patients. *Annals of oncology : official journal of the European Society for Medical Oncology*. 2018;29:966-72.
 19. Del Giudice I, Raponi S, Della Starza I, De Propriis MS, Cavalli M, De Novi LA, et al. Minimal Residual Disease in Chronic Lymphocytic Leukemia: A New Goal? *Frontiers in oncology*. 2019;9:689.
 20. Zhou Y, Jia E, Pan M, Zhao X, Ge Q. Encoding Method of Single-cell Spatial Transcriptomics Sequencing. *International journal of biological sciences*. 2020;16:2663-74.
 21. Moffitt JR, Hao J, Wang G, Chen KH, Babcock HP, Zhuang X. High-throughput single-cell gene-expression profiling with multiplexed error-robust fluorescence in situ hybridization. *Proceedings of the National Academy of Sciences of the United States of America*. 2016;113:11046-51.
 22. Khan S, Kaihara KA. Single-Cell RNA-Sequencing of Peripheral Blood Mononuclear Cells with ddSEQ. *Methods in molecular biology (Clifton, NJ)*. 2019;1979:155-76.
 23. Rai KR, Jain P. Chronic lymphocytic leukemia (CLL)-Then and now. *American journal of hematology*. 2016;91:330-40.
 24. Ruffo E, Wu RC, Bruno TC, Workman CJ, Vignali DAA. Lymphocyte-activation gene 3 (LAG3): The next immune checkpoint receptor. *Seminars in immunology*. 2019;42:101305.

25. Xia M, Luo TY, Shi Y, Wang G, Tsui H, Harari D, et al. Effect of Ibrutinib on the IFN Response of Chronic Lymphocytic Leukemia Cells. *Journal of immunology* (Baltimore, Md : 1950). 2020;205:2629-39.
26. Shapiro M, Herishanu Y, Katz BZ, Dezorella N, Sun C, Kay S, et al. Lymphocyte activation gene 3: a novel therapeutic target in chronic lymphocytic leukemia. *Haematologica*. 2017;102:874-82.
27. Kotaskova J, Tichy B, Trbusek M, Francova HS, Kabathova J, Malcikova J, et al. High expression of lymphocyte-activation gene 3 (LAG3) in chronic lymphocytic leukemia cells is associated with unmutated immunoglobulin variable heavy chain region (IGHV) gene and reduced treatment-free survival. *The Journal of molecular diagnostics : JMD*. 2010;12:328-34.
28. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al. Comprehensive Integration of Single-Cell Data. *Cell*. 2019;177:1888-902.e21.
29. Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, et al. Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. *Science* (New York, NY). 2014;344:1396-401.
30. Gaidano G, Rossi D. The mutational landscape of chronic lymphocytic leukemia and its impact on prognosis and treatment. *Hematology Am Soc Hematol Educ Program*. 2017;2017:329-37.
31. Burger JA, Landau DA, Taylor-Weiner A, Bozic I, Zhang H, Sarosiek K, et al. Clonal evolution in patients with chronic lymphocytic leukaemia developing resistance to BTK inhibition. *Nat Commun*. 2016;7:11589.
32. Brown JR, Tesar B, Hanna M, Ash M, Fernandes SM, Macconail L, et al. Amplification of 6p Associated with Familial CLL. *Blood*. 2010;116:1009-.
33. Qian S, Li J, Hong M, Zhu Y, Zhao H, Xie Y, et al. TIGAR cooperated with glycolysis to inhibit the apoptosis of leukemia cells and associated with poor prognosis in patients with cytogenetically normal acute myeloid leukemia. *J Hematol Oncol*. 2016;9:128.
34. Adekola KUA, Liu V, Ma S, Rosen ST, Shanmugam M. Targeting Glucose Metabolism and Compensatory Metabolic Resistance in Chronic Lymphocytic Leukemia. *Blood*. 2012;120.
35. Zhang W, Trachootham D, Liu J, Chen G, Pelicano H, Garcia-Prieto C, et al. Stromal control of cystine metabolism promotes cancer cell survival in chronic lymphocytic leukaemia. *Nat Cell Biol*. 2012;14:276-86.
36. Vangapandu HV, Ayres ML, Bristow CA, Wierda WG, Keating MJ, Balakrishnan K, et al. The Stromal Microenvironment Modulates Mitochondrial Oxidative Phosphorylation in Chronic Lymphocytic Leukemia Cells. *Neoplasia* (New York, NY). 2017;19:762-71.
37. Kurtova AV, Balakrishnan K, Chen R, Ding W, Schnabl S, Quiroga MP, et al. Diverse marrow stromal cells protect CLL cells from spontaneous and drug-induced apoptosis: development of a reliable and reproducible system to assess stromal cell adhesion-mediated drug resistance. *Blood*. 2009;114:4441-50.
38. Woyach JA, Johnson AJ. Targeted therapies in CLL: mechanisms of resistance and strategies for management. *Blood*. 2015;126:471-7.

39. Komarova NL, Burger JA, Wodarz D. Evolution of ibrutinib resistance in chronic lymphocytic leukemia (CLL). *Proc Natl Acad Sci U S A*. 2014;111:13906-11.
40. Mendez-Huergo SP, Blidner AG, Rabinovich GA. Galectins: emerging regulatory checkpoints linking tumor immunity and angiogenesis. *Curr Opin Immunol*. 2017;45:8-15.
41. Do JS, Visperas A, Sanogo YO, Bechtel JJ, Dvorina N, Kim S, et al. An IL-27/Lag3 axis enhances Foxp3⁺ regulatory T cell-suppressive function and therapeutic efficacy. *Mucosal Immunol*. 2016;9:137-45.
42. Helfman DM, Kim EJ, Lukanidin E, Grigorian M. The metastasis associated protein S100A4: role in tumour progression and metastasis. *Br J Cancer*. 2005;92:1955-8.
43. Gao SP, Sun HF, Jiang HL, Li LD, Hu X, Xu XE, et al. Loss of COX5B inhibits proliferation and promotes senescence via mitochondrial dysfunction in breast cancer. *Oncotarget*. 2015;6:43363-74.
44. Zhou X, Liang S, Zhan Q, Yang L, Chi J, Wang L. HSPG2 overexpression independently predicts poor survival in patients with acute myeloid leukemia. *Cell Death Dis*. 2020;11:492.
45. DeBerardinis RJ, Chandel NS. Fundamentals of cancer metabolism. *Sci Adv*. 2016;2:e1600200.
46. Chowdhury SR, Banerji V. Targeting Mitochondrial Bioenergetics as a Therapeutic Strategy for Chronic Lymphocytic Leukemia. *Oxid Med Cell Longev*. 2018;2018.
47. Jitschin R, Hofmann AD, Bruns H, Giessel A, Bricks J, Berger J, et al. Mitochondrial metabolism contributes to oxidative stress and reveals therapeutic targets in chronic lymphocytic leukemia. *Blood*. 2014;123:2663-72.
48. Herold T, Jurinovic V, Metzeler KH, Boulesteix AL, Bergmann M, Seiler T, et al. An eight-gene expression signature for the prediction of survival and time to treatment in chronic lymphocytic leukemia. *Leukemia*. 2011;25:1639-45.
49. Astorgues-Xerri L, Riveiro ME, Tijeras-Raballand A, Serova M, Rabinovich GA, Bieche I, et al. OTX008, a selective small-molecule inhibitor of galectin-1, downregulates cancer cell proliferation, invasion and tumour angiogenesis. *European journal of cancer (Oxford, England : 1990)*. 2014;50:2463-77.
50. Xu W, Yang S, Zhou K, Pan L, Li Z, Zhou J, et al. Treatment of relapsed/refractory chronic lymphocytic leukemia/small lymphocytic lymphoma with the BTK inhibitor zanubrutinib: phase 2, single-arm, multicenter study. *Journal of hematology & oncology*. 2020;13:48.
51. Tam CS, Trotman J, Opat S, Burger JA, Cull G, Gottlieb D, et al. Phase 1 study of the selective BTK inhibitor zanubrutinib in B-cell malignancies and safety and efficacy evaluation in CLL. *Blood*. 2019;134:851-9.
52. Goud NS, Soukya PSL, Ghouse M, Komal D, Alvala R, Alvala M. Human Galectin-1 and Its Inhibitors: Privileged Target for Cancer and HIV. *Mini reviews in medicinal chemistry*. 2019;19:1369-78.
53. Thijssen VL, Heusschen R, Caers J, Griffioen AW. Galectin expression in cancer diagnosis and prognosis: A systematic review. *Biochimica et biophysica acta*. 2015;1855:235-47.
54. Nambiar DK, Aguilera T, Cao H, Kwok S, Kong C, Bloomstein J, et al. Galectin-1-driven T cell exclusion in the tumor endothelium promotes immunotherapy resistance. *The Journal of clinical investigation*.

2019;129:5553-67.

55. Zhang PF, Li KS, Shen YH, Gao PT, Dong ZR, Cai JB, et al. Galectin-1 induces hepatocellular carcinoma EMT and sorafenib resistance by activating FAK/PI3K/AKT signaling. *Cell death & disease*. 2016;7:e2201.
56. Croci DO, Morande PE, Dergan-Dylon S, Borge M, Toscano MA, Stupirski JC, et al. Nurse-like cells control the activity of chronic lymphocytic leukemia B cells via galectin-1. *Leukemia*. 2013;27:1413-6.
57. Asgarian-Omran H, Forghani P, Hojjat-Farsangi M, Roohi A, Sharifian RA, Razavi SM, et al. Expression profile of galectin-1 and galectin-3 molecules in different subtypes of chronic lymphocytic leukemia. *Cancer investigation*. 2010;28:717-25.
58. Gassner FJ, Zaborsky N, Catakovic K, Rebhandl S, Huemer M, Egle A, et al. Chronic lymphocytic leukaemia induces an exhausted T cell phenotype in the TCL1 transgenic mouse model. *British journal of haematology*. 2015;170:515-22.
59. Yu X, Huang X, Chen X, Liu J, Wu C, Pu Q, et al. Characterization of a novel anti-human lymphocyte activation gene 3 (LAG-3) antibody for cancer immunotherapy. *mAbs*. 2019;11:1139-48.
60. Wierz M, Pierson S, Guyonnet L, Viry E, Lequeux A, Oudin A, et al. Dual PD1/LAG3 immune checkpoint blockade limits tumor development in a murine model of chronic lymphocytic leukemia. *Blood*. 2018;131:1617-21.
61. Dickinson JD, Gilmore J, Iqbal J, Sanger W, Lynch JC, Chan J, et al. 11q22.3 deletion in B-chronic lymphocytic leukemia is specifically associated with bulky lymphadenopathy and ZAP-70 expression but not reduced expression of adhesion/cell surface receptor molecules. *Leukemia & lymphoma*. 2006;47:231-44.

Figures

Figure 1

Expression profiling of 42288 single cells from the PBMC of nine samples. A T-SNE plot of all single cell data (10× Genomics). R1-3 represents three IBR patients. S1-4 represents four IBS patients. N1-2 represents two normal donors. B Result of the unsupervised clustering of all cells. C Cell-type annotation in the t-SNE plot. D Highly expressed genes in different cell types. Dot size indicates the gene expression percentage in the same cell type. Color indicates the average gene expression in the same cell type. E PBMC components in nine samples. F Large-scale CNV in single cells inferred based on scRNA-seq data.

Figure 2

Distinct transcriptional pattern of IBR B cells and intercellular interactions among multiple cell types. **A** Left: Unsupervised clustering of B cells showed in t-SNE plot. Right: t-SNE plot displays B cells from different types of samples. Red dots represent B cells from IBR patients; Green dots represent B cells from IBS patients; Blue dots represent B cells from normal donors. **B** Stem score visualization for B cell in the t-SNE plot. **C** Comparison of the stem score of B cells from IBR patients, IBS patients and normal donors.

Significance was calculated using the Wilcoxon rank-sum test. D Upper: Proportion of cells derived from three sample types in each cluster. Red represents IBR patients. Green represents IBS patients. Blue represents normal donors. The dashed lines represent 0.3 and 0.5, respectively. Lower: Bar plot shows the number of cells in each cluster. The dashed line represents 200. E Metabolic pathway enrichment in each cluster. Red represents IBR clusters. Green represents IBS clusters. Orange represents shared clusters. F Graph showing interactions among different cell types. Line thickness represents the number of significant interactions. G Graph displaying the summary of selected ligand–receptor interactions between different cell types. Circle size represents p–values (permutation test). Color gradient indicates the score of interaction.

Figure 3

Identification of the potential drivers of IBR based on bulk and single cell transcriptomic data. **A** Left: Heatmap representing the differential gene expression between IBR and IBS patients based on bulk RNA-seq data. Red indicates IBR patients. Green represents IBS patients. Right: Volcano plot depicting DEGs. The significant up-regulated genes in IBR patients were shown with red, and the significant down-regulated genes in IBR patients were shown with green. **B** Developmental trajectory from IBS to IBR

cells based on scRNA-seq data. C Dynamic changes in gene expression from IBS to IBR. D Graph showing up-regulated genes in single-cell IBR clusters. E Graph showing dynamic changes in LAG3, LGALS1, and PTMS expression along the developmental trajectory from IBS to IBR. Color gradient indicates pseudo-time scores. F Heatmap visualization of genes co-expressed with LGALS1 based on bulk RNA-seq data. Patients were sorted by the expression of LGALS1. G Functional enrichment analysis of genes co-expressed with LGALS1. Dot size represents the number of enriched genes in the pathway. Color gradient represents adjusted p-value.

Figure 4

Increased LGALS1 and LAG3 expression correlates with poor survival in CLL. A Kaplan–Meier analysis of CLL patient samples based on LGALS1 and LAG3 expression (n = 59). *p ≤ 0.05, **p ≤ 0.01. B Survival analysis of patients from GSE22762 stratified based on LGALS1 and LAG3 expression. C LGALS1 and

LAG3 mRNA levels in CLL patients. Schematic diagram of diverse clinical features and prognostic patterns in CLL samples. β 2-MG, β 2-microglobulin; del, deletion; IGHV, immunoglobulin heavy variable gene. D Gal-1 and LAG3 protein levels in four IBS CLL patients and four IBR CLL patients. S, sensitive; R, resistant.

Figure 5

High LGALS1 and LAG3 expression in cells with acquired ibrutinib resistance. A Schematic diagram of ibrutinib-resistant CLL cell line (MEC1-IR) establishment and cell viability throughout. B Relative proliferation of MEC1 and MEC1-IR cells treated with different ibrutinib concentrations for 48 h. IC₅₀ values were calculated from dose-response curves for four biologically-independent samples. The IC₅₀ for each cell line is indicated in parentheses. C, D MEC1 and MEC1-IR cell apoptosis following exposure to ibrutinib (IBR) for 48 h. E-H LGALS1 and LAG3 RNA and protein levels in MEC1 and MEC1-IR cells estimated using qRT-PCR (E), western blot (F), and ELISA (G, H). * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$.

Figure 6

OTX008 inhibits CLL-cell growth and exhibits efficacy against IBR CLL cells. A Parental (MEC1) and acquired ibrutinib-resistant (MEC1-IR) CLL cell lines and primary CLL cells were treated with increasing OTX008 concentrations for 48 h. Relative LGALS1 and LAG3 expression in CLL cells at the indicated OTX008 concentration was measured using qRT-PCR. B, C Western blot and immunofluorescence for Gal-1 and LAG3 in OTX008-treated CLL cells. D Cell proliferation was analyzed using CCK8 assays with or without OTX008 treatment. E Cell viability was analyzed using flow cytometry as the percentage of Annexin-V and PI double-negative cells. F Freshly isolated PBMCs from CLL patients were incubated with

10 μ M ibrutinib and 30 μ M OTX008 for 48 h. Cell viability was assessed using trypan blue staining. Cells with > 20 % cytotoxicity at 10 μ M ibrutinib were regarded as IBS (blue, n = 6) and those with < 20 % as IBR (red, n = 9).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalData.pdf](#)
- [SupplementalTables.xlsx](#)