Characterization of Recombinant β Subunit of Human MUC4 Mucin (rMUC4 β)

Prakash G. Kshirsagar

University of Nebraska Medical Center

Mansi Gulati

University of Nebraska Medical Center

Wade M. Junker

University of Nebraska Medical Center

Abhijit Aithal

University of Nebraska Medical Center

Gaelle Spagnol

University of Nebraska Medical Center

Srustidhar Das

University of Nebraska Medical Center

Kavita Mallya

University of Nebraska Medical Center

Shailendra K. Gautam

University of Nebraska Medical Center

Sushil Kumar

University of Nebraska Medical Center

Paul Sorgen

University of Nebraska Medical Center

Krishan K. Pandey

Saint Louis University Health Sciences Center

Surinder K. Batra

University of Nebraska Medical Center

Maneesh Jain (✉ mjain@unmc.edu)

University of Nebraska Medical Center

Research Article

Keywords: Recombinant MUC4 β , CD and I-TASSER, Protein-protein interactions

Posted Date: August 23rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-812058/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on December 1st, 2021.

See the published version at <https://doi.org/10.1038/s41598-021-02860-5>.

Abstract

MUC4 is a transmembrane mucin expressed on various epithelial surfaces, including respiratory and gastrointestinal tracts, and helps in their lubrication and protection. MUC4 is also aberrantly overexpressed in various epithelial malignancies and functionally contributes to cancer development and progression. MUC4 is putatively cleaved at the GDPH site into a mucin-like α -subunit and a membrane-tethered growth factor-like β -subunit. Due to the presence of several functional domains, the characterization of MUC4 β is critical for understanding MUC4 biology. We developed a method to produce and purify multi-milligram amounts of recombinant rMUC4 β (rMUC4 β). Purified rMUC4 β was characterized by Far-UV CD and I-TASSER-based protein structure prediction analyses, and its ability to interact with cellular proteins was determined by the pull-down assay. Two of the three EGF-like domains exhibited typical β -fold, while the third EGF-like domain and vWD domain were predominantly random coils. We observed that rMUC4 β physically interacts with Ezrin and EGFR family members. Overall, this study describes an efficient and simple strategy for the purification of biologically-active rMUC4 β that can serve as a valuable reagent for a variety of biochemical and functional studies to elucidate MUC4 function and generating domain-specific antibodies and vaccines for cancer immunotherapy.

Introduction

Mucins comprise a family of 21 high molecular weight secretory or membrane-bound glycoproteins that provide the structural framework of a 'mucus-like' layer covering the epithelial surfaces within our body. Their main roles are to protect epithelial surfaces from various insults and promote their regeneration and repair¹⁻⁴. One of the mucins, MUC4, is a large, multi-domain, transmembrane glycoprotein involved in multiple physiological and pathological processes⁵. Full-length MUC4 is synthesized as a single polypeptide that is putatively cleaved in an auto-catalytic manner at the Gly-Asp-Pro-His (GDPH) site located within the vWD domain resulting in two subunits, MUC4 α and MUC4 β ⁵⁻⁷. The large extracellular N-terminal subunit MUC4 α contains a characteristic tandem repeat (TR) domain along with NIDO and AMOP domains. The smaller, membrane-tethered, C-terminal subunit MUC4 β includes a truncated vWD domain, three (EGF)-like domains, and a cytoplasmic tail^{8,9}. Due to its aberrant overexpression in multiple carcinomas (pancreas, lung, cervix, ovary, colon, skin), MUC4 has emerged as a promising biomarker and therapeutic target¹⁰⁻¹³. Furthermore, MUC4 is also a potential target for cancer immunotherapy, owing to its aberrant glycosylation and the existence of several splice variants^{12,14}. The large, heavily glycosylated MUC4 α subunit plays a dominant role in modulating the adhesive properties of epithelial cells¹⁵. However, the MUC4 β subunit harboring three EGF-like domains mediates interactions between MUC4 and EGFR-family proteins (EGFR-1/HER-2/HER-3)¹⁶⁻²⁰. These interactions potentiate a diverse array of signaling pathways that result in aggressive growth and enhanced chemoresistance of cancer cells^{6,15,16,18}.

We have previously cloned MUC4 "minigene" containing all the MUC4 domains but only 10% of the TR domain and used it to discern the functional role of MUC4 in cancer cell lines using overexpression

studies^{8,21}. However, direct structure-function studies of MUC4 have been hampered by its large molecular size, lack of domain-specific reagents, and complexities associated with the existence of diverse glycoforms and splice variants in cancer. Purification of the mega-Dalton-sized, 'full-length' MUC4 by presently available molecular biology techniques is extremely challenging. Due to its potential retention on the cell membrane post-cleavage and the presence of several functional domains, we directed our efforts to clone, produce, and purify a recombinant MUC4 β (rMUC4 β) subunit using a bacterial expression system. The purified rMUC4 β has recently been evaluated for vaccine development^{22,23} and the generation of domain-specific monoclonal antibodies (mAbs)^{24,25}. Herein, we optimized the parameters for the production and purification of rMUC4 β and undertook its preliminary biophysical characterization. We also tested the utility of purified rMUC4 β for identifying MUC4-interacting partners. The optimized production and purification method yielded multi-milligram amounts of pure and soluble rMUC4 β . In affinity pull-down assays, rMUC4 β recognized known (EGFR, HER2, and HER3) and novel (Ezrin) MUC4-interacting partners. Our studies suggest that rMUC4 β produced and purified using the optimized method described herein, retains the characteristics of native MUC4, and can serve as a valuable tool to identify MUC4-interacting partners and undertake structure-function studies.

Materials And Methods

Materials

C41(DE3) and BL21(DE3) *E. coli* strains were purchased from Sigma-Aldrich. C41pLysS, C43pLysS, and Rosetta™ 2 *E. coli* strains were procured from Thermo scientific. Bacto™ tryptone, Bacto™ yeast, and Bacto™ Agar were purchased from BD Biosciences. Anti-His tag Ab (Clone 27E9, Cat. No. 2366) and anti-EGFR mAb (Clone D38B1, Cat. No. 4267) were purchased from Cell Signaling, while anti-Ezrin Ab (Clone H-276, Cat No. SC-20773) was bought from Santa Cruz Biotechnologies. HisTrap™ HP affinity Ni-NTA column was purchased from GE healthcare. Anti-His tag Ab Magnetic bead conjugate (Cat. No. 8811) and Protein A/G agarose beads (Cat. No. 9863) were purchased from Cell Signaling. Anti-MUC4 β mAb (Clone 6E8) was generated using recombinant rMUC4 β protein as an immunogen. More details on the construction of recombinant pET-28a-MUC4 β are mentioned in supporting information.

Optimization of culture parameters to enhance rMUC4 β yield

Selection of *E. coli* host for optimal production. Five different chemically competent *E. coli* (DE3) strains (C41, BL21, C41pLysS, C43pLysS, and Rosetta™ 2) were tested for optimal expression of rMUC4 β protein. Briefly, selected strains transformed with pET28a-MUC4 β plasmid were grown in LB/antibiotics media overnight at 37°C. A freshly-made pre-inoculum culture (2%) from each cell type was inoculated into 20 mL LB media and incubated at 37°C until the mid-exponential growth phase. When OD_{600 nm} reached ~0.6, cultures were induced with 0.5 mM IPTG for 4 h at 37°C (a set of negative control was incubated under similar conditions without IPTG). Next, both induced and uninduced fractions were harvested and disrupted in an appropriate volume of lysis buffer (500 mM Tris HCl, 2 mM MgCl₂, 10 mM NaCl, pH 8.0)

to adjust the OD_{600 nm} to ~1.2/mL. Equivalent amounts of crude lysate were loaded, ran on SDS-PAGE, and stained with Coomassie brilliant blue.

Optimization of rMUC4 β expression. C41(DE3) and Rosetta™ 2(DE3) competent cells transformed with pET28a-MUC4 β plasmid were grown in LB culture at 37°C. After reaching the mid-exponential phase, all bacterial cultures were separately induced at variable IPTG concentrations (0, 0.25, 0.5, and 1.0 mM) for 4 h at 37°C. Uninduced controls were prepared without adding IPTG. The effect of three different post-induction temperatures on the enrichment of rMUC4 β expression was evaluated in pET-28a-MUC4 β transformed Rosetta™ 2(DE3) cells⁶⁰. Bacterial culture was induced at ~0.6 OD by adding 0.5 mM of IPTG. After induction, each tube was incubated separately at 16°C, 25°C, and 37°C, respectively, for 4 h. To investigate the effects of different culture media on the expression efficiency of rMUC4 β in *E. coli*, freshly-made pre-inoculum culture (2%) of Rosetta™ 2(DE3) cells transformed with pET-28a-MUC4 β plasmid were added to 20 mL of culture media (LB and TB) and grown at 37°C until medium exponential phase. One set of grown cultures was induced with 0.5 mM IPTG and incubated for 4 h at 37°C. The second control set was grown without IPTG addition. A sample from each culture condition was pelleted and lysed in an appropriate volume of lysis buffer to maintain equal ODs (~1.2/mL). After lysis, SDS-PAGE and immunoblotting analysis were performed using equal amounts of lysates to identify IPTG concentration, temperature, and culture media for optimal production.

Scale-up production of rMUC4 β protein in Rosetta™ 2(DE3)

Large-scale production of rMUC4 β protein was undertaken using optimized culture conditions. The transformed cells were grown on LB agar plates containing kanamycin (50 μ g/mL) and chloramphenicol (40 μ g/mL). After overnight incubation at 37°C, a single colony was picked and grown overnight into 10 mL of LB/antibiotics culture in a shaking incubator at 37°C. Following day, 50 mL of LB/antibiotics culture was inoculated by 500 μ L of overnight-grown culture (1%) and incubated overnight at 37°C, 250 rpm. The next day, overnight grown pre-inoculum (15.0 mL) was inoculated into 1 L of LB/antibiotics media. Subsequently, the culture was incubated under continuous agitation at 37°C until OD_{600nm} reached ~0.5-0.6. MUC4 β expression was induced by adding IPTG [0.5 mM] and growing for another 4 h at 37°C. Cells were harvested by centrifugation (6000 rpm, for 30 min, at 4°C), washed once with ice-cold 1 $\times$ PBS, and stored overnight at -80°C⁶¹. The bacterial pellet was thawed and resuspended in ice-cold lysis buffer [50 mM Tris-HCl (pH 8.0), 2 mM MgCl₂, and 150 mM NaCl, EDTA-free protease inhibitor cocktail (250 μ L/10 g of the bacterial pellet) and benzonase nuclease (~10 U/mL pre-lysis culture). Following 15 min incubation on ice, cells were lysed with three passages through an EmulsiFlex-C3 (Avestin) at ~15,000 psi. Cell debris was removed by centrifugation (6000 rpm for 30 min at 4°C). The supernatant was treated again with benzonase nuclease (10 U/mL cell lysate) and incubated for 2 h at 4°C^{43,62}. The rMUC4 β protein present in the insoluble inclusion bodies was collected by high-speed centrifugation (18,000 rpm for 60 min at 4°C). The recovered pellet was resuspended in 40 mL of buffer A (1 $\times$ PBS, 6 M urea, 20 mM imidazole, 300 mM NaCl, 0.5% CHAPS, and 2 mM β -mercaptoethanol (BME), pH 8.0), incubated overnight at 4°C and again treated with Benzonase nuclease. Insoluble debris was

removed by centrifugation (18,000 rpm, 60 min, at 4°C), and the pre-ÄKTA supernatant was passed through a 0.22 µm Steriflip® vacuum filtration unit (Millipore). More details on the purification and refolding of rMUC4β protein can be found in supporting information.

Circular dichroism (CD) spectroscopy

Circular dichroism (CD) analysis of rMUC4β was performed on a Jasco J-815 spectrometer in the Far-UV spectral region ($\lambda = 190$ to 260 nm) using a 0.1 mm path length quartz cell. CD spectra of rMUC4β at a concentration of 39 µmol/L in 1× PBS were recorded at 7°C, 25°C and 37°C using a scan rate of 50 nm/min, an integration time of 1 second, and a bandwidth of 1.0 nm. The final spectrum was obtained by averaging five scans and was corrected by subtracting the solvent spectrum acquired under identical conditions and plotted using the GraphPad Prism8 software. CD data were processed using the spectra analysis function of Jasco Manager Version 2. Analysis of the CD spectrum was performed separately using two methods, CDSSTR and Contin/LL provided by DichroWeb (<http://dichroweb.cryst.bbk.ac.uk/>)
30.

Iterative Threading ASSEmbly Refinement (I-TASSER)

I-TASSER is a bioinformatics program that predicts the three-dimensional (3-D) model structure of the desired protein using amino acid (AA) sequences. We used this intelligent program to model the 3-D structure of the MUC4β protein by submitting the corresponding AA sequences to the I-TASSER online server (<https://zhanglab.ccmb.med.umich.edu/I-TASSER/>). After finding five different predicted models of MUC4β, the structure with the highest confidence score was selected for comparative analysis with CD data.

Affinity pull-down assays

High-affinity Ni-NTA resin (GenScript, Cat. No. L00250) was washed and equilibrated with ice-cold wash buffer (1× PBS, containing 1% CHAPS, 20 mM imidazole, and 2 mM BME, 300 mM NaCl, pH 8.0). The equilibrated Ni-NTA slurry was incubated with either CD18/HPAF or PANC-1 cell for 4 h at 4°C to pre-clear the cell lysates. Ni-NTA resin, after pre-clearing for each sample, was used as a “non-specific binding control.” A portion of each pre-cleared lysate supernatant was saved and used as ‘Input’ control. The lysates were incubated with 60 µg of rMUC4β protein [rMUC4β (+)] or without rMUC4β protein [rMUC4β (-)], in total 500 µL reaction volume overnight at 4°C with end-over-end mixing. The samples were centrifuged (2000 rpm, for 2 min at 4°C), and the supernatant containing unbound proteins was designated as “flow-through” (FT) and saved for further analysis. The Ni-resin was washed five times (2000 rpm, for 2 min at 4°C) with ice-cold wash buffer (1× PBS, containing 20 mM imidazole, 1% CHAPS, 2 mM BME, 300 mM NaCl, pH 8.0) and the last wash (Wash 5) was saved for analysis. Finally, the ‘bait-prey’ complexes were eluted with elution buffer (1× PBS supplemented with 500 mM imidazole, 1% CHAPS, and 2 mM BME, 300 mM NaCl, pH 8.0) ice for 15 min. The eluted proteins (Eluate) were recovered in the supernatant following centrifugation. An aliquot of rMUC4β-bound Ni-resin was also eluted with elution buffer to serve as a positive control (i.e., Eluted Recombinant Protein, “ERP (+)” control). All

collected samples were solubilized in 6× Laemmli sample buffer, heated at 95°C for 5 min, and subjected to SDS-PAGE and immunoblot analyses briefly described in supporting information.

Mass spectrometry (MS/MS) characterization of excised protein band

The protein profile of pull-down eluted complexes in two different cell lysates (CD18/HPAF and PANC-1) were visualized by Coomassie staining and silver staining. The distinct protein band is present in the eluted fraction of the rMUC4β (+) panel but absent in the eluted fraction rMUC4β (-) panel in both CD18/HPAF and PANC-1 lysates were destained and stored in 50% methanol overnight at 4°C. The selected band (MW ~80 kDa) was excised and digested with trypsin by the UNMC Mass Spectrometry and Proteomics Core. Following digestion, samples were dried by SpeedVac, resuspended in 20 µl of 0.5% trifluoroacetic acid, and were cleaned up using ZipTip µC18 columns (Millipore). Further, the samples were resuspended in 0.1% formic acid and injected onto an Eksigent cHiPLC column (75 µm x 15 cm ChromXP C18-CL 3 µm 120 Å), and 6600 TripleTOF instruments (gradient 2-60% ACN in 60 min).

Results

Cloning and expression of MUC4β

The schematic structure of MUC4 and the overall scheme of cloning strategy for the construction of the pET28a-MUC4β expression plasmid is described in **Fig. 1A**. The final construct was sequenced to confirm in-frame insertion of MUC4β with N-terminal hexahistidine tag (**Fig. S1**). The overall schema for the expression and purification of rMUC4β is described in **Fig. S2A**. Initially, C41(DE3) cells transformed with pET28a-MUC4β were grown until the mid-log phase (**Fig. S2B**) and induced by the addition of IPTG. The SDS-PAGE and immunoblot analyses of culture supernatants indicated a distinct ~72 kDa rMUC4β band following IPTG induction (**Fig. 1B**, lane 7 and 8), whereas negative control lanes did not show any expression (**Fig. 1B**, lanes 1-6). The initial purification yield of rMUC4β from a 4 L culture of C41(DE3) cells was found to be low (~2-3 mg/L). Hence, we optimized various culture growth and purification conditions (**Fig. S2C-D**) to improve the yield and purity of rMUC4β protein.

Optimization of culture parameters for rMUC4β expression

To improve the yield of purified rMUC4β, we optimized various parameters and examined their effect on protein expression. Five different, chemical competent *E. coli* (DE3) strains, namely C41, BL21, C41pLysS, C43pLysS, and Rosetta™ 2, were transformed and tested for rMUC4β expression. Of all the strains tested, the Rosetta™ 2(DE3) exhibited the highest rMUC4β expression, followed by C41(DE3), while the expression was very low in other strains (**Fig. S3A-B**). Comparative assessment of rMUC4β expression at variable IPTG concentrations showed higher rMUC4β expression in Rosetta™ 2(DE3) as compared to C41(DE3) (**Fig. 1C**). We also examined the effect of different culture media (i.e., Terrific Broth (TB) and LB) on the expression of rMUC4β in Rosetta 2(DE3) at different IPTG concentrations. We observed a ~1.3-fold higher expression of rMUC4β in the TB compared to LB (data not shown). However, the cost/benefit provided by the TB media was not substantial enough to justify its use, and thus LB was

employed for all further studies. A distinctly higher expression of rMUC4 β was observed at 25°C and 37°C following IPTG induction, while a relatively low expression was seen at 16°C (**Fig. 1D**). To minimize nucleic acid contamination, the Benzonase nuclease treatment was incorporated in the purification scheme (**Fig. S2A**). Treatment of Rosetta™ 2(DE3) lysates with Benzonase nuclease for 30 min at 25°C reduced the viscosity in a dose-dependent manner (**Fig. S3C**) and hydrolysis of nucleic acids into smaller fragments (<100 bp) (**Fig. S3D, lane 3**)^{26, 27}. Nucleic acids were undetectable in the purified rMUC4 β fraction recovered from Benzonase nuclease treated lysate (**Fig. S3D, lane 10**) compared to untreated lysate (**Fig. S3D, lane 11**).

Scale-up production and purification of MUC4 β

Using the optimized conditions (i.e., Rosetta™ 2 strain, 0.5 mM IPTG, 37°C, 4 h) for growth and induction, we scaled up the production to obtain a multi-milligram amount of rMUC4 β . A major fraction of rMUC4 β was observed in the inclusion bodies and was relatively pure (**Fig. 1E, Lane 2**). Thus, inclusion bodies were isolated by centrifugation, solubilized in 6M urea, and subjected to purification as outlined in **Fig. S2A**. The eluted protein was soluble in the 6M urea but exhibited significant precipitation when urea was removed following dialysis (**Fig. S2D**). To enhance the protein solubility, a zwitterionic detergent CHAPS was used instead of Triton X-100 in both wash and elution buffers during purification. This led to increased solubility of rMUC4 β during the refolding steps and minimized the residual detergent mass in the final lyophilized protein. Using the optimized expression and purification conditions, we accomplished a ~9-fold increase (i.e., ~18 mg/L, +50% recovery) in the total yield of purified-refolded rMUC4 β (**Table 1**). The purified rMUC4 β was observed as a single 72 kDa band in SDS-PAGE (**Fig. 1E, Lane 6**). Immunoblot analysis of purified protein was also performed for various fractions and conditions to validate using anti-His tag antibody (**Fig 1B-D**). Interestingly, immunoblot analysis of the purified protein following BN-PAGE indicated the tendency of rMUC4 β to form discrete covalent dimer a ~150 kDa under non-reducing conditions, which was resolved into a monomeric form in the presence of BME (**Fig. S3E**).

Prediction of MUC4 β secondary structure by CD spectroscopy

To evaluate the stability, folding, and structural integrity of the renatured rMUC4 β protein, we performed CD analysis and i-TASSER-based structure prediction. The Far-UV spectrum collected at 7°C and higher temperatures did not show significant differences (**Fig. 2A, S3F**). Analysis of the secondary structure recorded at 7°C is demonstrated in **Fig. 2B**. The CD spectrum (λ =190-260 nm) of refolded rMUC4 β (**Fig. 2A**) is characterized by one broad minima around 215 nm typical of β -strand structure and a lower than expected maxima at about 195 nm that is most likely caused by a significant amount of random coil content (typically shows as a pronounced minima at 197nm)²⁸. Further, the assessment of the CD spectrum (**Fig. 2B-D**) was performed separately using the CDSSTR and Contin/LL (Provencher & Glockner Method) programs provided by the online analysis tool DichroWeb^{29, 30}. Both analyses revealed an average secondary structure composition of ~10% for α -helical structure, ~48% for β -strand, and ~41% for random coil (**Fig. 2C-D**). Comparison of the secondary structure prediction data from the amino acid

sequence (GOR IV), I-TASSER, and CD (average composition from the CDSSTR or CONTIN/LL) analyses are shown in **Fig. 2B**.

3-D structure prediction of MUC4 β by I-TASSER

To predict the tertiary structure of MUC4 β , the amino acid sequence was submitted for 3-D structure prediction using an online server i-TASSER³¹⁻³³. The server-generated five different models were analyzed for the composition of the secondary structure. The percentage of α -helical structures in the 5 predicted models ranged between 1.7 to 6%, while extended β -strands ranged from 10.1 to 29.4%, and random coil structure ranged from 68 to 84% (PDB code: 5G56). Model 1 was chosen for further analysis as it has the highest confidence score (**Fig. 2E**). The remaining predicted structures were arranged according to their decreasing confidence scores (**Fig. S4B-E**). The tertiary structure of MUC4 β (Model 1) is displayed in a rainbow-colored spectrum (**Fig. 2E**), with the N-terminus and C-terminus denoted in blue and red color, respectively. To better visualize the EGF-like domains contained in MUC4 β , the structure was rotated 180 degrees. The model revealed that the EGF-like domain 1 (Cyan) and EGF-like domain 2 (Magenta) are composed of the expected typical β folds (anti-parallel b-sheet), while the EGF-like domain 3 (red) shows as a mostly random coil structure (**Fig. 2E**, right panel). Meanwhile, the vWD domain (Blue) appears as a highly random coiled structure; three small α -helix and three β -strands forming a parallel b-sheet) are observed.

Affinity pull-down and mass spectrometry to detect MUC4 interacting partners

To determine if the rMUC4 β exhibits a similar conformation to native MUC4, we tested its ability to interact with known and potentially novel MUC4-interacting partners using a His-affinity pull-down assay as outlined in **Fig. 3A**. We observed that rMUC4 β pulled down a ~80 kDa band from the lysates of both MUC4-expressing (CD18/HPAF) and non-expressing (PANC-1) cell lines (**Fig. 3B-C, lane 13**) which was not detected in the pull-down eluates when beads alone were used (**Fig. 3B-C, lane 6**). This band was excised, in-gel trypsin digested, and the resulting peptides were analyzed by tandem MS/MS. The most significant top hits were matched to Ezrin in CD18/HPAF and PANC-1 cells (**Fig. S5A-B**). Analysis of the pulled-down complexes by immunoblotting with the antibodies of known MUC4-interacting proteins (EGFR family receptors) indicated that rMUC4 β interacts with EGFR, HER2, and HER3 in both MUC4-expressing and non-expressing cells (**Fig. 4A-B**). Interaction with Ezrin was confirmed by a pull-down assay using an anti-His tag Ab-coupled to magnetic beads to immobilized rMUC4 β (**Fig. 4C**). Finally, the ability of endogenous MUC4 β to interact with Ezrin was further confirmed by CO-IP (i.e., immunoprecipitation with anti-MUC4 β mAb 6E8 (**Fig. 4D**)). Overall, these results suggest that the rMUC4 β recapitulates the functional conformation of native MUC4 β and can recognize its natural interacting partners.

Discussion

Due to their differential expression in various malignancies, mucins have been extensively investigated as determinants of disease initiation and progression and as diagnostic and therapeutic targets^{1, 34-36}.

MUC4 is a high molecular weight, multi-domain, and heavily glycosylated protein that is differentially overexpressed in multiple cancers, including pancreatic, breast, lung, and head and neck cancers, where it functionally contributes to disease initiation, progression, metastasis, and chemoresistance^{5, 9, 11, 13, 37, 38}. It is also a potential biomarker for the diagnosis and prognosis of various cancers and a novel target for cancer therapeutics¹¹⁻¹³.

Most information regarding MUC4 function has been discerned using cell lines engineered for MUC4 overexpression or knockdown, while its pathobiological significance has been determined from correlative studies involving the analysis of biological samples for MUC4 transcripts and protein. However, direct structure-function studies for human MUC4 have not been extensively performed due to its large genomic, molecular size, and extensive glycosylation. Due to the large size of the MUC4 gene and transcript, cloning and expressing a full-length MUC4 protein is practically impossible. We have previously cloned the MUC4 “minigene” in a eukaryotic expression system, containing only 10% of the tandem repeat sequence but with all other domains intact^{8, 21}. While this construct has been useful for studying the impact of MUC4 overexpression in non-expressing cells to discern MUC4 function, it is unsuitable for large-scale purification. An alternative approach is to clone, express, and purify smaller fragments containing functional domains. We have previously cloned and expressed various domains of the MUC4 α subunit in developing domain-specific anti-MUC4 monoclonal antibodies³⁹. Due to the presence of three EGF-like domains and a vWD domain, and its proximity to the cell membrane, several functions of MUC4, particularly the ones involving interactions with the EGF receptor family^{18, 19, 38}, have been attributed to MUC4 β subunit.

Further, our recent analysis suggests that the immunogenic peptides in MUC4 are enriched in the β -subunit¹². Due to these attributes, the MUC4 β subunit is a potential target for immunotherapy and targeted therapies^{11, 12, 29}. Thus, we directed our efforts to clone, express, and purify the MUC4 β subunit with a goal to utilize the purified recombinant protein to develop domain-specific antibodies and pancreatic cancer immunotherapy. Recently, Liberelle *et al.* generated fusion constructs containing GFP fused to either intact human MUC4 β or truncated mutants containing only EGF-like domains. These constructs were expressed in CHO cells and used to characterize MUC4-ErbB2 interactions. However, the studies were performed using CHO cell lysates and recombinant ErbB2, and no efforts were made to purify MUC4 β -GFP¹⁹.

Initially, the MUC4 β sequence was cloned into the pET-28a (+) expression vector. Initial scale-up attempts using 1L culture of pET28a-MUC4 β plasmid transformed in C41(DE3) cells resulted in a low yield of purified MUC4 β (i.e., ~1.8 mg/L of LB media). Analysis of MUC4 β sequence, with a rare codon calculator (RaCC), indicated the presence of 32 rare or minor codons including AGA, AGG, CGA, CGG, CUA, CCC, and AUA (<https://web.expasy.org/protparam/>). Rare or minor codons are underutilized codons by the bacterial translational machinery, resulting in low protein yield. We screened five different genetically engineered *E. coli* strains for MUC4 β expression, of which RosettaTM 2(DE3) provided the maximum improvement (>10 fold than the C41 strain) in protein yield (**Table 1**). This specific strain expresses the less utilized tRNAs to

compensate for rare codons that are generally used in eukaryotes but rarely used in *E. coli*⁴⁰⁻⁴². Prior to scaling up the production, we optimized various parameters, including post-induction incubation temperature, the concentration of IPTG, and culture media composition, to improve the rMUC4 β yield. Under conditions optimized for high expression (IPTG- 0.5 mM; growth medium-LB broth), rMUC4 β accumulated in inclusion bodies. We developed a comprehensive purification strategy that included cell lysis, removal of nucleic acids, isolation and solubilization of inclusion bodies, immobilized metal affinity chromatography, and dialysis to facilitate urea removal and protein refolding. Due to the high amount of nucleic acids, the high viscosity of the bacterial lysates can potentially interfere with downstream processing and impact protein yield. Reduction in cell lysate viscosity can be achieved by supplementing bacterial lysate with Benzonase nuclease^{26, 28, 43}. The use of CHAPS instead of non-ionic Triton X-100 in the solubilization and elution steps prevented protein aggregation during urea removal and contributed to the high solubility of rMUC4 β in water attributed to the high CMC value and zwitterionic nature of CHAPS⁴⁴. Immunoblotting analysis following BN-PAGE under reducing and non-reducing conditions suggested the tendency of purified rMUC4 β to form covalent dimers via disulfide bonds. The possibility of higher-order soluble aggregates can be ruled out as we did not observe higher molecular weight bands even by immunoblotting.

To gain insight into protein folding, we employed a GOR IV server to predict the secondary structure by amino acid sequences. Further, we performed structural characterization of the purified rMUC4 β protein by conducting CD spectroscopy. The CDSSTR and Contin/LL methods were employed to analyze the Far-UV CD spectrum^{29, 30, 45}. CD data demonstrated the presence of α -helix, β -strand, and random coils. While understanding the secondary structure formation, we observed structural disparity in the β -structure derived from CD and other secondary structure sequence prediction methods. Supportively, some reports have shown some discrepancies in CD analysis and sequence-based structure predictions regarding the quantification for β -structure because of their morphological and structural diversity⁴⁶⁻⁴⁸. To gain further insights into the tertiary structure and understand the folding of the rMUC4 β , we performed structure prediction analysis using an online-based I-TASSER server.

The server-generated five models. The model with the highest confidence score was selected to understand the folding pattern and its interacting partners. The predicted tertiary structure showed typical β -folds for two EGF-like domains, while the third EGF-like domain showed a random coiled structure. The vWD domain is majorly disordered, but we did observe a parallel β -sheet comprised of three β -strands (3-4 residues each). Three small α -helical structures (4 residues each) were also observed in the vWD domain. The presence of a high percentage of random coils (disordered structures) in MUC4 can be attributed to the existence of a high proportion of charged or polar residues. The decreased hydrophobic residues make the protein less susceptible to present a hydrophobic ordered core^{49, 50}. A previous study analyzing the intrinsically disordered structures of mucins has shown that MUC4 is 77% disordered; it also highlighted the vWD domain as a highly disordered domain⁵¹. Although the recombinant protein is non-glycosylated, protein-protein interaction studies using *in vitro* and *in silico* analysis showed MUC4 interaction with EGFR and HER2, as previously reported by native protein^{17-19, 52}. We observed some

differences in the predicted structures by GOR IV and i-TASSER; these differences can be attributed to the different algorithms employed by these two servers. GOR IV is a secondary structure prediction method that uses a probability-based algorithm and performs the jack-knife method to determine the secondary structure. This method aligns the target sequence with the GOR database of 267 sequences and has a prediction accuracy of about 64%⁵³. The tertiary structure predicted by i-TASSER involves structure prediction by identifying the structure templates by Local Meta-Threading Server (LOMETS) and other threading software, which generate thousands of structure alignments⁵⁴. The structures with the highest significance are ranked based on Z-scores. Both methods utilize different approaches to predict the structure of the target sequence; therefore, we observe differences in the predicted structures. The structural and functional characterization of the rMUC4 β paves a path to further explore its potential as a therapeutic agent.

MUC4 plays multifaced roles in cell adhesion, migration, proliferation to promote tumorigenesis and metastasis in pancreatic cancer and other epithelial malignancies^{5, 6, 37, 38}. One of the ways by which it mediates these roles is by interacting with cell surface molecules such as HER2, HER3, and integrins and soluble mediators in the extracellular microenvironment such as galectins^{16, 18, 20, 55}. To determine whether rMUC4 β achieves a folding state and is biologically active, we performed an affinity pull-down assay utilizing rMUC4 β as the 'bait' to isolate putative interacting partners from RIPA extracted lysates of a MUC4 expressing (CD18/HPAF) and a MUC4 non-expressing (PANC-1) cell line. A ~80 kDa band was observed in the eluted fraction that was identified as 'Ezrin' by MS/MS analysis. Ezrin is a cytoskeletal protein belonging to the Ezrin-Radixin-Moesin (ERM) family that plays an essential role in cell adhesion, motility, invasion, cancer progression, and metastasis⁵⁶⁻⁵⁸. Recently, the TCGA and CCLE datasets were used to identify genes for which expression is correlated with MUC4. EZRIN was one of the 178 reported genes with a Pearson correlation higher than 0.3¹⁷. However, to the best of our knowledge, no study has yet provided any experimental evidence pointing to the possible physical interaction between MUC4 and Ezrin. Interestingly, it has been shown that the cytoplasmic tail of another mucin, "MUC16," interacts with the ERM family of proteins via its characteristic polybasic sequence⁵⁹, connecting it to the actin cytoskeleton in the cytoplasm. It is possible that mucin-ERM interactions might be an important phenomenon and have functional relevance yet to be described.

We also confirmed the interaction of known interacting partners of MUC4 by immunoblotting the eluted fraction with specific antibodies. Both HER-2 and HER-3 were shown previously to interact with MUC4¹⁷⁻¹⁹ and are detected in the eluted fraction following multiple pull-down assays described with rMUC4 β . This further confirms that the interaction of these receptors with MUC4 lies in its β -subunit domain and is independent of glycosylation. Interestingly, we could also detect EGFR in the eluted fraction. This can either be due to direct interaction with the protein or indirectly because of its association with HER2. Given the folding pattern observed in our structure prediction analysis, it is possible that the three EGF-like domains of MUC4 bind to the EGF binding pocket of EGFR. Further studies are needed to confirm or rule out these observations. It will be interesting to determine if exogenously added rMUC4 β can potentially compete with the endogenous MUC4 for binding partners and serve as a potential therapeutic agent.

In summary, we successfully developed an expression and purification system to produce multi-milligram amounts of rMUC4 β . The recombinant protein fragment retained the binding characteristics of the native MUC4 β and can be a valuable tool to undertake MUC4 structure-function analysis. Further, purified rMUC4 β has been successfully formulated and evaluated as a nanovaccine^{22, 23} and used as an immunogen for the development of domain-specific anti-MUC4 antibodies^{24, 25}.

Abbreviations

Ab: antibody; EGFR: epidermal growth factor receptor; FPLC: fast protein liquid chromatography; MW: Molecular weight; MUC4: Mucin 4; MWCO: Molecular weight cut-off; PCR: polymerase chain reaction; TB: terrific broth; LB: Luria broth, BME: β -mercaptoethanol, vWD: Von Willebrand factor type D domain, PIC: Protease inhibitor cocktails, FPLC: Fast pressure liquid chromatography.

Declarations

Author contribution statement

P.G.K.: Conceptualization, Methodology, Validation, Formal analysis, Writing-Original draft, Prepared figures, Writing- Review & Editing. **M.G.:** Methodology, Validation, Formal analysis, Prepared figures, Writing- Review & Editing. **W.M.J.:** Methodology, Validation, Formal analysis, Writing- Review & Editing. **A.A.:** Methodology, Validation, Formal analysis, Writing- Review & Editing. **G.S.:** Formal analysis, Writing- Review & Editing. **S.D.:** Formal analysis, Writing- Review & Editing. **K.M.:** Formal analysis, Writing- Review & Editing. **S.K.G.:** Formal analysis, Writing- Review & Editing. **S.K.:** Formal analysis, Writing- Review & Editing. **P.S.:** Formal analysis, Resources, Writing- Review & Editing. **K.K.P.:** Formal analysis, Writing- Review & Editing. **S.K.B.:** Conceptualization, Resources, Supervision. **M.J.:** Conceptualization, Resources, Supervision.

Acknowledgments

This work and the authors of this manuscript are supported, in parts, by grants from the National Institutes of Health (U01 CA213862, R01 CA195586, R01 CA247471, and R44 DK117472). The authors acknowledge the invaluable technical support from Nicole Japp.

Conflicts of Interest

SKB is one of the co-founders of Sanguine Diagnostics and Therapeutics, Inc. The other authors disclosed no potential conflicts of interest.

References

1. Kaur, S., Kumar, S., Momi, N., Sasson, A.R. & Batra, S.K. Mucins in pancreatic cancer and its microenvironment. *Nature reviews. Gastroenterology & hepatology***10**, 607-620 (2013).

2. Kufe, D.W. Mucins in cancer: function, prognosis and therapy. *Nature Reviews Cancer***9**, 874-885 (2009).
3. Hollingsworth, M.A. & Swanson, B.J. Mucins in cancer: protection and control of the cell surface. *Nature Reviews Cancer***4**, 45-60 (2004).
4. Aithal, A. et al. MUC16 as a novel target for cancer therapy. *Expert Opinion on Therapeutic Targets***22**, 675-686 (2018).
5. Chaturvedi, P., Singh, A.P. & Batra, S.K. Structure, evolution, and biology of the MUC4 mucin. *The FASEB Journal***22**, 966-981 (2008).
6. Moniaux, N. et al. Alternative splicing generates a family of putative secreted and membrane-associated MUC4 mucins. *European journal of biochemistry***267**, 4536-4544 (2000).
7. Moniaux, N. et al. Complete sequence of the human mucin MUC4: a putative cell membrane-associated mucin. *J Biochemical Journal***338**, 325-333 (1999).
8. Bafna, S. et al. MUC4, a Multifunctional Transmembrane Glycoprotein, Induces Oncogenic Transformation of NIH3T3 Mouse Fibroblast Cells. *Cancer Research***68**, 9231-9238 (2008).
9. Carraway, K.L., Theodoropoulos, G., Kozloski, G.A. & Carothers Carraway, C.A. Muc4/MUC4 functions and regulation in cancer. *Future Oncol***5**, 1631-1640 (2009).
10. Chakraborty, S., Jain, M., Sasson, A.R. & Batra, S.K. MUC4 as a diagnostic marker in cancer. *Expert Opinion on Medical Diagnostics***2**, 891-910 (2008).
11. Gautam, S.K. et al. MUC4 mucin-a therapeutic target for pancreatic ductal adenocarcinoma. *Expert opinion on therapeutic targets***21**, 657-669 (2017).
12. Gautam, S.K. et al. in *Seminars in Immunology* 101391 (Elsevier, 2020).
13. Singh, A.P., Chaturvedi, P. & Batra, S.K. Emerging Roles of MUC4 in Cancer: A Novel Target for Diagnosis and Therapy. *Cancer Research***67**, 433-436 (2007).
14. Kumar, S. et al. Genetic variants of mucins: unexplored conundrum. *Carcinogenesis***38**, 671-679 (2017).
15. Albrecht, H. & Carraway, K.L. MUC1 and MUC4: Switching the Emphasis from Large to Small. *Cancer Biotherapy & Radiopharmaceuticals***26**, 261-271 (2011).
16. Chaturvedi, P. et al. MUC4 Mucin Interacts with and Stabilizes the HER2 Oncoprotein in Human Pancreatic Cancer Cells. *Cancer research***68**, 2065-2070 (2008).
17. Jonckheere, N. et al. The Mucin MUC4 and Its Membrane Partner ErbB2 Regulate Biological Properties of Human CAPAN-2 Pancreatic Cancer Cells via Different Signalling Pathways. *PLoS ONE***7**, e32232

(2012).

18. Lakshmanan, I. et al. Novel HER3/MUC4 oncogenic signaling aggravates the tumorigenic phenotypes of pancreatic cancer cells. *Oncotarget***6**, 21085-21099 (2015).
19. Liberelle, M. et al. MUC4-ErbB2 Oncogenic Complex: Binding studies using Microscale Thermophoresis. *Scientific Reports***9**, 16678 (2019).
20. Ponnusamy, M.P. et al. MUC4 activates HER2 signalling and enhances the motility of human ovarian cancer cells. *British Journal of Cancer***99**, 520-526 (2008).
21. Moniaux, N. et al. Human MUC4 mucin induces ultra-structural changes and tumorigenicity in pancreatic cancer cells. *British journal of cancer***97**, 345-357 (2007).
22. Banerjee, K. et al. Amphiphilic polyanhydride-based recombinant MUC4 β -nanovaccine activates dendritic cells. *Genes & Cancer***10**, 52-62 (2019).
23. Liu, L. et al. Polyanhydride nanoparticles stabilize pancreatic cancer antigen MUC4 β . *Journal of Biomedical Materials Research Part A*, 1-10 (2020).
24. Aithal, A. et al. in *PANCREAS*, Vol. 48 1401-1402 (Lippincott Williams & Two Commerce SQ, 2001 Market St, Philadelphia., 2019).
25. C. Orzechowski, A.A., W. Junker, P. Kshirsagar, S. Batra, M. Jain. Generation and characterization of novel antibodies against the β subunit of MUC4. *Cancer Research* 78 (13 Supplement), 3829-3829. (
26. Boynton, Z.L. et al. Reduction of Cell Lysate Viscosity during Processing of Poly(3-Hydroxyalkanoates) by Chromosomal Integration of the Staphylococcal Nuclease Gene in *Pseudomonas putida*. *Applied and Environmental Microbiology***65**, 1524-1529 (1999).
27. Rodríguez Gamero, J.E. et al. Nuclease expression in efficient polyhydroxyalkanoates-producing bacteria could yield cost reduction during downstream processing. *Bioresource Technology***261**, 176-181 (2018).
28. Manavalan, P. & Johnson, W.C. Sensitivity of circular dichroism to protein tertiary structure class. *Nature***305**, 831-832 (1983).
29. Abdul-Gader, A., Miles, A.J. & Wallace, B.A. A reference dataset for the analyses of membrane protein secondary structures and transmembrane residues using circular dichroism spectroscopy. *Bioinformatics***27**, 1630-1636 (2011).
30. Whitmore, L. & Wallace, B.A. DICHROWEB, an online server for protein secondary structure analyses from circular dichroism spectroscopic data. *Nucleic Acids Research***32**, W668-W673 (2004).

31. Roy, A., Kucukural, A. & Zhang, Y. I-TASSER: a unified platform for automated protein structure and function prediction. *Nature Protocols***5**, 725-738 (2010).
32. Yang, J. et al. The I-TASSER Suite: protein structure and function prediction. *Nature Methods***12**, 7-8 (2015).
33. Yang, J. & Zhang, Y. I-TASSER server: new development for protein structure and function predictions. *Nucleic acids research***43**, W174-W181 (2015).
34. Senapati, S., Das, S. & Batra, S.K. Mucin-interacting proteins: from function to therapeutics. *Trends in biochemical sciences***35**, 236-245 (2010).
35. Bafna, S., Kaur, S. & Batra, S.K. Membrane-bound mucins: the mechanistic basis for alterations in the growth and survival of cancer cells. *Oncogene***29**, 2893-2904 (2010).
36. Bhatia, R. et al. Cancer-associated mucins: role in immune modulation and metastasis. *Cancer metastasis reviews***38**, 223-236 (2019).
37. Chaturvedi, P. et al. MUC4 Mucin Potentiates Pancreatic Tumor Cell Proliferation, Survival, and Invasive Properties and Interferes with Its Interaction to Extracellular Matrix Proteins. *Molecular Cancer Research***5**, 309 (2007).
38. Mukhopadhyay, P. et al. MUC4 Overexpression Augments Cell Migration and Metastasis through EGFR Family Proteins in Triple Negative Breast Cancer Cells. *PLoS ONE***8**, 1-14 (2013).
39. Jain, M. et al. Monoclonal Antibodies Recognizing the Non-Tandem Repeat Regions of the Human Mucin MUC4 in Pancreatic Cancer. *PLoS ONE***6**, e23344 (2011).
40. Tegel, H., Tourle, S., Ottosson, J. & Persson, A. Increased levels of recombinant human proteins with the Escherichia coli strain Rosetta(DE3). *Protein Expression and Purification***69**, 159-167 (2010).
41. Kim, S. & Lee, S.B. Rare codon clusters at 5'-end influence heterologous expression of archaeal gene in Escherichia coli. *Protein Expression and Purification***50**, 49-57 (2006).
42. Kleber-Janke, T. & Becker, W.-M. Use of Modified BL21(DE3) Escherichia coli Cells for High-Level Expression of Recombinant Peanut Allergens Affected by Poor Codon Usage. *Protein Expression and Purification***19**, 419-424 (2000).
43. Snyder, R.O. & Flotte, T.R. Production of clinical-grade recombinant adeno-associated virus vectors. *Current Opinion in Biotechnology***13**, 418-423 (2002).
44. Kalipatnapu, S. & Chattopadhyay, A. Membrane protein solubilization: recent advances and challenges in solubilization of serotonin1A receptors. *IUBMB Life***57**, 505-512 (2005).

45. Sreerama, N. & Woody, R.W. Estimation of protein secondary structure from circular dichroism spectra: comparison of CONTIN, SELCON, and CDSSTR methods with an expanded reference set. *Analytical biochemistry***287**, 252-260 (2000).
46. Micsonai, A. et al. Accurate secondary structure prediction and fold recognition for circular dichroism spectroscopy. *Proceedings of the National Academy of Sciences of the United States of America***112**, E3095-3103 (2015).
47. Sreerama, N. & Woody, R.W. Structural composition of betal- and betall-proteins. *Protein Sci***12**, 384-388 (2003).
48. Manavalan, P. & Johnson, W.C., Jr. Variable selection method improves the prediction of protein secondary structure from circular dichroism spectra. *Analytical biochemistry***167**, 76-85 (1987).
49. Lise, S. & Jones, D.T. Sequence patterns associated with disordered regions in proteins. *Proteins***58**, 144-150 (2005).
50. Uversky, V.N. What does it mean to be natively unfolded? *European journal of biochemistry***269**, 2-12 (2002).
51. Carmicheal, J. et al. Presence and structure-activity relationship of intrinsically disordered regions across mucins. *The FASEB Journal***34**, 1939-1957 (2020).
52. Bhatia, R. et al. MUC4 interacts and stabilize EGFR1 in a ligand-dependent manner leading to sustained oncogenic signaling. *The FASEB Journal***33**, 631.633-631.633 (2019).
53. Kloczkowski, A., Ting, K.-L., Jernigan, R.L. & Garnier, J. Combining the GOR V algorithm with evolutionary information for protein secondary structure prediction from amino acid sequence. **49**, 154-166 (2002).
54. Wu, S. & Zhang, Y. LOMETS: A local meta-threading-server for protein structure prediction. *Nucleic Acids Research***35**, 3375-3382 (2007).
55. Senapati, S. et al. Novel INTERaction of MUC4 and galectin: potential pathobiological implications for metastasis in lethal pancreatic cancer. *J Clinical cancer research***17**, 267-274 (2011).
56. Arpin, M., Chirivino, D., Naba, A. & Zwaenepoel, I. Emerging role for ERM proteins in cell adhesion and migration. *Cell Adhesion & Migration***5**, 199-206 (2011).
57. Clucas, J. & Valderrama, F. ERM proteins in cancer progression. *Journal of Cell Science***127**, 267 (2014).
58. Meng, Y. et al. Ezrin promotes invasion and metastasis of pancreatic cancer cells. *Journal of Translational Medicine***8**, 61 (2010).

59. Blalock, T.D. et al. Functions of MUC16 in corneal epithelial cells. *Investigative ophthalmology visual science***48**, 4509-4518 (2007).
60. Koopaei, N.N. et al. Optimization of rPDT fusion protein expression by Escherichia coli in pilot scale fermentation: a statistical experimental design approach. *AMB Express***8**, 135 (2018).
61. Aithal, A. et al. Development and characterization of carboxy-terminus specific monoclonal antibodies for understanding MUC16 cleavage in human ovarian cancer. *PLoS ONE***13**, e0193907 (2018).
62. Ye, G. et al. Clearance and characterization of residual HSV DNA in recombinant adeno-associated virus produced by an HSV complementation system. *Gene therapy***18**, 135 (2011).

Tables

Due to technical limitations, table 1 is only available as a download in the Supplemental Files section.

Figures

Figure 1

Plasmid cloning strategy, expression, and purification of the human rMUC4 β . (A) Schematic representation of the complete structure of MUC4 protein and construction of pET-28a-MUC4 β plasmid: MUC4 β sequence (from GDPH to C-terminal) was amplified with primers incorporated with 'NdeI' and 'XhoI' restriction sites and cloned in-frame with the His6x tag and thrombin site (TS) encoded by pET-28a(+) vector. (B) Expression profile of total cellular proteins from E. coli C41(DE3) strain: The pET-28a

(empty vector) and pET-28a-MUC4 β transformed C41(DE3) cells were induced with an indicated amount of IPTG. SDS-PAGE and immunoblotting analyses were performed on the uninduced (U) and IPTG-induced (I) culture supernatant. (C) Comparative assessment of rMUC4 β expression profile in C41(DE3) and RosettaTM 2(DE3) competent cells at different IPTG concentrations using SDS-PAGE and an anti-His tag immunoblotting. (D) Effect of different post-induction incubation temperatures on rMUC4 β expression in RosettaTM 2(DE3): Total protein expression from each temperature condition was assessed by SDS-PAGE and immunoblotting. (D) Isolation and ÄKTA-FPLC affinity purification of rMUC4 β . The quality of pre-and-post-ÄKTA fractions was assessed by Coomassie staining and immunoblot analyses.

Figure 2

Biophysical characterization and secondary structure analysis of MUC4β. (A) Far-UV CD spectra of rMUC4β. (B) Comparison of the percentage of secondary structure content estimated or predicted from CD spectrum, amino acid sequence, and I-TASSER analyses. The 2D multicolor pie chart showing the structure compositions estimated using (C) CDSSTR and (D) Contin-LL (Provencher and Glockner) methods. (E) Predicted structure of rMUC4β generated with the I-TASSER tool. The cartoon structure of

MUC4 β indicating N-terminus (blue) and C-terminus (red). Right panel: I-TASSER predicted structure of MUC4 β rotated 180° to visualize the three EGF-like domains. EGF-like domain I (Cyan) and EGF-like domain II (Magenta) appear to be typical β -strand structures, while EGF-like domain III (Red) is a loop.

Figure 3

Affinity pull-down and tandem mass spectrometry to determine the interacting partners of MUC4. (A) Schematic outline of in vitro affinity pull-down assays (I) pre-clearing of total lysate [input] of MUC4

expressing (CD18/HPAF) and non-expressing (PANC-1) pancreatic cancer cell lines. (II) Affinity pull-down reactions were performed in the presence [rMUC4β (+)] and absence [rMUC4β (-), i.e., RIPA buffer] rMUC4β (bait) added with washed Ni-NTA beads. Respective supernatants [i.e., Flow-through (FT) and last wash (W5)] were collected before the final elution. (III) An aliquot of MUC4β-bound Ni-resin was eluted separately to serve as a positive control. All saved fractions (as shown in pink color, italic font) from CD18/HPAF (B) and PANC1 (C) were resolved on SDS-PAGE gel and stained with Coomassie Blue. The ~80 kDa band (arrowed) was identified, excised, and assessed by the tandem MS/MS and proteomic analyses.

Figure 4

Detection and confirmation of the new and known interacting partners of MUC4 via immunoblotting and IP analyses. (A) Expression profile of MUC4 in CD18/HPAF and PANC-1 cells determined by immunoblotting following 2% agarose gel electrophoresis and 10% SDS-PAGE. (B) Immunoblot analyses of Ni-Ni-NTA pull-down fractions from CD18/HPAF or PANC-1 cell lysates in the presence (+) or absence (-) of rMUC4β. The blots were probed with the indicated antibodies. Input, flow-through (FT), wash, and eluate are the same as described in Fig. 3A. (C) Immunoprecipitation using an anti-His tag Ab (clone 27E8) conjugated to magnetic beads following incubation of CD18/HPAF cell lysate in the presence and

absence of rMUC4 β . (D) Interaction of endogenous MUC4 with Ezrin. Immunoprecipitation was performed with protein A/G agarose beads on CD18/HPAF lysates following incubation with anti-MUC4 β Ab.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryInformation.docx](#)
- [Graphicalabstract.tif](#)
- [Table1.pdf](#)