Secular trends in cryoglobulinemia mortality in the United States in the era of direct-acting antivirals

Qianyu Guo

Shanxi Medical University

Jinfang Gao

Shanxi Medical University

Jiaoni Duan

Shanxi Medical University

Ruihong Hou

Shanxi Medical University

Liyun Zhang

Shanxi Medical University

Tsung-Hsueh Lu (✉ robertlu@mail.ncku.edu.tw)

National Cheng Kung University College of Medicine <https://orcid.org/0000-0003-4715-3787>

Research Article

Keywords: Cryoglobulinemia, Direct-acting antivirals, Hepatitis C virus, Mortality trends

Posted Date: September 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-814408/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

Hepatitis C virus (HCV) is the main etiology of cryoglobulinemia with mortality around 25%. Little is known on the changes in cryoglobulinemia mortality after the introduction of direct-acting antivirals (DAA) for treatment of HCV in 2014 in the US.

Methods

We calculated cryoglobulinemia mortality from 1999 to 2018. The proportionate mortality ratio (PMR) of cryoglobulinemia cases with HCV and those with autoimmune diseases were computed to assess the impact of introduction of DAA.

Results

We identified 1299 people aged ≥ 20 years who died with cryoglobulinemia between 1999 and 2018. The cryoglobulinemia mortality (deaths per million) declined from 1999 (0.4) to 2010 (0.22) and mildly increased from 2010 to 2014 (0.26), and then decreased abruptly from 2014 to 2018 (0.19). The proportion of cryoglobulinemia patients with HCV was 39% (118/302) in 2009–2013 and 26% (81/310) in 2014–2018, with a PMR of 0.67 (95% CI 0.50–0.89). By contrast, the proportion of cryoglobulinemia patients with systemic autoimmune diseases was 2.6% (8/302) in 2009–2013 and 4.2% (13/310) in 2014–2018, with a PMR of 1.58 (95% CI 0.66–3.82).

Conclusion

The changes in cryoglobulinemia mortality during the past two decades are mainly related to the aging and dying of the “baby boomer” cohort who had a high HCV prevalence and to the introduction of a DAA in 2014.

Introduction

Cryoglobulins are immunoglobulins that precipitate in vitro at temperatures below 37°C, and cryoglobulinemia is defined as the persistent presence of cryoglobulins in serum. Hepatitis C virus (HCV) is the main etiology accounting for 80–90% of cryoglobulinemia cases.¹ Although a review of a cryoglobulinemia case series conducted in 2008 indicated heterogeneous mortality results from 16–66%.² Later prognosis studies of patients with cryoglobulinemia and HCV suggested more consistent mortality results (ranging from 21–29%) relating to life-threatening organ damage involving renal failure, pulmonary hemorrhage, gastrointestinal ischemia, and cerebral ischemia or hemorrhage.^{3–7}

Several novel direct-acting antivirals (DAAs) for treatment of HCV were approved by the US Food and Drug Administration in late 2013, and many clinical trials have indicated a good response from patients with HCV and cryoglobulinemia.^{8–10} However, changes in the epidemiology of cryoglobulinemia before and after the introduction of DAAs have not been extensively examined in the United States. Therefore, this study was conducted to examine secular trends in cryoglobulinemia mortality within the United States from 1999 to 2018, with a particular focus on changes occurring after the introduction of DAAs in 2014.

Methods

Data sources and case definition

We used the multiple-cause mortality files compiled by the National Center for Health Statistics (NCHS) to identify decedents aged 20 years and above who died between 1999 and 2018 and whose death certificates mentioned cryoglobulinemia.¹¹ The multiple-cause mortality files are available in two data formats: entity axis and record axis codes.¹¹ We used record axis codes in this study because they are edited for NCHS core multi-cause tabulations.

The International Classification of Diseases, Tenth Revision (ICD-10) was adopted by the United States of America in 1999, and the ICD-10 codes associated with relevant diseases are as follows: D891 for cryoglobulinemia, B171 or B182 for HCV, M30–M36 for systemic autoimmune disease, M32 for systemic lupus erythematosus (SLE), and M350 for Sjogren syndrome.

Mortality rates

The cryoglobulinemia mortality rate was calculated by counting cases in which cryoglobulinemia was recorded either as the underlying or contributory cause of death on the death certificate as the numerator and general population as denominator. We calculated the age-standardized mortality rate using the age structure of the United States for the year 2000 as the standard population. The proportionate mortality rates (%) related to cryoglobulinemia with HCV and cryoglobulinemia with autoimmune diseases were also computed.

Statistical analysis

A joinpoint regression analysis was used to assess yearly changes in the secular trends of cryoglobulinemia mortality rates.¹² The annual percentage changes (APCs) were computed for each segment of the trends, and the trends in the statistical tests were examined if they differed from zero. To assess whether there was a significant change in the proportionate mortality rate following introduction of the DAA agents, we calculated the 2014–2018/2009–2013 proportionate mortality rate ratio (PMR) and the 95% confidence intervals (95% CIs).

Results

We identified 1299 people aged ≥ 20 years who died between 1999 and 2018 and whose death certificates mentioned cryoglobulinemia. The number of deaths decreased from 80 in 1999 to 58 in 2018, and the age-adjusted cryoglobulinemia mortality rate (deaths per million) declined from 0.40 in 1999 to 0.19 in 2018. Three mortality trend segments for were identified for age-adjusted mortality rate: the first was related to the years 1999 to 2010 (APC: -5.6%), the second to the years 2010 to 2014 (APC: 3.8%), and the third to the years 2014 to 2018 (APC: -8.9% ; Fig. 1A). The mortality rate for different age groups are illustrated in Fig. 1B, C, and D, respectively.

With regard to the demographic characteristics of decedents with cryoglobulinemia, a large decrease in the proportion of decedents aged 20–54 years was noted, from 30% (114/382) in 1999–2003 to 9% (27/310) in 2014–2018, whereas the proportion of decedents aged 65 years and above increased from 54% (207/382) in 1999–2003 to 60% (187/310) in 2014–2018 (Table 1).

Table 1
Number of deaths and mortality rates (deaths per million) with cryoglobulinemia in the United States, 1999–2018

	1999–2003			2004–2008			2009–2013			2014–2018			2014–2018/2009–2013	
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	Rate ratio	(95% CI)
Total*	382	(100)	0.37	305	(100)	0.28	302	(100)	0.24	310	(100)	0.21	0.91	(0.76–1.05)
Sex														
Men	168	(44)	0.34	136	(45)	0.26	149	(49)	0.27	141	(45)	0.24	0.89	(0.68–1.10)
Women	214	(56)	0.41	169	(55)	0.30	153	(51)	0.26	169	(55)	0.27	1.05	(0.81–1.27)
Age (years)														
20–54	114	(30)	0.16	94	(31)	0.13	48	(16)	0.06	27	(9)	0.04	0.56	(0.29–0.82)
55–64	61	(16)	0.48	69	(23)	0.43	114	(38)	0.61	96	(31)	0.46	0.77	(0.55–0.97)
65+	207	(54)	1.17	142	(46)	0.76	140	(46)	0.67	187	(60)	0.76	1.13	(0.88–1.38)
*Rate of total was age-adjusted using age structure of US population of 2000 as reference														
95% CI = 95% confidence interval														

The cryoglobulinemia mortality rates by sex and age in the four study periods are presented in Table 1. Two prominent reductions in mortality rates occurred in decedents aged 20–54 years: the first from 0.13 in 2004–2008 to 0.06 in 2009–2013 (50% decrease) and the second from 0.06 in 2009–2013 to 0.04 in 2014–2018 (44% decrease). By contrast, for decedents aged 55–64 years, an increase in mortality of 0.43 in 2004–2008 to 0.61 in 2009–2013 (40% increase) was noted, followed by a decrease in mortality from 0.61 in 2009–2013 to 0.46 in 2014–

2018 (24% decrease). A minor increase in the mortality of those aged 65 years and over was noted: 0.67 in 2009–2013 to 0.76 in 2014–2018 (13% increase).

The proportion of decedents with cryoglobulinemia and HCV was 39% (118/302) in 2009–2013, but this decreased to 26% (81/310) in 2014–2018 with a PMR of 0.67 (95% CI 0.50–0.89) (Table 2). The PMR for each age group was lower than one, but it was not statistically significant. The proportion of decedents with cryoglobulinemia and systemic autoimmune diseases was 2.6% (8/302) in 2009–2013, but this increased to 4.2% (13/310) in 2014–2018, with a PMR of 1.58 (95% CI 0.66–3.82). SLE (n = 15) and Sjogren syndrome (n = 20) were the two main systemic autoimmune diseases associated with cryoglobulinemia.

Table 2
Numbers of people who died with mention of cryoglobulinemia and hepatitis C virus and those died with mention of cryoglobulinemia and systemic autoimmune disease in the United States, 1999–2018

	1999–2003		2004–2008		2009–2013		2014–2018		2014–2018/2009–2013	
	No.	%	No.	%	No.	%	No.	%	PMR	95% CI
Cryoglobulinemia with hepatitis C virus										
Total	125	32.7	99	32.5	118	39.1	81	26.1	0.67	(0.50–0.89)
Sex										
Men	62	36.9	60	44.1	67	45.0	50	35.5	0.79	(0.55–1.14)
Women	63	29.4	39	23.1	51	33.3	31	18.3	0.55	(0.35–0.86)
Age (years)										
20–54	74	64.9	55	58.5	23	47.9	11	40.7	0.85	(0.41–1.74)
55–64	19	31.1	28	40.6	70	61.4	41	42.7	0.70	(0.47–1.02)
65+	32	15.5	16	11.3	25	16.7	29	15.5	0.93	(0.55–1.59)
Cryoglobulinemia with systemic autoimmune disease (ICD-10 codes)										
Autoimmune disease (M30-M36)	16	4.2	13	4.3	8	2.6	13	4.2	1.58	(0.66–3.82)
Systemic lupus erythematosus (M32)	6	1.6	3	1.0	2	0.7	4	1.3	1.95	(0.36–10.64)
Sjogren's syndrome (M350)	6	1.6	5	1.6	3	1.0	6	1.9	1.95	(0.49–7.79)
Other autoimmune disease	5	1.3	6	2.0	3	1.0	5	1.6	1.62	(0.39–6.79)
Sex										
Men	4	2.4	1	0.7	4	2.7	2	1.4	0.53	(0.10–2.88)
Women	12	5.6	12	7.1	4	2.6	11	6.5	2.49	(0.79–7.82)
Age (years)										
20–54	4	3.5	4	4.3	2	4.2	3	11.1	2.67	(0.45–15.96)
55–64	5	8.2	3	4.3	4	3.5	4	4.2	1.19	(0.30–4.75)
65+	7	3.4	6	4.2	2	1.3	6	3.2	2.41	(0.49–11.92)
PMR = proportionate mortality ratio; 95% CI = 95% confidence interval;										
ICD-10 codes = <i>International Classification of Diseases, Tenth Revision</i>										

Discussion

The findings of this national population-based study indicate that a persistent decline in the cryoglobulinemia mortality rates from 1999 to 2010, followed by a mild increase from 2010 to 2014, and then an abrupt decline from 2014 to 2018. The age-specific cryoglobulinemia mortality trend patterns correspond with the age-specific HCV mortality trend patterns.¹³ The age differences in mortality trend patterns reflect the aging and dying of “baby boomers” (born between 1945 and 1965 in the United States) with a high prevalence of HCV.¹⁴ The increased

cryoglobulinemia mortality among middle-age adults aged 55–64-years from the early 2000s and older adults aged 65-years and above from the late 2000s can be mainly attributed to the aging of “baby boomers.”

However, the abrupt decline in cryoglobulinemia mortality since 2014 among decedents aged 55–64 years and 65 years and above can be mainly attributed to the introduction of DAAs in 2014. The proportion of patients with cryoglobulinemia and HCV reported in a French study was 63% in 2011, 60% in 2015, and 33% in 2018.¹⁰ In this study, the proportion of decedents was 37% in 2011, 30% in 2015, and 26% in 2018.

With regard to the autoimmune diseases, the proportion of decedents with cryoglobulinemia who had systemic autoimmune diseases increased from 17% in 2011 to 36% in 2018 in the French study.¹⁰ Similar increases were observed in this study: the proportion was 2.6% in 2009–2013 and increased to 4.2% in 2014–2018.

One strength of the present study is that it used national population-based mortality data to examine the long-term cryoglobulinemia mortality trends and changes in association with HCV. To our knowledge, this study included a greater number of cryoglobulinemia cases and a larger observational year span than previous studies. However, several limitations should be noted when interpreting the findings of this study. First, some physicians may underreport either cryoglobulinemia, HCV, or systemic autoimmune diseases on death certificates. The comparison between proportions represented in this study and those in the French study suggest that the extent of underreporting may differ depending on the disease reported. For example, a study of death certificates in Connecticut revealed that among deaths for which hepatitis B, hepatitis C, and alcoholic liver disease were identified during the medical record review, the specified ICD codes were only recorded on 8.6%, 45.4%, and 36.5%, of the death certificates, respectively.¹⁵ However, our primary aim was to examine mortality trends, and it is unlikely that there would have been substantial changes in the underreporting behaviors of certifiers during the study period.

Second, it was not possible to further analyze mortality trends by cryoglobulinemia type, because the ICD-10 code for cryoglobulinemia does not further classify types. Third, as no information on clinical laboratory data were available from mortality data, it was not possible to stratify the decedents into different clinical profiles. Fourth, because only a small number of deaths had occurred, we did not further analyze data by race and place of residence. Fifth, information relating to whether the patients received DAA treatment was not available. These limitations show that further studies linking health care services data with mortality data are required.

Conclusion

In conclusion, a decline in cryoglobulinemia mortality occurred in the United States that began in the early 2000s, levelled off between 2010 and 2014, and then drastically declined between 2014 and 2018. The changes in mortality trends patterns are considered to be mainly attributed to the aging and dying of a “baby boomer” cohort who had a high prevalence of HCV and to the introduction of DAAs in 2014.

Abbreviations

APC, annual percent change; ASDR, age-standardized death rate; DAA, direct-acting antivirals; HCV, hepatitis C virus; ICD-10, International Classification of Diseases Tenth Revision; MCOD, multiple cause of death; NCHS, National Center for Health Statistics; PMR, proportionate mortality ratio; SLE, systemic lupus erythematosus; 95% CI, 95% confidence intervals.

Declarations

Acknowledgment

None.

Authors' contributions

QG, JG, and THL conceived and designed the study. All authors contributed to the acquisition of data. Analysis and interpretation of data were performed by QG, JG, JD, RH, and LZ. The first draft of the manuscript was written by QG and all authors critically revised it for important intellectual content. THL supervised the all investigation. All authors read and approved the final manuscript and agreed to be held accountable for all aspects of the work.

Funding

There was no funding support this study.

Availability of data and materials

The data are available upon the request to the corresponding author.

Ethics approval and consent to participate

This study was approved by the Institute Review Board of National Cheng Kung University Hospital. No participants were involved in this study.

Consent for publication

All authors have approved the final version of the manuscript and consented for publication.

Competing interests

The authors declare that they have no competing interests.

Authors details

¹Department of Rheumatology, Third Hospital of Shanxi Medical University, Shanxi Bethune Hospital, Shanxi Academy of Medical Sciences, Taiyuan, Shanxi, China

²Department of Public Health, National Cheng Kung University, Tainan, Taiwan.

References

1. Fuentes A, Mardones C, Burgos PI. Understanding the cryoglobulinemias. *Curr Rheumatol Rep*. 2019;21(11):60.
2. Della Rossa A, Marchi F, Catarsi E, Tavoni A, Bombardieri S. Mixed cryoglobulinemia and mortality: a review of the literature. *Clin Exp Rheumatol*. 2008;26(5 Suppl 51):105–8.
3. Landau DA, Scerra S, Sene D, Resche-Rigon M, Saadoun D, Cacoub P. Causes and predictive factors of mortality in a cohort of patients with hepatitis C virus-related cryoglobulinemic vasculitis treated with antiviral therapy. *J Rheumatol*. 2010;37(3):615–21.
4. Rossa AD, Tavoni A, D'Ascanio A, et al. Mortality rate and outcome factors in mixed cryoglobulinaemia: the impact of hepatitis C virus. *Scand J Rheumatol*. 2010;39(2):167–70.
5. Terrier B, Semoun O, Saadoun D, Sène D, Resche-Rigon M, Cacoub P. Prognostic factors in patients with hepatitis C virus infection and systemic vasculitis. *Arthritis Rheum*. 2011;63(6):1748–57.
6. Retamozo S, Díaz-Lagares C, Bosch X, et al. Life-threatening cryoglobulinemic patients with hepatitis C: clinical description and outcome of 279 patients. *Medicine*. 2013;92(5):273–84.
7. Lauletta G, Russi S, Conteduca V, Sansonno L, Dammacco F, Sansonno D. Impact of cryoglobulinemic syndrome on the outcome of chronic hepatitis C virus infection: a 15-year prospective study. *Medicine*. 2013;92(5):245–56.
8. Saadoun D, Pol S, Ferfar Y, et al. Efficacy and safety of sofosbuvir plus daclatasvir for treatment of HCV-associated cryoglobulinemia vasculitis. *Gastroenterology*. 2017;153(1):49–52.e5.
9. Cacoub P, Ahmed SNS, Ferfar Y, et al. Long-term efficacy of interferon-free antiviral treatment regimens in patients with hepatitis C virus-associated cryoglobulinemia vasculitis. *Clin Gastroenterol Hepatol*. 2019;17(3):518–26.
10. Boleto G, Ghillani-Dalbin P, Musset L, et al. Cryoglobulinemia after the era of chronic hepatitis C infection. *Semin Arthritis Rheum*. 2020;50(4):695–700.
11. National Center for Health Statistics. Public Use Data File Documentation: Mortality Multiple Cause-of-Death. https://www.cdc.gov/nchs/nvss/mortality_public_use_data.html Accessed November 4, 2019.
12. The National Cancer Institute. The Joinpoint Trend Analysis Software. <https://surveillance.cancer.gov/joinpoint/> Accessed November 4, 2019.
13. Sheu MJ, Liang FW, Lu TH. Hepatitis C virus infection mortality trends according to three definitions with special concern for the baby boomer birth cohort. *J Viral Hepat*. 2021;28(2):317–25.
14. Armstrong GL, Wasley A, Simard EP, et al. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann Intern Med*. 2006;144(10):705–14.
15. Ly KN, Speers S, Kleven RM, Barry V, Vogt TM. Measuring chronic liver disease mortality using an expanded cause of death definition and medical records in Connecticut, 2004. *Hepatol Res*. 2015;45(9):960–68.

Figures

Figure 1

Cryoglobulinemia mortality rates in the United States, 1999–2018. (Modelled rates were estimated according to joinpoint regression analysis.)