The Importance of Calibration Method in Determining the Association between Central Blood Pressure with Left Ventricular and Left Atrial Strain.

Satish Ramkumar (✉ satishramkumar2@gmail.com)

Baker Heart and Diabetes Institute

Hong Yang

University of Tasmania Menzies Research Institute: University of Tasmania Menzies Institute for Medical Research

Mark Nolan

Baker Heart Research Institute: Baker Heart and Diabetes Institute

Tomoko Negishi

University of Tasmania Menzies Research Institute: University of Tasmania Menzies Institute for Medical Research

James E Sharman

University of Tasmania Menzies Research Institute: University of Tasmania Menzies Institute for Medical Research

Thomas Hugh Marwick

Baker Heart Research Institute: Baker Heart and Diabetes Institute

Kazuaki Negishi

University of Sydney

Research Article

Keywords: strain, left atrium, atrial function, central blood pressure

Posted Date: August 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-815454/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at The International Journal of Cardiovascular Imaging on October 21st, 2021. See the published version at <https://doi.org/10.1007/s10554-021-02444-4>.

Abstract

Purpose

Afterload is an important determinant of left ventricular(LV) and atrial(LA) function, including myocardial strain. Central blood pressure(CBP) is the major component of cardiac afterload and independently associated with cardiovascular risk. However, the optimal means of calibrating CBP is unclear - standard CBP assessment uses systolic(SBP) and diastolic blood pressure(DBP) from brachial waveforms, but calibration with mean pressure(MAP) and DBP purports to be more accurate. Therefore, we sought to determine which CBP is best associated with LA and LV strain.

Methods

CBP was measured using both standard and MAP based calibration methods in 546 participants(age 70.7 ± 4.7 years, 45% male) with risk factors for heart failure. Echocardiography was performed in all patients and strain analysis conducted to assess LA/LV function. The associations of CBP with LA and LV strain were assessed using linear regression.

Results

MAP-derived CSBP(150 ± 20 mmHg) was higher than standard CSBP(128 ± 15 mmHg) and brachial SBP(140 ± 17 mmHg, $p < 0.001$), whereas DBPs were similar(84 ± 10 , 83 ± 10 , and 82 ± 10 mmHg). MAP-derived CSBP was not independently associated with LV strain($p > 0.05$), however was independently associated with LA reservoir strain($p < 0.05$). Brachial and central DBP were more strongly associated with LA reservoir/conduit and LV strain than brachial and central SBP. LA pump strain was not independently associated with any SBP or DBP parameter($p > 0.05$). MAP-derived CBP was more accurate in identifying patients with abnormal LA and LV strain than brachial SBP and standard CBP calibration.

Conclusion

CBP calibrated using MAP and DBP may be more accurate in identifying patients with abnormal LA and LV function than standard brachial calibration methods.

Introduction

Hypertension is a leading cause of morbidity and mortality[1, 2]. Recent changes in clinical practice guidelines have advocated for an aggressive approach in the diagnosis and treatment of hypertension[1]. However, there remains contention regarding the use of the most appropriate blood pressure (BP) marker[3], which is easily measurable, reproducible, and correlates with clinical outcomes. Some studies demonstrated that central blood pressure (CBP) is more closely associated with cardiovascular outcomes compared with peripheral brachial BP[4-7], although some did not[8]. There are multiple non-invasive methods of estimating central (aortic) pressure[9]. Most commonly, a waveform calibration method is to use brachial systolic blood pressure (SBP) and diastolic blood pressure (DBP). This method may be

associated with inaccuracy leading to underestimation of the systolic CBP and overestimation of diastolic CBP[10]. The use of mean arterial pressure (MAP) along with DBP may be a more accurate calibration method of CBP[9]. We previously reported better associations between MAP-derived CBP and cardiac anatomy as a marker of end-organ damage[11] but little is known about relationships with cardiac function. Identifying the BP parameter most closely associated with altered cardiac function is important because it may indicate those with early signs of end-organ damage related to hypertension.

Strain analysis using speckle tracking allows for quantitative assessment of left atrial (LA) and left ventricular (LV) function. Hypertension has a direct impact on both LV and LA function – the latter through LV stiffness contributing to increased LA pressure leading to LA dilatation and fibrosis[12]. In this study, we aimed to elucidate the associations between the BPs and myocardial deformation parameters with the strongest correlation with target organ functional alteration.

Methods

Study population. This is a cross-sectional study using baseline data from a large community-based study of stage A heart failure (HF), with the primary objective of early detection of left ventricular dysfunction using strain imaging (Tas-ELF study, ACTRN12614000080628). Asymptomatic participants ≥ 65 years were recruited if they had 1 or more risk factors, including hypertension (SBP ≥ 140 mmHg, DBP ≥ 90 mmHg or pre-existing use of anti-hypertensive medications), type 2 diabetes mellitus (T2DM, based on self-report of diagnosis or the current use of diabetic medications), or obesity (defined as a body mass index ≥ 30 kg/m²). Exclusion criteria included: (1) inability to provide written consent to participate in the study, (2) history of moderate or greater valvular disease, (3) known history of HF, (4) reduced LV systolic function on baseline echocardiography (LVEF $<40\%$), (5) contraindications to beta blockers or angiotensin converting enzyme inhibitors (ACE-I), (6) expected life expectancy of less than one year or (7) inability to perform strain analysis or acquire interpretable images from baseline echocardiogram. All patients with a known history of AF or documented AF on the baseline ECG were excluded from the study. All patients were provided written informed consent and approval was obtained from the institution's Human Research Ethics Committee (University of Tasmania HREC project number H0013333)

Baseline data collection. All participants undertook a clinical history and answered questionnaires to assess overall health status at the start of the study. Information regarding demographics, past medical history, medication history as well as baseline examination data (height, weight, body mass index (BMI)) was recorded for all participants. Baseline electrocardiography (ECG) and echocardiography was conducted in all participants.

Blood Pressure measurement. Peripheral BP (3x measurements) was measured in the supine position following a minimum of 10 minutes in a quiet room without auditory stimuli. A validated oscillometric device (Mobil-O-Graph, IEM, Stolberg, Germany) was used for all measurements. CBP was measured using two calibration methods. Standard CBP was measured from calibration of the brachial BP

waveforms using oscillometric SBP and DBP. Using an automated batch method, each brachial BP waveform was then recalibrated to derive MAP based CBP using oscillometric MAP and DBP measurements. Cut-offs for hypertension using CBP and brachial BP was determined as BP $\geq$ 140/80 mmHg.

Echocardiography. All echocardiograms were performed by qualified sonographers who were blinded to clinical information using the same equipment (Siemens ACUSON SC2000, Siemens Healthcare, Mountain View, CA) and transducers (4V1c, 1.25-4.5 MHz; 4Z1c, 1.5-3.5 MHz). Two-dimensional, M-mode, and Doppler measures were obtained using standard techniques outlined by the American Society of Echocardiography. LV dimensions were calculated in both diastole and systole in parasternal long axis views. LV hypertrophy (LVH) was defined as LVM index >115 g/m² in men and >95 g/m² in women. LV and LA volumes were indexed to body surface area (LAVi) and calculated by the biplane method of discs. Abnormal LAVi was defined as ≥ 34 ml/m².

Global longitudinal strain (GLS) was calculated in apical views using speckle tracking imaging. Manual tracing of the endocardial border of the LV was performed in end-systole and this was tracked during the cardiac cycle. Abnormal GLS was defined as $>-16\%$ [13]. LA reservoir, conduit and pump strain were assessed using speckle tracking imaging by an external third-party software program (ImageArena, Tomtec, Munich, Germany). Apical four and two chamber images were selected with a frame rate of 60-80 frames/sec. The endocardial border of the LA was manually traced and strain analysis was performed using the LV strain algorithm, utilizing the average of both the four- and two-chamber values. The reference point for image analysis was taken at the onset of the QRS complex (R-R gating). Abnormal LA strain was defined as LA reservoir strain $<38\%$, LA conduit $< 21\%$ and LA pump strain $<16\%$. Patients with poor image quality, where strain analysis could not be performed, were excluded. All strain measurements were performed by two investigators. Reproducibility was assessed using a random sample of 20 patients and mean percentage difference was calculated.

Statistical analysis. All categorical variables are presented as frequencies/percentages and continuous variables presented as means/standard deviation (if normally distributed) or medians/IQR (if non-parametric). Clinical and echocardiographic characteristics were compared in participants with normal/abnormal GLS. Baseline characteristics were compared using the chi-square test for categorical data, and Student t-test for continuous data or the Mann-Whitney U test as appropriate. The correlation between standard and MAP-derived CBP was assessed using scatterplots and Pearson's r. Scatterplots were also used to compare both standard and MAP-derived CBP based on the presence of reduced GLS/LA strain. Multivariable linear regression analysis was used to assess the independent association between BP parameters and strains by adjusting BMI, T2DM, family history of HF, use of angiotensin-converting enzyme inhibitors/angiotensin receptor blockers, beta blockers, calcium channel blockers and diuretics. Correlation co-efficients between standard and MAP-derived CBP were compared using Steiger's Z test. Analyses were considered statistically significant if 2-tailed p-values were <0.05 . Statistical analysis was performed using SPSS v.22 (SPSS, Chicago, IL), R version 3.5.0 [<https://www.r-project.org/>] and Stata v.13 (StataCorp, College Station, Texas).

Results

Baseline characteristics. A summary of baseline patient characteristics is shown in **Table 1**. A total of 546 patients were included in the study (age 70.7 ± 4.7 years, male 45%). Most patients had T2DM (52%), obesity (43%), hypercholesterolemia (54%) and hypertension (79%). The mean GLS, LA reservoir, conduit, and pump strain of the overall cohort was $-18.6 \pm 2.5\%$, $39.3 \pm 6.8\%$, $19.9 \pm 5.4\%$, and $19.4 \pm 5.0\%$, respectively.

87 participants (16%) of the cohort had abnormal GLS (GLS $-14.6 \pm 1.3\%$ vs. $-19.3 \pm 1.9\%$ in those with normal GLS, $p < 0.001$). Those with abnormal GLS had higher BMI, were more likely to have a diagnosis of diabetes mellitus and had reduced exercise capacity based on the six-minute walk test ($p < 0.05$). SBPs were not significantly different in those with normal and abnormal GLS ($p > 0.05$), however, those with abnormal GLS had higher DBP compared with those with normal GLS (85.2 ± 10.3 vs. 81.1 ± 10.2 mmHg for brachial DBP, 86.7 ± 10.4 vs. 82.1 ± 9.9 mmHg for standard central DBP and 87.7 ± 10.9 vs. 83.0 ± 10.0 mmHg for MAP-derived central DBP, $p \leq 0.001$). LA strains were lower in those with abnormal GLS ($36.7\% \pm 7.9\%$ vs. $39.7\% \pm 6.5\%$, $p = 0.001$ for LA reservoir strain; and $18.1\% \pm 5.9\%$ vs. $20.2\% \pm 5.2\%$, $p = 0.002$ for LA conduit strain), except for LA pump strain, which was similar among the groups ($p > 0.05$).

Correlation between different CBP calibration methods. **Figure 1** illustrates a scatterplot showing the correlation among the three BP methods. MAP-derived central SBP was higher than standard systolic CBP and brachial SBP (150 ± 20 mmHg vs. 128 ± 15 mmHg vs. 140 ± 17 mmHg, both $p < 0.001$). Whereas MAP-derived central DBP was similar to standard central DBP and brachial DBP (84 ± 10 mmHg vs. 83 ± 10 mmHg vs. 82 ± 10 mmHg, $p > 0.05$). There was a modest correlation between MAP-derived central SBP and standard central SBP (Pearson's $r = 0.74$, $p < 0.001$). A stronger correlation was observed between standard central SBP with brachial SBP than that between MAP-derived central SBP and brachial SBP ($p = 0.018$). DBP, MAP-derived central DBP and standard central DBP are very closely correlated to each other (Pearson's r between 0.93-0.98, all $p < 0.001$).

Association between BP and LA/LV strain. **Table 2** summarizes the associations between BPs and strains. DBP (both brachial and central) were independently associated with GLS, LA reservoir, and LA conduit strain ($p < 0.05$) but not with LA pump strain. Brachial and standard central SBPs were independently associated with GLS ($\beta = -0.09$, $p = 0.04$ for brachial SBP and $\beta = -0.15$, $p = 0.001$ for standard central SBP). MAP-derived central SBP was independently associated with LA reservoir strain ($\beta = -0.12$, $p = 0.01$) but not with GLS or LA conduit strain ($p > 0.05$). No BP parameter was independently associated with LA pump strain ($p > 0.05$).

Discrimination of abnormal LA and LV function. **Figures 2 and 3** demonstrate scatterplots among brachial BP, MAP-derived or standard central SBP with abnormal GLS and LA reservoir strain being color-coded (cut-offs reported in methods section). These plots were divided into 4 subsections based on cutoffs for SBP (140 mmHg for both central and brachial SBP). Based on these cut-off values for central BPs, MAP-derived central SBP identified an additional 32 (37%) patients with abnormal GLS (total $n = 87$

for abnormal GLS) who were classified as normotensive based on standard central SBP criteria (**Figure 2A**). For LA strain, MAP-derived central SBP identified an additional 101 (47%) patients with abnormal LA reservoir strain (total n=213) (**Figure 2C**), 157 (49%) patients with abnormal LA conduit strain (total n=321) (**Figure 3A**); and 63 (48%) patients with abnormal LA pump strain (total n=130) who were classified as normotensive based on standard central SBP (**Figure 3C**). On the contrary, standard central SBP identified no additional patients with abnormal GLS and only 2 additional patients with abnormal LA strain (1 LA reservoir and 1 LA conduit) who were classified as normotensive based on MAP-derived central SBP. Collectively, net increments of patients with abnormal GLS, LA reservoir, conduit, and pump strain were 32 (37%), 100 (47%), 156 (49%), and 63 (48%), respectively.

Compared with brachial SBP, MAP-derived central SBP identified an additional 16 (18%) patients with abnormal GLS (**Figure 2B**), 36 (17%) patients with abnormal LA reservoir strain (**Figure 2D**), 57 (18%) patients with abnormal LA conduit strain (**Figure 3B**), and 31 (24%) patients with abnormal LA pump strain who were classified as normotensive based on brachial SBP criteria (**Figure 3D**). On the contrary, brachial SBP identified an additional 6 patients with abnormal GLS and 17 patients with abnormal LA strain (4 patients with LA reservoir/10 patients with LA conduit and 3 patients with LA pump strain) who were classified as normotensive based on MAP-derived central SBP. Collectively, net increments of patients with abnormal GLS, LA reservoir, conduit, and pump strain were 10 (11%), 32 (15%), 47(15%), and 28(22%), respectively.

Reproducibility. Reproducibility was assessed by blinded strain measurements in a random sample of 20 patients. All measurements were done by the same two investigators (SR and TN) and the mean of the absolute value of differences between measurements was calculated. For GLS the mean $\pm$ SD difference was 0.7 $\pm$ 0.7%. For LA strain the mean difference was 8.0 $\pm$ 7.0% for reservoir strain, 5.4 $\pm$ 4.1% for conduit strain and 5.6 $\pm$ 4.6% for pump strain. Intra-observer variability was assessed by one investigator (SR) who repeated LA strain measurements in the same 20 patients at a different timepoint. The mean difference for LA strain was 3.8 $\pm$ 2.9% for reservoir strain, 2.7 $\pm$ 1.4% for conduit strain and 2.7 $\pm$ 1.4% for pump strain.

Discussion

This study demonstrated that MAP-derived central SBP identified more patients with abnormal LA and LV strain (i.e. those with subclinical LV/LA dysfunction) compared with standard central SBP. Strong correlations among DBP (both brachial and central) measurements were observed whilst only moderate correlations in SBP measurements were observed. MAP-derived central SBP was higher than standard central SBP. CBP calibrated using MAP and DBP was more closely associated with LA reservoir strain compared with standard CBP, which is based on SBP and DBP. Compared with SBP, DBP was more strongly associated with LA reservoir/conduit strain and GLS.

Clinical Relevance of CBP. Clinical assessment of BP is an essential component of patient assessment. Use of peripheral (brachial) BP is convenient, easily reproducible and non-invasive. Clinical practice guidelines and cardiovascular risk assessment currently depend on brachial BP recordings[1]. However

brachial BP only provides a rough estimate of central (aortic) pressure[9]. Previously CBP was only able to be measured invasively, having a limited clinical role. Newer methods of measuring CBP non-invasively using waveform analysis have several advantages. As noninvasive CBP may be more closely representative of invasive aortic pressure than brachial BP[9, 10, 14], it may have a useful role in assessment of cardiovascular outcomes[4-7], having a prognostic role[7] and potentially associated with improved hypertension management[15].

Association of BP with LA/LV strain. Strain analysis using speckle tracking provides quantitative information of LA and LV function. BP has a strong impact on LA and LV function. Elevated BP leads to LV hypertrophy, reduced compliance and increased stiffness. The increase in LA pressure contributes to LA dilatation and may lead to fibrosis[12, 16]. There is interdependence on the three components of LA strain. LV function has a direct influence on LA reservoir and conduit function[17], where LA pump function has much less association with LV function[17]. This is important in the pathogenesis in conditions such as HF with preserved ejection fraction as well as AF.

In our study, DBPs were more strongly associated with GLS and LA strain than SBPs. This is an interesting finding because SBP is also more commonly used in clinical practice to guide diagnosis. In addition, previous studies have not demonstrated a stronger association between DBP and cardiovascular outcomes compared with SBP[1, 18]. Our finding may have important clinical implications and highlights the potential importance in monitoring DBP in patients with hypertension and the use of DBP in identifying those with subclinical LA/LV dysfunction. Very strong mutual correlations were noted among all three methods of DBP, with little difference in measured values. Thus, there may not be additional utility in measurement of central DBP compared with brachial DBP or standard central DBP.

Furthermore, we observed that mean MAP-derived central SBP was higher than brachial SBP and standard central SBP. A weaker correlation between MAP-derived central SBP and brachial SBP compared with standard central SBP, suggesting MAP-derived central SBP may possess different diagnostic signals to brachial BP. Indeed, MAP-derived central SBP identified more patients with abnormal LV and LA strain than standard central SBP/brachial SBP did. These observations have important clinical implications. Noninvasive BP are different among calibration methods. MAP-derived CBP may help identify those with end organ functional alteration earlier, where subclinical LV and LA dysfunction are risk factors for developing overt HF or atrial fibrillation (AF)[19-21]. Therefore, MAP-derived central SBP may be utilized as a gate keeper to rule-out low risk patients and identify those who would benefit from further advanced echocardiographic assessment. This finding is also consistent with our previous findings showing that MAP-derived central SBP had increased discriminatory power compared with other SBP markers in detecting those with LVH and LA enlargement in patients with stage A HF[11].

Use of CBP in hypertension management. CBP may be a more appropriate BP marker to use in the diagnosis of hypertension. MAP-derived central SBP was noted in our study to be higher than brachial SBP and standard central SBP, which has been observed in other studies[11, 22]. This suggests an underestimate of central SBP using brachial SBP calibration. Given recent guidelines have advocated for

earlier recognition of patients with hypertension and more aggressive treatment of BP[1], the use of a more sensitive marker will aid diagnosis and monitoring. As anti-hypertensive drugs have different effects on CBP than peripheral BP, the use of CBP may result in improved drug titration and may represent a more accurate reflection on treatment response.[7, 9, 23-26]

Comparison of MAP-derived and standard CBP calibration methods. There are multiple different calibration methods used to calculate CBP non-invasively[9]. Previously there were no standardized guidelines and different algorithms and calibrations were utilized leading to discrepancy in the accuracy of CBP measurements. A recent taskforce has been established to help create more uniform standards[9]. The use of different calibration standards has potentially a large impact on clinical outcomes. We noted a large discrepancy in SBP recordings using the two most common CBP calibration methods and brachial BP. DBPs were much more strongly correlated with each other. Future research is required in determining the most appropriate calibration method which should be used in measuring CBP. Currently brachial BP offers a simple, effective and reproducible assessment of BP in clinical practice. However, with further improvements in determining CBP accurately using non-invasive, cost effective methods, may translate into a more feasible assessment tool for clinical practice.

Limitations: This was a cross sectional study so that causality is unable to be established. CBP was not assessed invasively and we used a single CBP device for all measurements. Although this device has been validated[22], it will be important to replicate these results across other device vendors to ensure no variability exists. LA and LV strain were both measured using speckle tracking. This was performed offline using an LV algorithm. There are currently differences in the algorithm used amongst software vendors as well as differences in nomenclature and ECG gating. Despite recent attempts at creating normal reference values for LA strain[27] and standard consensus guidelines[28], further studies are warranted to assess the generatability of our results.

Conclusion

CBP calibrated using MAP and DBP may be more closely associated with LA and LV function than standard brachial calibration methods. MAP-derived central SBP can be used as a sensitive screening tool to detect functional target organ damages from hypertension.

Abbreviations

ACE-I - Angiotensin Converting Enzyme Inhibitors

AF – Atrial Fibrillation

BMI – Body Mass Index

BP – Blood Pressure

CBP – Central Blood Pressure

DBP – Diastolic Blood Pressure

ECG – Electrocardiograph

GLS – Global Longitudinal Strain

HF – Heart Failure

LA – Left Atrium

LV – Left Ventricle

LVEF – Left Ventricular Ejection Fraction

LVH – Left Ventricular Hypertrophy

LVM – Left Ventricular Mass

MAP – Mean Arterial Pressure

SBP – Systolic Blood Pressure

T2DM – Type II Diabetes Mellitus

Tas-ELF – Tasmanian AF Screening Cohort

TTE – Transthoracic echocardiography

Declarations

Disclosures: No conflicts of interest

Funding: S.R is supported by a research scholarship by the National Heart Foundation of Australia. This study was partially supported by Diabetes Australia, the Tasmanian Community Fund and Siemens Healthcare Australia. ECG equipment and software support were provided by Semacare Inc, a manufacturer of handheld ECG devices. The sponsors had no role in the design and conduct of the study, in the collection, analysis, and interpretation of the data, and in the preparation, review, or approval of the manuscript.

Conflicts of Interest – none to declare

References

- [1] Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. 2018;71:e127-e248.
- [2] Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224-60.
- [3] Picone DS, Schultz MG, Otahal P, Aakhus S, Al-Jumaily AM, Black JA, et al. Accuracy of Cuff-Measured Blood Pressure: Systematic Reviews and Meta-Analyses. *Journal of the American College of Cardiology*. 2017;70:572-86.
- [4] Cheng HM, Chuang SY, Sung SH, Yu WC, Pearson A, Lakatta EG, et al. Derivation and validation of diagnostic thresholds for central blood pressure measurements based on long-term cardiovascular risks. *Journal of the American College of Cardiology*. 2013;62:1780-7.
- [5] Huang CM, Wang KL, Cheng HM, Chuang SY, Sung SH, Yu WC, et al. Central versus ambulatory blood pressure in the prediction of all-cause and cardiovascular mortalities. *Journal of Hypertension*. 2011;29:454-9.
- [6] Kollias A, Lagou S, Zeniodi ME, Boubouchairopoulou N, Stergiou GS. Association of central versus brachial blood pressure with target-organ damage: systematic review and meta-analysis. *Hypertension*. 2016;67:183-90.
- [7] McEniery CM, Cockcroft JR, Roman MJ, Franklin SS, Wilkinson IB. Central blood pressure: current evidence and clinical importance. *European Heart Journal*. 2014;35:1719-25.
- [8] Dart AM, Gatzka CD, Kingwell BA, Willson K, Cameron JD, Liang Y-L, et al. Brachial blood pressure but not carotid arterial waveforms predict cardiovascular events in elderly female hypertensives. *Hypertension*. 2006;47:785-90.
- [9] Sharman JE, Avolio AP, Baulmann J, Benetos A, Blacher J, Blizzard CL, et al. Validation of non-invasive central blood pressure devices: ARTERY Society task force consensus statement on protocol standardization. *European Heart Journal*. 2017;38:2805-12.
- [10] Kobayashi H, Kinou M, Takazawa K. Correlation between the brachial blood pressure values obtained using the cuff method and the central blood pressure values obtained invasively. *Internal Medicine*. 2013;52:1675-80.
- [11] Negishi K, Yang H, Wang Y, Nolan MT, Negishi T, Pathan F, et al. Importance of calibration method in central blood pressure for cardiac structural abnormalities. *American Journal of Hypertension*. 2016;29:1070-6.

- [12] Lau DH, Mackenzie L, Kelly DJ, Psaltis PJ, Brooks AG, Worthington M, et al. Hypertension and atrial fibrillation: evidence of progressive atrial remodeling with electrostructural correlate in a conscious chronically instrumented ovine model. *Heart Rhythm*. 2010;7:1282-90.
- [13] Mor-Avi V, Lang RM, Badano LP, Belohlavek M, Cardim NM, Derumeaux G, et al. Current and evolving echocardiographic techniques for the quantitative evaluation of cardiac mechanics: ASE/EAE consensus statement on methodology and indications endorsed by the Japanese Society of Echocardiography. *European Journal of Echocardiography*. 2011;12:167-205.
- [14] Millasseau S, Agnoletti D. Non-invasive estimation of aortic blood pressures: a close look at current devices and methods. *Current Pharmaceutical Design*. 2015;21:709-18.
- [15] Sharman JE, Marwick TH, Gilroy D, Otahal P, Abhayaratna WP, Stowasser M. Randomized trial of guiding hypertension management using central aortic blood pressure compared with best-practice care: principal findings of the BP GUIDE study. *Hypertension*. 2013;62:1138-45.
- [16] Vaziri SM, Larson MG, Lauer MS, Benjamin EJ, Levy D. Influence of blood pressure on left atrial size: the Framingham Heart Study. *Hypertension*. 1995;25:1155-60.
- [17] Ramkumar S, Yang H, Wang Y, Nolan M, Negishi T, Negishi K, et al. Association of the active and passive components of left atrial deformation with left ventricular function. *Journal of the American Society of Echocardiography*. 2017;30:659-66.
- [18] Benetos A, Thomas F, Bean K, Gautier S, Smulyan H, Guize L. Prognostic value of systolic and diastolic blood pressure in treated hypertensive men. *Archives of Internal Medicine*. 2002;162:577-81.
- [19] Ramkumar S, Ochi A, Kawakami H, Yang H, Potter EL, D'Elia N, et al. Echocardiographic Risk Assessment to Guide Screening for Atrial Fibrillation. *Journal of the American Society of Echocardiography*. 2019;32:1259-67.
- [20] Kawakami H, Ramkumar S, Nolan M, Wright L, Yang H, Negishi K, et al. Left Atrial Mechanical Dispersion Assessed by Strain Echocardiography as an Independent Predictor of New-Onset Atrial Fibrillation: A Case-Control Study. *Journal of the American Society of Echocardiography*. 2019;32:1268-76.e3.
- [21] Kawakami H, Ramkumar S, Pathan F, Wright L, Marwick TH. Use of echocardiography to stratify the risk of atrial fibrillation: comparison of left atrial and ventricular strain. *European Heart Journal Cardiovascular Imaging*. 2020;21:399-407.
- [22] Weber T, Wassertheurer S, Rammer M, Maurer E, Hametner B, Mayer CC, et al. Validation of a brachial cuff-based method for estimating central systolic blood pressure. *Hypertension*. 2011;58:825-32.
- [23] Kelly R, Gibs H, O'ROURKE M, Daley J, Mang K, Morgen J, et al. Nitroglycerin has more favourable effects on left ventricular afterload than apparent from measurement of pressure in a peripheral artery.

European Heart Journal. 1990;11:138-44.

[24] Matsui Y, Eguchi K, O'Rourke MF, Ishikawa J, Miyashita H, Shimada K, et al. Differential effects between a calcium channel blocker and a diuretic when used in combination with angiotensin II receptor blocker on central aortic pressure in hypertensive patients. *Hypertension*. 2009;54:716-23.

[25] Manisty CH, Zambanini A, Parker KH, Davies JE, Francis DP, Mayet J, et al. Differences in the magnitude of wave reflection account for differential effects of amlodipine-versus atenolol-based regimens on central blood pressure: an Anglo-Scandinavian Cardiac Outcome Trial substudy. *Hypertension*. 2009;54:724-30.

[26] Agabiti-Rosei E, Mancia G, O'Rourke MF, Roman MJ, Safar ME, Smulyan H, et al. Central Blood Pressure Measurements and Antihypertensive Therapy. *Hypertension*. 2007;50:154-60.

[27] Pathan F, D'Elia N, Nolan MT, Marwick TH, Negishi K. Normal Ranges of Left Atrial Strain by Speckle-Tracking Echocardiography: A Systematic Review and Meta-Analysis. *Journal of the American Society of Echocardiography*. 2017;30:59-70 e8.

[28] Badano LP, Koliass TJ, Muraru D, Abraham TP, Aurigemma G, Edvardsen T, et al. Standardization of left atrial, right ventricular, and right atrial deformation imaging using two-dimensional speckle tracking echocardiography: a consensus document of the EACVI/ASE/Industry Task Force to standardize deformation imaging. *European Heart Journal-Cardiovascular Imaging*. 2018;19:591-600.

Tables

Table 1 –Baseline characteristics of the overall cohort and participants grouped by GLS (cut-off -16%).

Baseline Patient Characteristics	Entire cohort	Subjects with normal GLS ($\leq -16\%$)	Subjects with abnormal GLS ($> -16\%$)	P value
Demographics	n=546	n=459	n=87	
Age - years (SD)	70.7 (4.7)	70.6 (4.7)	71.3 (4.6)	0.20
Male n(%)	247 (45)	194 (42)	53 (61)	0.001
BMI (kg/m ²) (SD)	29.4 (5.2)	29.2 (5.1)	30.5 (6.1)	0.03
Current Smoking n(%)	12 (2)	11 (2)	1 (1)	0.47
Diabetes Mellitus n(%)	284 (52)	224 (49)	60 (69)	0.001
Obesity n(%)	234 (43)	192 (42)	42 (48)	0.27
Hypercholesterolemia n(%)	281/533 (54)	234/442 (53)	47/80 (59)	0.34
Hypertension n(%)	429 (79)	360 (78)	69 (79)	0.86
Previous history of IHD n(%)	429 (79)	29 (6)	9 (10)	0.18
Median Six Minute Walk Test m(IQR)	38 (7) 485 (115.5)	490 (111.8)	469 (129.0)	0.02
Medications				
Beta blockers n(%)	34 (6)	24 (5)	10 (12)	0.03
ACE Inhibitor/Angiotensin Receptor Blocker n(%)	363 (67)	309 (67)	54 (62)	0.34
Calcium Channel Blocker n(%)	117/503 (23)	91/425 (21)	26/78 (33)	0.02
Lipid lowering agents n(%)	278/505 (55)	231/427 (54)	47/78 (60)	0.32
Anti-platelet agents n(%)	189/502 (38)	159/426 (37)	30/76 (40)	0.72
Blood Pressure				
Brachial Systolic BP mmHg (SD)	140.0 (16.6)	139.7 (15.7)	141.8 (20.6)	0.38
Brachial Diastolic BP mmHg (SD)	81.7 (10.3)	81.1 (10.2)	85.2 (10.3)	0.001
Standard Systolic Central BP mmHg (SD)	128.2 (15.2)	127.9 (14.6)	130.0 (18.0)	0.30
Standard Diastolic Central BP mmHg (SD)	82.8 (10.2)	82.1 (9.9)	86.7 (10.4)	<0.001
		150.0 (18.7)	151.7 (26.5)	0.59
		83.0 (10.0)	87.7 (10.9)	<0.001

MAP derived Systolic Central BP mmHg (SD)	150.3 (20.2)			
MAP derived Diastolic Central BP mmHg (SD)	83.7 (10.3)			
Echocardiographic Parameters	Mean (SD)			
Ejection Fraction % (SD)	63.7 (5.9)	64.4 (5.2)	60.0 (7.6)	<0.001
Global Longitudinal Strain % (SD)	-18.6 (2.5)	-19.3 (1.9)	-14.6 (1.3)	<0.001
E/e' (Average of lateral and septal) (SD)	8.9 (2.6)	8.9 (2.6)	8.8 (2.8)	0.66
Left atrial volume - indexed ml/m ² (SD)	31.6 (9.2)	31.5 (9.1)	32.4 (9.9)	0.38
LV mass index g/m ² (SD)	91.5 (22.7)	90.0 (21.6)	99.2 (26.8)	0.003
LA Reservoir Strain % (SD)	39.3 (6.8)	39.7 (6.5)	36.7 (7.9)	0.001
LA Conduit Strain % (SD)	19.9 (5.4)	20.2 (5.2)	18.1 (5.9)	0.002
LA Pump Strain % (SD)	19.4 (5.0)	19.5 (4.9)	18.6 (5.5)	0.14

Table 2: Association between brachial/CBP and LV/LA strain.

		LV Strain		LA Strain					
		GLS		Reservoir		Conduit		Pump	
Variables		β	P value	β	P value	β	P value	β	P value
SBP	Brachial	-0.09	0.04	-0.09	0.04	-0.13	0.004	0.01	0.78
	Standard Central	-0.15	0.001	-0.06	0.21	-0.12	0.006	0.06	0.23
	MAP Derived Central	-0.05	0.23	-0.12	0.01	-0.07	0.12	-0.08	0.09
DBP	Brachial	-0.20	<0.001	-0.14	0.002	-0.20	<0.001	0.02	0.70
	Standard Central	-0.22	<0.001	-0.12	0.01	-0.19	<0.001	0.04	0.36
	MAP Derived Central	-0.20	<0.001	-0.14	0.002	-0.19	<0.001	0.01	0.87

Multivariable model adjusted for body mass index, type 2 diabetes mellitus, family history, angiotensin-converting enzyme/angiotensin receptor blocker, beta blocker, calcium channel blocker and diuretics.

The above table shows that DBP is more closely associated with LA and LV function than SBP. Standard CBP and brachial SBP was more closely associated with LV function than MAP derived CBP calibration. Steiger's Z test (comparing correlation co-efficient of MAP central SBP and Standard central SBP):

(LA Reservoir - Z score = 0.96, p=0.36, GLS – Z score = -1.1, p=0.27)

Figures

Figure 1

Scatterplot comparing correlation between brachial, standard and MAP-derived CBP measurements

Figure 2

Scatterplots comparing MAP-derived central SBP with standard central SBP and brachial SBP based on presence of reduced GLS and LA reservoir strain (cut-off LA reservoir 38%, GLS -16%).

Figure 3

Scatterplots comparing MAP-derived central SBP with standard central SBP and brachial SBP based on presence of reduced LA conduit and pump strain (cut-off LA conduit 21%, LA pump 16%).