Genetic diversity of Brazilian *Bacillus thuringiensis* isolates with toxicity against *Aedes aegypti* (Diptera: Culicidae)

Geysla Costa Fernandes

State University of Maranhão

Dalton Kaynnan Prado Costa

State University of Maranhão

Nayanne Santos Oliveira

State University of Maranhão

Emanuelle Cristine Pereira Sousa

State University of Maranhão

Déborah Heloísa Bittencourt Machado

São Paulo State University (Unesp)

Antônio Ricardo Polanczyk

São Paulo State University (Unesp)

Herbert Álvaro Abreu Siqueira

Federal Rural University of Pernambuco

Maria Cleoneide Silva (✉ cleouema@hotmail.com)

State University of Maranhão

Research Article

Keywords: solates, btma, toxic, aegypti, standard, subspecies

Posted Date: August 24th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-817179/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Genetic diversity of Brazilian *Bacillus thuringiensis* isolates with toxicity against *Aedes aegypti* (Diptera: Culicidae)

Geysla da Costa Fernandes¹; Dalton Kaynnan de Prado Costa²; Nayanne Santos de Oliveira²; Emanuelle Cristine Pereira de Sousa³; Déborah Heloísa Bittencourt Machado⁴; Ricardo Antônio Polanczyk⁵; Herbert Álvaro Abreu de Siqueira⁶; Maria Cleoneide da Silva^{7*}

¹ PostGraduate Program in Biodiversity, Environment and Health, Center for Higher Studies of Caxias, State University of Maranhão, Praça Duque de Caxias, Morro do Alecrim, S/N, Caxias, MA, 65604-380, Brazil

² Laboratory of Entomopathogenic Bacteria and Molecular Markers, Center for Higher Studies of Caxias, State University of Maranhão, Praça Duque de Caxias, Morro do Alecrim, S/N, Caxias, MA, 65604-380, Brazil

³ Postgraduate Program in Animal Science, State University of Maranhão, University City Paul VI, Avenida Lourenço Vieira da Silva 1000, Jardim São Cristóvão, São Luís, MA, 65055-310, Brazil

⁴ Postgraduate Program in Agronomy (Agricultural Entomology), São Paulo State University (Unesp), School of Agricultural and Veterinarian Sciences, Jaboticabal 14884-900, Brazil

⁵ Department of Plant Health, São Paulo State University (Unesp), School of Agricultural and Veterinarian Sciences, Jaboticabal 14884-900, Brazil

⁶Department of Agronomy (Entomology), Federal Rural University of Pernambuco, Recife, PE,
52171-900, Brazil

^{7*}Department of Chemistry and Biology, Center for Higher Studies of Caxias, State University
of Maranhão, Praça Duque de Caxias, Morro do Alecrim, S/N, Caxias, MA, 65604-380, Brazil

Correspondence: Maria Cleoneide Silva, Department of Chemistry and Biology, Laboratory
of Entomopathogenic Bacteria and Molecular Markers, Center for Higher Studies of Caxias,
State University of Maranhão, Praça Duque de Caxias, Morro do Alecrim, S/N, Caxias, MA,
65604-380, Brazil, cleouema@hotmail.com, <https://orcid.org/0000-0002-5273-8125>

Abstract

Bacillus thuringiensis (Bt) isolates native to Maranhão (BtMA) that are highly toxic to *Aedes aegypti* larvae and seven standard subspecies of Bt were analyzed for genetic diversity using the rep-PCR technique with BOX, ERIC, REP, MB1, and GTG₅ markers. The rep-PCR technique is considered an extremely reliable, reproducible, fast and highly discriminatory technique that may be used even among populations of the same species. These five markers revealed a total of 38 polymorphic DNA fragments for 30 BtMA isolates. Eight groups were obtained with the dendrogram generated through Pearson's correlation analysis, with four groups formed only with BtMA isolates and four comprised of isolates of BtMA and the standard subspecies toxic to dipterans and lepidopterans. Despite the high genetic diversity of BtMA, a low correlation between the collection site, gene content and mortality against *A. aegypti* larvae was evidenced. The clustering of the standard subspecies of Bt that were toxic against dipterans with BtMA isolates confirm the mosquitocidal action of the native isolates

from Maranhão, and they can be used as an alternative for *A. aegypti* control and other insects of medical importance and for the control of agricultural pests.

Introduction

Bacillus thuringiensis (Bt) is a gram-positive, aerobic, rod-shaped and spore-forming entomopathogenic bacterium that is highly effective in controlling immature forms of mosquitoes, since several strains produce protein crystals containing δ -endotoxin (Cry and Cyt) with mosquitocidal action^{16, 3, 11, 55, 25}.

Several subspecies of Bt have been reported to be highly toxic against dipteran insects, such as *Bt israelensis* (Bti), *Bt entomocidus*, *Bt sotto*, *Bt jegathesan*, *Bt darmstadiensis*, *Bt medellin*, *Bt fukuokaensis*, and *Bt higo*^{40, 5, 14, 60, 29, 49, 41, 24}. Bti is considered the most toxic to dipteran larvae^{16, 13, 3}. The mosquitocidal activity of Bti is due to insecticidal proteins encoded by the *cry4Aa*, *cry4Ba*, *cry11Aa*, *cyt1Aa*, *cry10Aa* and *cyt2Ba* genes^{4, 3, 55}.

Molecular typing has been used to identify and classify species belonging to the genus *Bacillus*, in addition to monitoring the evolution and phylogeny of the group^{30, 15, 28, 39}. Thus, the rep-PCR technique is widely used to investigate specific genetic relationships between Bt isolates^{9, 18, 19, 37, 53, 35}.

The rep-PCR technique uses primers specific to the *Bacillus cereus* group and is considered an extremely reliable, reproducible, fast and highly discriminatory technique that may be used even among populations of the same species^{58, 45, 1, 23, 50} or even among microorganisms^{58, 23, 9, 35}.

The objective of the current study was to estimate the genetic diversity of Bt isolates from the Entomopathogenic Bacteria Collection of Maranhão (CBENMA) and the Entomopathogenic Bacilli Bank of Maranhão (BBENMA) that are toxic to *A. aegypti* larvae^{52, 34, 59} using the repetitive sequences BOX, ERIC, REP, MB1, and GTG₅. Furthermore, the study

aimed to evaluate the association between Bt isolate clusters and standard subspecies of this bacterium, mortality, and amplification of dipteran-specific genes.

Results

The diversity of *Bacillus thuringiensis* isolates native to Maranhão (BtMA) was determined by rep-PCR, and a complex fingerprint pattern was obtained for the 30 BtMA isolates with the BOX, ERIC, REP, MB1 and GTG₅ primers. The results revealed a total of 38 polymorphic DNA fragments (Fig. 1).

The number of fragments per isolate for the BOX primer ranged from one to seven, with the smallest fragment of 100 bp and the largest exceeding 10.000 bp, with a total of 24 electrophoretic patterns (Fig. 1a). The ERIC and REP primers provided fragments ranging from one to six per isolate. For the ERIC primer, the base pairs ranged from 250 to 1,250 bp, and for REP, they ranged from 250 to 2,000 bp, with a total of 13 and 12 electrophoretic patterns, respectively (Fig. 1b and Fig. 1c). The MB1 primer provided between one to eight fragments and a total of 31 electrophoretic patterns with fragment sizes between 250 and 6,000 bp (Fig. 1d). Regarding the GTG₅ primer, the BtMA isolates provided fragments of repetitive sequences that ranged from two to 11 bands, with the majority being similar, from 200 to 3,000 bp, with a total of 11 electrophoretic patterns (Fig. 1e). Although some isolates shared the same bands, most of them presented different fragment profiles.

A dendrogram was generated through Pearson's correlation for genetic diversity analyses using the sequences BOX, ERIC, REP, MB1, and GTG₅, with an average distance of 45% (cutoff point), in which eight groups (designated I, II, III, IV, V, VI, VII and VIII) were obtained, which comprised 67,5% of the isolates (Fig. 2). The other Bt isolates showed an average distance greater than the cutoff point.

Groups I, V, VI and VIII were formed only with BtMA isolates (Fig. 2). Group I (BtMA-212, BtMA-527 and BtMA-459) and group V (BtMA-687, BtMA-694 and BtMA-703) included isolates obtained from different biomes. The BtMA-212 isolate obtained from Amazonia grouped with the BtMA-527 and BtMA-459 isolates from the Cerrado. These three isolates, in addition to presenting 100% mortality to *A. aegypti* larvae in the second evaluation (48 h), amplified the *cry* and *cyt* genes.

Groups VI and VIII were formed by isolates from the same biome, with isolates of group VI (BtMA-679, BtMA-684, BtMA-685, BtMA-688 and BtMA-689) all from the Cerrado and those from group VIII (BtMA-229 and BtMA-237) from the Amazon. All isolates in group VI caused 100% mortality within a 24-hour period and amplified all analyzed dipteran-specific genes, with the exception of the BtMA-689 isolate, which did not present the *cry11Ba* gene. For the genetic profile of isolates from group VIII, it was found that they did not share any of the *cry* and *cyt* genes analyzed. An *A. aegypti* larval mortality rate of 100% was caused by isolates of BtMA in this group at 48 h.

Groups II, III, IV and VII comprised isolates of BtMA and the standard subspecies. The BtMA-179 isolate that grouped with the subspecies *Bacillus thuringiensis* subsp. *aizawai* (Bta) (group II) was isolated from the soil of the Amazon biome, caused 100% mortality in 48 h and amplified only the dipteran-specific *cry4Ba* gene, and the BtMA-233 and BtMA-241 isolates grouped with subspecies *Bacillus thuringiensis* subsp. *yunnanensis* (Bty) (IV) were isolated from the soil of the Amazon and Cerrado biomes, respectively. These three isolates were toxic to *A. aegypti* larvae over a period of 48 h, and isolates BtMA-233 and BtMA-241 presented the *cyt2Aa* gene in common. The subspecies Bta and Bty are toxic to insects of the order Lepidoptera, which suggests that isolates from groups II and IV may also be active against insects of that order.

The BtMA-215 isolate, obtained from Amazonian soil, did not amplify any of the analyzed genes, caused 100% mortality in *A. aegypti* larvae in 48 hours, and grouped with two subspecies, *Bacillus thuringiensis* subsp. *israelensis* (Bti) and *Bacillus thuringiensis* subsp. *sotto* (Bts) (group III). These subspecies possibly confirm the mosquitocidal action of the isolate BtMA-215 for having grouped with Bti and possibly have potential for the control of lepidopterans because of being grouped with Bts.

The isolates BtMA-682, BtMA-690 and BtMA-691 were found in group VII, all from the Cerrado soil, with different mortality rates and genetic profiles, and grouped with the subspecies *Bacillus thuringiensis* subsp. *entomocidus* (Bte), which is at the same time effective against insects of the orders Diptera and Lepidoptera. It is worth mentioning that the BtMA-690 isolate amplified all the *cry* and *cyt* genes analyzed and provided 100% mortality of *A. aegypti* larvae in 24 h.

Discussion

The repetitive element palindromic-polymerase chain reaction (rep-PCR) technique has been used to investigate the interspecific and intraspecific genetic diversity of Bt isolates, allowing us to correlate them with the geographic regions of collection, the origin of the substrate, morphological, biochemical and genetic characteristics, and rates of mortality and toxicity against insect pests and vectors^{32, 50, 21, 19, 35}.

In this research, the genetic diversity of BtMA isolates collected in different biomes of Maranhão, with a high rate of mortality against larvae of *A. aegypti* and amplified for several *cry* and *cyt* dipteran-specific genes, was studied using group analysis with data obtained with the BOX, ERIC, REP, MB1 and GTG₅ primers.

The BOX and MB1-PCRs were the most informative, producing the most complex fragment patterns^{7, 9, 53}. Robust and, commonly, highly complex fingerprints are obtained using

the BOX primer^{46, 44}. The MB1 primer generated extremely complex fingerprints, showing that almost all 30 BtMA isolates had their own unique MB1-PCR standard, suggesting high genetic variability among the isolates^{7, 18}.

Bacillus thuringiensis is a genetically diverse species, so its large number of strains can form several different profiles according to their genetic profile⁴⁸. The high genetic variability of Bt may be related to both the influence of geographic and ecological factors^{19, 37, 35} because this bacterium is isolated from several substrates and has a wide range of target hosts.

Several studies have reported that ERIC markers are more informative than REP markers in discriminating Bt strains^{32, 26, 9, 35}, as was also verified in this research, suggesting that ERIC-like sequences may be more widely distributed than REP-like sequences in Bt²⁶.

Among all the markers, GTG₅ was not as effective in discriminating between isolates of BtMA, indicating low genetic diversity. A similar result was verified when this marker was used to determine the diversity among the different *Bacillus* species^{17, 53}.

It was possible to verify that groups VI, VII and VIII included isolates of BtMA from the same biome. The similarity of the BtMA isolates from the Cerrado (groups VI and VII) and the Amazon (VIII) biomes may be related to the geographic location. However, groups I, IV and V included isolates of BtMA from the Cerrado with other biomes. The tendency to form groups or subgroups in relation to the similarity of Bt according to geographic origin, using the Rep-PCR technique, is quite controversial.

Huerta et al.²¹, using REP primers, evaluated the genetic diversity of 53 Bt isolates collected in soil from different geographical regions of Peru and correlated the similarity of these isolates according to geographic origin. Similarly, Da Silva and Valicente⁹ pointed out that the similarity of Bt isolates may be related to geographic location when they used the ERIC, REP and BOX primers simultaneously. Katara et al.²⁶ observed that the REP-PCR and

ERIC-PCR patterns, generated with 113 Bt native isolates from diverse habitats of India, formed four main groups with isolates of diverse origin.

All isolates of BtMA from the VI group (BtMA-684, BtMA-685, BtMA-688, BtMA-689, and BtMA-679) presented the same mortality rate (100% in 24 h), and all isolates of BtMA from the I, II and VIII groups caused 100% mortality within 48 h. Groups V and VII included isolates with different mortality rates (24 and 48 h). These results suggest that there was no genetic relationship between BtMA isolates and their *A. aegypti* larvae. When correlating the mortality of the BtMA isolates and the site of origin, only group VI, constituted by the isolates BtMA-684, BtMA-685, BtMA-688, BtMA-689 and BtMA-679, showed a genetic relationship.

This low correlation was also verified by Huerta et al.²¹ when verifying subgroups of Bt formed according to the geographical origin and potential use against *A. aegypti*. Likewise, Da Silva and Valicente⁹ and Machado et al.³⁵, comparing the genetic relationship of Bt isolates toxic to *Spodoptera* spp. and the collection sites of these isolates reported a low correlation. In addition, in agreement with Reyes-Ramirez and Ibarra⁴⁸ and Da Silva and Valicente⁹, the genomic relationship between Bt strains is not defined only by their specific toxicity but by several characteristics, such as the content of the *cry* gene, crystal morphology and plasmid pattern. Bt larvicidal activity comes from crystal protoxins encoded by *cry* and *cyt* genes.

The BtMA isolates showed a great diversity of the diptera-specific genes *cry4*, *cry10*, *cry11*, *cyt1* and *cyt2*^{52, 34, 59}. Groups V and VI included BtMA isolates with all the investigated genes, with the exception of the BtMA-694 isolate from group V, which did not amplify the *cyt2Ba* gene, and the BtMA-689 isolate from group VI, which did not amplify the *cry11Ba* gene. The great diversity of genes encoding mosquito-specific toxins in BtMA isolates, presenting the same genetic content as Bti, represents a great opportunity to control vector insects and enables strategies to manage the evolution of insect resistance to Bt insecticidal proteins.

One factor that may or may not explain the correlation with the Bt collection sites is that the diversity may also be associated with the region's climate, soil conditions where the samples were collected, and the influence of other microbial communities present in the region³⁷.

Interestingly, the isolates of group V (BtMA-687 BtMA-694 BtMA-703) did not show any relation to the geographic origin and mortality of the *A. aegypti* larvae, and the isolates of VI (BtMA-684, BtMA-685, BtMA-688, BtMA-689 and BtMA-679) were all from the same biome and caused 100% mortality to *A. aegypti* larvae in the first evaluation (24 h). Based on these results, the correlation between the genetic relationship of BtMA isolates and their investigated gene content is very low. Lima et al.³² highlighted that repetitive sequences showed no relation to the types of insecticidal protein produced by Bt, which demonstrates that the majority of clusters that occur between isolates that do not carry the same combinations of dipteran-specific genes.

For the groups formed with the standard subspecies and the BtMA isolates (groups II, III, IV and VII), it was possible, in addition to confirming the toxic activity of these isolates against *A. aegypti* larvae, to suggest the potential of the BtMA isolates for the control of insects of the order Lepidoptera. However, it will be necessary to carry out bioassays with larvae of pest lepidopterans and to characterize these isolates with lepidopteran-specific genes of Bt.

An interesting fact was the grouping of the isolate BtMA-215 with the subspecies Bti (group III). Unlike the subspecies Bti, this isolate caused mortality to *A. aegypti* larvae within 48 h and did not amplify any of the diptera-specific *cry* and *cyt* genes selected for this study. However, Soares-da-Silva et al.⁵² reported that the BtMA-215 isolate amplified the *cry32* gene. There are 29 Cry32 proteins⁸, and according to Van Frankenhuyzen⁵⁶ and Rajchanuwong et al.⁴⁷, these proteins are toxic to insects of the order Diptera, especially mosquito larvae.

In this work, we provided evidence of high genetic variability of *Bacillus thuringiensis* isolates native to Maranhão (BtMA) and toxic to *A. aegypti*, providing important information about their phylogenetic relationships and similarity with the standard subspecies (Bti, Bta, Bte, Bty, and Bts).

In addition, the clusters with the standard subspecies of Bt toxic to dipterans with BtMA isolates confirm the mosquitocidal action of the native isolates from Maranhão and that they can be used as an alternative for control of *A. aegypti* and other insects of medical importance. This fact also demonstrated that isolates that grouped with subspecies toxic to lepidopterans, such as Bta and Bty, could constitute new tools in integrated pest management, providing different Cry insecticidal proteins than those expressed in ordinary Bt formulations⁴³.

We must note that detailed information on the Bt *cry* gene profile is highly important not only for approaches to insect resistance management and host spectrum^{56, 31, 61} but also for providing important insights into Bt formulation because different insecticidal proteins can require specific fermentation conditions, such as dissolved oxygen and temperature^{42, 20}.

Methods

***Bacillus thuringiensis* isolates.** Thirty *Bacillus thuringiensis* isolates native to Maranhão (BtMA) obtained from soil samples and with high toxicity to *A. aegypti* larvae amplified for *cry* (*cry4Aa*, *cry4Ba*, *cry10Aa*, *cry11Aa* and *cry11Ba*) and *cyt* (*cyt1Aa*, *cyt1Ab*, *cyt2Aa* and *cyt2Ba*)^{52, 34, 59} and seven standard subspecies of Bt were used in this survey (Tables 1 and 2). All these isolates are stored in CBENMA located in the Laboratory of Entomopathogenic Bacteria and Molecular Markers (BEMMOL) at the Centro de Estudos Superiores de Caxias of Universidade Estadual do Maranhão (CESC/UEMA).

DNA extraction and PCR conditions. Genomic DNA from BtMA isolates and standard subspecies of Bt was extracted using the Instagene matrix (Bio-Rad) according to the manufacturer's recommendations. PCR was performed in final volume of 20 μ L. Each reaction mixture contained 50 ng of genomic DNA of Bt isolates, 1x buffer, 2 mM $MgCl_2$, each dNTP at a final concentration of 200 μ M, 1 U of Taq DNA Polymerase and 1 μ M each primer (BOX, ERIC, REP, MB1, and GTG₅) (Table 3). Amplification was accomplished with the Gencycler-G96G thermocycler (Biosystems), following the standard program at 94 °C for 5 min; 36 cycles at 94 °C for 1 min; annealing (with temperatures varying for each primer) for 1 min; extension at 72 °C for 2 min; and final extension at 72 °C for 7 min. The PCR products were visualized in a 1,5% agarose gel using a photo documentation system (L-PIX EX Loccus).

Data analysis. The Rep-PCR profiles (banding patterns) obtained with the 30 BtMA isolates and the seven standard subspecies of Bt generated binary matrices that were used as input data into Bionumerics software (Applied Maths, Belgium) after Pearson's correlation analysis. The Dice similarity coefficient was used to calculate the similarity matrix from binary data. Clustering analysis was performed using this coefficient and UPGMA (unweighted pair-group method with arithmetic mean) with 1,000 bootstrapping replicates to evaluate the consistency of the group, and Bionumerics was used to produce both the similarity matrix and dendrogram.

Acknowledgments

We wish to thank FAPEMA (Maranhão Research and Development Support Foundation) for financially supporting this research and UEMA (State University of Maranhão) for providing the studentships to the students.

Conflict of interest

The authors declare that there are no conflicts of interest.

References

1. Abriouel, H. *et al.* Differentiation and characterization by molecular techniques of *Bacillus cereus* group isolates from potto potto and dégué, two traditional cereal-based fermented foods of Burkina Faso and Republic of Congo. *J. Food Prot.* **70**, 1165–1173. <https://doi.org/10.4315/0362-028x-70.5.1165> (2007).
2. Balasubramanian, P. *et al.* Cloning and characterization of the crystal protein-encoding gene of *Bacillus thuringiensis* subsp. *yunnanensis*. *Appl. Environ. Microbiol.* **68**, 408–411. <https://doi.org/10.1128/AEM.68.1.408-411.2002> (2002).
3. Ben-Dov, E. *Bacillus thuringiensis* subsp. *israelensis* and its dipteran-specific toxins. *Toxins* **6**, 1222–1243. <https://doi.org/10.3390/toxins6041222> (2014).
4. Berry, C. *et al.* Complete sequence and organization of pBtoxis, the toxin-coding plasmid of *Bacillus thuringiensis* subsp. *israelensis*. *Appl. Environ. Microbiol.* **68**, 5082–5095. <https://doi.org/10.1128/AEM.68.10.5082-5095.2002> (2002).
5. Bourgouin, C., Delécluse, A., Ribier, J., Klier, A. & Rapoport, G. A *Bacillus thuringiensis* subsp. *israelensis* gene encoding a 125-kilodalton larvicidal polypeptide is associated with inverted repeat sequences. *J. Bacteriol.* **170**, 3575–3583. <https://doi.org/10.1128/jb.170.8.3575-3583.1988> (1988).

6. Bourgouin, C., Klier, A. & Rapoport, G. Characterization of the genes encoding the haemolytic toxin and the mosquitocidal delta-endotoxin of *Bacillus thuringiensis israelensis*. *Mol. Gen. Genet.* **205**, 390–397. <https://doi.org/10.1007/BF00338072> (1986).
7. Brumlik, M. J. *et al.* Genetic diversity among *Bacillus anthracis*, *Bacillus cereus* and *Bacillus thuringiensis* strains using repetitive element polymorphism-PCR. *Pol. J. Microbiol.* **53**, 215–225. (2004).
8. Crickmore, N. *et al.* Bacterial Pesticidal Protein Resource Center. <https://www.bpprc.org/>. (2020).
9. Da Silva, R. B. & Valicente F. H. Molecular characterization of *Bacillus thuringiensis* using rep-PCR. *SpringerPlus* **2**, 641. <https://doi.org/10.1186/2193-1801-2-641> (2013).
10. Dankocsik, C., Donovan, W. P. & Jany, C. S. Activation of a cryptic crystal protein gene of *Bacillus thuringiensis* subspecies *kurstaki* by gene fusion and determination of the crystal protein insecticidal specificity. *Mol. Microbiol.* **4**, 2087–2094. <https://doi.org/10.1111/j.1365-2958.1990.tb00569.x>. (1990).
11. De Bortoli, C. P. & Jurat-Fuentes J. L. Mechanisms of resistance to commercially relevant entomopathogenic bacteria. *Curr. Opin. Insect Sci.* **33**, 56–62. <https://doi.org/10.1016/j.cois.2019.03.007> (2019).
12. Dequech, S. T. B., Fiuza, L. M., Silva, R. F. P. & Zumba, R. C. Histopatologia de lagartas de *Spodoptera frugiperda* (Lep., Noctuidae) infectadas por *Bacillus thuringiensis*

aizawai e com ovos de *Campoletis flavicincta* (Hym., Ichneumonidae). *Cienc. Rural* **37**, 273–276. <https://doi.org/10.1590/S0103-84782007000100045> (2007).

13. Diaz-Mendoza, M., Bideshi, D. K. & Federici, B. A. A 54-kilodalton protein encoded by *pBtoxis* is required for parasporal body structural integrity in *Bacillus thuringiensis* subsp. *israelensis*. *J. Bacteriol.* 194:1562–1571. <https://doi.org/10.1128/JB.06095-11> (2011).

14. Drobniewski, F. A. & Ellar, D. J. Purification and properties of a 28-kilodalton hemolytic and mosquitocidal protein toxin of *Bacillus thuringiensis* subsp. *darmstadiensis* 73-E10-2. *J. Bacteriol.* **171**, 3060–3067. <https://doi.org/10.1128/jb.171.6.3060-3067.1989> (1989).

15. Ehling-Schulz, M. & Messelhäusser, U. *Bacillus* “next generation” diagnostics: moving from detection toward subtyping and risk-related strain profiling. *Front. Microbiol.* **4**, 1–8. <https://doi.org/10.3389/fmicb.2013.00032> (2013).

16. Federici, B. A., Park H.-W. & Bideshi, D. K. Overview of the basic biology of *Bacillus thuringiensis* with emphasis on genetic engineering of bacterial larvicides for mosquito control. *Open Toxinology J.* **3**, 154–171. <https://doi.org/10.2174/1875414701003010083> (2010).

17. Freitas, D. B. *et al.* Genotypic and phenotypic diversity of *Bacillus* spp. isolated from steel plant waste. *BMC Res. Notes* **1**, 92. <https://doi.org/10.1186/1756-0500-1-92> (2008).

18. Galvis, F. & Moreno, L. Y. Caracterización molecular mediante rep-PCR de aislados nativos de *Bacillus thuringiensis*, obtenidos de muestras de suelo. *Agron. Costarric.* **38**, 223–229. (2014).
19. García, K. *et al.* Variability of *Bacillus thuringiensis* strains by ERIC-PCR and biofilm formation. *Curr. Microbiol.* **70**, 10–18. <https://doi.org/10.1007/s00284-014-0675-8> (2015).
20. Ghribi, D., Zouari, N., Trabelsi, H. & Jaoua, S. Improvement of *Bacillus thuringiensis* delta-endotoxin production by overcome of carbon catabolite repression through adequate control of aeration. *Enzyme Microb. Tech.* **40**, 614–622. <https://doi.org/10.1016/j.enzmictec.2006.05.015> (2007).
21. Huerta, D., Acosta, O. & Chang, M. Diversidad genético molecular de cepas de *Bacillus thuringiensis* con potencial tóxico contra *Aedes aegypti*. *An. Fac. Med.* **73**, 205–210. (2012).
22. Ishii, T. & Ohba, M. Diversity of *Bacillus thuringiensis* environmental isolates showing larvicidal activity specific for mosquitoes. *J. Gen. Microbiol.* **139**, 2849–2854. <https://doi.org/10.1099/00221287-139-11-2849> (1993).
23. Ishii, S. & Sadowsky, M. J. Applications of the rep-PCR DNA fingerprinting technique to study microbial diversity, ecology and evolution. *Environ. Microbiol.* **11**, 733–740. <https://doi.org/10.1111/j.1462-2920.2008.01856.x> (2009).

24. Ito, T., Ikeya, T., Sahara, K., Bando, H. & Asano, S. Cloning and expression of two crystal protein genes, *cry30Ba1* and *cry44Aa1*, obtained from a highly mosquitocidal strain, *Bacillus thuringiensis* subsp. *entomocidus* INA288. *Appl. Environ. Microbiol.* **72**, 5673–5676. <https://doi.org/10.1128/AEM.01894-05> (2006).
25. Jurat-Fuentes, J. L., Heckel, D. G. & Ferre, J. Mechanisms of resistance to insecticidal proteins from *Bacillus thuringiensis*. *Ann. Rev. Entomol.* **66**, 121–140. <https://doi.org/10.1146/annurev-ento-052620-073348> (2021).
26. Katara, J., Deshmukh, R., Singh, N. K. & Kaur, S. Molecular typing of native *Bacillus thuringiensis* isolates from diverse habitats in India using REP-PCR and ERIC-PCR analysis. *J. Gen. Appl. Microbiol.* **58**, 83–94. <https://doi.org/10.2323/jgam.58.83> (2012).
27. Koeuth, T., Versalovic, J. & Lupski, J. R. Differential subsequence conservation of interspersed repetitive *Streptococcus pneumoniae* BOX elements in diverse bacteria. *Genome Res.* 5:408–418. <https://doi.org/10.1101/gr.5.4.408> (1995).
28. Kumar, A., Kumar, A. & Pratush, A. Molecular diversity and functional variability of environmental isolates of *Bacillus* species. *SpringerPlus* **3**, 312. <https://doi.org/10.1186/2193-1801-3-312> (2014).
29. Lee, H.-K. & Gill, S. S. Molecular cloning and characterization of a novel mosquitocidal protein gene from *Bacillus thuringiensis* subsp. *fukuokaensis*. *Appl. Environ. Microbiol.* **63**, 4664–4670. <https://doi.org/10.1128/aem.63.12.4664-4670.1997> (1997).

30. Lee, J. *et al.* A RAPD-PCR method for the rapid detection of *Bacillus cereus*. *J. Microbiol. Biotechnol.* **21**, 274–276. <https://doi.org/10.4014/jmb.1008.08031> (2011).
31. Liang, H. *et al.* Characterization of *cry2-type* genes of *Bacillus thuringiensis* strains from soil-isolated of Sichuan basin, China. *Braz. J. Microbiol.* **42**, 140–146. <https://doi.org/10.1590/S1517-83822011000100018> (2011).
32. Lima, A. S. G., Guidelli, A. M., Abreu, I. L. & Lemos, M. V. F. Identification of new isolates of *Bacillus thuringiensis* using rep-PCR products and δ -endotoxin electron microscopy. *Genet. Mol. Biol.* **25**, 225–229. <https://doi.org/10.1590/S1415-47572002000200017> (2002).
33. Lima, M. P. L., Oliveira, J. V., Gondim Junior, M. G. C., Marques, E. J. & Correia, A. A. Bioatividade de formulações de NIM (*Azadirachta indica* A. Juss, 1797) e de *Bacillus thuringiensis* subsp. *aizawai* em lagartas de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). *Ciênc. Agrotec.* **34**, 1381–1389. <https://doi.org/10.1590/S1413-70542010000600004> (2010).
34. Lobo, K. S. *et al.* Isolation and molecular characterization of *Bacillus thuringiensis* found in soils of the Cerrado region of Brazil, and their toxicity to *Aedes aegypti* larvae. *Rev. Bras. Entomol.* **62**, 5–12. <https://doi.org/10.1016/j.rbe.2017.11.004> (2018).
35. Machado, D. H. B. *et al.* Molecular characterization of *Bacillus thuringiensis* strains to control *Spodoptera eridania* (Cramer) (Lepidoptera: Noctuidae) population. *Rev. Bras. Entomol.* **64**, e201947. <https://doi.org/10.1590/1806-9665-RBENT-2019-47> (2020).

36. Magda, S. & Moharam, M. E. Comparing the effect of seven isolated *Bacillus thuringiensis* against *Tuta absoluta* infesting in laboratory and field condition. *Int. J. Sci. Res.* **4**, 458–462. <https://doi.org/10.21275/v4i11.01111502> (2015).
37. Mishra, P. K. *et al.* Genetic diversity and functional characterization of endophytic *Bacillus thuringiensis* isolates from the North Western Indian Himalayas. *Ann. Microbiol.* **67**, 143–155. <https://doi.org/10.1007/s13213-016-1244-0> (2017).
38. Moar, W. J., Masson, L., Brousseau, R. & Trumble, J. T. Toxicity to *Spodoptera exigua* and *Trichoplusia ni* of individual P1 protoxins and sporulated cultures of *Bacillus thuringiensis* subsp. *kurstaki* HD-1 and NRD-12. *Appl. Environ. Microbiol.* **56**, 2480–2483. <https://doi.org/10.1128/aem.56.8.2480-2483.1990> (1990).
39. Mohkam, M., Nezafat, N., Berenjian, A., Mobasher, M. A. & Ghasemi, Y. Identification of *Bacillus* probiotics isolated from soil rhizosphere using 16S rRNA, *recA*, *rpoB* gene sequencing and RAPD-PCR. *Probiotics Antimicrob. Proteins* **8**, 8–18. <https://doi.org/10.1007/s12602-016-9208-z> (2016).
40. Ohba, M., Saitoh, H., Miyamoto, K., Higuchi, K. & Mizuki, E. *Bacillus thuringiensis* serovar *higo* (flagellar serotype 44), a new serogroup with a larvicidal activity preferential for the anopheline mosquito. *Lett. Appl. Microbiol.* **21**, 316–318. <https://doi.org/10.1111/j.1472-765x.1995.tb01068.x> (1995).
41. Ohgushi, A., Saitoh, H., Wasano, N., Uemori, A. & Ohba, M. Cloning and characterization of two novel genes, *cry24B* and *slorf2*, from mosquitocidal strain of

Bacillus thuringiensis serovar *sotto*. *Curr. Microbiol.* **51**, 131–136.
<https://doi.org/10.1007/s00284-005-7529-3> (2005).

42. Özkan, M., Dilek, F. B., Yetis, Ü. & Özcengiz, G. Nutritional and cultural parameters influencing antidipteran delta-endotoxin production. *Res. Microbiol.* **154**, 49–53.
[https://doi.org/10.1016/s0923-2508\(02\)00006-2](https://doi.org/10.1016/s0923-2508(02)00006-2) (2003).

43. Polanczyk, R. A., Van Frankenhuyzen, K & Pauli, G. The American *Bacillus thuringiensis* based biopesticides market (eds. Fiuza, L. M., Polanczyk, R. A. & Crickmore, N.) 173–184 (Springer International Publishing, 2017).

44. Rademaker, J. L. W., Aaris, H. J. M. & Vinuesa, P. Molecular typing of environmental isolates (eds. Osborn, A. C. & Smith, C. J.) 97–134 (Taylor & Francis Group, 2005).

45. Rademaker, J. L. W. *et al.* Comparison of AFLP and rep-PCR genomic fingerprinting with DNA–DNA homology studies: *Xanthomonas* as a model system. *Int. J. Syst. Evol. Microbiol.* **50**, 665–677. <https://doi.org/10.1099/00207713-50-2-665> (2000).

46. Rademaker, J. L. W., Louws, F. J., Versalovic, J. & de Bruijn, F. J. Characterization of the diversity of ecological important microbes by rep-PCR genomic fingerprinting. (eds. Kowalchuk, G. A., de Bruijn, F. J., Head, I. M., Akkermans, A. D. & van Elsas, J. D.) 611–644 (Kluwer Academic Publishers 2004).

47. Rajchanuwong, P., Chanpaisaeng, J. & Kaewsompong, S. Characterization and toxicity of *Bacillus thuringiensis* serovar *chanpaisis* (H46): A serovar from Thailand.

Songklanakarin J. Sci. Technol. **41**, 804–812.

<https://doi.org/10.14456/sjst-psu.2019.103> (2019).

48. Reyes-Ramirez, A. & Ibarra, J. E. Fingerprinting of *Bacillus thuringiensis* type strains and isolates by using *Bacillus cereus* group-specific repetitive extragenic palindromic sequence-based PCR analysis. *Appl. Environ. Microbiol.* **71**, 1346–1355. <https://doi.org/10.1128/AEM.71.3.1346-1355.2005> (2005).

49. Ruiz, L. M., Segura, C., Trujillo, J. & Orduz, S. In vivo binding of the Cry11Bb toxin of *Bacillus thuringiensis* subsp. *medellin* to the midgut of mosquito larvae (Diptera: Culicidae). *Mem. Inst. Oswaldo Cruz* **99**, 73–79. <https://doi.org/10.1590/S0074-02762004000100013> (2004).

50. Sauka, D. H., Basile, J. I. & Benintende, G. Evidence of *Bacillus thuringiensis* intra-serovar diversity revealed by *Bacillus cereus* group-specific repetitive extragenic palindromic sequence-based PCR genomic fingerprinting. *J. Mol. Microbiol. Biotechnol.* **21**, 184–190. <http://doi.org/10.1159/000335532> (2011).

51. Shin, B.-S. *et al.* Distribution of *cryV*-type insecticidal protein genes in *Bacillus thuringiensis* and cloning of *cryV*-Type genes from *Bacillus thuringiensis* subsp. *kurstaki* and *Bacillus thuringiensis* subsp. *entomocidus*. *Appl. Environ. Microbiol.* **61**, 2402–2407. <http://doi.org/10.1128/aem.61.6.2402-2407> (1995).

52. Soares-da-Silva, J. *et al.* Molecular characterization of the gene profile of *Bacillus thuringiensis* Berliner isolated from brazilian ecosystems and showing pathogenic

activity against mosquito larvae of medical importance. *Acta Trop.* **176**, 197–205.
<http://doi.org/10.1016/j.actatropica.2017.08.006> (2017).

53. Subbanna, A. R. N. S. *et al.* Interspecies diversity of *Bacillus thuringiensis* isolates native
 from North Western Indian Himalayas. *J. Environ. Biol.* **39**, 306–313.
<http://doi.org/10.22438/jeb/39/3/MRN-574> (2018).

54. Thorne, L. *et al.* Structural similarity between the Lepidoptera- and Diptera-specific
 insecticidal endotoxin genes of *Bacillus thuringiensis* subsp. "*kurstaki*" and "*israelensis*".
J. Bacteriol. **166**, 801–811. <http://doi.org/10.1128/jb.166.3.801-811.1986> (1986).

55. Valtierra-de-Luis, D., Villanueva, M., Lai, L., Williams, T. & Caballero, P. Potential of
 Cry10Aa and Cyt2Ba, two minority δ -endotoxins produced by *Bacillus thuringiensis* ser.
israelensis, for the control of *Aedes aegypti* larvae. *Toxins* **12**, 355.
<https://doi.org/10.3390/toxins12060355> (2020).

56. Van Frankenhuyzen, K. Insecticidal activity of *Bacillus thuringiensis* crystal proteins. *J.*
Inverteb. Pathol. **101**, 1–16. <https://doi.org/10.1016/j.jip.2009.02.009> (2009).

57. Versalovic, J., Koeuth, T. & Lupski, J. R. Distribution of repetitive DNA sequences in
 eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Res.* **19**,
 6823–6831. <https://doi.org/10.1093/nar/19.24.6823> (1991).

58. Versalovic, J., Schneider, M., de Bruijn, F. J. & Lupski, J. R. Genomic fingerprinting of bacteria using repetitive sequence-based polymerase chain reaction. *Methods Mol. Cell Biol.* **5**, 25–40. (1994).
59. Vieira-Neta, M. R. A. *et al.* Estirpe de *Bacillus thuringiensis* da restinga, tóxico ao *Aedes* (*Stegomyia*) *aegypti* (Linnaeus) (Diptera, Culicidae). *Braz. J. Biol.* **81**, 872–880. <https://doi.org/10.1590/1519-6984.228790> (2021).
60. Wirth, M. C., Delécluse, A., Federici B.A. & Walton, W. E. Variable cross-resistance to Cry11B from *Bacillus thuringiensis* subsp. *jegathesan* in *Culex quinquefasciatus* (Diptera: Culicidae) resistant to single or multiple toxins of *Bacillus thuringiensis* subsp. *israelensis*. *Appl. Environ. Microbiol.* **64**, 4174–4179. <https://doi.org/10.1128/AEM.64.11.4174-4179.1998> (1998).
61. Zorzetti, J. *et al.* Isolation, morphological and molecular characterization of *Bacillus thuringiensis* strains against *Hypothenemus hampei* Ferrari (Coleoptera: Curculionidae: Scolytinae). *Rev. Bras. Entomol.* **62**, 198–204. <https://doi.org/10.1016/j.rbe.2018.07.002> (2018).

Figures

Figure 1

Rep-PCR fingerprint showing the amplification of the BOX (a), ERIC (b), REP (c), MB1 (d) and GTG5 (e) molecular markers in 30 isolates of *Bacillus thuringiensis* and seven standard subspecies. MM: Molecular marker; Bta: *B. thuringiensis* subsp. *aizawai*; Bte: *B. thuringiensis* subsp. *entomocidus*; Btf: *B.*

thuringiensis subsp. fukuokaensis; Bti: *B. thuringiensis* subsp. israelensis; Btk: *B. thuringiensis* subsp. kurstaki; Bts: *B. thuringiensis* subsp. sotto; Bty: *B. thuringiensis* subsp. yunnanensis; BtMA: *Bacillus thuringiensis* from Maranhão; NC: negative control.

Figure 2

Dendrogram produced by cluster analysis (UPGMA) based on the Jaccard coefficient by using the BOX, ERIC, REP, MB1 and GTG5 molecular markers and molecular fingerprint of 30 isolates of *Bacillus thuringiensis* from diverse locations in Maranhão and seven standard subspecies of *B. thuringiensis*.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [ArticleFERNANDESetal2021.Tables.pdf](#)