Maternal Social Support and Determinants of Health Facility Delivery in Rural Ethiopia

Abebe Mamo (✉ abemam24@gmail.com)

Faculty of Public Health, Institute of Health, Jimma University <https://orcid.org/0000-0002-7118-1932>

Muluemebet Abera

Jimma University

Lakew Abebe

Jimma University

Nicole Bergen

Ottawa University

Shifera Asfaw

Jimma University

Gebeyehu Bulcha

Jimma zone health office

Yisalemush Asefa

Jimma University

Endale Erko

Addis Ababa City Administration Health Bureau

Kunuz Haji Bedru

Jimma Zone Health office

Mihiretu Lakew

Jimma Zone Health office

Jaameeta Kurji

Ottawa University

Manisha A. Kulkarni

Ottawa University

Ronald Labonté

Ottawa University

Zewdie Birhanu

Jimma University

Sudhakar Morankar

Jimma University

Keywords: Maternal and child health, social support, health facility delivery, Ethiopia.

Posted Date: August 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-819229/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Maternal mortality continues to decrease in the world but remains the most important health problems in low-income countries. Although evidence indicates that social support is important factor influencing health facility delivery, it has not been extensively studied in Ethiopia. Therefore, this study aimed to assess social support in influencing health facility childbirth in Ethiopia.

Methods: A cross-sectional survey data on 3304 women aged 15–49 years from an Innovating for Maternal and Child Health in Africa project in three districts of Ethiopia, were analyzed, including data on social support and important factors. Using multivariate logistic regression, we examined the association between health facility birth, social support, and included variables. Adjusted odds ratios with 95% confidence intervals were used to identify statistically significant associations at 5% alpha level.

Result: Overall, 46.9% of women delivered at health facility in their last pregnancy. Average travel time from closest health facility (AOR: 1.51, 95% CI 1.21 to 2.90), mean perception score of health facility use (AOR: 1.83, 95% CI 1.44 to 2.33), involvement in final decision to identify their place of childbirth (AOR: 2.12, 95% CI 1.73 to 2.58) had significantly higher odds of health facility childbirth. From social support variables, women who perceived there were family members and husband to help them during childbirth (AOR: 3.62, 95% CI 2.74 to 4.79), women who received continuous support (AOR: 1.97, 95% CI 1.20 to 3.23), women with companions for facility visits (AOR: 1.63, 95% CI 1.34 to 2.00) and women who received support from friends (AOR: 1.62, 95% CI 1.16 to 3.23) had significantly higher odds of health facility childbirth.

Conclusions: Social support was critical to enhance health facility delivery, especially if women's close ties help facility delivery. An intervention to increase facility delivery uptake should target not only the women's general social supports, but also continuous support during childbirth from close ties including family members and close friends as these are influential in place of childbirth and also actions that increase women's healthcare decision could be effective in improving health facility delivery.

Background

Pregnancy and childbirth are important moments in the lives of our families and our communities, and most pregnancies end with the birth of live babies to healthy mothers. However, for some, pregnancy and childbirth is an unsafe journey and it is a time of pain, fear, suffering, and even death (1–3). In 2017, the World Health Organization (WHO) estimated that 295 000 women die annually from preventable problems during and following pregnancy and childbirth (4). To address these devastating maternal and child health (MCH) issues, WHO and allies identified the reduction of maternal mortality as a public health priority, and further agreed on a global target for a maternal mortality ratio (MMR) of less than 70/100,000 live births by 2030, with no single country having an MMR greater than 140(4). However, worldwide more than half a million women and girls die each year and almost all (94%) of maternal deaths occur in low and middle-income countries (LMICs), with sub-Saharan African (SSA) countries

accounting for around two thirds of the global burden (62%) (4–7). Most maternal deaths occur from preventable and treatable conditions associated with pregnancy and childbirth, including hypertensive disorders, postpartum hemorrhaging, sepsis, complications from prior childbirth or unsafe abortions, and indirect conditions such as malaria and anemia, and the majority of which occur during the first 24 hours of birth (8).

According to The Lancet newborn survival series, the most efficient and cost-effective strategy to reduce maternal mortality and newborn deaths is for every woman to use health facility delivery and to be assisted by a skilled birth attendant (2). Success of this strategy is dependent not only upon the availability health facilities but also the use of these facilities during pregnancy and childbirth. Although, studies reported around 67% of newborn deaths could be prevented with health facility delivery, data from recent studies suggest uptake is low even in settings where services are available. In Africa, including Ethiopia over half of all births occur outside of health facilities (6–10), and the utilization of the existing maternal and child health services is very low (11). The health status of Ethiopia by most indicators ranks amongst the worst globally, even when compared to those for most SSA countries (12–15). According to the 2019 Ethiopian mini Demographic and Health Survey (EDHS), 72% of births among urban populations in Ethiopia occurred in health facilities, compared to only 40% of births among rural populations where over 80% of Ethiopians living, which highlights the need to improve use of facility delivery in underutilization (9).

Major factors that have been studied as contributing factors to the low utilization of existing maternal health services in SSA countries like Ethiopia are related to distance to a facility, means of transportation, maternal socio-demographics (e.g. age, parity, education, and wealth), health decision-making autonomy, and perceived needs/benefits of facility birth and attitude (6, 16). Most of these factors have been studied extensively in Ethiopia and elsewhere (10, 15, 16). Further, recent studies examined the influence of social support during pregnancy and childbirth on birth weight, length of gestation including fewer pregnancy complications for women with high levels of stress, reduce fear/anxiety, improve outcomes, and stimulate healing but not for those with low levels of stress (17, 18), however, these studies do not fully account for the interpersonal, family and community interactions thought to facilitate or constrain service use as well as decision making; a social context in which individual determinants are known to operate. For example, the importance of social networks in Ethiopia have long recognized and exhibit different kinds of social interaction and rely more on family and friend ties, and often consult family and friends in case of illness and service use (17–20). However, maternal social support as direct determinants of health facility delivery including the different dimensions of maternal social support and its complex interactions have not been extensively studied in Ethiopia. Therefore, the aim of the present study was to assess the maternal social support and related factors on health facility childbirth, and investigating the effects of social support on health facility childbirth may provide insights into additional areas of interventions to increase uptake in facility delivery.

Social support including family and community contexts are mini universes of complex social, political, associational, economic, power and cultural dynamics, providing a different direction for encouraging the

use of health services and facilitating behavior change than in a health facility (21–24). Social support is considered a complex and dynamic process that often used in a broad sense, including social networks and social capital, which also involves individuals and their social networks, working to satisfy their needs, provide and complement the resources they have, and thus cope with new situations (7, 17). One frequent criticism of research in this area is the lack of consensus about social support in terms of its definition and how best to measure it (7, 17–19). However, many scholars and theorists conceptualized social support as a contextual phenomenon in the sense that it is both individual and a community characteristic reflecting the daily interaction between social relationships/neighbors that may benefit health through interpersonal trust and norms of mutual aid, promoting collective efficacy and neighborhood cohesion (1, 25).

Further, studies have specified the common dimensions or forms of social support including functional and structural supports. Structural social support refers to the structure and quantity of social relationships, such as the size/density of networks and the frequency of interaction. Functional Social support, in contrast, refers to the function and quality of social relationships, such as perceived availability of help and the resources exchanged among individuals in a relationship such as informational support (advice/suggestions), instrumental support (aid or assistance), emotional support (empathy, care and trust), and accompaniment (7, 10, 26). Different studies including systematic reviews reported the association between social support and access to health care behaviors(7, 16, 17), and absence of social support was associated with increased maternal morbidity and complication (17, 18, 25).

Furthermore, studies in Ethiopia, Kenya and Bangladesh found that social support from a spouse or partner and a social network of family and friends has been found to influence women's decisions regarding obtaining prenatal care, childbirth, and breastfeeding (17, 18, 27–29), and other studies also found that female relatives and friends accompanying laboring women to maternity units (13, 15, 18), and the presence of a female relative during labor, is associated with improved labor outcomes (30). Various studies also acknowledged the diversity and uniqueness of each social support characteristics as a major source of strength, and identified that people who receive higher levels of social support tend to be less stressed and have better health status than more socially isolated people (1, 25, 31). Moreover, maternal social support can also be received in different contexts which leads to different healthcare seeking behaviors (32). For example, studies in SSA countries reported that pregnant women initiate ANC early and felt the importance of health facility childbirth, only when they are sick or experience any pregnancy-related health problems(32–35). Early initiators emphasized that they go to the ANC and started to think health facility childbirth and need different supports from family and relatives if the pregnancy is under threat and or they are sick in the following comments. Moreover, if pregnant women perceived pregnancy to be a normal health condition and with previous successful home delivery, they felt seeking health care or giving birth at facilities was unnecessary; hence develop positive attitude towards home delivery and placed a low value on using antenatal care and health facility delivery (32–34). Thus, maternal social support plays a crucial role in improving maternal health outcomes, first, in

recognizing the need for health services, and, second, in facilitating or constraining the use of those services (10, 30, 36).

About one-third of currently married women in Ethiopia, however, have no final say concerning their own healthcare decisions, with recent studies finding that 32.3% report that their husbands make the final decision (15, 27). According to the 2016 EDHS, more than 70% of women report social barriers to accessing MCH care, including getting money for advice or treatment (55%), long distance to a health facility (50%), lack of accompaniment (42%), and getting permission to go for treatment (32%) (6). Furthermore, in traditional and patriarchal societies like Ethiopia, where restrictions are placed on a woman's freedom of movement and contact with unrelated men, the influence of the social environment through maternal social support factors may be important factors in determining whether MCH care, particularly health facility delivery is used or not (15, 37). Therefore, assessing the distinctions among different dimensions and complex interactions of social support factors may provide a comprehensive view of how social support influences health facility delivery, alongside other important factors.

Methods

Data source

The data for this study was obtained from a household survey conducted prior to delivering interventions (baseline survey) being evaluated in three districts in Jimma Zone, Ethiopia using a cluster randomized controlled trial (RCT) design. The intervention components under study in the cluster RCT were upgraded maternity waiting homes and local leader training that were hypothesized to improve levels of health facility delivery. Women reporting a pregnancy outcome (livebirth, stillbirth or abortion) up to 12 months prior to commencement of the baseline survey in January 2017 were eligible to be interviewed. A two-stage sampling strategy was employed. First, 24 primary health care units (PHCUs) within the three districts were randomly selected for the trial, then, 160 women per PHCU were randomly selected from community-based lists of pregnant women generated as part of health post records. The questionnaire, administered in a face-to-face interview with women by trained research assistants in a private space at women's homes. Questionnaire contained modules on socio-demographics, maternal health service use, danger sign awareness, social support, decision-making practices and sources of information. At baseline, interviews were conducted with 3,784 (~99%) women of the 3,840 targeted sample size calculated to meet trial requirements. A protocol with detail methodology of the trial has been published previously and (14) and the trial identifier is NCT03299491.

Study setting

Jimma Zone is located 356 km from Addis Ababa in Southwest Ethiopia, Oromia National Regional State. Jimma Zone has 21 districts, and 42 urban and 513 rural kebeles.^[1] The total population of Jimma Zone is estimated to be 3.2 million with the majority of the population living in rural areas (38). The three

study districts selected for the cluster RCT were Gomma, Seka Chekorsa and Kersa, which had populations ranging from 180 000 to 270 000 in 2016 (39).

Design and analytic approach

A community based cross-sectional study design was used to assess association between health facility delivery and social support variables described in subsections below. The sample for this work included women of reproductive age group (15–49 years) who had uncomplicated pregnancy and delivery resulting in a live birth during the year preceding the survey. Women who faced pregnancy and delivery related complications such as bleeding, pre-eclampsia, eclampsia, mental health problems, infections, severe headache, water breaks without labour, persistent vomiting and related complications were more likely to receive immediate supports from family or relatives or close friends . Therefore, we excluded women who reported experiencing these complications during antepartum or intrapartum periods. Among the 3784 women surveyed at baseline, 480 reported experiencing antepartum or intrapartum complications and were excluded from the final analysis. Thus, data from 3304 respondents were analyzed.

The primary dependent variable was the place of delivery of the most recent child, dichotomized as health facility or not health facility (home, in ambulance, on the road). Health facility delivery included those at the health center, hospitals and private hospital/clinic.

Variables of interest

Outcome variable

The primary dependent variable was the place of delivery of the latest child, dichotomized as health facility or not health facility (home, in ambulance, on the road). Health facility deliveries included those at the health center, hospitals, and private hospital/clinic.

Independent variables

The main independent variables of interest included perceived social support, structural and functional social support characteristics. In addition to main variables we also included numerous independent variables that might be associated with place of delivery including age, education level, occupation, marital status, age at first pregnancy, household income, availability of health facility, types of health facility, the time required to reach the nearest delivery facility, number of family members, social group participation, participated a programme that promote MCH cares, attitude towards health facility delivery, perception of health care services provision, planning to give birth at health facility, sources of information and decision-making autonomy (10,13,15,16,29,37). So, we developed social support

questions on the basis of the result of the pre-intervention study which was conducted to explore major areas of maternal social supports, roles of community health actors and common sources of information in the same settings (15,33), and adopted social support questions from existing measures and review of the network literature (10,36–38). So, we included key social support domains such as perceived social support, structural support and functional supports.

The perceived social support variable consisted of two items; first, women were asked whether they had someone that they could depend on during their last pregnancy and childbirth when they were in need. Responses were recorded as 1 = yes and 0 = no, and if the response was 'yes', it followed by asking the relationship of the women with the perceived individual (s) to provide supports. Responses were lists of options from 1 to 11 including husband/partner, parents, siblings, friends/neighbours, in-laws, religious groups, women developmental army [WDA], health workers, health extension workers (HEWs) and other to specify. Finally, these responses were collapsed into four categories: 1. Partner/family members (include partner, children or any family member in the household), 2. Relatives (like sibling, parent and in-law), 3. Friends/neighbors, 4. Health workers (midwives, nurses, health extension workers). Functional social support had five domains, including a) practical help with routine activities, b) material aid (financial or in-kind assistance), c) emotional support (empathy, care and trust), d) informational support (advice/suggestions), and e) accompaniment supports. For each section, subjects were first asked whether they had received that type of functional support in the last pregnancy and childbirth. For example, to assess practical support, women were asked, "for your last pregnancy and childbirth, did you receive any practical help with routine activities?" (Example: help with child care, house chores, food preparation, cattle herding, etc). Responses were recorded as (1=yes and no = 0). If women answered yes, it was followed by "when did you receive this help?" Responses were recorded as 1-during pregnancy, 2-during labour and childbirth, and 3- during postpartum. This question was also followed by structural support [types of network members] questions "from whom did you receive this help?" Responses were lists of options from 1 to 11 including husband/partner, parents, siblings, friends/neighbours, in-laws, religious groups, WDA, health workers, HEW and other to specify. These responses were collapsed into four categories: 1. Partner/family members (include partner, children or any family member in the household), 2. Relatives (like sibling, parent and in-law), 3. Friends/neighbors, 4. Health workers (midwives, nurses, health extension workers). To provide an overall support size of each woman received during pregnancy, labour, and delivery, receipt of support was counted with 0 indicating no support received, 1 indicating received one type of support, 2 indicating received two types of support, and so on. Then, the total number of received supports was summed and divided by the number of maximum received supports, where 1 is the maximum score or women received highest number of different supports over the course of pregnancy and delivery. Density was conceptualized as the number of supporters/ ties that the women had during pregnancy and delivery. so, the density for each of the five types of support was counted with 0 indicating do not receive any support from anyone, 1 indicating support received from one person, 2 indicating any support received from two persons, and so on. Therefore, the total number of ties/supporters was summed and divided by the number of maximum supporters, where 1 is the maximum score indicated that women have highest ties of social support over

the course of pregnancy and delivery. For receiving continuous social support, we created a variable from the frequency of periods the women received all types of social supports from network members. A simple summative score from one open ended item, “when did you receive this help”, with the responses of during pregnancy, during childbirth, and during postnatal period. Finally, those who reported they received supports during both pregnancy and delivery were considered as receiving continuous support/resulting in a final continuous support score with values between 0 (not continuous) and 1 (received continuous support).

From socio-economic and healthcare variables, educational level was assessed based on women’s responses on highest level of education completed, and categorized as; 1) no formal education, 2) primary school, 3) high school and 4) above high school. Women’s responses on their primary occupation were collapsed into a nominal variable to reflect the main occupations listed; 1) housewife, 2) farmer, 3) traders, 4) others (private workers, unemployed, employee and student). Household income was created using information on the total household income of last year, which were then grouped into quintiles; the first quintile (lowest quintile) corresponded to the lowest household income and the fourth quintile corresponding to the highest household income. Travel time to closest health facility was assessed as continuous variable based on women’s estimated time (one way walk in hours) required reaching the nearest health facility. Women’s responses on their final decision/ involvement in decision making on place of childbirth was assessed based a question whether they made decisions on their own, jointly with someone else or someone else. Then response was categorized as: 1= involved (if decision made by women/their own and jointly with someone else), 0 = not involved (if the decision made by someone else or not included the women). Attitude towards health facility delivery was measured through seven items using Likert scale [with response ranged from 1= strongly disagree to 5= strongly agree] resulted with final score ranged from 7 to 35. Finally, the mean score was calculated with minimum score of 0.20 to maximum score of 1, in which the higher scores indicated having high attitude towards health facility delivery. Perception on healthcare provision was assessed based on evaluation scores that ranged from 1-5 to evaluate the service provided in the closest health facility; with response 5= very good and 1 = very poor. Finally, the mean score was calculated with minimum score of 0.32 to maximum score of 1, in which the higher scores indicated having highest perception on healthcare provision. Social group membership was assessed based on women’s responses on membership of any social group, organization or association, and the responses were categorized as (1=yes, 0=no). Participation on programme that promote MCH cares was assessed based on women’s responses on participation on any programme that promote MCH cares, and categorized as (1=yes, 0=no). Age and age at first pregnancy were assessed based on interval age in years and number of household members was also assessed on interval numbers of household members. Whether the women planned to give birth at health facility was assessed by binary responses (1=yes, 0=no). Antenatal care use was assessed based on the number of visits during pregnancy as ANC visits and collapsed to binary responses as (1= ANC four or plus visits and 0= none/ less than four visits).

Analysis

First, descriptive analysis was carried out to summarize the socio-demographic and health service utilization characteristics of the respondents using frequencies and proportions. We also conducted bivariate and multivariate analyses of the association between each variable included in the study and health facility birth. We also ran separate logistic regression models to test two-way interactions, specifically whether structural social support moderated the relationship between functional social support and facility birth. To identify variables associated with health facility childbirth, multivariable regression analysis was used. All independent variables and interaction terms that associated at bivariate levels (occupation, household income, number of household members, travel hour from closer health facility, participated on role that promotes maternal healthcare, involved in decisions on place of childbirth, attitude to facility delivery and perception on healthcare provision) were included in multivariable regression. From maternal social support variables perceived partner/family members to help women during labour and childbirth, women who received practical support, accompanied to health facility during last child birth, size of received supports, received continuous supports, received from partners/family members, from relatives/siblings and from friends/neighbors were also included in multivariate analysis. From two-way interactions or correlation between social support variables all tested interactions between each structural social support and functional social support variable, were not significantly associated with facility delivery. Age was not included because it is highly correlated with number of family members ($r = .82$), and from perceived social support, perceived relatives to help women during labour and childbirth was highly correlated with perceived partner/family members to help them ($r = .89$). All analyses were performed using SPSS software version 20.0 (SPSS Inc., 2008), and an alpha level of 0.05 was selected.

Results

Socio-demographic factors and MCH service utilization

Table 1 showed the descriptive analysis of data from 3304 eligible respondents. Participants' age ranged from 15–47 (with mean 28 and $SD \pm 6.0$) years, the majority (30.8%) belongs to age category of 25 to 29 years and 5.6% were younger than 19 years. The majority of women were married (97.9%), 56.2% of women had no formal education and 77.1% of them were housewife. In relation to household income, a larger proportion belonged to first (poorest) quartile (31.2%), and the majority of women's partners were farmers (72.7%) (table 1).

Healthcare related variables and health facility delivery

The majority of women (76.2%) used four or more antenatal care services and 46.9% of women delivered at health facility. Most women mentioned they had health facilities closer to their community (95.7%) with the average travel time (one way walk) from home to the closest health facility was around 1 hour. About 46% of women mentioned that they planned to give birth at the current place of delivery. Half of the women (50%) participated in a programme that promotes MCH cares and only 17.2% of women

mentioned that they were a member of any social group. Almost half of the women (49%) mentioned that they involved in deciding place of delivery for their last childbirth. The mean score of age at first pregnancy, average number of household members, mean score of attitude towards health facility delivery and perception on healthcare provision were: 19.04 (± 2.61), 3.72 (± 2.20), 0.38 (± 0.28) and 0.26 (± 0.29), respectively (table2).

Maternal social support and health facility Delivery

From women who delivered at health facility, 45.2% of them perceived that they had someone to help them during their last pregnancy and childbirth. Most of the women (81.4%) perceived their partners including family members to help them during labour and childbirth, while 58.3% of women perceived friends/neighbors and 55.2% any health workers (55.2) to help them during their last childbirth. In regard to functional social support, women who had health facility birth had a higher mean size of received supports (0.6 ± 0.2), a higher mean of received continuous support (0.4 ± 0.2). Additionally, from women who delivered at health facility, 57.7% of them received practical support with routine activities, 54.7% of them accompanied during last childbirth, 47.9% of women received emotional support, 48.2% of women received material support and 45.2% of women received informational support from different network members. Further, women identified they received these supports from partner/family members, relatives/siblings, friends/neighbors and health workers with the proportions of 55.3%, 55.2%, 58.3% and 45.6% respectively. The number of network members (density of ties) among women who delivered at health facility consisted of the same mean score (0.4 ± 0.2) with women who delivered out of health facility (table 3).

Multivariate logistic regression analysis of social support and related factors with health facility delivery

After controlling for associated variables at bivariate levels, three socio-demographic and healthcare related factors including travel time from closest health facility, perception towards health facility use and decision in using health facility delivery resulted in statistically significant associations with health facility delivery. Therefore, the multivariate analysis showed that per unit increase in total score of travel time to health facilities the likelihood of health facility delivery also increased by 1.5 (AOR: 1.51, 95% CI 1.21 to 2.90), and women who were involved in the final decision to identify their place of childbirth had twice the odds of using health facility delivery than women who did not deliver at a health facility (AOR: 2.12, 95% CI 1.73 to 2.58). Similarly, this study identified that with each unit increase in total score of perception towards the benefits of health facility use the likelihood of health facility delivery also increased (AOR: 1.83, 95% CI 1.44 to 2.33).

Further, from social support variables our study revealed that women's perception that there were partner or family members to help them during labour and childbirth; women who received continuous support,

women who were accompanied during childbirth, and women who received any support from their friends were all significantly associated with health facility delivery.

Women who perceived there were partner/family members to help them during labour and childbirth were more than three times more likely to have a health facility delivery than women who did not perceived there were partner/family members to help (AOR: 3.62, 95% CI 2.74 to 4.79). Women who described having companions to accompany them for health facility visits during labour and delivery had 1.6 times the odds of having used health facility delivery than women who did not this kind of support (AOR: 1.63, 95% CI 1.34 to 2.00).

Women who received continuous support during pregnancy, labour and delivery had almost twice the odds of having used a health facility for delivery than women who did not deliver at a health facility (AOR: 1.97, 95% CI 1.20 to 3.23), and women who received any type of social support from friends/neighbors during pregnancy and delivery were 1.6 times more likely to use health facility delivery than women who did not deliver at a health facility (AOR: 1.62, 95% CI 1.16 to 3.23) (Table 4). We also ran separate logistic regression models to test two-way interactions, specifically whether density of ties moderated the relationship between each network members (family members, relatives, friends and health workers) and health facility birth, as well as whether each network members moderated the relationship between each functional supports and health facility birth. At bivariate analysis, the result showed that except the interaction between emotional support and support received from relatives, all other interactions were not significant, but at multivariate analysis; all interactions were not significantly associated with health facility delivery (table 4).

Discussion

In order to understand culturally and socially relevant types of maternal social support, and to identify which measures may provide the best fit for our specific setting, we assessed the patterns of maternal social support and health facility delivery including the interactive effect of structural and functional social supports. In this study we found that around 47% of women delivered at health facility, consistent with recent studies in the country including data from the 2019 Mini EDHS and others which reported a proportion between 43%–52% (9,40,41). Previous reports in Ethiopia and in other low-income countries like Ghana, Kenya, and Bangladesh (7,10,16,42) showed much higher rates of facility deliveries (between 70%–74%), a discrepancy that could be explained by lower health facility delivery in rural areas, such as the study area of our analysis. The other explanation could be that our analysis considered women with uncomplicated pregnancy and childbirth, who may be less likely to delivery at a health facility because pregnancy and labour are seen as normal human events and not considered as a problem unless the women encountered complications and related health problems (15,42,43).

The current findings showed that perceived support, types of support and the people providing this support were associated with place of delivery. Multivariable logistic regression analysis revealed that women who perceived that their partner /family members helped them during childbirth were more likely

to deliver at health facilities, although this association should be interpreted with caution due to the possibility of reverse causality possible with cross-sectional data. Consistent with previous research (7,10,17) women who perceived that their husbands and family members were in favour of health facility deliveries were more likely to have given birth at one; receiving advice from family members during pregnancy further strengthened this likelihood.

In terms of functional social support, we found that women who were accompanied during labour and childbirth had a significantly higher odd of facility delivery than women without such support. The finding was supported by other studies which find that women strongly prefer to have a birth companion (22,44), which is also associated with increased satisfaction with healthcare services. A Cochrane effectiveness review suggested that having a labour companion enhances utilization of MCH services, and improves outcomes for women(45,46). Further, in addition to enhancing health facility delivery, women who have companions to accompany them during labour and childbirth may also benefit from more rapid diffusion of health information, and through practical and affective support in accessing to local services and facilities (13,45–47).

This study also supported the hypothesis that women who received continuous social support are more likely to use health facility delivery, which is in agreement with a previous study conducted in different LMICs (15,18,46,48). This suggests that women appreciate the continuous presence of maternal social supporters. Studies including systematic review recognizes the significance of continuous support of the woman's choice from her social network (such as her husband, partner, mother, or friend) during childbirth and WHO also encourages different organizations to issue practical guidelines promoting continuous support (4,49), as well as efforts to reduce the rate of dropout in the continuum of MCH care (1,25,50). A Cochrane effectiveness review on women's birth experience collected data from 11 RCTs comparing the impact of continuous and intermittent support; it reported that negative feelings about the childbirth experience and home delivery were significantly lower among women who received continuous support (44,46,48,51,52). The result of this study affirms these earlier findings and recommendations of the positive role of continuous support during pregnancy and childbirth in enhancing health facility delivery, when the supporter is part of the woman's social network.

By companions of the woman's choice from her social network (such as her husband, partner, mother, or friend). Women who received continuous labour support were more likely to give birth 'spontaneously'

From structural social support, the use of health facility delivery was positively and significantly associated when women received supports from close friends/neighbors. Although different studies report that friendship can have both negative and positive health impacts, many scholars report that the benefits of friends are greater than those from relatives. For example, in terms of predicting health, friendship occasionally predicts health to an equivalent and, in some cases, larger degree compared to spousal and parent–child relationships (26,29,50). Different social support variables highlighted by different researchers as predictors of the health service utilizations, such as number of ties/density, emotional supports, interaction between functional and structural dimensions, participation in roles that

promote healthcare, and member of any social groups, were not associated with health facility delivery in our study. Similarly, some studies argue that, instead of counting number of ties, amount of received supports and participation in activities that have different objectives than MCH care, it would be better to assess the quality of relationship and consistency of support from network members (24,50,53). Further, different studies on social support recommended caution during subgroup analyses and interpretation of social support variables (10,31,54), due to the fact that consistent patterns about the effectiveness of different social supports during pregnancy and childbirth may be enhanced or reduced by contexts in the birth setting, type of provider, and timing of onset of support (29,50,55). So, this result may imply about the main advantage of getting local context-based labour support which may target underserved pregnant women, provides service at no cost to the client, and seeks to employ supporters who are from the same local contexts in which they are living. In Ethiopia, receiving such in-kind social support during labour and childbirth is congruent with the social and cultural norms, where relationships and support from family and extended family, in-laws, close friends and customs are highly valued, including input and advice with regard to the decision about where and with whom to give birth.

Consistent with previous studies in Ethiopia, Ghana, Kenya, and Nepal (13,15,17,43), our analysis identified a significant positive association between women who are involved in deciding their place of childbirth and health facility delivery. Different social support studies and theories have suggested that interactions between network structure and perceptions of advice from family members and close friends would help explain pregnant women's decision to utilize health facility birth (15,18,56). Further, this study indicated that women who had a positive perception towards the benefits of health facility delivery were significantly more likely to seek health facility delivery. Several studies corroborated this finding, and the positive experiences of women in their childbirth helped them to have a positive perception, hence determining the childbirth place in their subsequent deliveries. Further, women whose families and close friends had positive perceptions about health facility birth were more likely to deliver in a health facility (57,58).

Our findings also revealed that women who lived further from the health facility were more likely to have used a health facility for delivery than those women who lived closer. In this regard, studies in Africa and Southeast Asia reported different findings on distance from health facility as an influencing factor for women's decisions to use MNCH care (13,57,59). Qualitative and quantitative studies from the same setting in Ethiopia reported that women living a greater distance to the maternity facility were more likely to utilize MNCH services including maternity waiting homes (15,16,37,45). Within the study setting, there is a freely available ambulance transport service during labour and the availability of maternity waiting homes may help women from remote areas use childbirth care (13,15,16,37). The other reason could be providing women with supports and transportation funds before they go to a facility for delivery and managing transportation options, as well as using maternity waiting home may increase service utilization (3,13,28,42,60).

The findings of this study, however, must be considered in light of various limitations. Women were only asked about current patterns of social supports, while the childbearing was done in the past, again

leading to issues about the direction of the relationships as well as the issue of whether current patterns of maternal social support consistently reflect the future patterns. Again it is not clear whether participant's perceptions of network approval of facility delivery may have been influenced by their actual experiences, rather than simple explanations of maternal support for facility delivery; mixed research methods including longitudinal studies could be important here in order to establish causality. Further, women could have responded with socially desirable answers such as reporting that they considered facility delivery in a positive light. In addition, ascertaining support for practical and advice regarding facility delivery does not provide the whole picture of social support. We therefore measured only some of the instrumental and informational support available to women. Despite these limitations, this study makes valuable contributions to the existing literature on social support in low resource settings. We also sampled women who had given birth 12 months prior to the survey which helped to limit recall bias

Conclusion

In conclusion, this study underscores the importance of accounting for the roles of network members in supporting women during pregnancy and childbirth when designing maternal and child health interventions to promote use of health facility delivery care. Community health education and mobilization interventions attempting to increase demand for health facility delivery should be tailored to incorporate social supports, as it could reflect the importance of supports in the forms of continuous supports and companionship accompanying women to health facility visits from different network members, particularly family members and close friends to close the gaps of distance and related constraints. In countries like Ethiopia, where social support is considered as a norm, such work may have immediate and positive influence by informing ongoing national level interventions to improve health services throughout pregnancy, delivery and early postnatal periods. Further, exploring different dimensions of maternal social support may be an existing opportunity for closing the gaps and complete the continuum of MCH utilization by combining effective cares that begins in the home, community, continues to labour and childbirth time, and follows woman back home with her baby in the early postpartum period. Therefore, future research should explore why and how social support can be considered as a pragmatic, problem-solving approach and an effective strategy in closing the gaps of completing continuum of MCH cares.

Abbreviations

ANC - Antenatal care; EDHS – Ethiopian demographic and health survey; cRCT - cluster-randomized control trial; FMoH – Federal Ministry of Health; HEWs - Health Extension Workers; IEC – Information, Education and Communication; LMIC -low and middle income countries; MCH – Maternal and child health; MNCH – Maternal, neonatal and child health; MMR - Maternal Mortality Rate; MWH - Maternity Waiting Home ; PNC – Postnatal care; PHCU – primary health care unit; RCT - randomized control trial; SSA – sub-Saharan African; SS – social support.

Declarations

Ethics approval and consent to participate

Ethical approval was obtained from Jimma University Institute of Health Institutional Review Board (Ref No: RPGE/449/2016) and from the University of Ottawa Health Sciences and Science Research Ethics Board (File No: H10-15-25B) and documented at both Jimma University and University of Ottawa. For written consent, participants were given time to read the consent forms and to ask questions before signing. For oral consent, the objectives of the research and ethical issues related to safety, confidentiality, and privacy were explained to uneducated participants in the local language of women's choice (Amharic or Afan Oromo) before asking if they agreed to the interview or focus group.

Consent to publish

Not applicable.

Availability of data and materials

Data used for this analysis (copy of the questionnaire used in the trial) will be provided by the authors upon reasonable request.

Competing Interests

The authors declare that they have no competing interests.

Funding

This work was carried out with grants from the Innovating for Maternal and Child Health in Africa initiative - a partnership of Global Affairs Canada (GAC), the Canadian Institutes of Health Research (CIHR) and Canada's International Development Research Centre (IDRC); it does not necessarily reflect the opinions of these organizations. The funder had no role in the study design, in the collection, analysis or interpretation of data, or in writing the manuscript.

Authors' contributions

AM, MS, ZB, SA, MA, MAK and LA contributed to conception and design of the study, collection, analysis of data, and were involved in drafting the manuscript. LA, RL, MS, NB, JK, MA, GK, ZB, MAK, YA, ML, GB, KHB and EE participated in interpretation of the results and contributed to development and finalization of the manuscript. All authors approved the final manuscript for publication.

Acknowledgements

The authors gratefully acknowledge the study participants for their time and effort. We wish to acknowledge Innovating for Maternal and Child Health in Africa project for funding this study through International Development Research Centre (IDRC) the government of Canada. The authors deeply acknowledge our research team in Jimma and Ottawa Universities for their dedication and contribution to complete this manuscript. We deeply appreciate Jimma University who provided logistical support throughout the data collection period, as well as Jimma zone Health offices for their support too.

References

1. Votano J, Parham M, Hall L. Celebrating Birth - Exploring the Role of Social Support in Labour and Delivery for First Nations Women and Families [Internet]. Chemistry & National Aboriginal Health Organization; 2008. 31 p. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/cbdv.200490137/abstract>
2. Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J, et al. Global causes of maternal death: A WHO systematic analysis. *Lancet Glob Heal* [Internet]. 2014 [cited 2021 May 20];2(6).
3. Arba MA, Darebo TD, Koyira MM. Institutional delivery service utilization among women from rural districts of Wolaita and Dawro zones, southern Ethiopia; a community based cross-sectional study. *PLoS One*. 2016;11(3).
4. Statistics World Health. Monitoring health for the SDGs, sustainable development goals [Internet]. Vol. 4, Who. 2019. p. 75–84.
5. Sofer D. Why are women still dying of pregnancy and childbirth? [Internet]. Vol. 118, *American Journal of Nursing*. Lippincott Williams and Wilkins; 2018 [cited 2021 May 20]. p. 12.
6. Ethiopia Demographic and Health Survey 2016. Central Statistical Agency (Ethiopia)ICF International. Ethiopia Demographic and Health Survey 2016.
7. Cofie LE, Barrington C, Singh K, Sodzi-Tettey S, Ennett S, Maman S. Structural and functional network characteristics and facility delivery among women in rural Ghana. *BMC Pregnancy Childbirth*. 2017 Dec 19;17(1).
8. World Health Statistics. Monitoring health for the SDGs, sustainable development goals [Internet]. Vol. 4, Who. 2019 [cited 2021 May 15]. p. 75–84.
9. Ethiopian Public Health Institute (EPHI) and ICF. 2019. Mini Demographic and Health Survey 2019: key Indicators. Rockville, Maryland, USA: EPHI and ICF. 2019. 35 p.

10. Edmonds JK, Hruschka D, Bernard HR, Sibley L. Women's social networks and birth attendant decisions: Application of the Network-Episode Model. *Soc Sci Med*. 2012 Feb;74(3):452–9.
11. UNICEF. Maternal mortality rates and statistics - UNICEF DATA [Internet]. [cited 2021 May 23]. Available from: <https://data.unicef.org/topic/maternal-health/maternal-mortality/>
12. Central Statistical Agency (Ethiopia), ICF International. Ethiopia Demographic and Health Survey 2016. Fairfax, United States of America: ICF International, 2017.
13. Kurji J, Gebretsadik LA, Wordofa MA, Sudhakar M, Asefa Y, Kiros G, et al. Factors associated with maternity waiting home use among women in Jimma Zone, Ethiopia: A multilevel cross-sectional analysis. *Vol. 9, BMJ Open*. BMJ Publishing Group; 2019. p. 1–9.
14. Kurji J, Kulkarni MA, Gebretsadik LA, Wordofa MA, Morankar S, Bedru KH, et al. Effectiveness of upgraded maternity waiting homes and local leader training in improving institutional births among women in the Jimma zone , Ethiopia : study protocol for a cluster-randomized controlled trial. 2019;1–11.
15. Mamo A, Morankar S, Asfaw S, Bergen N, Kulkarni MA, Abebe L, et al. How do community health actors explain their roles? Exploring the roles of community health actors in promoting maternal health services in rural Ethiopia. *BMC Health Serv Res*. 2019;19(1):1–12.
16. Kurji J, Gebretsadik LA, Wordofa MA, Morankar S, Bedru KH, Bulcha G, et al. Effectiveness of upgraded maternity waiting homes and local leader training on improving institutional births: a cluster-randomized controlled trial in Jimma, Ethiopia. *BMC Public Health*. 2020;20(1):1–15.
17. Ono M, Matsuyama A, Karama M, Honda S. Association between social support and place of delivery: A cross-sectional study in Kericho, Western Kenya [Internet]. *Vol. 13, BMC Pregnancy and Childbirth*. 2013. Available from: <http://www.biomedcentral.com/1471-2393/13/214>
18. Afulani P, Kusi C, Kirumbi L, Walker D. Companionship during facility-based childbirth: Results from a mixed-methods study with recently delivered women and providers in Kenya. *BMC Pregnancy Childbirth*. 2018 May 10;18(1).
19. Abosse Z, Woldie M, Ololo S. Factors Influencing Antenatal Care Service Utilization in Hadiya Zone [Internet]. *Vol. 20, Ethiopian Journal of Health Sciences*. 2011 [cited 2021 May 18].
20. Bergen N, Zhu G, Yedenekal SA, Mamo A, Abebe Gebretsadik L, Morankar S, et al. Promoting equity in maternal, newborn and child health—how does gender factor in? Perceptions of public servants in the Ethiopian health sector. *Glob Health Action*. 2020 Dec 31;13(1).
21. Leban K. How Social Capital in Community Systems Strengthens Health Systems: People, Structures, Processes. 2011;(October).

22. Andersen R, Newman JF. Societal and individual determinants of medical care utilization in the United States. *Milbank Q* [Internet]. 2005 Dec 10 [cited 2021 May 18];83(4).
23. Leahy-Warren P, McCarthy G, Corcoran P. First-time mothers: Social support, maternal parental self-efficacy and postnatal depression. *J Clin Nurs*. 2012 Feb;21(3–4):388–97.
24. Putnam RD. E pluribus unum: Diversity and community in the twenty-first century the 2006 Johan skytte prize lecture. *Scan Polit Stud*. 2007 Jun;30(2):137–74.
25. Levine S, Lipson J. Fostering Social Support: Accommodating Patients’ Family and Friends as Part of the Healing Process. unpublished [Internet]. 2010 [cited 2021 May 18];1–52.
26. Mattson M& JGH. Linking Health Communication with Social Support. *Health as Communication Nexus*. 2011. 181–218 p.
27. Bergen N, Hudani A, Asfaw S, Mamo A, Kiros G, Kurji J, et al. Promoting and delivering antenatal care in rural Jimma Zone, Ethiopia: A qualitative analysis of midwives’ perceptions. *BMC Health Serv Res*. 2019 Oct 21;19(1).
28. Kincaid DL. From innovation to social norm: Bounded normative influence. *J Health Commun*. 2004;9:37–57.
29. Edmonds JK, Paul M, Sibley LM. Type, content, and source of social support perceived by women during pregnancy: Evidence from Matlab, Bangladesh. *J Heal Popul Nutr*. 2011;29(2):163–73.
30. Rosen P. Supporting Women in Labor: Analysis of Different Types of Caregivers. *J Midwifery Womens Health*. 2004 Jan 2;49(1):24–31.
31. Pescosolido BA. Illness Careers and Network Ties: A Conceptual Model of Utilization and Compliance [Internet]. Vol. 2, *Advances in Medical Sociology*. 1991 [cited 2021 May 18]. p. 161–84. Available from: <https://www.google.com/search?q>
32. Kotoh AM, Boah M. “No visible signs of pregnancy, no sickness, no antenatal care”: Initiation of antenatal care in a rural district in Northern Ghana. *BMC Public Health*. 2019;19(1).
33. Sialubanje C, Massar K, Hamer DH, Ruiter RAC. Reasons for home delivery and use of traditional birth attendants in rural Zambia: A qualitative study. *BMC Pregnancy Childbirth*. 2015;15(1).
34. Warri D, George A. Perceptions of pregnant women of reasons for late initiation of antenatal care: A qualitative interview study. *BMC Pregnancy Childbirth*. 2020;20(1).
35. Bergen N, Zhu G, Yedenekal SA, Mamo A, Abebe Gebretsadik L, Morankar S, et al. Promoting equity in maternal, newborn and child health—how does gender factor in? Perceptions of public servants in the Ethiopian health sector. *Glob Health Action*. 2020 Dec 31;13(1).

36. Behrman JR, Kohler H-P, Watkins SC. Social networks and changes in contraceptive use over time: Evidence from a longitudinal study in rural Kenya. *Demography*. 2002 Nov 1;39(4):713–38.
37. Asfaw S, Morankar S, Abera M, Mamo A, Abebe L, Bergen N, et al. Talking health: Trusted health messengers and effective ways of delivering health messages for rural mothers in Southwest Ethiopia. *Arch Public Heal*. 2019 Feb 21;77(1).
38. zone Health Office 2016 J, 2019 JZHO. Jimma Zone Health Office. Jimma zone annual health bulletin. 2019.
39. Jimma zone Health Office 2016, Jimma Zone Health Office 2019, zone Health Office 2016 J, 2019 JZHO. Jimma Zone Health Office. Jimma zone annual health bulletin. 2019.
40. Emiru AA, Alene GD, Debelew GT. Women's retention on the continuum of maternal care pathway in west Gojjam zone, Ethiopia: Multilevel analysis. *BMC Pregnancy Childbirth*. 2020 Apr 29;20(1).
41. Haile D, Kondale M, Andarge E, Tunje A, Fikadu T, Boti N. Level of completion along continuum of care for maternal and newborn health services and factors associated with it among women in Arba Minch Zuria woreda, Gamo zone, Southern Ethiopia: A community based crosssectional study. *PLoS One*. 2020 Jun 1;15(6).
42. Bergen N, Abebe L, Asfaw S, Kiros G, Kulkarni MA, Mamo A, et al. Maternity waiting areas—serving all women? Barriers and enablers of an equity-oriented maternal health intervention in Jimma Zone, Ethiopia. *Glob Public Health*. 2019;14(10):1509–23.
43. Adhikari R. Effect of Women's autonomy on maternal health service utilization in Nepal: A cross sectional study. *BMC Womens Health*. 2016;16(1).
44. Story WT. Social capital and the utilization of maternal and child health services in India: A multilevel analysis. *Heal Place [Internet]*. 2014 [cited 2021 May 19];28:73–84.
45. Bohren MA, Berger BO, Munthe-Kaas H, Tunçalp Ö. Perceptions and experiences of labour companionship: A qualitative evidence synthesis. *Cochrane Database Syst Rev*. 2019;2019(3).
46. Hodnett ED, Gates S, Hofmeyr GJ, Sakala C. Continuous support for women during childbirth. Vol. 2013, *Cochrane Database of Systematic Reviews*. John Wiley and Sons Ltd; 2013.
47. MA CAB, Bohren MA, Hofmeyr GJ, Sakala C, Fukuzawa RK, Cuthbert A, et al. *Cochrane Database of Systematic Reviews Continuous support for women during childbirth (Review) Continuous support for women during childbirth [Internet]*. 2017 [cited 2021 May 19].
48. Agampodi TC, Rheinländer T, Agampodi SB, Glozier N, Siribaddana S. Social capital and health during pregnancy; An in-depth exploration from rural Sri Lanka. *Reprod Health*. 2017 Jul 27;14(1).

49. Lunda P, Minnie CS, Benadé P. Women's experiences of continuous support during childbirth: A meta-synthesis. *BMC Pregnancy Childbirth*. 2018;18(1):1–11.
50. Lu P, Oh J, Leahy KE, Chopik WJ. Friendship Importance Around the World: Links to Cultural Factors, Health, and Well-Being. *Front Psychol*. 2021;11.
51. Diamond-Smith N, Sudhinaraset M, Melo J, Murthy N. The relationship between women's experiences of mistreatment at facilities during childbirth, types of support received and person providing the support in Lucknow, India. *Midwifery*. 2016 Sep 1;40:114–23.
52. Owili PO, Muga MA, Chou YJ, Hsu YHE, Huang N, Chien LY. Associations in the continuum of care for maternal, newborn and child health: A population-based study of 12 sub-Saharan Africa countries. *BMC Public Health* [Internet]. 2016 [cited 2021 May 19];16(1).
53. Mary Ross-Davie PhD MA RMBA, Helen Cheyne PhD MSc RMRGN, Ross-Davie PhD, MA, RM, BA M, Cheyne PhD, MSc, RM, RGN H. Intrapartum support: what do women want? A literature review. *Evid Based Midwifery* [Internet]. 2014;12(2):52–8.
54. Fine B. Theories of social capital: Researchers behaving badly, *Political Economy and Development*, 2010, ISBN 978-1-84964-442-6, Pluto Press, London.
55. Bergen N, Mamo A, Asfaw S, Abebe L, Kurji J, Kiros G, et al. Perceptions and experiences related to health and health inequality among rural communities in Jimma Zone, Ethiopia: A rapid qualitative assessment. *Int J Equity Health*. 2018 Jun 18;17(1).
56. Speizer IS, Story WT, Singh K. Factors associated with institutional delivery in Ghana: The role of decision-making autonomy and community norms. *BMC Pregnancy Childbirth*. 2014 Nov 27;14(1).
57. Gao LL, Chan SWC, Mao Q. Depression, perceived stress, and social support among first-time Chinese mothers and fathers in the postpartum period [Internet]. Vol. 32, *Research in Nursing and Health*. 2009 [cited 2021 May 19]. p. 50–8.
58. Kapp M., M. K, Kapp M. Mothers' perceptions of confidence with self-care and infant care. *Journal of Perinatal Education*, 1998 - Google Search. *J Perinat Educ* [Internet]. [cited 2021 May 19];
59. Choulagai B, Onta S, Subedi N, Mehata S, Bhandari GP, Poudyal A, et al. Barriers to using skilled birth attendants' services in mid- and far-western Nepal: A cross-sectional study. *BMC Int Health Hum Rights*. 2013 Dec 23;13(1).
60. Ajzen I. The theory of planned behavior. *Organ Behav Hum Decis Process*. 1991;50(2):179–211.

Tables

Table 1. Characteristics of socio-demographic factors and health facility delivery of women's most recent childbirth in Jimma, Ethiopia, 2017(N = 3304)

Variables	Frequency	%
Age		
15 - 19	186	5.6
20 - 24	756	22.9
25 – 29	1016	30.8
30 - 34	765	23.2
35 - 39	482	14.6
40 - 44	91	2.8
45+	8	0.2
Marital status		
Married	3233	97.9
Not married	71	2.1
Women Education		
No formal education	1856	56.2
Primary school	1278	38.7
High school	143	4.3
Above high school	27	0.8
Women's occupation		
Housewife	2547	77.1
Farmer	495	15.0
Traders	169	5.1
Others(private workers, unemployed & student)	93	2.8
Household income per year		
1st quartile	922	31.2
2nd quartile	512	17.3
3rd quartile	769	26.0
4th quartile	756	25.5
Partner's occupation (N=495)		
Farmer	360	72.7
Trader	106	21.4
Others(private workers, employed & student)	29	5.9

Table 2 Characteristics of healthcare related attributes and health facility delivery of women's most recent childbirth in Jimma, Ethiopia, 2017(N = 3304)

Variables	Frequency	%
ANC 4+ visits		
No	957	29.0
Yes	2347	71.0
Health facility delivery		
No	1755	53.1
Yes	1549	46.9
Availability of health facility		
No	143	4.3
Yes	3161	95.7
Types of health facility		
Private health facilities	8	0.3
Health posts	1560	49.4
Government hospitals and health centers	1593	50.4
Planned place of delivery		
No	1274	38.6
Yes	2030	61.4
Member of any social group		
No	2736	82.8
Yes	568	17.2
Participated a programme that promote healthcare		
No	1653	50.0
Yes	1651	50.0
Involved in deciding place of delivery		
No	1594	50.4
Yes	1566	49.6

Table 3 Analysis of maternal social support and health facility delivery in most recent childbirth in southwest Ethiopia, 2017 (N = 3304).

Domains of Social support		Health facility delivery		P.value
		No (N=1755) N (%)	Yes(N=1549) N (%)	
Perceived social supports				
Perceived women had someone to help during childbirth	No	575(50.9)	555(49.1)	0.064
	Yes	1180(54.3)	994(45.7)	
Relationship to perceived source of help				
Partner/family	No	701(38.9)	1103(61.1)	0.001
	Yes	100(18.6)	439(81.4)	
Relatives	No	700(39.1)	1090(60.9)	0.001
	Yes	101(18.3)	452(81.7)	
Friends	No	1356(52.5)	1229(47.5)	0.149
	Yes	399(55.5)	320(44.5)	
Health workers	No	1227(52.2)	1123(47.8)	0.102
	Yes	528(55.3)	426(44.7)	
Functional supports/received supports				
Practical support with routine tasks	No	1486(54.8)	1224(45.2)	0.001
	Yes	269(45.3)	325(57.7)	
Accompanied in last childbirth	No	911(61.6)	568(38.4)	0.001
	Yes	269(45.3)	325(54.7)	
Emotional support	No	563(51.4)	532(48.6)	0.168
	Yes	1192(54.0)	1017(46.0)	
Material support	No	1079(52.1)	992(47.9)	0.129
	Yes	676(54.8)	557(45.2)	
Informational support	No	294(51.8)	274(48.2)	0.477
	Yes	1461(53.4)	1275(46.6)	
Size of received supports [<i>mean(±SD)</i>]		0.5(0.2)	0.6 (0.2)	0.001
Continuous support[<i>mean(±SD)</i>]		0.3(0.2)	0.4(0.2)	0.001
Structural supports/Relationship with supporters				
Partner/family members	No	1625(53.9)	1388(46.1)	0.002
	Yes	130(44.7)	161(55.3)	
Relatives/siblings	No	1557(54.4)	1305(45.6)	0.001
	Yes	198(44.8)	244(55.2)	
Friends/neighbors	No	1617(54.4)	1356(45.6)	0.001
	Yes	138(41.7)	193(58.3)	
Health workers	No	895(52.0)	827(48.0)	0.711
	Yes	860(54.4)	722(45.6)	
Density of ties [<i>mean (±SD)</i>]		0.4(0.2)	0.4(0.2)	0.230

Table 4. Multivariable regression analysis of social support and related factors with health facility delivery in most recent childbirth in southwest Ethiopia, 2017 (N = 3304).

Variables	Health Facility Delivery		OR (95%CI)	AOR (95%CI)
	No (N=1755) N (%)	Yes (N=1549) N (%)		
Women Occupation				
Housewife	1378(54.1)	1169(45.9)	1	1
Farmer	226(45.7)	269(54.3)	1.40 (1.16,1.70)*	0.97(0.73, 1.29)
Traders	97(57.4)	72(42.6)	0.88 (0.64, 1.20)	0.94(0.62, 1.41)
Others (private workers & student)	54(58.1)	39(41.9)	0.85 (0.56, 1.30)	0.92(0.51, 1.69)
Household income				
1st quartile	461(50.0)	461(50.0)	1	1
2nd quartile	280(54.7)	232(45.3)	1.20 (0.93,1.53)	0.71(0.53, 1.01)
3rd quartile	410(53.3)	359(46.7)	1.23 (0.99,1.54)	0.92(0.71, 1.20)
4th quartile	437(57.8)	319(42.2)	1.52 (1.23,1.88)*	0.80(0.62, 1.05)
Number of household members [<i>Mean (±SD)</i>]	5.6(1.9)	5.8(2.0)	1.04 (1.01,1.08)*	1.00(0.95, 1.06)
Travel time to closer health facility [<i>Mean(±SD)</i>]	0.5(0.4)	0.6 (0.5)	1.44 (1.24,1.67)*	1.51 (1.21,0.90)*
Participated on program that promote healthcare				
No	844(51.1)	809(48.9)	1	1
Yes	911(55.2)	740(44.8)	0.85(0.74,0.97)*	0.98(0.80, 1.20)
Involved in deciding place of delivery				
No	957(60.0)	637(40.0)	1	1
Yes	723(46.2)	843(53.8)	1.75(1.52, 2.02)*	2.12(1.73, 2.58)
Attitude to facility delivery [<i>Mean (±SD)</i>]	0.4(0.2)	0.4(0.2)	1.46 (1.14,1.88)*	0.94(0.65, 1.37)
Perception on provided MCH [<i>Mean (±SD)</i>]	0.3(0.2)	0.3(0.3)	1.83 (1.44,2.33)*	1.89 (1.31, 2.71)*
Perceived partner/family members to help them				
No	701(38.9)	1103(61.1)	1	1
Yes	100(18.6)	439(81.4)	2.82 (2.22, 3.45)*	3.62 (2.74, 4.79)*
Received practical support				
No	1486(54.8)	1224(45.2)	1	1
Yes	269(45.3)	325(57.7)	1.47 (1.23, 1.75)*	0.81(0.50, 1.40)
Accompanied to health facility				
No	911(61.6)	568(38.4)	1	1
Yes	269(45.3)	325(54.7)	1.86 (1.62, 2.14)*	1.63 (1.34, 2.00)*
Size of received supports(<i>Mean (±SD)</i>)	0.5(0.2)	0.6 (0.2)	1.91 (1.39, 2.62)*	1.30(0.94, 1.78)

Received continuous support (<i>Mean (±SD)</i>)	0.3(0.2)	0.4(0.2)	2.17 (1.52, 3.10)*	1.97 (1.20, 3.23)*
Received from partner/family members				
No	1625(53.9)	1388(46.1)	1	1
Yes	130(44.7)	161(55.3)	1.45(1.14, 1.85)*	1.20(0.73, 1.97)
Received from relatives/siblings				
No	1557(54.4)	1305(45.6)	1	1
Yes	198(44.8)	244(55.2)	1.47(1.20, 1.80)*	1.30(0.94, 1.78)
Received from friends/neighbors				
No	1617(54.4)	1356(45.6)	1	1
Yes	138(41.7)	193(58.3)	1.67(1.33, 2.10)*	1.62(1.16, 2.26)*

*significant variables