The Transcriptome-Wide N6-Methyladenosine (m⁶A) Map Profiling Reveals the Regulatory Role of m⁶A in the Yak Ovary

Shaoke Guo

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Xingdong Wang

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Mengli Cao

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Xiaoyun Wu

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Lin Xiong

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Pengjia Bao

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Min Chu

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Chunnian Liang

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Ping Yan

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Jie Pei

Key Laboratory of Yak Breeding Engineering of Gansu Province, Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences

Xian Guo (✉ guoxian@caas.cn)

Research Article

Keywords: mRNAs, MeRIP-seq, estrus, anestrus, pregnancy.

Posted Date: September 2nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-820378/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background and aim: Yak estrus is a seasonal phenomenon, probably involving epigenetic regulation of synthesis and secretion of sex hormones as well as growth and development of follicles. N6-methyladenosine (m⁶A) is the most common internal modification of the eukaryotic mRNA. However, there are no detailed reports on the m⁶A transcriptome map of yak ovary. Therefore, this study aimed to collect the yak ovarian tissues at three different states of anestrus (YO-A), estrus (YO-F), and pregnancy (YO-P), and obtained the full transcriptome m⁶A map in yak by MeRIP-seq.

Results: The HE staining revealed that the number of growing follicles and mature follicles in the ovary during the estrus period was relatively higher than those in the estrus period and the pregnancy period. The RT-qPCR showed that the expression of *METTL3*, *METTL14*, *FTO*, *YTHDC1* were significantly different across different periods in the ovaries, which suggests that m⁶A may play a regulatory role in ovarian activity. Next, we identified 20174, 19747 and 13523 m⁶A peaks in the three ovarian samples of YO-A, YO-F and YO-P using the methylated RNA immunoprecipitation sequencing (MeRIP-seq). The m⁶A peaks are highly enriched in the coding sequence (CDS) region and 3'untranslated region (3'UTR) as well as the conserved sequence of "RRACH." Functional analysis revealed the involvement of m⁶A in many physiological activities of the yak's ovary during reproductive cycle. The association analysis found that some genes such as *BNC1*, *HOMER1*, *BMP15*, *BMP6*, *GPX3*, and *WNT11* were related to ovarian functions.

Conclusions: The comparison of the distribution patterns of methylation peaks in the ovarian tissues across different periods further explored the m⁶A markers related to the regulation of ovarian ovulation and follicular development in the yak ovary. This comprehensive map provides a solid foundation for revealing the potential function of the mRNA m⁶A modification in the yak ovary.

Background

Yak (*Bos grunniens*), an endemic cattle species, is capable of adapting to extremely cold and low-oxygen environments and highly resistant to diseases. It is not only an important source of life and economy for herdsmen in the Qinghai Tibet Plateau but also indispensable livestock in the local animal husbandry economy[1]. Yak generally calve once in two years or twice in three years. Yak ovulations are usually manifested by seasonal physiological changes, and the time of ovulation and conception mainly ranging from August to September[2]. It is reported that epigenetic might regulate the pre-ovulatory follicular growth and development, as well as the follicular selection[3]. In the cold season, yak undergo ovarian dysfunction due to the accumulation of the antral follicles and dysfunction of the granulosa cells. This process may be related to steroid hormone biosynthesis and estrogen signaling pathway, or regulated by epigenetic mechanisms[4]. Xia et.al found that a decreased m⁶A level in the zebrafish (*Danio rerio*) not only disrupted the sex hormone synthesis and the gonadotropin signal transduction gene expression, but also impaired the meiosis process of oocytes[5].

N6-methyladenosine (m⁶A) discovered in 1974, constitute the most prominent internal RNA modification[6, 7]. Remarkably, numerous studies have reported m⁶A executed an important regulatory role in animal reproduction. The spermatocytes lacking METTL3 could not reach the pachytene stage of the meiotic prophase, inhibiting the spermatogonial differentiation and the normal initial meiosis[8], inactivating ALKBH5 and leading to sterility in male mice[9]. It was found in zebrafish that METL3 deletion can impair the meiotic process of oocytes[10]. YTHDF2 may regulate mRNA degradation during meiosis and maturation of oocytes in an m⁶A-dependent manner[11]. YTHDC2 plays an important role in the meiotic process of male and female germ cells, and the deletion may lead to infertility or infertility[12]. However, to the best of our knowledge, there are no reports on m⁶A in the yak ovarian tissues.

Despite the early discovery of m⁶A, studies on this modification were hindered by multiple factors. Until the emergence of new sequencing technology, Dominisini and Meyer [11, 13] combined immunoprecipitation technology with next generation high-throughput sequencing (m⁶A-Seq or MeRIP-Seq). Since then, researchers have successfully constructed the m⁶A map for a variety of animals and plants using the MeRIP-Seq. To study the regulatory mechanism of the yak ovary during estrus and pregnancy, and to promote the further study of yak m⁶A, the ovarian tissues of yak in three different states of anestrus, estrus (also known as follicular phase) and pregnancy, respectively, were collected and the novel full transcriptome m⁶A map in yak was obtained by MeRIP-Seq. By comparing the distribution patterns of the different m⁶A peaks in different ovarian tissues, the relationship between the regulation of the gene expression by m⁶A modification and ovarian function was further explored.

Methods

Sample collection

The ovarian tissues were collected from three healthy yaks of the same age during three different periods of the year. The samples were separately collected during the anestrus in April (A1, A2, A3), the follicular (also as estrus) phase in August (F1, F2, F3), and the pregnancy period in December (P1, P2, P3). Identification of the yak estrus was performed as described by Zi et al.[35]. The ovarian tissue was divided into two parts, one part was fixed in 10% neutral formaldehyde for making tissue sections, the other part was immediately stored in liquid nitrogen for RNA isolation. All the samples were collected from the Menyuan Hui Autonomous County, Haibei Tibetan Autonomous Prefecture, Qinghai Province, China (37°39′N 101°62′E). All ovarian samples were collected from euthanized yaks within 1 hour.

HE staining

The fixed-tissues were dehydrated, embedded, and sliced using a fully automatic dehydrator, the following operations are performed: hematoxylin staining for 10–20 min; tap water for 1–3min; hydrochloric acid alcohol differentiation for 5–10s; tap water for 1–3min; treated with 50°C warm water or weak alkaline aqueous solution until the blue color appears; rinsed with tap water for 1–3min; 85% alcohol for 3–5min; dye with eosin for 3–5 min; washed for 3–5s; gradient alcohol dehydration;

transparent xylene; sealed with neutral gum. The above specimens were carried out according to the pathological examination standard operation procedure (SOP). The dehydration, trimming, embedding, sectioning, staining, mounting, etc., and finally, the microscopic examination of the slice was captured using the Panoramic 250 digital slice scanner (DRNJIER, Ji Nan, CHINA).

RNA isolation and purification

The total RNA was isolated and purified using the TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The amount and purity of the RNA samples were quantified using the NanoDrop ND-1000 (NanoDrop, Wilmington, DE, USA). Then the concentration and integrity of each RNA sample was assessed by Bioanalyzer 2100 (Agilent, CA, USA) and verified by agarose electrophoresis. The RNA that meets the conditions of concentration > 50 ng/μL, Rin > 7.0, OD260 / 280 > 1.8, total RNA > 50 μg, is used for downstream experiments..

RT-qPCR

The extracted RNA was reverse transcribed to obtain cDNA using the *Evo M-MLV* RT Kit with gDNA Clean for qPCR II (AG, Hunan, China). The primer synthesis template of the target gene is shown in Additional file 8. The real-time quantitative real-time PCR (RT-qPCR) was performed using the SYBR Green master mix (Yeasen, Shanghai, China) and the LightCycler System (CFX96™ Optics Module, Singapore). The reaction volume of 20μL were used, including 10μL of 2x Precision PLUS Master Mix, 1μL of diluted cDNA (25 ng), each 1μL (300 nmol) of gene-specific forward and reverse primers, and 7μL of RNase/Dnase free water. The standard PCR reaction conditions for all transcripts are 95°C for 3min, 35 cycles of 95°C for 20 s, 52°C for 40 s, and 72°C for 4 min, followed by 72°C for 10 min. The data are expressed as mean ± SE (n = 3). The *Glyceraldehyde-3-phosphate dehydrogenase (GAPDH)* gene was used as an internal reference gene, and the expression of mRNA was normalized by the $2^{-\Delta\Delta CT}$ method. All the statistical analyses were calculated using a test in the SAS 9.4 statistical software. Statistically significant difference were considered as $P < 0.05$.

The methylated RNA immunoprecipitation sequencing (MeRIP-Seq)

The Poly (A) RNA was purified from 50 μg total RNA using Dynabeads Oligo (dT) 25-61005 (Thermo Fisher, CA, USA) through two rounds of purification. Then the captured RNA was fragmented into small pieces using the Magnesium RNA Fragmentation Module (NEB, cat.e6150, USA) for 7 min at 86°C. The fragmented RNA was premixed with the Dynabeads Antibody Coupling Kit (Thermo Fisher, CA, USA) and m⁶A antibody (No.202003, Synaptic Systems, Germany) in IP buffer (50 mM Tris-HCl, 750 mM NaCl and 0.5% Igepal CA-630) and incubated for 2 h at 4°C. The IP RNA was reverse-transcribed to synthesize the cDNA using the SuperScript™ II Reverse Transcriptase (Invitrogen, cat. 1896649, USA). Then the two-strand synthesis was performed with RNase H (NEB, product number m0297, USA) using the *E. coli* DNA polymerase I (NEB, product number m0209, USA) to synthesize U-labeled second-stranded DNAs. At the same time, the dUTP Solution (Thermo Fisher, Catalog No. R0133, CA, USA) was incorporated into the two strands, and the ends of the double-stranded DNA were filled with blunt ends. An A-base was added at

each end of the fragment to connect it to the T-base overhang, and the size of the fragment was screened and purified using the AMPureXP beads. The U-labeled second-stranded DNAs were digested using a heat-labile UDG enzyme (NEB, cat.m0280, USA), then amplified by PCR under the following conditions: Pre-denaturation at 95°C for 3 min, denaturation at 98°C (8 cycles) for 15 s, annealing at 60°C for 15 s, extending at 72°C for 30 s, and finally extending at 72°C for 5 min to form a cDNA library of 300 ± 50 BP. At last, the 2 × 150 bp paired-end sequencing (PE150) was performed on an Illumina Novaseq™ 6000 (LC-Bio Technology CO., Ltd., Hangzhou, China) following the vendor's recommended protocol.

Data Analysis

The Fastp software was used to remove the reads of the IP and input samples that contained splices, repetitive sequences and low-quality sequences[36] with default parameters. The map reads to the reference genome of *Bos grunniens* (Version: v101)[37] by HISAT2, and the file format is bam. The R package exomepeak[38] was used for peak calling analysis and gene difference peak analysis for the bam files obtained from the IP samples and input samples. The candidate peak region was extended on the genome to obtain a modeled region of a certain length. According to all the unique comparison reads in this region, the Poisson distribution model was used to test and calculate the p-value of the candidate peak region. The default p-value < 0.05, the area smaller than the p-value is considered to be a peak. The obtained m⁶A peaks with the bed or bigwig format are visually displayed using the IGV software. The called peaks were annotated by intersection with the gene architecture using R package ChIPseeker [39], MEME [40] and HOMER were used for motif analysis. The StringTie software[41] was used to perform gene assembly and quantification by using the quantification method FPKM (total exon fragments /mapped reads (millions) × exon length (kB)). The differentially expressed mRNAs were selected with log₂ (fold change) >0.58496 or log₂ (fold change) <-0.58496 and p value < 0.05 using the R package edgeR[42].

Results

Observation of ovarian tissue section

The ovarian tissue slices in the anestrus (YO-A), follicular (YO-F), and pregnancy (YO-P) phases were observed and scanned under the microscope using the CaseViewer software. Based on the classification of follicles on the ovarian surface by Yu Sijiu[14], the ovarian capsule was found to be intact, with primordial, primary follicles, secondary, and mature follicles evident in the cortex, and the development of the follicles were normal at all levels (Fig. 1). The fibrous connective tissue in the medullary area was arranged closely without obvious edema or necrosis, and a small number of lymphocytes were infiltrated. By comparison, it was found that there were more growth follicles and mature follicles in August, while in April and December, the number of mature follicles was less. The arrangement of follicular cells in the granulosa layer of primary or secondary follicles was disordered in some tissues. In terms of the volume of follicles, the volume of atresia follicles is larger in August. The diameter of atresia follicles is mostly 2–

3 mm, and the largest being up to 3.6 mm. The atretic follicles observed in April were slightly smaller, with a diameter of 0.5–1.5 mm. The diameter of atretic follicles evident in December was between 1–3 mm.

Comparison of the m⁶A enzyme expression level

In order to further determine whether m⁶A plays a regulatory role in yak ovarian activity, the gene expression of 13 m⁶A-related enzymes: METL3, METL14, WTAP, FTO, ALKBH5, YTHDF1-3, YTHDC1-2, RBM15, VIRMA, and ZC3H13 were detected using the RT-qPCR. Figure 2a depicted that the expression levels of METL3, FTO, and ALKBH5 in the anestrus ovaries were significantly higher than those of the ovaries during estrus and pregnancy. On the contrary, the expression levels of YTHDF1, YTHDF2, and YTHDC2 in the ovaries during the estrus period were significantly higher than those in other periods. The expression of METL14 and VIRMA in the ovaries was significantly higher during pregnancy than that in estrus and anestrus. It can be seen from the test results that there are significant differences in the expression of these methylases at different stages of the ovary. Therefore, we speculate that m⁶A plays an important role in ovarian activity in yaks.

Sequencing quality control and reference genome alignment

To obtain the m⁶A transcriptome map of the yak ovary, the ovarian tissues in three different periods- YO-A, YO-F, and YO-P were detected by MeRIP-Seq. The samples were obtained from three yaks during each period. The result of the principal component analysis showed that the YO-F2 had poor reproducibility for the same group of samples (Fig. 2b), therefore, the sample data would not be analyzed in the future. The reading with connector, low quality readings (the number of bases with a quality value of $Q \leq 10$ accounts for more than 20% of the total read), and low quality bases were filtered out by the Trimomatic v0.36 tool. In the MERIP-seq library, the three groups of YO-A, YO-F, and YO-P yielded an average of 51426911, 51743067, and 50341117 original data readings, and an average of 50785722, 51071937 and 49348875 valid data readings. The proportion of valid readings is above 90% (Supplementary file 1). Unlike m⁶A-seq, which was found in early mammals[11] (11 million to 24 million readings per sample) and rice[15] (23 million to 47 million readings per sample), the technology for m⁶A-seq in the current study was significantly improved and all the sequences have high depths.

The preprocessed valid data was compared to the reference genome (LU_Bosgru_v3.0) by the software Hisat2. The number of reads that can be matched to the genome accounts for about 90% of the data after quality control, among which the number of reads that can only be matched at one location of the genome accounts for 62–67%, while the number of reads that can be matched to multiple locations of the genome accounts for 22–29% (Supplementary file 1), as shown in Fig. 2c. According to the regional information of the reference genome, the valid data that can be matched to the reference genome can be defined as those matched to the exon, intron, and intergenic regions. It can be seen from Fig. 2d that the percentage content of sequencing the sequence localization in the exon region is the highest.

Identification of the genome-wide m⁶A peak

The readings obtained from the yak samples were compared with the yak reference genome. These peaks were annotated using the R package, ChIPseeker after peak scanning the whole gene (Supplementary file 2). Referring to the previous studies, highly credible peaks were determined for each experimental condition (one inbred line under one environmental condition). In short, by crossing the peak regions in pairs between all the three replicates, at least two overlapping regions in the three replicates were designated as the high confidence m⁶A peak regions. About 20,174, 19,747, and 13,523 m⁶A peaks were separately identified in the Yo-A, Yo-F, and Yo-P ovary samples. All the peaks had total lengths of 95,594,574 bp, 122,175,034 bp, and 53,590,770 bp, respectively, accounting for 3.37%, 4.31%, and 1.89% of the genome. The average lengths of m⁶A peaks in the YO-A, YO-F, and YO-P groups were 4738.50 bp, 6,187.02 bp, and 3962.94 bp, the length distribution has been shown in Figure 2e. The heat map shows the relative expression levels of IP in the three groups of ovarian samples, with similar performance in the three samples of the same group (Fig. 2f). Interestingly, the distribution trend of m⁶A methylation sites in the yak genome of the three different samples was consistent (Fig. 2g). Chr 20 contains more m⁶A sites, followed by chr 8, and chr 23 contains the least m⁶A sites. Due to this, this result is slightly different from the identification result obtained in the rat brain tissue[16]. Each gene or transcript contains 1–3 m⁶A peaks, and more than half of the methylated mRNAs contain only one m⁶A peak. To analyze the topological pattern of m⁶A in the yak ovary, the gene distribution of all identified m⁶A peaks in YO-A, YO-F, and YO-P transcripts was statistically analyzed. It was found that the m⁶A peak was highly enriched in the coding sequence (CDS) region and 3' untranslated region (3' UTR) (Fig. 2h). This feature indicates that the distribution of m⁶A methylation in the gene structure of organisms is highly conserved.

Identification of the differentially methylated RNAs

The ovarian samples obtained from the three yaks- YO-A, YO-F, and YO-P were divided into three comparison groups, YO-FvsA, YO-FvsP, and YO-PvsA. The comparative analysis of these three groups separately screened 1186, 1890, and 1528 differentially m⁶A peaks in the YO-FvsA, YO-FvsP, and YO-PvsA (Supplementary file 3). Compared to YO-A group, 574 m⁶A peaks were hypermethylated and 612 m⁶A peaks were hypomethylated in the YO-F group (Fig. 3a); compared to the YO-P group, 1611 m⁶A peaks were hypermethylated and 279 m⁶A peaks were hypomethylated in the YO-F group (Fig. 3b); compared to the YO-A group, 182 m⁶A peaks were hypermethylated, and 1346 hypomethylated m⁶A peaks in YO-P group (Fig. 3c). Overall, the differences of m⁶A in the ovarian mRNAs were the greatest between the follicular phase and pregnancy, and the smallest between the follicular phase and anestrus. The motif with high reliability in the peak region was analyzed using the motif analysis software, Homer, and the width, E-value, position frequency matrix (PFM), position-specific scoring matrix (PSSM) of each motif, and its total position information in each peak sequence were obtained. The sequence modified by m⁶A was identified as “RRACH” (R is A/G, A is m⁶A, H is A/C/U)[17, 18]. To determine whether the identified m⁶A peaks were enriched on the common sequence of RRACH, we performed the motif prediction for each group and the differential results, and found consistent results in the differential methylation peaks of YO-FvsA, YO-FvsP, and YO-PvsA (Fig. 3d).

The GO and KEGG analysis of differentially methylated RNAs

To estimate the potential biological significance of m⁶A methylation in the ovary, we divided the differential methylation peaks in YO-FvsA, YO-FvsP, and YO-PvsA into hypermethylated and hypomethylated DMRs (differentially methylated RNAs). Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) were used to analyze the gene function enrichment of these DMRs with significantly changes in the methylation level of each comparison group. The GO analysis of the hypermethylated and hypomethylated DMRs in YO-FvsA group (the number of genes involved in GO item > 2, P < 0.05) showed that in terms of molecular function, they were significantly correlated with protein binding, ion binding, catalytic activity, and transferase activity; In terms of biological processes, the hypermethylated and hypomethylated DMRs were significantly involved in cellular processes, metabolic processes, the regulation of biosynthesis and metabolic processes; In terms of cell components, it is mainly related to organelle organization (Supplementary file 4). The GO analysis of the YO-FvsP and YO-PvsA groups is similar to the results of the YO-FvsA group, as shown in the (Supplementary file 5 and 6) .

The KEGG analysis of the YO-FvsA group (Supplementary file 4) (the number of genes involved in the KEGG item > 2, P < 0.05) showed that the hypermethylated DMRs significantly participated in the selenocompound metabolism, ubiquinone and other terpenoid-quinone biosynthesis pathway, and the hypomethylated DMRs significantly participated in the Notch signal pathway, Parathyroid hormone synthesis, secretion and action, and NF-kappa B signaling pathway. The KEGG analysis results of the YO-FvsP group (Supplementary file 5) showed that hypermethylated DMRs peaks were significantly enriched in adipocyte factor signaling pathway, Th17 cell differentiation, notch signaling pathway, PPAR signaling pathway, AMPK signaling pathway, and hypomethylated DMRs were significantly enriched in pyruvate metabolism, dopaminergic synapse , sphingolipid signaling pathway. The KEGG analysis results of the YO-PvsA group (Supplementary file 6) showed that the hypermethylated DMRs were mainly related to AMPK signaling pathway and fatty acid metabolism, and the hypomethylated DMRs were mainly related to base excision repair and notch signaling pathway. These significantly enriched pathways may be related to special regulatory mechanisms such as hormone secretion on the ovary during estrus and follicular development. These results indicate that m⁶A is involved in multiple physiological activities of the ovary during estrus and pregnancy in yaks.

The association analysis between the differential m⁶A peaks and differentially expressed genes

To further correlate the m⁶A modification with gene expression, we combined the MERIP-seq and RNA-seq analyses to investigate whether the degree of m⁶A methylation was related to the mRNAs of differentially expressed genes (DEGs) (Supplementary file 7). To intuitively represent the association between gene expression and m⁶A methylation, we constructed a four-quadrant diagram of the differentially expressed genes and m⁶A methylation genes. The results showed that in the YO-FvsA group (Fig. 3e), 75 genes were identified as significantly co-differentially expressed genes, including 60 hypermethylated DMRs (37 mRNAs were upregulated, 23 mRNAs were downregulated) and 15 hypomethylated DMRs (10 mRNAs were upregulated, 5 mRNAs were downregulated); A total of 165 co-differentially expressed genes were

identified in the YO-FvsP group (Fig. 3f), including 139 hypermethylated DMRs (107 mRNAs were upregulated, 32 mRNAs were downregulated), and 26 hypomethylated DMRs (18 mRNAs were upregulated and 8 mRNAs were downregulated); A total of 63 co-differentially expressed genes were identified in the YO-PvsA group (Fig. 3g), including 25 hypermethylated DMRs (22 mRNAs were upregulated, 3 mRNAs were downregulated) and 38 hypomethylated DMRs (29 mRNAs were upregulated, 9 mRNAs were downregulated). We then found that the mRNAs related to ovarian and follicular development such as *BNC1*, *HOMER1*, *P4HA3*, *BMP15*, and *BMP6* in the YO-FvsA and YO-FvsP groups all showed significantly increased m⁶A levels compared to the comparison group. The transcriptome analysis showed that the *BNC1* transcription level was significantly upregulated in the estrus ovaries compared to the anestrus ovaries and pregnant ovaries, and the *BMP15* transcription level was also significantly upregulated in estrus ovaries compared to the pregnant ovaries. On the contrary, the transcription levels of *HOMER1* and *P4HA3* were significantly downregulated compared to the anestrus group, and the transcription of *BMP6* was significantly downregulated compared to the pregnant group. In the YO-PvsA group, we found the mRNA *GPX3* and *WNT11* with significantly lower m⁶A levels compared to the control group, while their transcription levels were significantly upregulated compared to the anestrus ovaries. Then, four key genes were selected from the three comparison groups, and the comparative analysis of gene qPCR results and transcriptome expression was carried out (Fig. 4). It was found that the expression trend of gene at qPCR level and transcription level was consistent, which also further verified the results of transcriptome.

Discussion

The m⁶A modification, evident in various tissues reportedly regulates the gene expression and executes corresponding biological functions by regulating the RNA metabolism, splicing, degradation and translation[19]. In recent years, reversible RNA methylation modifications have led to the third wave of epigenetic modifications based on DNA and protein modifications. Many studies related to m⁶A have been conducted in the model organisms such as humans, mice, and *Arabidopsis thaliana*, and common organisms such as pigs and poultry, but there are no reports on the map of m⁶A in yak. To our knowledge, this is the first comprehensive and high-throughput study on RNA methylation in the yak ovary. Our data indicate that there is a significantly large amount of m⁶A methylation in the yak ovary during the estrus and pregnancy stages. Through further analysis, we found that m⁶A modification may play an important role in ovarian activity.

Many studies have shown that the m⁶A peaks are mainly concentrated in long exons, stop codons and 3' UTR. Our results also showed that these peaks are mainly concentrated in the exon region. These peaks aligned to the intron and intergenic regions may be due to pre-mRNA, splicing events, incomplete genome annotation, and background noise. The methylation and demethylation of RNA start from multiple binding proteins bind to the motif at the site where the methylation occurs. The motif is essentially a biologically significant nucleic acid sequence pattern that these RNA methylation-related enzymes can recognize and bind to influence the gene expression. The study of gene expression regulation

mechanisms is the focus of biological research. Identifying these motifs is of great significance for the study of mechanisms regulating gene expression. According to the published data, the consistent motif “RRACH” was overexpressed in the m⁶A motif region, our results also confirm it.

As the basic functional unit of the ovary, the follicle is responsible for the growth and ovulation of high-quality oocytes and produces steroid and peptide hormones which are important for reproductive physiology. An enormous amount of primordial follicles are stored in the ovary of mammals from birth. In our results, the number of growing follicles in the ovaries of yaks was less in the anestrus period. With the arrival of the estrus season, the dominant follicles began to appear in the ovaries of yaks and secreted a lot of estrogens, hence, the yaks showed estrus symptoms and ovulated. There will still be a wave of non-ovulatory follicles in the ovary during pregnancy, but the number of such follicles is less than that in the non-pregnant state. The types of follicles present in the ovaries during pregnancy are similar to those in the anestrus period, mainly replacement follicles, and fewer selected follicles and dominant follicles. The corpus luteum is usually formed in the ovaries during pregnancy and used to synthesize estrogen and progesterone. However, as the fetus develops, the corpus luteum is gradually replaced by the placental tissue, and the corpus luteum begins to shrink.

The formation, growth and function of follicles are controlled by the ovarian endocrine signal network. In this process, the hormones and cytokines strictly regulate the secretion of the gonadal hormones and expression of steroid-producing proteins in the corpus luteum and granulosa cells through various signaling pathways such as the PI3K/AKT, mTOR, transforming growth factor β , and Notch signaling pathways[20–22]. Interestingly, we also found consistent results after performing KEGG on the hypermethylated genes and hypomethylated genes of each comparison group, the Notch signaling pathway appeared in all three comparison groups. The Notch signaling plays a significant role in ovarian function, such as follicle assembly and growth, oocyte meiotic maturation, ovarian angiogenesis and steroid production[21, 22]. The expression of the Notch signal components in the mammalian ovary is conservative, which is mainly accomplished by the interaction of one of the four Notch receptors (Notch1–4) with one of the five notch ligands Jagged1 (Jag1), Jagged2 (Jag2), Delta like 1 (Dll1), Dll3 and Dll4[23]. The Notch signal has both inhibitory and stimulating effects on steroid synthesis. Wang et al.[24] reported that Notch signaling plays an inhibitory role in progesterone biosynthesis and steroidogenic protein expression in GCs. Jag1 is the most abundant Notch ligand expressed in the mouse ovary. According to Prasya and Mayo[25], the absence of Jag1 in mouse GCs results in the inhibition of granulosa cell differentiation, which shows that the expression of enzymes and factors involved in steroid biosynthesis and steroid secretion decreases, indicating that Notch signal has a stimulating effect on steroid synthesis. Conditional knockout of Notch2 in the granulosa cells and Jag1 in the oocytes leading to the formation of multiple oocyte follicles and subsequently declines the fertility of the mice[26, 27].

The development of mammalian follicles is coordinated by a complex interaction of multiple signals, including gonadotropins, paracrine/autocrine, and paracrine signals. Through the analysis of transcriptome and differential peaks, some genes related to ovarian function were found, such as *BNC1*, *HOMER1*, *BMP15*, *BMP6*, *GPX3*, and *WNT11*. *BNC1* appears to be a member of the mammalian maternal

effect gene family. Previous studies have detected the BNC1 protein in adult ovarian tissue, especially in the oocytes that are in the secondary follicular phase and the ovulating oocytes with nuclear-cytoplasmic expression patterns[28]. The female mouse model with *BNC1* frame-shift mutation showed infertility, due to the significantly increased serum follicle-stimulating hormone, and reduced ovary with a declined number of follicles [29]. Our results showed that the *BNC1* m⁶A levels were significantly increased in the ovaries during estrus and the expression of the mRNAs was also significantly higher than those in the ovaries during anestrus and pregnancy. Therefore, we speculated that the mRNA expression of the m⁶A modified *BNC1* gene is upregulated during estrus and ovulation, which promotes oocyte development and follicle formation. A study by Zhang et al. speculated that BNC1 might affect oocyte genesis through the oocyte-derived growth factor/signal transduction and PI3K pathway [28]. The bone morphogenetic proteins 6 (BMP6) and 15 (BMP15) are transforming growth factors- β (TGF- β) superfamily, secreted by the oocytes[30]. BMP15 has been shown to play an important role in mammalian ovarian and follicular development, including promoting the proliferation of the granulosa cells (GCSs) and steroid production, controlling the ability of oocytes, and ovulation[31, 32]. BMP6 is expressed in the oocytes and granulosa cells of healthy follicles and inhibits the activity of FSH by inhibiting the activity of adenylate cyclase and lead to the “selection” of dominant follicles[33]. In addition, BMP6 also prevents bovine cumulus cell apoptosis and preserves the normal oocyte quality. In our study, the m⁶A and mRNA levels of *BMP15* in the estrus ovaries were significantly higher than those in pregnancy. Therefore, the m⁶A methylation of BMP15 may play a potential role in promoting the growth and maturation of yak follicles.

We found that the expression levels of *METTL3* and *FTO* in the anestrus ovary were significantly higher than those in the estrus and pregnancy ovaries, instead, the expression of readings *YTHDF1*, *YTHDF2*, and *YTHDC2* during the estrus and pregnancy were significantly higher than those in the anestrus. Studies have shown that *METTL3*, *FTO*, *YTHDF2*, and *YTHDC2* are related to regulating mRNA degradation, mRNA stability, and gene expression [8, 34]. In estrous ovaries, the level of m⁶A and transcripts of BNC1 and BMP15 were up-regulated, indicating that methylase promoted the expression of mRNAs. The m⁶A levels of *HOMER1*, *P4HA3*, and *BMP6* in the ovary during estrus were significantly up-regulated compared with the anestrus and pregnant ovary. However, in transcription level, the *HOMER1* and *P4HA3* were significantly down-regulated compared with the anestrus ovary, and the transcription level of *BMP6* was significantly lower than that of the pregnant ovary. This phenomenon might be related to the regulation of the degradation of mRNAs and the inhibition of gene expression levels by methylases. However, the specific methylase and the regulatory mechanism are still unclear.

Conclusions

In summary, this study analyzed the m⁶A methylation modification in the ovary tissues of the yak during the anestrus, estrus, and pregnancy. The differences in the expression of *METTL3*, *METTL14*, *FTO*, and other enzymes and the functional enrichment analysis of the methylated genes suggest that m⁶A methylation might influence the regulation of the yak follicles development and the expression of sex hormone secretion-related genes. This study shows that BNC1, BMP15, and BMP6 are most likely to play

a key role in regulating the growth and development of follicles. In addition, the data obtained by the high-throughput sequencing provides a foundation for future research on the role of m⁶A methylation in the process of animal estrus and ovulation.

Abbreviations

m⁶A: N6-methyladenosine; CDS: coding sequence; 3'UTR: 3'untranslated region; MeRIP-seq: methylated RNA immunoprecipitation sequencing; DMRs: differentially methylated RNAs; GO: Gene Ontology; KEGG: Kyoto Encyclopedia of Genes and Genomes; DEGs: differentially expressed genes

Declarations

Ethics approval and consent to participate

All the ovarian samples were collected from the euthanized yak within 1 h. All the procedures involving animals were performed according to the guidelines of the China Council on Animal Care and the Ministry of Agriculture of the People's Republic of China. The Animal Care and Use Committee of the Lanzhou Institute of Husbandry and Pharmaceutical Sciences Chinese Academy of Agricultural Sciences approved all yak handling procedures (Permit No: SYXK-2014-0002).

Consent for publication

Not applicable.

Availability of data and material

The dataset supporting the conclusions of this article is available in the GEO repository, unique persistent identifier and hyperlink to datasets in <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE180401>. The data supporting the conclusions of this study are available within the Additional files.

Competing interests

The authors declare no competing interests.

Funding

This study was financially supported by the China Agriculture Research System of MOF and MARA (CARS-37) and Innovation Project of Chinese Academy of Agricultural Sciences (25-LZIHPS-01).

Authors' Contributions

Conceptualization, X.G. and P.Y.; Investigation, X.W.2 and L.X.; Methodology and software, M.C.1, M.C.2 and C.L.; Data curation, S.G. and X.W.1; Writing—original draft, S.G.; Writing—review& editing, P.B. and J.P.;

funding acquisition, X.G and P.Y. All authors contributed to the interpretation of the results, and writing of the article.

Acknowledgements

The authors would like to thank the technical assistance in transcriptome sequencing from LC-Bio Company (Hangzhou, China).

References

1. Qiu Q, Zhang G, Ma T, Qian W, Wang J, Ye Z, Cao C, Hu Q, Kim J, Larkin DM *et al*: **The yak genome and adaptation to life at high altitude**. *Nature genetics* 2012, **44**(8):946-949.
2. Zi XD: **Reproduction in female yaks (*Bos grunniens*) and opportunities for improvement**. *Theriogenology* 2003, **59**(5-6):1303-1312.
3. Gershon E, Dekel N: **Newly Identified Regulators of Ovarian Folliculogenesis and Ovulation**. *International journal of molecular sciences* 2020, **21**(12) :4565.
4. Ding C, Zou Q, Ding J, Ling M, Wang W, Li H, Huang B: **Increased N6-methyladenosine causes infertility is associated with FTO expression**. *Journal of cellular physiology* 2018, **233**(9):7055-7066.
5. Xia H, Zhong C, Wu X, Chen J, Tao B, Xia X, Shi M, Zhu Z, Trudeau VL, Hu W: **Mettl3 Mutation Disrupts Gamete Maturation and Reduces Fertility in Zebrafish**. *Genetics* 2018, **208**(2):729-743.
6. Desrosiers R, Friderici K, Rottman F: **Identification of methylated nucleosides in messenger RNA from Novikoff hepatoma cells**. *Proc Natl Acad Sci U S A* 1974, **71**(10):3971-3975.
7. Perry RP, Kelley DE: **Existence of methylated messenger RNA in mouse L cells**. *Cell* 1974, **1**(1):37-42.
8. Xu K, Yang Y, Feng GH, Sun BF, Chen JQ, Li YF, Chen YS, Zhang XX, Wang CX, Jiang LY *et al*: **Mettl3-mediated m⁶A regulates spermatogonial differentiation and meiosis initiation**. *Cell research* 2017, **27**(9):1100-1114.
9. Tang C, Klukovich R, Peng H, Wang Z, Yu T, Zhang Y, Zheng H, Klungland A, Yan W: **ALKBH5-dependent m⁶A demethylation controls splicing and stability of long 3'-UTR mRNAs in male germ cells**. *Proc Natl Acad Sci U S A* 2018, **115**(2):E325-e333.
10. Haussmann IU, Bodi Z, Sanchez-Moran E, Mongan NP, Archer N, Fray RG, Soller M: **m⁶A potentiates Sxl alternative pre-mRNA splicing for robust *Drosophila* sex determination**. *Nature* 2016, **540**(7632):301-304.
11. Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, Cesarkas K, Jacob-Hirsch J, Amariglio N, Kupiec M *et al*: **Topology of the human and mouse m⁶A RNA methylomes**

revealed by m6A-seq. *Nature* 2012, **485**(7397):201-206.

12. Alam H, Maizels ET, Park Y, Ghaey S, Feiger ZJ, Chandel NS, Hunzicker-Dunn M: **Follicle-stimulating hormone activation of hypoxia-inducible factor-1 by the phosphatidylinositol 3-kinase/AKT/Ras homolog enriched in brain (Rheb)/mammalian target of rapamycin (mTOR) pathway is necessary for induction of select protein markers of follicular differentiation.** *The Journal of biological chemistry* 2004, **279**(19):19431-19440.
13. Meyer KD, Saletore Y, Zumbo P, Elemento O, Mason CE, Jaffrey SR: **Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons.** *Cell* 2012, **149**(7):1635-1646.
14. Cui Y, Yu SJ: **An anatomical study of the internal genital organs of the yak at different ages.** *Veterinary journal (London, England : 1997)* 1999, **157**(2):192-196.
15. Li Y, Wang X, Li C, Hu S, Yu J, Song S: **Transcriptome-wide N⁶-methyladenosine profiling of rice callus and leaf reveals the presence of tissue-specific competitors involved in selective mRNA modification.** *RNA biology* 2014, **11**(9):1180-1188.
16. Yu J, Zhang Y, Ma H, Zeng R, Liu R, Wang P, Jin X, Zhao Y: **Epitranscriptomic profiling of N⁶-methyladenosine-related RNA methylation in rat cerebral cortex following traumatic brain injury.** *Molecular brain* 2020, **13**(1):11.
17. Wang X, Lu Z, Gomez A, Hon GC, Yue Y, Han D, Fu Y, Parisien M, Dai Q, Jia G *et al*: **N⁶-methyladenosine-dependent regulation of messenger RNA stability.** *Nature* 2014, **505**(7481):117-120.
18. Csepány T, Lin A, Baldick CJ, Jr., Beemon K: **Sequence specificity of mRNA N⁶-adenosine methyltransferase.** *The Journal of biological chemistry* 1990, **265**(33):20117-20122.
19. Fu Y, Dominissini D, Rechavi G, He C: **Gene expression regulation mediated through reversible m⁶A RNA methylation.** *Nature reviews Genetics* 2014, **15**(5):293-306.
20. Stocco C, Telleria C, Gibori G: **The molecular control of corpus luteum formation, function, and regression.** *Endocrine reviews* 2007, **28**(1):117-149.
21. Vanorny DA, Mayo KE: **The role of Notch signaling in the mammalian ovary.** *Reproduction (Cambridge, England)* 2017, **153**(6):R187-r204.
22. Knight PG, Glister C: **TGF-beta superfamily members and ovarian follicle development.** *Reproduction (Cambridge, England)* 2006, **132**(2):191-206.
23. Boggs K, Henderson B, Reisman D: **RBP-Jkappa binds to and represses transcription of the p53 tumor suppressor gene.** *Cell biology international* 2009, **33**(3):318-324.

24. Wang Y, Lu E, Bao R, Xu P, Feng F, Wen W, Dong Q, Hu C, Xiao L, Tang M *et al*: **Notch signalling regulates steroidogenesis in mouse ovarian granulosa cells.** *Reproduction, fertility, and development* 2019, **31**(6):1091-1103.
25. Prasasya RD, Mayo KE: **Notch Signaling Regulates Differentiation and Steroidogenesis in Female Mouse Ovarian Granulosa Cells.** *Endocrinology* 2018, **159**(1):184-198.
26. Xu J, Gridley T: **Notch2 is required in somatic cells for breakdown of ovarian germ-cell nests and formation of primordial follicles.** *BMC biology* 2013, **11**:13.
27. Vanorny DA, Prasasya RD, Chalpe AJ, Kilen SM, Mayo KE: **Notch signaling regulates ovarian follicle formation and coordinates follicular growth.** *Molecular endocrinology (Baltimore, Md)* 2014, **28**(4):499-511.
28. Zhang D, Liu Y, Zhang Z, Lv P, Liu Y, Li J, Wu Y, Zhang R, Huang Y, Xu G: **Basonuclin 1 Deficiency is a Cause of Primary Ovarian Insufficiency.** *Human Molecular Genetics* 2018, **27** (21):3787-3800.
29. Ma J, Zeng F, Schultz RM, Tseng H: **Basonuclin: a novel mammalian maternal-effect gene.** *Development (Cambridge, England)* 2006, **133**(10):2053-2062.
30. Zhai B, Liu H, Li X, Dai L, Gao Y, Li C, Zhang L, Ding Y, Yu X, Zhang J: **BMP15 prevents cumulus cell apoptosis through CCL2 and FBN1 in porcine ovaries.** *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology* 2013, **32**(2):264-278.
31. Fabre S, Pierre A, Mulsant P, Bodin L, Di Pasquale E, Persani L, Monget P, Monniaux D: **Regulation of ovulation rate in mammals: contribution of sheep genetic models.** *Reproductive biology and endocrinology : RB&E* 2006, **4**:20.
32. Hussein TS, Thompson JG, Gilchrist RB: **Oocyte-secreted factors enhance oocyte developmental competence.** *Developmental biology* 2006, **296**(2):514-521.
33. Otsuka F, Moore RK, Shimasaki S: **Biological function and cellular mechanism of bone morphogenetic protein-6 in the ovary.** *The Journal of biological chemistry* 2001, **276**(35):32889-32895.
34. Sui X, Hu Y, Ren C, Cao Q, Zhou S, Cao Y, Li M, Shu W, Huo R: **METTL3-mediated m⁶A is required for murine oocyte maturation and maternal-to-zygotic transition.** *Cell cycle (Georgetown, Tex)* 2020, **19**(4):391-404.
35. Zi XD, He SM, Lu H, Feng JA, Lu JY, Chang S, Wang X: **Induction of estrus in suckled female yaks (*Bos grunniens*) and synchronization of ovulation in the non-sucklers for timed artificial insemination using progesterone treatments and Co-Synch regimens.** *Anim Reprod Sci* 2006, **92**(1-2):183-192.
36. Chen S, Zhou Y, Chen Y, Gu J: **fastp: an ultra-fast all-in-one FASTQ preprocessor.** *Bioinformatics* 2018, **34**(17):i884-i890.

37. Kim D, Langmead B, Salzberg SL: **HISAT: a fast spliced aligner with low memory requirements.** *Nat Methods* 2015, **12**(4):357-360.
38. Meng J, Lu Z, Liu H, Zhang L, Zhang S, Chen Y, Rao MK, Huang Y: **A protocol for RNA methylation differential analysis with MeRIP-Seq data and exomePeak R/Bioconductor package.** *Methods* 2014, **69**(3):274-281.
39. Yu G, Wang LG, He QY: **ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization.** *Bioinformatics* 2015, **31**(14):2382-2383.
40. Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS: **MEME SUITE: tools for motif discovery and searching.** *Nucleic Acids Res* 2009, **37**(Web Server issue):W202-208.
41. Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL: **StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.** *Nat Biotechnol* 2015, **33**(3):290-295.
42. Robinson MD, McCarthy DJ, Smyth GK: **edgeR: a Bioconductor package for differential expression analysis of digital gene expression data.** *Bioinformatics* 2010, **26**(1):139-140.

Figures

Figure 1

The microscopic observation of the tissue sections of the yak ovaries. YO-A, YO-F, and YO-P separately represent the yak ovary in the anestrus period, follicular (estrus) period, and pregnancy period.

Figure 2

The results of m6A methylase expression and genome sequencing. a. The gene expression levels of 13 common m6A methylases on the three groups of YO-A, YO-F and YO-P. b. The principal component analysis of the three groups of samples. c. The distribution of different types of reads in each sample was counted. d. The valid data were matched to the reference genome defined as matched to exon, intron and intergenic. e. The length distribution of m6A peaks in the YO-A, YO-F, and YO-P groups. f. The relative

expression levels of IP in the three groups of ovarian samples. g. The distribution of m6A peak sites in the yak genome was obtained from three different samples. The outermost circle represents the chromosome distribution on the genome, the red circle represents the YO-A group, the purple circle represents the YO-F group, and the blue circle represents the YO-P group. h. Density distribution of the m6A peaks in YO-A, YO-F and YO-P transcripts in the different gene structures.

Figure 3

Overview of the differential methylation peaks in the YO-FvsA, YO-FvsP, and YO-PvsA comparison groups. a. The number of hypermethylation and hypomethylation in the YO-FvsA group. b. The number of hypermethylation and hypomethylation in the YO-FvsP group. c. The number of hypermethylation and hypomethylation in the YO-PvsA group. d. The motif sequences “RRACH ” were identified in the YO-FvsA, YO-FvsP, and YO-PvsA groups. e. The four-quadrant diagram of the differentially expressed genes and m6A methylation genes in the YO-FvsA group. f. The four-quadrant diagram of the differentially expressed genes and m6A methylation genes in the YO-FvsP group. g. The four-quadrant diagram of differentially expressed genes and m6A methylation genes in the YO-PvsA group.

Figure 4

The gene qPCR results and transcriptome expression levels of BNC1, HOMER1, MMP16, P4HA3, BMP15, BMP6, ISG15, CCRL2, GPX3, WNT11, CYP1B1, and ITGA2 in the YO-A, YO-F and YO-P groups.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.xlsx](#)
- [Additionalfile2.xlsx](#)
- [Additionalfile3.xlsx](#)
- [Additionalfile4.tif](#)
- [Additionalfile5.tif](#)
- [Additionalfile6.tif](#)
- [Additionalfile7.xlsx](#)
- [Additionalfile8.xlsx](#)