PSP2 , a gene encoding RGG-motif protein, is a novel suppressor of clathrin heavy chain deficiency

Debadrita Roy

Indian Institute of Science

Mani Garg

Indian Institute of Science <https://orcid.org/0000-0003-3794-418X>

Purusharth I Rajyaguru (✉ rajyaguru@iisc.ac.in)

<https://orcid.org/0000-0003-1478-1787>

Research article

Keywords: Clathrin heavy chain, translation repressor, SCD6, PSP2, SBP1, DED1, DHH1, GBP2 and RGG-motif

Posted Date: November 20th, 2019

DOI: <https://doi.org/10.21203/rs.2.17572/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

***PSP2*, a gene encoding RGG-motif protein, is a novel suppressor of clathrin heavy chain deficiency**

Debadrita Roy, Mani Garg and Purusharth I Rajyaguru*

Department of Biochemistry, Indian Institute of Science, C V Raman road, Bangalore – 560012

Author for correspondence: *Purusharth I Rajyaguru, Department of Biochemistry, Indian Institute of Science, C V Raman road, Bangalore – 560012; email: rajyaguru@iisc.ac.in

Running title: RGG-motif proteins suppress clathrin heavy chain deficiency.

Keywords: Clathrin heavy chain, translation repressor, *SCD6*, *PSP2*, *SBP1*, *DED1*, *DHH1*, *GBP2* and RGG-motif.

Abstract

Clathrin, made up of the heavy- and light-chains, constitutes one of the most abundant protein in vesicles involved in intracellular protein trafficking and endocytosis. *YPR129W*, which encodes RGG-motif containing translation repressor was identified as a part of multi-gene construct (*SCD6*) that suppressed clathrin deficiency. However, the contribution of *YPR129W* alone in suppressing clathrin deficiency has not been documented. In this study we identify *YPR129W* as a necessary and sufficient gene in a multigene construct *SCD6* that suppresses clathrin deficiency. Importantly, we identify cytoplasmic RGG-motif protein encoding gene *PSP2* as a novel suppressor of clathrin deficiency. Three other RGG-motif protein encoding genes *SBP1*, *DED1* and *GBP2* do not suppress clathrin deficiency. *DHH1*, a DEAD-box RNA helicase with translation repression activity also fails to rescue clathrin deficiency. α -factor secretion assay suggests that suppression of clathrin deficiency by *SCD6* and *PSP2* is not mediated by the rescue of the trans-Golgi network (TGN) protein sorting defect observed in the absence of *CHC1*. Detailed domain analysis of the two suppressors reveals that the RGG-motif of both *Scd6* and *Psp2* is important for suppressing clathrin deficiency. Additionally, the Lsm domain deletion as well as the arginine to alanine mutation in the arginine methylation defective (AMD) mutant render *Scd6* defective in suppressing clathrin deficiency. Overall based on our results using *SCD6* and *PSP2* proteins, we identify a novel role of RGG-motif in suppressing clathrin deficiency. Since both the suppressors are RNA-binding granule-resident proteins, this study opens an exciting avenue for exploring the connection between clathrin function and cytoplasmic RNA metabolism.

Introduction

Clathrin is a protein of vital importance in cells. Clathrin coated vesicles (CCVs) play a key role in endocytosis and various intracellular trafficking. The triskelion, which is made up of the clathrin light chain and heavy chain monomers, forms a typical polygonal surface lattice on CCVs. Clathrin mediated endocytosis, which is highly conserved from yeast to humans, is the major pathway for the internalization of external and membrane molecules into the cell. There are clathrin-independent pathways, however without clathrin-mediated endocytosis, uptake is greatly reduced (Bitsikas et al. 2014; Kaksonen and Roux 2018; Mayor and Pagano 2007; Prosser et al. 2011). CCVs are also involved in the intracellular trafficking pathway. Deficiency of clathrin in unicellular organisms like yeast, amoebae and protozoa are associated with defects in intracellular trafficking and receptor-mediated endocytosis (Allen et al. 2003; Seeger and Payne 1992a).

In certain yeast strains, the deletion of the clathrin heavy chain-1 gene (*CHC1*) renders cells inviable, whereas in others, cells are viable. This is owing to the presence of an independently segregating gene – suppressor of clathrin deficiency 1 (*SCD1*). Presence of the *scd1-i* allele leads to lethality in *chc1* deficient cells, whereas the *scd1-v* allele allows growth, albeit poorly. Along with slow growth, *chc1Δ scd1-v* cells have abnormal morphology, genetic instability, reduced endocytosis and mis-localization of endoproteases from the *trans*-Golgi network to the cell membrane (Lemmon et al. 1990; Lemmon and Jones 1987; Nelson et al. 1996; Payne and Schekman 1985; Payne and Schekman 1989; Seeger and Payne 1992b). *SCD1* was recently identified to encode *PAL2*, a protein involved in localizing to cortical patches with other endocytic factors, thereby facilitating endocytosis. It was found that *scd1-i* allele has a premature stop codon which results in a truncated version of wild type *PAL2*, encoded by *scd1-v* allele (Moorthy et al. 2019). YEpSCD6 was one of the six multi-gene multicopy suppressors identified in a genetic screen (Nelson and Lemmon 1993). This suppressor contains a total of 8 genes along with *YPR129W* which encodes an RGG-motif translation repressor protein.

RGG-motif proteins contain single or multiple repeats (Table 1) of RGG/RGX ('X' being any residue). This motif is known to be involved in protein-nucleic acids and protein-protein interactions (Thandapani et al. 2013). The RGG-motif containing proteins function in processes like transcription, cell cycle, apoptosis and synaptic plasticity. The RGG-motif constitutes the second most common class of domains/motifs that contribute to RNA-binding (Ozdilek et al. 2017). Recently, a subset of yeast RGG-motif proteins (such as Scd6, Sbp1, Npl3 and Ded1) have been implicated in translation control (Hilliker et al. 2011; Rajyaguru et al. 2012) as these proteins bind conserved translation initiation factor eIF4G through their RGG-motif to repress translation. Scd6 and Sbp1 are translation repressors and decapping activator proteins. Consistent with their role, both Scd6 and Sbp1 localize to RNA granules that are sites of translation repression and mRNA decay (Mitchell et al. 2013; Rajyaguru et al. 2012; Segal et al. 2006). *PSP2* encodes a cytosolic RGG-motif (Table 1) containing protein originally identified as a suppressor of intron-splicing defects of a mutation in *MRS2* and subsequently as a suppressor of conditional mutation in DNA POL I in yeast (Formosa and Nittis 1998; Waldherr et al. 1993). Interestingly, Psp2 localizes to processing bodies during glucose deprivation (Mitchell et al. 2013). *DED1* encodes a DEAD-box RNA helicase and a RNA granule resident protein that can both promote and inhibit translation initiation in yeast (Beckham et al. 2008; lost et al. 1999). *GBP2* is nucleo-cytoplasmic shuttling protein involved in telomere length modification (Pang et al. 2003), mRNA export (Häcker and Krebber 2004; Hurt et al. 2004;

Windgassen and Krebber 2003) and mRNA quality control (Hackmann et al. 2014). Despite being a nuclear protein, *GBP2* associates with polysomes and is a component of stress granules (Buchan et al. 2008; Windgassen et al. 2004) the significance of which remains to be understood. *DHH1* is a DEAD-box RNA helicase that acts as a translation repressor and decapping activator (Coller and Parker 2005). It localizes to RNA granules and regulates P-body assembly in ATP-dependent manner (Mugler et al. 2016). In this work, we report that *YPR129W* is a bonafide suppressor of clathrin deficiency. We identify *PSP2* a gene encoding RNA-binding and RGG-motif protein as a novel suppressor of clathrin deficiency. The RGG-motif of both *Scd6* and *Psp2* is important for suppression of clathrin deficiency. These results establish an unprecedented link between clathrin heavy chain function and granule-resident RGG-motif proteins. Our observations pave the way for exploring the mechanistic basis underlying this link in the future.

Table 1: Number of RGG/RGX in different proteins tested in this study

Protein (# of amino acids)	# of RGG-/RGX-	Location of RGG-/RGX- motifs
Scd6 (349)	1/7	C-terminal
Psp2 (593)	4/10	C-terminal
Sbp1 (294)	8/5	Central
Ded1 (604)	4/4	Dispersed
Gbp2 (427)	4/6	Predominantly towards N-terminal

RESULTS

***YPR129W* is necessary and sufficient to suppress clathrin deficiency**

YE Δ SCD6, a high copy number multi-gene plasmid was reported to be capable of rescuing inviability of clathrin heavy chain-deficient yeast strain expressing endogenous *CHC1* under a galactose-inducible promoter (Nelson and Lemmon 1993). This multi-gene construct contained 7 other genes along with *YPR129W*. A personal communication on *Saccharomyces* Genome Database (SGD) implicated *YPR129W* as a suppressor of clathrin deficiency (Gelperin et al. 1995, personal communication to SGD), however evidence supporting it is not published. We were intrigued by the possible role of an RNA-binding translation repressor protein in suppressing clathrin heavy chain requirement and therefore wanted to indeed confirm the role of *YPR129W* in clathrin function. To test in our hands, the necessity and sufficiency of *YPR129W*, we created YE Δ SCD6 (lacking *YPR129W*) and YEonlySCD6 (YEpoSCD6; expressing only *YPR129W*) by site directed mutagenesis. These plasmids were transformed into *GAL1:CHC1* strain and the transformants were assayed for growth on glucose and galactose. We observed that YE Δ SCD6 transformant grew as poorly as the empty vector control on glucose plate (Figure 1A) indicating that in the absence of *SCD6* none of the other 7 genes could suppress the clathrin deficiency growth defect. Thus, *SCD6* was necessary in the multi-gene construct to suppress clathrin deficiency mediated growth defect. YEpoSCD6 transformed cells grew in manner comparable to YE Δ SCD6 transformed cells suggesting that *SCD6* was sufficient to suppress the clathrin deficiency mediated growth defect (Figure 1A). In order to quantify the extent of rescue, a plating assay was carried out on YEP media, wherein after depleting *Chc1*, cells were plated on galactose and glucose followed by counting the colonies. Consistent with the growth assay, cells expressing YE Δ SCD6 and YEpoSCD6 but not YE Δ scd6 could suppress clathrin deficiency.

(Figure 1B). To further confirm its role, we cloned *YPR129W* in a 2u plasmid pGP564 with a C-terminal His-tag and checked its ability to suppress clathrin deficiency. We observed that *SCD6* suppressed the clathrin deficiency-mediated growth defect in a manner comparable to *YE_pSCD6* (Figure 1C). Based on these results we confirmed and established that *YPR129W* is a suppressor of clathrin deficiency.

***PSP2* is a novel multicopy suppressor of clathrin deficiency**

Scd6 is an RGG-motif containing RNA granule-resident protein (Rajyaguru et al. 2012). We tested *PSP2* that encodes another granule-resident protein with RGG-motif for its ability to suppress clathrin deficiency. *PSP2* was expressed from 2μ plasmid in a strain expressing clathrin under galactose-inducible promoter. Strikingly, we observed that *PSP2* suppressed clathrin deficiency mediated growth defect (Figure 2A). Plating analysis followed by colony count confirmed the results obtained using growth assay (Figure 2B). The percentage rescue by *PSP2* improved when the plating assay was performed on selective synthetic media plates likely due to increased retention of the plasmid (Supp. Figure 1). Based on these results we conclude that *PSP2* is a novel suppressor of clathrin deficiency.

***DED1*, *SBP1*, *DHH1* and *GBP2* do not suppress clathrin deficiency**

SCD6 and *PSP2* encode RGG-motif containing proteins whereas *Scd6* is also a translation repressor. We wanted to check whether other genes encoding RGG-motif proteins and/or translation repressors can suppress clathrin deficiency. We therefore tested *DED1*, *SBP1*, *DHH1* and *GBP2*. *Sbp1*, *Ded1* and *Dhh1* can act as translation repressor proteins. *Sbp1* and *Ded1* (but not *Dhh1*) contains RGG-/RG- motifs which vary in number of repeats and location in the protein sequence (Table 1). *Gbp2* is a shuttling RGG-motif protein which predominantly localizes to the nucleus. Interestingly none of the four genes suppressed the clathrin deficiency mediated growth defect (Figure 3A). Plating assay further confirmed the growth assay results (Figure 3B). These results indicate that genes encoding RGG-motif containing or translation repressor proteins in general do not suppress clathrin deficiency highlighting the specificity of suppression of clathrin deficiency by *SCD6* and *PSP2*.

SCD6* and *PSP2* do not rescue the trans-Golgi network (TGN) sorting function of the *CHC1

We next tested the ability of *SCD6* and *PSP2* to rescue the TGN sorting defects of *CHC1* depleted cells. MATα cells are defective in the secretion of mature form of the mating pheromone α-factor, in the absence of clathrin. This is due to the mis-localization of α-factor processing enzymes from the trans-Golgi network to the plasma memberane (Payne and Schekman 1989). As a result, MATα cells with defective clathrin function do not inhibit the growth of MATα cells otherwise evident by a zone of growth inhibition (halo) in the halo assay. Since *SCD6* and *PSP2* rescue the growth defect phenotypes upon depletion of *Chc1*, we decided to test their contribution to α-factor secretion. *GAL1:CHC1* MATα cells overexpressing plasmid encoding *CHC1*, *SCD6*, *SBP1*, *PSP2*, *GBP2* or *DED1* were depleted of *Chc1* by growing on glucose and spotted on a lawn of MATα cells expressing the *sst1-2* allele (BJ3556 strain in Table 2). *GAL1:CHC1* MATα cells overexpressing *CHC1* from a plasmid form a clear zone of growth inhibition (halo) however such a zone of inhibition is absent for cells overexpressing any of the tested genes including *SCD6* and *PSP2* (Figure 4). We interpret this result to suggest that the suppression of clathrin deficiency growth defect by *SCD6* and *PSP2* does not involve complementation of the TGN function.

RGG-motif plays a vital role in suppressing clathrin-deficiency

Both *SCD6* and *PSP2* contain RGG-motif rich C-terminal region. *SCD6* also contains an Lsm domain at its N-terminus and a central FDF domain. *PSP2* does not contain any other canonical domain/motif other than the RGG-motif. We investigated which domain of these proteins was responsible for the observed suppression of growth defect. To this end, we generated domain deletion constructs and assayed them for suppression of clathrin deficiency mediated growth defect. The RGG-motif deletion construct of *SCD6* was highly defective in suppressing clathrin deficiency (Figure 5A). The Lsm domain deletion construct was also defective in suppressing clathrin deficiency albeit to slightly lesser extent than the RGG-motif deletion construct. Deletion of the FDF motif weakly affected the ability of the mutant to suppress clathrin deficiency. The RGG-motif (282-348 residues) of *SCD6* as annotated in literature (Anantharaman and Aravind 2004) comprises of RGG-rich sequences (282-310 residues) and QN-rich sequences (311-348 residues). Deletion of just the RGG-rich sequences (282-310) (*scd6RGG'*) resulted in a defect that was comparable to the deletion of 282-348 residues indicating that the *SCD6* RGG-rich sequence was indeed important for the rescue of clathrin deficiency. Based on these results we conclude that all the three domains affect the ability of *SCD6* to suppress clathrin deficiency. RGG-motif is the least dispensable motif and FDF-domain is the most dispensable domain for suppressing clathrin deficiency. Plating experiments followed by colony counting confirmed the growth assay results (Figure 5B). Interestingly the FDF domain deletion mutant was also defective in suppressing clathrin deficiency in the plating assay which was not evident as much in the growth assay (Figure 5A). The differential behaviour of this mutant in two different assays is interesting however the basis for this is unclear. Some of the arginine residues in the RGG-motif of *Scd6* are methylated and this modification promotes the repression activity (Poornima et al. 2016). We tested the role of arginine residues of the RGG-motif in suppressing clathrin deficiency. We observed that the arginine methylation defective mutant (AMD; with 9 R to A substitutions) failed to suppress the clathrin deficiency both in growth and plating assay (Figure 5C & D). It is possible that arginine methylation of *SCD6* RGG-motif could play a role in suppression of clathrin deficiency. Endogenous *Scd6* is expressed in low copy number [1280 molecules/cell (Ghaemmaghami et al. 2003)] and we believe that its expression is highly regulated. Detection of *Scd6* and its mutants expressed from pGP564 (Figure 5) using an anti-His antibody has been technically challenging. We however know that expression of both RGG-deletion and AMD mutant of *SCD6* is not compromised when expressed under galactose-inducible promoter from a 2u plasmid (BG1805 vector) and detected using PAP reagent that detects the C-terminal ZZ-tag (Poornima et al. 2016). We also know that expression of Lsm domain deletion mutant is not compromised when expressed in the BG1805 vector (Parbin et al., under review). These observations indicate that the inability of RGG-motif deletion, Lsm-domain deletion and AMD mutants to suppress clathrin deficiency is likely not due to compromised protein levels.

Apart from the RGG-motif, no other known domain/motif has been reported in *PSP2*. An RGG-deletion mutant of *PSP2* was created and tested for its ability to suppress clathrin deficiency. We observed that, like in the case of *SCD6*, the RGG-motif deletion mutant of *PSP2* was defective in suppressing clathrin deficiency (Figure 6A). Plating assays followed by colony counting further confirmed that the RGG-motif deletion mutant of *PSP2* was defective in rescuing growth on the glucose media (Figure 6B and Supp. Figure 1). We conclude that the RGG-motif of *PSP2* plays an important role in suppressing clathrin deficiency.

Discussion

We provide evidence suggesting that two genes encoding RGG-motif proteins function as suppressors of clathrin deficiency. To the best of our knowledge this is the first peer-reviewed report implicating RGG-motif proteins in clathrin function. Our conclusion is based on the following observations, a) *SCD6* acts as a suppressor of clathrin deficiency (Figure 1) b) *PSP2* is a novel suppressor of clathrin deficiency (Figure 2) c) RGG-motifs of Scd6 and Psp2 are required for suppressing clathrin deficiency (Figure 5 and 6) d) Arginine residues of the Scd6 RGG-motif play an important role in suppressing clathrin deficiency (Figure 5C and D).

SCD6 was identified as one of the genes in a multi-gene construct that suppressed clathrin heavy chain deficiency, however the specific contribution of *SCD6* remained to be confirmed. Our results prove that in the reported multi-gene construct, *SCD6* is both necessary and sufficient to suppress clathrin deficiency mediated growth defect. This along with the observation that *SCD6* cloned in a different vector also suppresses clathrin deficiency establishes it as a genuine suppressor of clathrin deficiency. Reported roles of *SCD6* orthologs point towards a role of this family of proteins in endocytosis and/or trafficking. *CAR-1* (worm ortholog) plays a role in maintaining ER organization as upon *CAR-1* knockdown, ER is disorganized into large patches and thick strands (Squirrell et al. 2006). Trailer hitch (*Drosophila* ortholog) localizes to ER-exit sites, which persist as large foci upon *Tral* knockdown (Wilhelm et al. 2005). Whether *SCD6* functions along similar lines as its worm and fly ortholog is unclear.

Suppression of clathrin deficiency by *PSP2* is a striking result. Psp2 is known to localize to RNA granules (Mitchell et al. 2013) and upon overexpression, can rescue P-body formation in *edc3Δlsm4Δc* strain (Rao and Parker 2017). Scd6 binds *TIF4631* (eIF4G1) to repress translation and interestingly Psp2 is reported to physically interact with its paralog *TIF4632* (Krogan et al. 2006). Although the functional relevance of Psp2-eIF4G2 interaction is not yet clear, we hypothesize that Psp2 could play a role in mRNA translation and decay like Scd6. Requirement of the Psp2 RGG-motif in suppressing clathrin deficiency (Figure 6B) points toward a mechanism similar to that of Scd6. Lack of suppression by other RGG-motif proteins and translation repressors (Sbp1, Ded1, Dhh1 and Gbp2) highlights the specificity of Scd6 and Psp2 in clathrin function. It is unlikely that the contribution of Scd6 and Psp2 in clathrin function is dependent on the number of RGG-/RGX- repeats as Sbp1 contains more repeats than Scd6 (Table 1) but fails to suppress. Interestingly both Scd6 and Psp2 harbor RGG-motifs at its C-terminal but none of the other tested proteins including Sbp1 have RGG-motifs at their C-terminus.

Clathrin deficiency suppression mechanisms might likely fall into one of the following two categories: a) Suppressors that directly contribute to clathrin function in vesicular transport and endocytosis and b) Suppressors that indirectly increase tolerance to clathrin deficiency. *SCD4* (Huang et al. 1997) and *SCD5* (Nelson et al. 1996) belong to the former category that contribute directly to vesicular transport. *SCD2* on the other hand belongs to latter category of suppressors. It was identified as *UBI4* (Nelson and Lemmon 1993) which likely suppresses clathrin deficiency defect by accelerating clearance of mis localized proteins We hypothesize that *SCD6* and *PSP2* might act in an indirect manner by repressing translation to reduce the load of newly synthesized proteins on the trafficking machinery. This hypothesis is based on the following observations: 1) RGG-motif, which is required for translation repression activity of *SCD6* (Poornima et al. 2019; Poornima et al. 2016;

Rajyaguru et al. 2012) is also required for suppressing clathrin deficiency defect by *SCD6* and *PSP2* 2) *SCD6* AMD mutant is defective in repressing translation as well as in suppressing clathrin deficiency 3) Lsm domain is required for the translation repression activity of Scd6 (Parbin et al., *communicated*) and also for suppressing clathrin deficiency. Further experiments will be required to test this hypothesis. It is also possible that translation control of specific mRNAs involved in endocytosis and/or trafficking by Scd6 and perhaps Psp2 contributes to the suppression of clathrin deficiency. Identification of specific mRNA targets of Scd6 and Psp2 will provide important insights in this regard. Since Scd6 localizes to RNA granules in RGG-motif dependent manner and RNA granules are sites of translation repression and mRNA decay, we tested the localization of Scd6 upon clathrin depletion. We did not observe any significant difference in localization of Scd6 to foci (data not shown) suggesting that suppression is likely not mediated by changes in localization to foci.

LSM14 is the human ortholog of Scd6 (Anantharaman and Aravind 2004; Marnef et al. 2009). The role of Scd6 identified in mRNA fate determination in yeast is conserved in humans. It is possible that LSM14 could contribute to clathrin mediated endocytosis and trafficking. This exciting possibility will be tested in future. Overall our results establish a new link between RGG-motif containing RNA-binding proteins and clathrin function. Analyzing the mechanistic basis of this link could lead to unraveling of the role of mRNA translation control pathway in clathrin mediated endocytosis. It will also be pertinent to test if the players contributing to endocytic/trafficking pathway including Chc1 are involved in modulating cytoplasmic mRNA fate. Our current work thus highlights several interesting research questions and addressing these questions will be an exciting future endeavor.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: All authors have consented for this manuscript.

Competing Interests: There are no conflict of interests.

Author's Contribution

PIR and DR designed the experiments, DR and MG performed the experiments, PIR wrote the first draft of the manuscript. PIR and MG revised and finalized the manuscript.

Acknowledgements

This work was predominantly supported by Department of Science and Technology grant (EMR/2017/001332) from the government of India. It was also supported by India Alliance DBT-Wellcome trust and DBT-IISc partnership program. DR was supported by DST-INSPIRE fellowship; MG is supported by University Grants Commission (UGC) fellowship. We are indebted to Sandra Lemmon (University of Miami) for helping us with various strains and reagents. We thank her for providing inputs at different stages of this work. We also sincerely thank Roy Parker (University of Colorado, Boulder) and Kenji Irie (University of Tsukuba) for sharing plasmids and strains.

Materials and methods

Table 2: List of strains used in this study

Name	Genotype	Description	Source
<i>GAL:CHC1</i>	<i>MATα GAL1:CHC1 scd1-i leu2 ura3-52 trp1 his1 GAL2</i>	Wild type <i>Saccharomyces cerevisiae</i> with inviable allele of <i>SCD1</i> and <i>CHC1</i> under galactose inducible promoter	(Nelson and Lemmon 1993)
BJ3556	<i>MATα sst1-2 ade2-1 his6 met1 cyh2 rme1 ura1 can1</i>	<i>Saccharomyces cerevisiae</i> with defective adaptation to alpha factor response pathway	(Newpher et al. 2006)
MAT α	<i>MATα leu2 ura3-52 trp1 his3-200</i>	Wild type <i>Saccharomyces cerevisiae</i> with 'a' mating type	(Newpher et al. 2006)
MAT α	<i>MATα leu2 ura3-52 trp1 his3-200</i>	Wild type <i>Saccharomyces cerevisiae</i> with 'alpha' mating type	(Newpher et al. 2006)

Table 3: List of plasmids used in this study

Name	Description	Vector	Source
YEp24 (empty vector)	YEp24 library empty vector, multicopy 2 μ plasmid with <i>URA3</i> and ampicillin resistance genes	YEp24	(Nelson and Lemmon 1993)
YEpSCD6	YEp24 cloned with a fragment from digested yeast genome containing <i>YPR129W</i> along with other flanking genes	YEp24	(Nelson and Lemmon 1993)
YEpoSCD6	Plasmid with only <i>SCD6</i> gene from YEpSCD6, other flanking genes deleted	YEp24	This study
YEp Δ scd6	Plasmids retaining all genes except <i>SCD6</i> from YEpSCD6	YEp24	This study
pGP564 (empty vector)	Yeast shuttle vector, multicopy 2 μ plasmid with <i>LEU2</i> and kanamycin resistance genes	pGP564	YGTC library
pGP <i>SCD6</i>	pGP564 containing <i>SCD6</i> under its own promoter. Along with the ORF 550 nucleotides upstream and 225 nucleotides downstream are present. A 6X-His tag is present before the stop codon of <i>SCD6</i> .	pGP564	This study
pGP <i>scd6ΔLsm</i>	pGP <i>SCD6</i> with Lsm domain deletion: amino acids deleted: 2-93	pGP564	This study
pGP <i>scd6ΔFDF</i>	pGP <i>SCD6</i> with FDF domain deletion; amino acids deleted: 199-281	pGP564	This study
pGP <i>scd6ΔRGG</i>	pGP <i>SCD6</i> with RGG domain (including QN rich part) deletion;	pGP564	This study

	amino acids deleted: 282-348		
pGP <i>scd6ΔRGG'</i>	pGP <i>SCD6</i> with RGG domain (excluding QN rich part) deletion, amino acids deletion: 282-310	pGP564	This study
pGP <i>scd6AMD</i>	pGP <i>SCD6</i> with R288A, R292A, R294A, R298A, R301A, R304A, R306A, R310A, R316A	pGP564	This study
pRS426 (empty vector)	Yeast shuttle vector, multicopy 2μ plasmid with <i>URA3</i> and ampicillin resistance genes	pRS426	Roy Parker, University of Colorado
pRS426 <i>PSP2</i>	pRS426 expressing <i>PSP2</i> under its own promoter	pRS426	(Rao and Parker 2017)
pRS426 <i>psp2ΔRGG</i>	pRS426 <i>PSP2</i> with RGG domain deletion; amino acids deleted: 418-577	pRS426	This study
pRS426 <i>DED1</i>	pRS426 expressing <i>DED1</i> under its own promoter	pRS426	(Hilliker et al. 2011)
pRS426 <i>SBP1</i>	pRS426 expressing <i>SBP1</i> under its own promoter	pRS426	(Segal et al. 2006)
YEplac195 (empty vector)	Yeast shuttle vector, multicopy 2μ plasmid with <i>URA3</i> and ampicillin resistance genes	YEplac195	Kenji Irie; University of Tsukuba
YEplac195 <i>DHH1</i>	YEplac195 expressing <i>DHH1</i> under its own promoter	YEplac195	Kenji Irie; University of Tsukuba
<i>Gbp2-GFP</i>	pPS1372 expressing <i>GBP2-GFP</i> under its own promoter, multicopy 2μ plasmid with <i>LEU</i> and ampicillin resistance genes	pPS1372	(Hackmann et al. 2014)

Growth Assay

Growth assays were performed to assess the ability of plasmids expressing desired genes to rescue clathrin deficiency mediated growth defect of the *GAL:CHC1* strain on glucose media. In *GAL:CHC1* strain, *CHC1* is under a galactose inducible promoter (Nelson and Lemmon 1993) which results in a growth defect on glucose media. Transformants grown on galactose containing media (10 OD₆₀₀) were, serially diluted and spotted on the respective minimal media containing glucose and galactose plates. All the plated were incubated at 30°C and imaged after 2 days (galactose media) or 4-5 days (glucose media). Growth assays were performed at least three times with comparable results.

Plating Assay

After growing cells in YEP-glucose for more than 16 -20h, different *GAL:CHC1* transformants were diluted to 0.001 OD₆₀₀. Three different volumes (50ul, 100ul and 200ul) were plated on YEP-galactose and glucose plates. The galactose containing plates were incubated for 4 days and glucose plates for 7 days before imaging. Colonies were counted and plotted as percentage rescue which

was calculated as [(no. of colonies on glucose/no. of colonies on galactose) *100]. Plating assays were performed at least three times with comparable results which were quantitated.

Halo Assay

The background tester MATa strain (BJ3556) carrying the *sst1-2* allele and the different *GAL:CHC1* transformants were grown on YEP-glucose for 10-12 h and 16-20 h respectively. The tester strain was diluted to 0.1 OD₆₀₀ and spread onto YEP-glucose plate. Subsequently different *GAL:CHC1* transformant cultures were spotted onto the plate which were incubated for 2-3 days before imaging. MATa and MAT α strains labelled in the Figure 4 are wild type strain controls.

References

1. Allen CL, Goulding D, Field MC (2003) Clathrin - mediated endocytosis is essential in *Trypanosoma brucei*. The EMBO journal 22: 4991-5002
2. Anantharaman V, Aravind L (2004) Novel conserved domains in proteins with predicted roles in eukaryotic cell-cycle regulation, decapping and RNA stability. BMC genomics 5: 45
3. Beckham C, Hilliker A, Cziko A-M, Noueiry A, Ramaswami M, Parker R (2008) The DEAD-box RNA helicase Ded1p affects and accumulates in *Saccharomyces cerevisiae* P-bodies. Molecular biology of the cell 19: 984-993
4. Bitsikas V, Corrêa Jr IR, Nichols BJ (2014) Clathrin-independent pathways do not contribute significantly to endocytic flux. Elife 3: e03970
5. Buchan JR, Muhlrud D, Parker R (2008) P bodies promote stress granule assembly in *Saccharomyces cerevisiae*. The Journal of cell biology 183: 441-455
6. Collier J, Parker R (2005) General translational repression by activators of mRNA decapping. Cell 122: 875-886
7. Formosa T, Nittis T (1998) Suppressors of the temperature sensitivity of DNA polymerase α mutations in *Saccharomyces cerevisiae*. Molecular and General Genetics MGG 257: 461-468
8. Ghaemmaghami S, Huh W-K, Bower K, Howson RW, Belle A, Dephoure N, O'Shea EK, Weissman JS (2003) Global analysis of protein expression in yeast. Nature 425: 737
9. Häcker S, Krebber H (2004) Differential export requirements for shuttling serine/arginine-type mRNA-binding proteins. Journal of Biological Chemistry 279: 5049-5052
10. Hackmann A, Wu H, Schneider U-M, Meyer K, Jung K, Krebber H (2014) Quality control of spliced mRNAs requires the shuttling SR proteins Gbp2 and Hrb1. Nature communications 5: 3123
11. Hilliker A, Gao Z, Jankowsky E, Parker R (2011) The DEAD-box protein Ded1 modulates translation by the formation and resolution of an eIF4F-mRNA complex. Molecular cell 43: 962-972
12. Huang KM, Gullberg L, Nelson KK, Stefan CJ, Blumer K, Lemmon SK (1997) Novel functions of clathrin light chains: clathrin heavy chain trimerization is defective in light chain-deficient yeast. Journal of cell science 110: 899-910
13. Hurt E, Luo M-j, Röther S, Reed R, Sträßer K (2004) Cotranscriptional recruitment of the serine-arginine-rich (SR)-like proteins Gbp2 and Hrb1 to nascent mRNA via the TREX complex. Proceedings of the National Academy of Sciences 101: 1858-1862
14. Iost I, Dreyfus M, Linder P (1999) Ded1p, a DEAD-box protein required for translation initiation in *Saccharomyces cerevisiae*, is an RNA helicase. Journal of Biological Chemistry 274: 17677-17683
15. Kaksonen M, Roux A (2018) Mechanisms of clathrin-mediated endocytosis. Nature Reviews Molecular Cell Biology 19: 313

16. Krogan NJ, Cagney G, Yu H, Zhong G, Guo X, Ignatchenko A, Li J, Pu S, Datta N, Tikuisis AP (2006) Global landscape of protein complexes in the yeast *Saccharomyces cerevisiae*. *Nature* 440: 637
17. Lemmon SK, Freund C, Conley K, Jones EW (1990) Genetic instability of clathrin-deficient strains of *Saccharomyces cerevisiae*. *Genetics* 124: 27-38
18. Lemmon SK, Jones EW (1987) Clathrin requirement for normal growth of yeast. *Science* 238: 504-509
19. Marnef A, Sommerville J, Lodomery MR (2009) RAP55: insights into an evolutionarily conserved protein family. *The international journal of biochemistry & cell biology* 41: 977-981
20. Mayor S, Pagano RE (2007) Pathways of clathrin-independent endocytosis. *Nature reviews Molecular cell biology* 8: 603
21. Mitchell SF, Jain S, She M, Parker R (2013) Global analysis of yeast mRNPs. *Nat Struct Mol Biol* 20: 127-133
22. Moorthy BT, Sharma A, Boettner DR, Wilson TE, Lemmon SK (2019) Identification of Suppressor of Clathrin Deficiency-1 (SCD1) and Its Connection to Clathrin-Mediated Endocytosis in *Saccharomyces cerevisiae*. *G3: Genes, Genomes, Genetics* 9: 867-877
23. Mugler CF, Hondele M, Heinrich S, Sachdev R, Vallotton P, Koek AY, Chan LY, Weis K (2016) ATPase activity of the DEAD-box protein Dhh1 controls processing body formation. *Elife* 5: e18746
24. Nelson KK, Holmer M, Lemmon SK (1996) SCD5, a suppressor of clathrin deficiency, encodes a novel protein with a late secretory function in yeast. *Molecular biology of the cell* 7: 245-260
25. Nelson KK, Lemmon S (1993) Suppressors of clathrin deficiency: overexpression of ubiquitin rescues lethal strains of clathrin-deficient *Saccharomyces cerevisiae*. *Molecular and cellular biology* 13: 521-532
26. Newpher TM, Idrissi F-Z, Geli MI, Lemmon SK (2006) Novel function of clathrin light chain in promoting endocytic vesicle formation. *Molecular biology of the cell* 17: 4343-4352
27. Ozdilek BA, Thompson VF, Ahmed NS, White CI, Batey RT, Schwartz JC (2017) Intrinsically disordered RGG/RG domains mediate degenerate specificity in RNA binding. *Nucleic acids research* 45: 7984-7996
28. Pang T-L, Wang C-Y, Hsu C-L, Chen M-Y, Lin J-J (2003) Exposure of single-stranded telomeric DNA causes G2/M cell cycle arrest in *Saccharomyces cerevisiae*. *Journal of Biological Chemistry* 278: 9318-9321
29. Payne GS, Schekman R (1985) A test of clathrin function in protein secretion and cell growth. *Science* 230: 1009-1014
30. Payne GS, Schekman R (1989) Clathrin: a role in the intracellular retention of a Golgi membrane protein. *Science* 245: 1358-1365
31. Poornima G, Mythili R, Nag P, Parbin S, Verma PK, Hussain T, Rajyaguru PI (2019) RGG-motif self-association regulates eIF4G-binding translation repressor protein Scd6. *RNA biology*
32. Poornima G, Shah S, Vignesh V, Parker R, Rajyaguru PI (2016) Arginine methylation promotes translation repression activity of eIF4G-binding protein, Scd6. *Nucleic acids research* 44: 9358-9368
33. Prosser DC, Drivas TG, Maldonado-Báez L, Wendland B (2011) Existence of a novel clathrin-independent endocytic pathway in yeast that depends on Rho1 and formin. *J Cell Biol* 195: 657-671
34. Rajyaguru P, She M, Parker R (2012) Scd6 targets eIF4G to repress translation: RGG motif proteins as a class of eIF4G-binding proteins. *Molecular cell* 45: 244-254
35. Rao BS, Parker R (2017) Numerous interactions act redundantly to assemble a tunable size of P bodies in *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences* 114: E9569-E9578

36. Seeger M, Payne GS (1992a) A role for clathrin in the sorting of vacuolar proteins in the Golgi complex of yeast. *The EMBO Journal* 11: 2811-2818
37. Seeger M, Payne GS (1992b) Selective and immediate effects of clathrin heavy chain mutations on Golgi membrane protein retention in *Saccharomyces cerevisiae*. *The Journal of Cell Biology* 118: 531-540
38. Segal SP, Dunckley T, Parker R (2006) Sbp1p affects translational repression and decapping in *Saccharomyces cerevisiae*. *Molecular and cellular biology* 26: 5120-5130
39. Squirrell JM, Eggers ZT, Luedke N, Saari B, Grimson A, Lyons GE, Anderson P, White JG (2006) CAR-1, a protein that localizes with the mRNA decapping component DCAP-1, is required for cytokinesis and ER organization in *Caenorhabditis elegans* embryos. *Molecular biology of the cell* 17: 336-344
40. Thandapani P, O'Connor TR, Bailey TL, Richard S (2013) Defining the RGG/RG motif. *Molecular cell* 50: 613-623
41. Waldherr M, Ragnini A, Jank B, Teply R, Wiesenberger G, Schweyen RJ (1993) A multitude of suppressors of group II intron-splicing defects in yeast. *Current genetics* 24: 301-306
42. Wilhelm JE, Buszczak M, Sayles S (2005) Efficient protein trafficking requires trailer hitch, a component of a ribonucleoprotein complex localized to the ER in *Drosophila*. *Developmental cell* 9: 675-685
43. Windgassen M, Krebber H (2003) Identification of Gbp2 as a novel poly (A)⁺ RNA - binding protein involved in the cytoplasmic delivery of messenger RNAs in yeast. *EMBO reports* 4: 278-283
44. Windgassen M, Sturm D, Cajigas LJ, González CI, Seedorf M, Bastians H, Krebber H (2004) Yeast shuttling SR proteins Npl3p, Gbp2p, and Hrb1p are part of the translating mRNPs, and Npl3p can function as a translational repressor. *Molecular and cellular biology* 24: 10479-10491

Figures

Figure 1

YPR129W is necessary and sufficient to rescue clathrin deficiency (A) Growth assay of *GAL1 CHC1* cells (endogenous *CHC1* is under galactose inducible promoter) transformed with YEp 24 (empty vector), YEp SCD 6 YEp Δ scd 6 (lacking YPR 129 W and YEp oSCD 6 YEp only SCD 6 expressing only YPR 129 W 10 OD 600 cells were serially diluted and spotted on glucose or galactose containing selective media plates. These plates were incubated at 30 C for 2 days (galactose media) or 4 5 days (glucose media) before imaging. (B) Quantification of the percentage rescue of *GAL1 CHC1* transformants upon clathrin deficiency indicates $p < 0.05$. Percentage rescue calculated as $(\text{number of colonies on glucose} / \text{number of colonies on galactose}) \times 100$. (C) Growth assay of *GAL1 CHC1* transformants. SCD 6 panel refers to the transformant expressing SCD 6 from pGP 564 vector with C terminal His tag. SCD 6 expressed from pGP 564 vector was spotted on the same plate as the rest of the three spottings. Image was spliced to remove other spottings not relevant to this figure. Note: Because of the difference between the selection marker between Yep and pGP 564, this assay was done on YEP plates. Plates were incubated at 30 C for 2 days (galactose media) or 4 5 days (glucose plate) before imaging.

Figure 2

PSP2 is a novel suppressor of clathrin deficiency (A) GAL 1 CHC 1 cells were transformed with pRS 426 (empty vector) or pRS 426 PSP 2 10 OD 600 cells were serially diluted and spotted on glucose or galactose containing selective media plates These plates were incubated at 30 C for 2 days (galactose media) or 4 5 days (glucose media) before imaging B) Quantification of the percentage rescue by PSP 2 upon clathrin deficiency using plating assay and indicates $p < 0.01$ and $p < 0.001$ respectively) Percentage rescue was calculated as the number of colonies on glucose plate/number of colonies on galactose plate* 100

Figure 3

DED1, SBP1, DHH1 and GBP2 fail to rescue clathrin deficiency growth defect. A) Growth assay of GAL1:CHC1 cells transformed with different plasmids as labeled. 10 OD600 cells were serially diluted and spotted on glucose or galactose containing selective media plates. These plates were incubated at 30°C for 2 days (galactose media) or 4-5 days (glucose media) before imaging. B) Quantification of plating assay to measure percentage rescue by GBP2, DED1 and DHH1 (** indicates $p < 0.01$). Percentage rescue was calculated as the number of colonies on glucose plate/number of colonies on galactose plate*100.

Figure 4

SCD 6 and PSP 2 fail to rescue the trans Golgi network (sorting function of clathrin heavy chain MAT α strain expressing endogenous CHC 1 under galactose inducible promoter was transformed with plasmids expressing SCD 6 SBP 1 DED 1 PSP 2 GBP 2 DHH 1 or CHC 1 Respective empty vectors were used as control Transformants were spotted on a lawn of MAT α cells carrying the sst 1 2 allele and incubated on YEP glucose plate until 2 3 days) the appearance of a zone of clearance (YEp 24 is the empty vector control for SCD 6 and CHC 1 pRS 426 is the empty vector control for DED 1 SBP 1 and PSP 2 pGP 564 is the empty vector control for GBP 2 and YEplac 195 is the empty vector control for DHH 1 Wild type MAT α and MAT α strains act as negative and positive controls, respectively pRS426 SCD6 GBP2 pGP564 SBP1 PSP2 DED1 YEp24 YEplac195 CHC1 MAT α MAT α DHH1

Figure 5

RGG motif and Lsm domain are important for suppress clathrin deficiency growth defect by Scd 6 (A) Growth assay of GAL 1 CHC 1 cells transformed with plasmids expressing wild type, Lsm FDF, RGG or RGG' domain/motif deletion mutants of Scd 6 10 OD 600 cells were serially diluted and spotted on glucose or galactose containing selective media plates These plates were incubated at 30 C for 2 days (galactose media) or 4 5 days (glucose media) before imaging B) Quantification of the percentage rescue of GAL 1 CHC 1 by SCD 6 mutants upon clathrin depletion ..(Growth assay to test the ability of a rginine methylation defective (mutant of SCD 6 to suppressing clathrin deficiency ..(Quantification of the percentage rescue of GAL 1 CHC 1 transformants by the AMD mutant of SCD 6 upon clathrin deficiency

and indicates $p < 0.05$ and $p < 0.01$ respectively) Percentage rescue was calculated as the number of colonies on glucose plate/number of colonies on galactose plate* 100

Figure 6

Deletion of RGG-motif compromises the ability of Psp2 to suppress clathrin deficiency growth defect. (A) Growth assay of GAL1:CHC1 cells transformed with wild type and RGG-motif deletion mutants of Psp2. 10 OD600 cells were serially diluted and spotted on glucose or galactose containing selective media plates. These plates were incubated at 30°C for 2 days (galactose media) or 4-5 days (glucose media) before imaging. (B) Quantification of the percentage rescue of GAL1:CHC1 transformants with wild type or RGG-motif deletion mutant of PSP2 upon clathrin deficiency (* indicates $p < 0.05$). Percentage rescue was calculated as the number of colonies on glucose plate/number of colonies on galactose plate*100.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupFigure1.jpg](#)