Combined use of Specific Length Amplified Fragment Sequencing (SLAF-seq) And Bulk Segregant Analysis (BSA) For Rapid Identification of Genes Influencing Fiber Content of Hemp (*Cannabis sativa* L.)

Yue Zhao

Daqing Branch of Heilongjiang Academy of Sciences

Yufeng Sun

Daqing Branch of Heilongjiang Academy of Sciences

Kun Cao

Daqing Branch of Heilongjiang Academy of Sciences

Xiaoyan Zhang

Daqing Branch of Heilongjiang Academy of Sciences

Jing Bian

Daqing Branch of Heilongjiang Academy of Sciences

Chengwei Han

Daqing Branch of Heilongjiang Academy of Sciences

Ying Jiang

Daqing Branch of Heilongjiang Academy of Sciences

Lei Xu

Daqing Branch of Heilongjiang Academy of Sciences

Xiaonan Wang (✉ wxn_fern@163.com)

Daqing Branch of Heilongjiang Academy of Sciences

Research Article

Keywords: Hemp fiber content, Specific length amplified fragment sequencing (SLAF-seq), Bulk Segregant analysis (BSA), Candidate gene, SNP analysis

Posted Date: August 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-820517/v1>

Abstract

Hemp (*Cannabis sativa* L.), an ancient crop, is a significant source of high-quality fiber that primarily caters to the textile industry worldwide. Compared to other fiber crops, hemp is an environmentally friendly fiber crop which requires fewer pesticides and purifies the soil during cultivation. Fiber content is a crucial quantitative trait for evaluating fiber yield in hemp. Understanding the genetic mechanisms involved in hemp breeding is essential for improving yield. In this study, to investigate these underlying mechanisms, we developed 660 F1 plants from a cross between Jindao-15 (high fiber content fiber-use variety) and Fire No.1 (low fiber content fiber-use variety), and thirty plants each with high and low fiber content were selected from 305 monoecious plants of this population according to 5 %-10 % of population size for quantitative traits. The DNA from these plants was extracted to establish two bulk DNA pools. The DNA was then subjected to the restriction digestion by the enzymes RsaI and HaeIII to obtain 314–364 bp digestion fragments and subjected to sequencing using specific length amplified fragment sequencing (SLAF-seq). Then, each fragment was labeled with specific SLAF tags, and single nucleotide polymorphisms (SNPs) were identified to determine the candidate regions related to fiber content through marker association analysis of genotype frequency differences between the mixed pools of DNA. The gene annotation results identified the candidate genes responsible for the fiber content of hemp. In this study, we successfully developed 368,404 SLAF tags, which led to the detection of 389,687 SNPs. These SNPs were then subjected to the SNP-Index correlation algorithm, which revealed four candidate regions related to fiber content traits on Chromosome 1, with a length of 8.68 Mb and containing 389 annotated genes. We found that 199 non-synonymous mutations which change the amino acids sequences existed in the parent population. The annotation information and the comparison results identified 15 genes that were highly likely to modulate the fiber content of hemp. Further, qPCR validation identified six genes (LOC115705530, LOC115705875, LOC115704794, LOC115705371, LOC115705688 and LOC115707511) that were highly positively correlated with influencing the hemp fiber content. These genes were involved in the transcription regulation, auxin and water transportation, one carbon and sugar metabolism. Thus, our study highlights the importance of the combined use of SLAF-Seq and Bulk Segregant analysis (BSA) to locate genes related to hemp fiber content rapidly. Hence, our study provides novel mechanistic inputs for the fast identification of genes related to important agronomic traits of hemp and other crops catering to the textile industry.

Introduction

Cannabis sativa L. is an annual herb that can be grouped as hemp or marijuana based on its tetrahydrocannabinol (THC) content, of which hemp contains less than 0.3 % and marijuana contains more than 0.3 % [1]. Active research on hemp mainly focuses on determining the plant's fiber, grain, and cannabidiol (CBD) content. It is known that plant fibers have great potential to be used in various innovative applications as biodegradable, ecological, and renewable resources with unique properties [2, 3], with hemp bast fiber being one of the best. The unique structure of hemp fiber makes it superior to cotton fiber due to its comprehensive moisture absorption, dispersion rate, and heat-resisting

temperature. Other superior qualities of the hemp fiber include its antibacterial effects, the ability to prevent damage due to exposure to radiation and ultraviolet rays, and the prevention of sound absorption and static. Due to its capability to cater to military and civil needs, the textile market has an ever-increasing demand for hemp bast fiber. However, the price of hemp textiles is relatively high due to the production process and other factors. It is vital to increase the fiber output to meet the production demand and reduce the production cost.

It is reported that the probable measure to increase fiber yield through improving cultivation conditions [4, 5], but it has limited effects. Since a sole improvement of the hemp germplasm can fundamentally increase the yield, in this study, we investigate the crucial involvement of genetic factors in hemp production, apart from the roles played by the natural environment and the cultivation conditions. At present, hemp fiber improvement mainly relies on the traditional time-consuming method of phenotypic selection. Here, we highlight how delineating the molecular and genetic basis for hemp breeding can have an improved efficiency. However, it is imperative to identify the genes affecting hemp fiber yield to proceed with our proposed line of study.

The current methods of gene mining mainly include reverse genetics that differ phenotype from genotype and positive genetics that identifies the reverse [6]. The development of sequencing techniques and bioinformatics has provided many valuable strategies for discovering new genes related to crop yields [7]. SLAF-seq combined with BSA is one of the positive genetics methods. Specific length amplified fragment sequencing (SLAF-seq) is a technique that uses high-depth sequencing to identify specific unique fragments in the genome and thereby help to locate molecular markers affecting the traits of interest. This method compares the differences in the SNP markers' occurrence frequency between two mixed pools with different genotypes [8]. The advantages of such techniques include a high rate of identification of molecular marker density, low cost, high efficiency, and high accuracy [9]. The SLAF-seq technology has been implicated in multiple studies that include gene mapping of wheat, cucumber, pepper, melon, and other crops [10–13]. However, the use of SLAF-seq in determining hemp fiber yield remains largely unexplored. Fiber content has always been an important index to evaluate the hemp fiber yield, which is governed by multiple genetic components. It is known that the genetic component regulating hemp bast fiber content is significant, the gene-environment (G-E) interaction is low, and the generalized heritability is high [14]. The high heterozygosity and variability of hemp cause the separation of hemp traits in the F1 generation. Lavery et al. have shown that the physical and genetic map of hemp can be constructed using the F1 generation population [15]. Such studies indicate the feasibility and importance of identifying genes regulating fiber yield by studying the quantitative traits of the F1 generation through SLAF and BSA technology.

This study used the isolated population constructed by crossing Jindao-15 and Fire No.1, which have a significant difference in their fiber content, to perform our investigations. We also coupled bioinformatic analysis to rapidly identify the genes regulating hemp fiber content with the SLAF-seq and BSA technologies. It greatly saved cost and improved gene mining efficiency. Our investigation provides a novel theoretical basis for improved breeding of hemp using molecular markers and genetic studies.

Materials And Methods

Plant materials used and construction of two distinct pools of plants

The selection of plant varieties for this study depended on the total fiber rate (calculated after retting) which can indicate the fiber content. Jindao-15 (Fig. 1A) was produced in the Ukrainian Academy of Agricultural Sciences in 2014, and its total fiber rate is 34.4 %. Fire No.1 (Fig. 1B, C) was bred by Daqing Branch of Heilongjiang Academy of Sciences in 2015, with a total fiber rate of 22.9 %. These cultivars were provided by the Daqing Branch of Heilongjiang Academy of Sciences. Jindao-15 and Fire No.1 were crossed as the male and female parents respectively to achieve the F1 generation group. The monoecious hemp cultivar Jindao-15 and the dioecious Fire No.1 were grown in the greenhouse of Chunlei farm (Daqing, Heilongjiang, China) in the winter, 2018. The male plants of Fire No.1 were uprooted when the buds appeared. A total of 660 seeds were obtained by crossing the two cultivars in the spring of 2019. The seeds and their parents were then grown in an experimental base of Dongfeng farm under natural conditions. The leaves of 305 monoecious plants of the F1 population were collected and immediately immersed into liquid nitrogen during budding. The leaves were stored at -80°C. The 305 plants were harvested during the initiation of flowering. The weights of fresh plant stems and bast fibers (Fig. 1D) peeled from the stem were recorded. Phenotypic data were statistically analyzed to calculate the fiber content (Fiber content = bast fiber weight / stem weight × 100 %) using Microsoft Excel (Microsoft Office, Microsoft, 2003). Thirty of each high and low fiber-containing plants were selected for the extreme bulk construction. Total genomic DNA was isolated from tender leaves of both parental lines and every plant of the two segregating bulks using a Plant Genomic DNA Kit (Tiangen Biotech, Beijing, China) following the manufacturer's protocol. DNA concentration and quality were measured by 1 % agarose gel electrophoresis. Thus, considering equal amounts of DNA from thirty high fiber-containing plants and thirty low fiber-containing plants, two separate and distinct gene pools were made.

Construction of SLAF-seq library and high-throughput sequencing

The entire genome of hemp (*Cannabis sativa* L.) (https://www.ncbi.nlm.nih.gov/genome/11681?genome_assembly_id=455372, version 1) was chosen as the reference genome for this study. The SLAF-predict software was used to predict the digestion pattern of the hemp reference genome using restriction endonucleases. Based on this analysis, *Rsa*I and *Hae*III were chosen to get the SLAF tag fragments of length 314 ~ 364 bp for SLAF library construction. Single-nucleotide A overhangs and dual-index sequencing adaptors were ligated to these fragments obtained by digestion of the hemp genomic DNA [16]. The modified fragments were then amplified by PCR, purified, pooled, and screened to construct the SLAF library. A detailed description of SLAF library construction and screening processes is given by Sun et al. [17]. Finally, the SLAF library was sequenced at an Illumina HiSeq 2500 platform (Illumina, Inc., San Diego, CA, USA) at Biomarker Technologies Corporation in Beijing. The *Oryza sativa*

genome(<http://www.ncbi.nlm.nih.gov/genome/?term=Arabidopsis%20thaliana>, version 7.0) was used as a positive control to evaluate the quality of the experiment and the enzyme digestion reaction. The control sequencing reads were compared with the reference genome by Burrows-Wheeler Alignment tool (BWA) software(<https://nchc.dl.sourceforge.net/project/bio-bwa/bwa-0.7.15.tar.bz2>)[18].

Identification of high-quality SNPs

To ensure high quality of data, low-quality raw reads and reads with adapter were filtered to retain only the clean reads. The BWA software was used to compare the clean reads with the reference genome to locate their positions. The samples' sequencing depth, genome coverage, and other crucial information were recorded, followed by detection of variations.

According to the localization of the clean reads in the reference genome, the Mark Duplicate tool of Picard (<http://sourceforge.net/projects/picard/>) was used to remove duplicates to negate the impact of PCR duplication. Base recalibration and variant calling were performed using Genome Analysis Toolkit (GATK) software (<https://gatk.broadinstitute.org/hc/en-us>) [19]. The SNPs were filtered strictly to get the final set of SNPs. First, SNPs with multiple genotypes were filtered out, then SNPs with read support less than four were filtered out, and then SNPs with consistent genotypes between mixing pools and SNPs with recessive mixing pool genes that did not come from recessive parents were filtered out. All the variable sites between the test and the reference genomes were identified based on a comparison between the two. All the variations in SNP annotation and effectiveness prediction were identified using the SnpEff software (<https://sourceforge.net/projects/snpeff/>).

Association analysis

SNP-index is a marker correlation analysis method that relies on the differences in the genotype frequency between the mixing pools [20, 21]. This analysis mainly looks for significant differences in genotype frequency between the mixed pools with Δ (SNP-index) statistics. The closer a marker is associated with a trait, the closer the Δ (SNP-index) value reaches. Table 1 shows the calculation formulae.

The process of elimination of false-positive sites mainly involves the identification of different markers located in the genome. The SNPNUM method(a script written by Biomarker Technologies Corporation)was adopted to plot the Δ (SNP-index) values. The target locus and its adjacent linkage loci should approach the threshold which normally is 99 % when a high peak should occur near the significantly associated region. The result was calculated by using a computer simulation experiment [22].

Gene annotation

The coding genes in the candidate area of confidence interval were accurately annotated using Non-redundant(NR)[23], Swiss-prot [24], Gene Ontology (GO) [25], Kyoto Encyclopedia of Genes and Genomes (KEGG) [26], and Cluster of Orthologous Groups of proteins (COG) [27] by BLAST software [28]. Such detailed annotation helps in the rapid screening of candidate genes.

2.6 Validation of Gene Expression by Real-Time quantitative PCR (qPCR)

The expression levels of candidate genes were validated by the Real-time quantitative PCR (qPCR) method. Total RNA was extracted from stalks of 3 varieties of hemp with different total fiber rates (Fire No.1 (22.9 %), Han Ma No.10 (27.1 %), and Jindao-15 (34.4 %)) at the seedling stage and the technical maturity stage using RNA Extraction Kit (Tiangen, China). The RNA was dissolved in Ultra Pure™ DNase/RNase-free distilled water (Invitrogen, USA). The total RNA was reverse-transcribed using FastKing RT Kit(KR116)reagent. Primer sequences were designed using Primer 7.0 and screened using SeqHunter 1.0 (Supplemental Table 1). The qPCRs were performed using Power qPCR PreMix (Genecopoeia, USA) according to the manufacturer's instructions. SYBR Green PCR cycling was performed in an IQTM5 Multicolor Real-Time PCR Detection System (Bio-Rad, USA) using 20 µl reaction volumes. The reaction conditions are as follows: 95 °C for 10 min, followed by 40 cycles of 95 °C for 10 s, 60 °C for 40 s, and 60 °C for 15 s. The relative quantitation of gene expression was calculated and normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH), used as a housekeeping gene. Three biological replicates from each condition were used for the qPCRs.

Results

Construction of SLAF library using restriction digestion of the reference hemp genome

Initially, the high total fiber-containing hemp variety Jindao-15 as the male parent was crossed with the low total fiber-containing Fire No.1 as the female parent to obtain the F1 generation group, in which the highest fresh fiber content was 45.56 %, and the lowest fiber content was 10.72 % (Supplemental Table 2). From the F1 population, thirty plants with high fiber content (35.78 %-45.56 %) and thirty with low fiber content (10.72 %-26.55 %) were selected for further studies. An equal amount of genomic DNA was taken from each of the two groups and mixed to prepare a pool. It is already known that the size of the hemp reference genome is 876.148 Mb and the GC content is 34 %. This led to the possibility of the formation of 105823 SLAF tags from the hemp reference genome distributed uniformly on the chromosomes when subjected to restriction digestion (Fig. 2). Thus, the SLAF tag library was prepared by subjecting the DNA to RsaI and HaeIII-mediated restriction digestion for further analysis.

Analysis of sequence data

Our analysis led to the generation of 36.45 Gb clean data developed from the parents through resequencing, with the average sequencing depth of the individual populations being 10.40X. To further characterize the genetic elements affecting hemp fiber production, we developed 368,404 SLAF tags from the DNA pool created by mixing the two populations using the simplified SLAF genome sequencing method (Supplemental Table 3). An average Q30(Q30 indicates a quality score of 30, indicating a 0.1 %

chance of error and thus 99.9 % confidence) ratio of 93.41 % and 37.94 % GC content (Table 2) revealed the number of reads and the quality of the data, respectively, from the sequencing analysis. The numbers of SLAF tags in the high and low fiber-containing pools were 225,318 and 320,016, respectively. Moreover, the sequencing depth of the former pool was 41.66X, while that of the latter was 48.19X. BWA software analysis of the sequencing reads showed a 90.88 % efficiency (within the normal range) of the double end analysis compared to the reference rice genome. We also observed that the actual lengths of the SLAF tags calculated according to the pair-end mapped reads of contrast locations in the genome correspond to the expected fragment size. The statistical analysis of the SLAF tags on each chromosome according to their positioning is shown in Table 3.

Identification of SNPs based on SLAF sequencing

Next, the sequencing reads obtained from each of the mixed pools were clustered for analysis. The clustered reads originating from the same SLAF fragment were defined as a particular SLAF tag. Each label, which showed differences in the sequence and SNPs between the different samples, was a polymorphic SLAF label. Our study identified 4, 295,852 SNPs between the two parent populations, 389,687 SNPs from the bulked pools, and 102,964 polymorphic SLAF labels between the mixed pools (Supplemental Table 3). The results of the statistical analysis for SNP filtering are shown in Table 4. The comprehensive analysis finally identified 25,133 high-quality SNP sites in the hemp genome.

SNP-index association analysis

Next, the sites showing a difference between the mixed pools were used for the SNP-index analysis. A computer simulation experiment revealed no correlation with relevant candidate regions when the confidence was 0.90. Ideally, it is expected that the target site and its adjacent linkage sites should be close to the threshold, resulting in a high peak near the significant correlation region. However, due to the lack of regions exceeding the theoretical threshold, no significant position-related results were observed. To fully utilize the data, we lowered the theoretical threshold to locate possible areas of interest. We used the 99 percentile for fitting the Δ (SNP-index) (Fig. 3), namely 0.10, which revealed a total length of 8.68 Mb of four candidate regions, including 397 genes. The information regarding the correlation region and the statistical analysis is shown in Table 5.

Association of regional Gene annotation

Our study found that 389 genes encoded in the candidate regions identified from the previous analysis were annotated in multiple databases (Supplemental Table 4). Out of the total 389 annotated genes from the candidate regions, 199 harbored non-synonymous mutations between the parent populations, suggesting that a minimal number of these might play a functional role in regulating hemp fiber content. However, the involvement of these genes in affecting the hemp fiber content remains uninvestigated to date. A comparative homological analysis between the hemp genes and those of *Arabidopsis*, flax, and cotton genes and a thorough literature review helped us narrow down the scope of candidate genes.

Finally, we chose 15 interesting candidate genes that might be involved in hemp fiber content regulation (LOC115705530, LOC115706200, LOC115707511, LOC115706733, LOC115704794, LOC115707202, LOC115707643, LOC115705371, LOC115705688, LOC115705010, LOC115705568, LOC115705891, LOC115706691, LOC115708167 and LOC115705875) for further analysis.

Validation of Gene Expression by Real-Time quantitative PCR (qPCR)

The stalks of Fire No. 1, Han Ma No. 10, and Jindao-15 were named H1J1, H10J1, and JD15J1, respectively, at the seedling period, while at maturity period, they were named H1J2, H10J2, and JD15J2, respectively. The gene expression analysis revealed an increase in the expression levels of LOC115705530, LOC115707511, LOC115704794, LOC115705371, and LOC115708688 with an increase in the total fiber content. In contrast, the expression levels of LOC115705875 decreased with the increase in total fiber content. The expression level of LOC115705875 in the seedling period was higher than that in the technical maturity period. The expression level of the five other genes in the seedling period did not increase to the extent of expression level in the technical maturity period (Fig. 4). This analysis identified a differential expression pattern of the different candidate genes during the seedling and the mature phase, suggesting a growth phase-dependent involvement of multiple genes in regulating hemp fiber content.

Discussion

Fiber yield from multiple plants is a crucial parameter that affects the commercial production of textiles and has been studied in multiple crops [29, 30]. It is challenging to identify the key genetic drivers regulating this complex yet essential trait. The genes regulating the fiber content of hemp remain unexplored to date. Hence, in this study, we leveraged the potential of next-generation sequencing (NGS), a highly efficient, cost-effective, and accurate method compared with conventional methods, to develop new genetic markers modulating essential traits such as fiber content of crops [31–33]. Recently, SLAF-seq has emerged as a technique with exceptionally high resolution and efficiency for identifying SNP markers in specific populations. Molecular markers can be directly developed from paired-end analysis of the sequence-specific restriction fragment lengths. Our study has identified 389,687 SNP locus that can be used to develop SNP molecular markers. KEGG analysis of the identified genes from our study showed the enrichment of candidate regions closely associated with of amino sugar and nucleotide sugar metabolism, glycosphingolipid biosynthesis of both globo and ganglio series, one carbon pool by folate, basal transcription factors, lysine biosynthesis, photosynthesis, glycosaminoglycan degradation, and starch and sucrose metabolism pathways (Fig. 5). Moreover, quantitative analysis showed six genes (LOC115705530, LOC115705875, LOC115704794, LOC115705371, LOC115705688 and LOC115707511) that revealed their essential role in hemp fiber content regulation.

One of the candidate genes identified by our analysis, LOC115705530, is located in region 1. A comparative analysis revealed that this gene has the highest similarity with the gene At3G11280, which encodes the superfamily protein of replica-like homologous domain in *Arabidopsis thaliana*, and encodes

for a protein that has a typical MYB domain. Studies involving the novel regulators of vascular development in *Arabidopsis thaliana* show that MYB has 40 interacting molecules [34]. Also, it has been reported that the gene FSM1, encoding an atypical MYB-like domain short protein found in tomato, negatively regulates the expansion of cells in the vascular bundle of fruit pericarp [35]. These studies collectively indicated an essential role of the MYB domain in regulating vascular development in multiple plants. MYB is also known to regulate lignin biosynthesis by recognizing AC elements in promoters of many lignin monomer biosynthesis genes [36] and is highly implicated in the regulation of secondary cell wall biosynthesis [37–39]. MYB, a transcription factor, is also closely associated with lignification in jute and ramie [40, 41], while MYB46-1, another MYB family transcription factor, regulates secondary cell wall and lignin biosynthesis in hemp [42]. However, further investigation is required to conclude the actual role of the gene identified in our study concerning hemp.

Another gene identified in our study was LOC115705875, located in region 2, which had only one candidate gene. The protein encoded by this gene had the highest homology with AUXIN1 (AUX1). AUX1 had been reported to encode a high-affinity auxin influx vector. In *Arabidopsis thaliana*, AUX1 belongs to the AUX/LAX multigene family, consisting of four highly conserved genes AUX1 and Like AUX1 (LAX) genes LAX1, LAX2, and LAX3. All four AUX/LAX family members are known to have the auxin uptake function [43]. Auxin is an essential hormone for plant growth and development. Auxin flow carrier AUX1/LAX transports auxin into cells and promotes xylem differentiation in stem and root tissues by increasing cytoplasmic auxin signaling to regulate vascular patterns and differentiation [44].

Our RT-qPCR analysis showed that LOC115705875, the expression pattern of which shows a negative regulation during the seedling and the technical maturity period, differed in terms of function, with respect to the other genes mentioned. Low fiber-containing hemp varieties required more auxin, which could be attributed to the fact that a lower fiber content requires more auxin for xylem differentiation. The expression of this gene at the technical maturity stage was significantly lower than that at the seedling period, which might be because of the increased requirement of auxin by hemp during early development. Guerriero et al. found that the expression of genes related to auxin metabolism was higher in the older stem nodes at the 6-week seedling stage [45]. Studies on the secondary growth stage of hemp showed a high biological activity of auxin during the deposition and remodeling of the primary cell walls [46]. In other words, there is an increased demand for auxin in the early development of secondary phloem fibers of hemp. These reports collectively validate our findings based on the expression pattern of this gene and suggest a probable mechanism by which auxin might regulate the fiber content of hemp.

The next candidate gene, LOC115704794, was located in region 3 and had the highest homology with AT4G00430, encoding aquaporin protein AtPIP1;4. Aquaporins(AQPs) are transmembrane channel proteins that regulate the intracellular and intercellular diffusion of water and other uncharged solutes such as glycerin, hydrogen peroxide, ammonia, small organic acids, urea, and metallic substances. AQPs are essential for maintaining water composition, osmotic regulation, signal transduction, detoxification processes, and the acquisition and transport of nutrients in various organisms [47, 48]. Plasma membrane intrinsic proteins (PIP) are one of five AQP subfamilies that have attracted particular attention

for their potential to improve water retention and photosynthesis in plants [49]. PIPs can be divided into the subtypes PIP1 and PIP2 [50], which have an 80 % amino acid sequence homology. The main differences between the two groups lie at the N- and C-terminal ends, the ring A length, and the amino acid composition [51]. The PIP1 subfamily was initially thought to be nonfunctional due to its failure to localize to the plasma membrane [52]. However, later experiments have shown that PIP1 does have a functional role inside the cell. For example, it was found that the stem parenchymal cells' response to drought stress was significantly upregulated by the PIP1 subfamily of water channels, rather than the PIP2 subfamily [53].

Moreover, it is known that AtPIP1;2 in *Arabidopsis thaliana* promotes the water conductivity of roots and rosette leaves [54], while AtPIP1;4 mediates the transport of CO₂, an essential regulator of photosynthesis [55]. However, a recent study has reported negligible changes in photosynthetic efficiency and mesophyll conductivity of *Arabidopsis* aquaporin knockout mutants (PIP1;2, PIP1;3, PIP2;6), compared to the control group [56]. A study based on cotton plants showed that GhPIP1-2, which belongs to the PIP1 family, is mainly expressed during the fiber extension period of cotton. The gene expression was recorded to be the highest at five days post-flowering, suggesting a vital role in supporting the rapid water flow into the vacuoles during cell elongation in cotton [57]. The transcriptional abundance of the PIP gene family in *Calotropis procera* fiber cells is greater than that of cotton. Studies on long thorns, which are adapted to survive in harsh environmental conditions such as drought and salty and alkaline conditions, also verified the role of PIP aquaporin in the elongation of fiber cells. However, the study suggests a more critical role of PIP2 than that of PIP1 [58]. Therefore, we conclude that LOC115704794 may affect the fiber yield by regulating the CO₂ transport for photosynthesis. We also suggest a similar role of LOC115704794 to that of GhPIP1-2, whereby it adjusts the length and width of fibers to affect the fiber content. However, further experiments are needed to determine the exact mechanism by which LOC115704794 functions in regulating the hemp fiber content.

The following candidate gene, LOC115705371, also located in region 3, had the highest similarity to the *Arabidopsis* gene AT2G28760, which encodes the protein UXS6. There are six UXS genes in the *Arabidopsis* genome, among which UXS3, UXS5, and UXS6 have the highest expression in the stem, mainly in the xylem cells and the interfascicular fibers. The proteins encoded by these genes, which regulate secondary wall formation, are directly regulated by the secondary wall NAC transcription factors. The simultaneous down-regulation/mutation of UXS3, UXS5, and UXS6 results in a significant decrease in the primary wall xyloglucan content, the thickening of the secondary wall, the content of xylan, and severe deformation of xylem vessels. Xylan and xyloglucan are the two main hemicelluloses in plant cell walls. Xylan is the main hemicellulose in the primary wall of dicotyledonous plants. Their biosynthesis requires a stable supply of sugar donors like UDP-xylose, which is synthesized by converting UDP-glucuronic acid via the activity of UDP-xylose synthase. UXS3, UXS5, and UXS6 play a significant role in the supply of UDP-xylose for the biosynthesis of xylan and xyloglucan [59]. Hence, we conclude that LOC115705371 may play a possible role in affecting the fiber content by regulating the hemicellulose content in hemp phloem fibers.

The following candidate gene, LOC115705688, located in region 3, showed the highest similarity with AT2G35040. This gene encodes the protein phosphoribosylaminimidazole formamide formyltransferase, which belongs to the AICARFT/IMPCHase two-enzyme family proteins [60]. GO analysis showed that this gene was involved in nucleotide transport and metabolism, while KEGG pathway analysis showed the involvement of this gene in one-carbon (C1) metabolism. C1 metabolism is closely related to lignin biosynthesis [61–63]. Hemp lignification is one of the differentiation processes of bast fiber and core fiber. A comparative gene expression analysis of hemp bast fiber and core fiber showed that most of the coding proteins were involved in regulating C1 metabolism and lignin biosynthesis [64]. We conclude that LOC115705688 may participate in the lignification process to regulate hemp fiber content.

The final candidate gene, LOC115707511, located in region 4, had the highest similarity with WRKY transcription factor WRKY70. WRKY transcription factors belong to one of the largest transcription factor families discovered to date and participate in regulating development, signal transduction, and stress defense processes in various plants. WRKY executes transcriptional activation or inhibition in either a homodimeric or a heterodimeric form [65]. WRKY70 has also been reported to regulate jasmonic acid and salicylic acid signaling [66]. A study of cotton plants showed that the laccase gene GhLac1 regulates fiber initiation and elongation by coordinating jasmonic acid and flavonoid metabolism [67]. WRKY transcription factors are also found to be up-regulated in jute during early fiber development [68]. The jasmonic acid biosynthesis-related gene expression in adult stem nodes was higher in the phloem [69]. Therefore, we conclude collectively from all the studies that WRKY70 may affect fiber development by regulating the jasmonic acid pathway.

Our study highlights our novel identification of genes involved in regulating the fiber content trait of hemp by using the integrated SLAF-seq and BSA methods. The findings of our study have the potential to lay a good foundation for determining regulators of hemp breeding via molecular marker-assisted selection. However, further in-depth analysis and functional characterization of these candidate genes by transformation or assessment of mutation are required to delineate their roles in regulating hemp fiber content conclusively.

Conclusions

In our study, the F1 population was constructed by crossbreeding hemp parents with a significant difference in total fiber content. The mixing pool was established according to the fiber content, and SLAF database construction and sequencing were performed. A total of 368,404 SLAF tags were developed, and 389,687 SNP sites were detected. According to the SNP-Index correlation algorithm, four candidate regions related to fiber content traits were obtained on chromosome 1, with a length of 8.68 Mb and 389 annotated genes, among which 199 genes were non-synonymous mutants between the parental populations. According to the annotation and comparison results, the 15 genes (LOC115705530, LOC115706200, LOC115707511, LOC115706733, LOC115704794, LOC115707202, LOC115707643, LOC115705371, LOC115705688, LOC115705010, LOC115705568, LOC115705891, LOC115706691, LOC115708167 and LOC115705875) were highly significantly positively correlated candidate genes for

hemp fiber content. The quantitative analysis validated that the genes LOC115705530, LOC115705875, LOC115704794, LOC115705371, LOC115705688 and LOC115707511 were indeed positively correlated with fiber content. These studies indicated that SLAF-Seq and BSA methods could be used to locate genes related to important agronomic traits in hemp rapidly.

Declarations

DATA AVAILABILITY STATEMENT

The sequencing data associated with this study have been deposited in NCBI with accession number PRJNA749899. ([http://www.ncbi.nlm.nih.gov/sra/ PRJNA 749899](http://www.ncbi.nlm.nih.gov/sra/PRJNA749899))

AUTHOR CONTRIBUTIONS

YZ and YS designed the experiment; XZ, CH and YJ constructed the F1 population and collected the samples; KC, JB and LX contributed to data processing and analysis; YZ wrote this manuscript; XW checked and revised the manuscript; all authors reviewed and approved this submission.

ACKNOWLEDGMENT

The authors gratefully acknowledge funding from the Youth Innovation Program of the Heilongjiang Academy of Science (CXMS2019DQ01), President Foundation Program of the Heilongjiang Academy of Sciences (YZ2020DQ01).

References

1. Avico U, Pacifici R, Zuccaro P: **Variations of tetrahydrocannabinol content in cannabis plants to distinguish the fibre-type from drug-type plants.** *Bulletin on narcotics* 1985, **37**(4):61–65.
2. Sanjay MR, Arpitha GR, Yogesha B: **Study on Mechanical Properties of Natural - Glass Fibre Reinforced Polymer Hybrid Composites: A Review.** *Materials Today: Proceedings* 2015, **2**(4):2959–2967.
3. Pickering KL, Efendy MGA, Le TM: **A review of recent developments in natural fibre composites and their mechanical performance.** *Composites Part A: Applied Science and Manufacturing* 2016, **83**:98–112.
4. Struik PC, Amaducci S, Bullard MJ, Stutterheim NC, Venturi G, Cromack HTH: **Agronomy of fibre hemp (*Cannabis sativa* L.) in Europe.** *IND CROP PROD* 2000, **11**(2):107–118.
5. Amaducci S, Zatta A, Pelatti F, Venturi G: **Influence of agronomic factors on yield and quality of hemp (*Cannabis sativa* L.) fibre and implication for an innovative production system.** *FIELD CROP RES* 2008, **107**(2):161–169.
6. Takahashi J, Pinto L, Vitaterna M: **Forward and reverse genetic approaches to behavior in the mouse.** *SCIENCE* 1994, **264**(5166):1724–1733.

7. Varshney RK, Graner A, Sorrells ME: **Genomics-assisted breeding for crop improvement.** *TRENDS PLANT SCI* 2005, **10**(12):621–630.
8. Sun X, Liu D, Zhang X, Li W, Liu H, Hong W, Jiang C, Guan N, Ma C, Zeng H *et al.*: **SLAF-seq: an efficient method of large-scale de novo SNP discovery and genotyping using high-throughput sequencing.** *PLOS ONE* 2013, **8**(3):e58700.
9. Teitelbaum J: **A Hemp oil, CBD, and Marijuana Primer: Powerful Pain, Insomnia, and Anxiety-relieving Tools!.** *Altern Ther Health Med* 2019, **25**(S2):21–23.
10. Trick M, Adamski NM, Mugford SG, Jiang CC, Febrer M, Uauy C: **Combining SNP discovery from next-generation sequencing data with bulked segregant analysis (BSA) to fine-map genes in polyploid wheat.** *BMC PLANT BIOL* 2012, **12**:14.
11. Xu X, Lu L, Zhu B, Xu Q, Qi X, Chen X: **QTL mapping of cucumber fruit flesh thickness by SLAF-sEq.** *Sci Rep* 2015, **5**:15829.
12. Xu X, Chao J, Cheng X, Wang R, Sun B, Wang H, Luo S, Xu X, Wu T, Li Y: **Mapping of a Novel Race Specific Resistance Gene to Phytophthora Root Rot of Pepper (Capsicum annuum) Using Bulk Segregant Analysis Combined with Specific Length Amplified Fragment Sequencing Strategy.** *PLOS ONE* 2016, **11**(3):e151401.
13. Zhang H, Yi H, Wu M, Zhang Y, Zhang X, Li M, Wang G: **Mapping the Flavor Contributing Traits on "Fengwei Melon" (Cucumis melo L.) Chromosomes Using Parent Resequencing and Super Bulk Segregant Analysis.** *PLOS ONE* 2016, **11**(2):e148150.
14. Petit J, Salentijn E, Paulo MJ, Thouminot C, van Dinter BJ, Magagnini G, Gusovius HJ, Tang K, Amaducci S, Wang S *et al.*: **Genetic Variability of Morphological, Flowering, and Biomass Quality Traits in Hemp (Cannabis sativa L.).** *FRONT PLANT SCI* 2020, **11**:102.
15. Lavery KU, Stout JM, Sullivan MJ, Shah H, Gill N, Holbrook L, Deikus G, Sebra R, Hughes TR, Page JE *et al.*: **A physical and genetic map of Cannabis sativa identifies extensive rearrangements at the THC/CBD acid synthase loci.** *GENOME RES* 2019, **29**(1):146–156.
16. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD: **Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform.** *Appl Environ Microbiol* 2013, **79**(17):5112–5120.
17. Sun X, Liu D, Zhang X, Li W, Liu H, Hong W, Jiang C, Guan N, Ma C, Zeng H *et al.*: **SLAF-seq: an efficient method of large-scale de novo SNP discovery and genotyping using high-throughput sequencing.** *PLOS ONE* 2013, **8**(3):e58700.
18. Cingolani, P. E., Platts, A. E., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X.J., Ruden, D.M.: **A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3.** *Fly* 2012, **6**(2): 80–92.
19. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M *et al.*: **The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.** *GENOME RES* 2010, **20**(9):1297–1303.

20. Hill JT, Demarest BL, Bisgrove BW, Gorski B, Su YC, Yost HJ: **MMAPPR: mutation mapping analysis pipeline for pooled RNA-seq.** *GENOME RES* 2013, **23**(4):687–697.
21. Fekih R, Takagi H, Tamiru M, Abe A, Natsume S, Yaegashi H, Sharma S, Sharma S, Kanzaki H, Matsumura H *et al.* **MutMap+: genetic mapping and mutant identification without crossing in rice.** *PLOS ONE* 2013, **8**(7):e68529.
22. Takagi H, Abe A, Yoshida K, Kosugi S, Natsume S, Mitsuoka C, Uemura A, Utsushi H, Tamiru M, Takuno S *et al.* **QTL-seq: rapid mapping of quantitative trait loci in rice by whole genome resequencing of DNA from two bulked populations.** *PLANT J* 2013, **74**(1):174–183.
23. Deng,Y.Y.,Li, J.Q., Wu, S.F.,Zhu, Y.P.,Chen, Y.W.,He,F.C.: **Integrated nr Database in Protein Annotation System and Its Localization.** *Computer Engineering* 2006(05):71–73.
24. Soudy M, Anwar AM, Ahmed EA, Osama A, Ezzeldin S, Mahgoub S, Magdeldin S: **UniprotR: Retrieving and visualizing protein sequence and functional information from Universal Protein Resource (UniProt knowledgebase).** *J PROTEOMICS* 2020, **213**:103613.
25. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT *et al.* **Gene ontology: tool for the unification of biology.** The Gene Ontology Consortium. *NAT GENET* 2000, **25**(1):25–29.
26. Kanehisa M, Goto S, Kawashima S, Okuno Y, Hattori M: **The KEGG resource for deciphering the genome.** *NUCLEIC ACIDS RES* 2004, **32**(Database issue):D277-D280.
27. Tatusov RL, Galperin MY, Natale DA, Koonin EV: **The COG database: a tool for genome-scale analysis of protein functions and evolution.** *NUCLEIC ACIDS RES* 2000, **28**(1):33–36.
28. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: **Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.** *NUCLEIC ACIDS RES* 1997, **25**(17):3389–3402.
29. Fang L, Wang Q, Hu Y, Jia Y, Chen J, Liu B, Zhang Z, Guan X, Chen S, Zhou B *et al.* **Genomic analyses in cotton identify signatures of selection and loci associated with fiber quality and yield traits.** *NAT GENET* 2017, **49**(7):1089–1098.
30. Huang C, Nie X, Shen C, You C, Li W, Zhao W, Zhang X, Lin Z: **Population structure and genetic basis of the agronomic traits of upland cotton in China revealed by a genome-wide association study using high-density SNPs.** *PLANT BIOTECHNOL J* 2017, **15**(11):1374–1386.
31. Huang X, Feng Q, Qian Q, Zhao Q, Wang L, Wang A, Guan J, Fan D, Weng Q, Huang T *et al.* **High-throughput genotyping by whole-genome resequencing.** *GENOME RES* 2009, **19**(6):1068–1076.
32. Rubin BE, Ree RH, Moreau CS: **Inferring phylogenies from RAD sequence data.** *PLOS ONE* 2012, **7**(4):e33394.
33. Xie W, Feng Q, Yu H, Huang X, Zhao Q, Xing Y, Yu S, Han B, Zhang Q: **Parent-independent genotyping for constructing an ultrahigh-density linkage map based on population sequencing.** *Proc Natl Acad Sci U S A* 2010, **107**(23):10578–10583.
34. Smit ME, McGregor SR, Sun H, Gough C, Bagman AM, Soyars CL, Kroon JT, Gaudinier A, Williams CJ, Yang X *et al.* **A PXY-Mediated Transcriptional Network Integrates Signaling Mechanisms to Control**

Vascular Development in Arabidopsis. *PLANT CELL* 2020, **32**(2):319–335.

35. Machemer K, Shaiman O, Salts Y, Shabtai S, Sobolev I, Belausov E, Grotewold E, Barg R: **Interplay of MYB factors in differential cell expansion, and consequences for tomato fruit development.** *PLANT J* 2011, **68**(2):337–350.
36. Zhao Q, Dixon RA: **Transcriptional networks for lignin biosynthesis: more complex than we thought?** *TRENDS PLANT SCI* 2011, **16**(4):227–233.
37. McCarthy RL, Zhong R, Fowler S, Lyskowski D, Piyasena H, Carleton K, Spicer C, Ye ZH: **The poplar MYB transcription factors, PtrMYB3 and PtrMYB20, are involved in the regulation of secondary wall biosynthesis.** *PLANT CELL PHYSIOL* 2010, **51**(6):1084–1090.
38. Zhong R, Ye ZH: **MYB46 and MYB83 bind to the SMRE sites and directly activate a suite of transcription factors and secondary wall biosynthetic genes.** *PLANT CELL PHYSIOL* 2012, **53**(2):368–380.
39. Nakano Y, Yamaguchi M, Endo H, Rejab NA, Ohtani M: **NAC-MYB-based transcriptional regulation of secondary cell wall biosynthesis in land plants.** *FRONT PLANT SCI* 2015, **6**:288.
40. Chakraborty A, Sarkar D, Satya P, Karmakar PG, Singh NK: **Pathways associated with lignin biosynthesis in lignomaniac jute fibres.** *MOL GENET GENOMICS* 2015, **290**(4):1523–1542.
41. Pingan G, Bo W, Yancheng Z, Jie C, Wenlue L, Lijun L, Dingxiang P: **Transcriptome analyses provide insights into the effect of temperature change on fiber quality of ramie.** *Industrial Crops & Products* 2020, **152**.
42. Behr M, Legay S, Hausman J, Lutts S, Guerriero G: **Molecular Investigation of the Stem Snap Point in Textile Hemp.** *GENES-BASEL* 2017, **8**(12).
43. Benjamin P, Kamal S, Alison F, Malvika S, Yaodong Y, Stijn D, Nicholas J, Ilda C, Paula P, Adnan S *et al*: **AUX/LAX Genes Encode a Family of Auxin Influx Transporters That Perform Distinct Functions during Arabidopsis Development(C)(W).** *PLANT CELL* 2012, **24**(7).
44. Norma F, Pau F, Ana C, Riccardo S, Jose MA, Ranjan S, Malcolm JB, Ari PMHN, Ana ICO, Marta IE: **Auxin Influx Carriers Control Vascular Patterning and Xylem Differentiation in Arabidopsis thaliana.** *PLOS GENET* 2015, **11**(4).
45. Guerriero G, Behr M, Legay S, Mangeot-Peter L, Zorzan S, Ghoniem M, Hausman JF: **Transcriptomic profiling of hemp bast fibres at different developmental stages.** *Sci Rep* 2017, **7**(1):4961.
46. Behr M, Legay S, Zizkova E, Motyka V, Dobrev PI, Hausman JF, Lutts S, Guerriero G: **Studying Secondary Growth and Bast Fiber Development: The Hemp Hypocotyl Peeks behind the Wall.** *FRONT PLANT SCI* 2016, **7**:1733.
47. Bezerra-Neto JP, de Araujo FC, Ferreira-Neto J, Da SM, Pandolfi V, Aburjaile FF, Sakamoto T, de Oliveira SR, Kido EA, Barbosa AL *et al*: **Plant Aquaporins: Diversity, Evolution and Biotechnological Applications.** *Curr Protein Pept Sci* 2019, **20**(4):368–395.
48. Li G, Santoni V, Maurel C: **Plant aquaporins: roles in plant physiology.** *Biochim Biophys Acta* 2014, **1840**(5):1574–1582.

49. Groszmann M, Osborn HL, Evans JR: **Carbon dioxide and water transport through plant aquaporins.** *PLANT CELL ENVIRON* 2017, **40**(6):938–961.
50. Anderberg HI, Danielson JA, Johanson U: **Algal MIPs, high diversity and conserved motifs.** *BMC EVOL BIOL* 2011, **11**:110.
51. Chaumont F, Barrieu F, Wojcik E, Chrispeels MJ, Jung R: **Aquaporins constitute a large and highly divergent protein family in maize.** *PLANT PHYSIOL* 2001, **125**(3):1206–1215.
52. Yaneff A, Vitali V, Amodeo G: **PIP1 aquaporins: Intrinsic water channels or PIP2 aquaporin modulators?** *FEBS LETT* 2015, **589**(23):3508–3515.
53. Secchi F, Zwieniecki MA: **Patterns of PIP gene expression in *Populus trichocarpa* during recovery from xylem embolism suggest a major role for the PIP1 aquaporin subfamily as moderators of refilling process.** *PLANT CELL ENVIRON* 2010, **33**(8):1285–1297.
54. Postaire O, Tournaire-Roux C, Grondin A, Boursiac Y, Morillon R, Schaffner AR, Maurel C: **A PIP1 aquaporin contributes to hydrostatic pressure-induced water transport in both the root and rosette of *Arabidopsis*.** *PLANT PHYSIOL* 2010, **152**(3):1418–1430.
55. Li L, Wang H, Gago J, Cui H, Qian Z, Kodama N, Ji H, Tian S, Shen D, Chen Y *et al*: **Harpin Hpa1 Interacts with Aquaporin PIP1;4 to Promote the Substrate Transport and Photosynthesis in *Arabidopsis*.** *Sci Rep* 2015, **5**:17207.
56. Kromdijk J, Glowacka K, Long SP: **Photosynthetic efficiency and mesophyll conductance are unaffected in *Arabidopsis thaliana* aquaporin knock-out lines.** *J EXP BOT* 2020, **71**(1):318–329.
57. Liu D, Tu L, Wang L, Li Y, Zhu L, Zhang X: **Characterization and expression of plasma and tonoplast membrane aquaporins in elongating cotton fibers.** *PLANT CELL REP* 2008, **27**(8):1385–1394.
58. Aslam U, Khatoon A, Cheema HM, Bashir A: **Identification and characterization of plasma membrane aquaporins isolated from fiber cells of *Calotropis procera*.** *J Zhejiang Univ Sci B* 2013, **14**(7):586–595.
59. Zhong R, Teng Q, Haghighat M, Yuan Y, Furey ST, Dasher RL, Ye ZH: **Cytosol-Localized UDP-Xylose Synthases Provide the Major Source of UDP-Xylose for the Biosynthesis of Xylan and Xyloglucan.** *PLANT CELL PHYSIOL* 2017, **58**(1):156–174.
60. Peltier JB, Cai Y, Sun Q, Zabrouskov V, Giacomelli L, Rudella A, Ytterberg AJ, Rutschow H, van Wijk KJ: **The oligomeric stromal proteome of *Arabidopsis thaliana* chloroplasts.** *MOL CELL PROTEOMICS* 2006, **5**(1):114–133.
61. Srivastava AC, Palanichelvam K, Ma J, Steele J, Blancaflor EB, Tang Y: **Collection and Analysis of Expressed Sequence Tags Derived from Laser Capture Microdissected Switchgrass (*Panicum virgatum* L. Alamo) Vascular Tissues.** *BIOENERG RES* 2010, **3**(3):278–294.
62. Villalobos DP, Diaz-Moreno SM, Said E, Canas RA, Osuna D, Van Kerckhoven SH, Bautista R, Claros MG, Canovas FM, Canton FR: **Reprogramming of gene expression during compression wood formation in pine: coordinated modulation of S-adenosylmethionine, lignin and lignan related genes.** *BMC PLANT BIOL* 2012, **12**:100.

63. Tang HM, Liu S, Hill-Skinner S, Wu W, Reed D, Yeh CT, Nettleton D, Schnable PS: **The maize brown midrib2 (bm2) gene encodes a methylenetetrahydrofolate reductase that contributes to lignin accumulation.** *PLANT J* 2014, **77**(3):380–392.
64. **Proceedings of the VII. Alps-Adria Scientific Workshop, 28 April–2 May 2008, Stara Lesna, Slovakia. Part III.** *CEREAL RES COMMUN* 2008, **36**(5):1395–2094.
65. Eulgem T, Rushton PJ, Robatzek S, Somssich IE: **The WRKY superfamily of plant transcription factors.** *TRENDS PLANT SCI* 2000, **5**(5):199–206.
66. Ren CM, Zhu Q, Gao BD, Ke SY, Yu WC, Xie DX, Peng W: **Transcription factor WRKY70 displays important but no indispensable roles in jasmonate and salicylic acid signaling.** *J INTEGR PLANT BIOL* 2008, **50**(5):630–637.
67. Hu Q, Xiao S, Guan Q, Tu L, Sheng F, Du X, Zhang X: **The laccase gene GhLac1 modulates fiber initiation and elongation by coordinating jasmonic acid and flavonoid metabolism.** *The Crop Journal* 2020, **8**(4):522–533.
68. Pradipta S, Sanjoy S, Asitava B: **Identification of differentially expressed transcripts associated with bast fibre development in Corchorus capsularis by suppression subtractive hybridization.** *PLANTA* 2015, **241**(2).
69. Gea G, Marc B, Sylvain L, Lauralie M, Simone Z, Mohammad G, Jean-Francois H: **Transcriptomic profiling of hemp bast fibres at different developmental stages.** *SCI REP-UK* 2017, **7**(1).

Tables

Table 1 SNP-index Calculation formulae

Calculation formulae
$\text{SNPindex}_{aa} = \frac{Maa}{(Maa + Paa)}$ <p>Maa showed the female sample depth from aa pool while Paa indicate the male sample depth from the same pool.</p>
$\text{SNPindex}_{ab} = \frac{Mab}{(Mab + Pab)}$ <p>Mab stood for the female sample depth from ab pool while Pab was the male sample depth from the same pool</p>
$\Delta(\text{SNP-index}) = \text{SNPindex}_{aa} - \text{SNPindex}_{ab}$

Table 2 Statistic results of sequencing data from each sample

Sample	Sample ID	Clean-Read number	Clean-Base number	Q30 percentage(%)	GC percentage(%)
Male parent	R01	61,937,343	18,556,453,868	93.05	35.00
Female parent	R02	59,738,133	17,895,069,796	93.60	34.16
High-fiber content pool	aa	12,354,037	3,113,217,324	93.40	42.51
Low-fiber content pool	ab	19,527,628	4,920,962,256	93.60	40.08

Table 3 Number of all SLAFs on each chromosome

Chromosome_ID	Chromosome_length	Expected_SLAF_number	Average_SLAF_distance
NC_044370.1	104987320	13010.7	8069.31
NC_044371.1	101209240	11194.1	9041.34
NC_044372.1	94670641	11249.6	8415.47
NC_044373.1	91913879	11419.3	8048.97
NC_044374.1	88181582	11734.1	7514.98
NC_044375.1	96346938	11349.4	8489.18
NC_044376.1	61561104	7909.49	7783.19
NC_044377.1	79335105	10857.7	7306.82
NC_044378.1	71238074	8857.93	8042.29
NC_044379.1	64622176	8240.72	7841.81
Total	854066059	105823	8070.7

Table 4 SNP filtering Statistic

Total SNP	Locus of multiple alleles	Locus with Read support less than 4	Locus of mixed pool genotypic uniformity	Locus filtered by the parent	High quality SNP
7,331,769	44,040	6,773,604	122,330	366,662	25,133

Table 5 The statistic of associated region information

Chromosome_ID	Start×bp×	End×bp×	Size(Mb)	Gene_Number
Chr1	12,344,405	12,561,297	0.22	12
Chr1	14,324,437	14,342,707	0.02	1
Chr1	14,600,744	22,565,820	7.97	336
Chr1	248,400	718,823	0.47	48
Total	-	-	8.68	397

Figures

Figure 1

Experimental cultivars and materials (A) Monoecious Jindao-15. (B) Female plant of dioecious Fire No.1 (C) Male plant of Fire No.1. (D) Fresh bast fiber peeled from stem.

Figure 2

Distribution diagrams of all SLAFs (black lines) on each chromosome of hemp. The black lines indicate SLAF tags.

Figure 3

Distribution of SNP-index correlation values on chromosomes. The x-axis indicate chromosome name. Color dot represents the calculated SNPS - index (or Δ SNPS - index) values, the black line represents fitting SNP-index (or Δ SNPS - index) values.(A) SNP-index graphs of high fiber content mixed pool. (B) SNP-index graphs of low fiber content mixed pool. (C) Distribution diagram of Δ SNP-index value, of which the red dashed line represents the threshold line for the 99th percentile.

Figure 4

Relative expression level of six candidate genes in different hemp varieties. Each bar represents the average expression level of three independent biological replicates. Error bars show standard errors of the average values. (*P < 0.05, **P < 0.01)

Figure 5

Genes enrichment results in the candidate regions via KEGG analysis. The x-axis represents the enrichment factor and the y-axis represents Q-value. The smaller the Q-value, the higher the enrichment degree.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaltable1.xls](#)
- [Supplementaltable2.xls](#)
- [Supplementaltable3.xls](#)
- [Supplementaltable4.xls](#)