

Risk Factors of Metachronous Brain Metastasis in Patients with EGFR-Mutated Advanced Non-Small Cell Lung Cancer

Wen Ouyang

Wuhan University Zhongnan Hospital

Jing Yu

Wuhan University Zhongnan Hospital

Yan Zhou

Wuhan University Zhongnan Hospital

Jing Hu

Wuhan University Zhongnan Hospital

Zhao Huang

Wuhan University Zhongnan Hospital

Junhong Zhang

Wuhan University Zhongnan Hospital

Conghua Xie (✉ chxie_65@whu.edu.cn)

Wuhan University Zhongnan Hospital Department of Chemotherapy and Radiation Therapy

<https://orcid.org/0000-0003-2836-6373>

Research article

Keywords: non-small cell lung cancer, epidermal growth factor receptor, brain metastases, risk factor

Posted Date: May 18th, 2020

DOI: <https://doi.org/10.21203/rs.2.17604/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on July 30th, 2020. See the published version at <https://doi.org/10.1186/s12885-020-07202-8>.

Abstract

Background NSCLC patients with EGFR mutation were at a higher incidence of developing brain metastasis (BM). Patients with BM are associated with high mortality. Reducing BM incidence becomes increasingly significant for NSCLC patients to achieve prolonged survival. The aim of the study was to explore the possible risk factors of developing metachronous BM during EGFR-TKIs treatment, and to identify the potential candidates for prophylactic cranial irradiation (PCI) or the first-line osimertinib treatment.

Methods A total of 157 consecutive EGFR-mutated advanced NSCLC patients without BM at initial diagnosis in our institution from 2012 and 2018 were retrospectively reviewed. Comparisons of OS were performed based on BM status. The cumulative incidence of metachronous BM was calculated by the Kaplan-Meier method, and the independent risk factors of metachronous BM were investigated by multivariate analysis.

Results Patients developing metachronous BM had worse survival (mOS: 22.1 months) than patients not-developing BM (mOS: 44.8 months). Moreover, the multivariate analysis indicated that age ≤ 49 years ($P = 0.035$), number of extracranial metastases ($P = 0.013$), and malignant pleural effusion ($P = 0.002$) were independent risk factors of metachronous BM. Furthermore, the 1-year actuarial incidence of developing metachronous BM in patients with no risk factor ($n = 101$), 1 risk factor ($n = 46$), and 2 risk factors ($n = 10$) were 7.01%, 14.61%, and 43.75%, respectively ($P < 0.001$).

Conclusions Patients developing metachronous BM during EGFR-TKIs treatment have worse outcomes. Our results suggested that EGFR-mutated advanced NSCLC patients with ≥ 1 risk factors were candidates for PCI or the first-line Osimertinib treatment.

Introduction

Lung cancer is the leading cause of cancer death all over the world [1]. Among them, 80-85% of patients are diagnosed as non-small cell lung cancer (NSCLC) [2]. Despite the presence of the blood-brain barrier (BBB), brain is still a frequent site of NSCLC metastasis. 10% of NSCLC patients present brain metastasis (BM) at their initial diagnosis, and 40-50% of patients develop metachronous BM during the course of the disease [3]. The median survival in patients who suffer from BM without treatment is less than 1 month, and whole brain radiation therapy (WBRT), as the standard treatment for BM, also can bring about only 3-6 months median survival [4]. Patients with BM are associated with high morbidity, poor prognosis, neurocognitive and life quality deficits [5]. Therefore, reducing incidence of BM becomes increasingly significant for NSCLC patients to achieve prolonged survival.

Epidermal growth factor receptor (EGFR) mutations are observed in approximately 10-15% of the Caucasian population [6] and more than 50% of the Asian population [7] with non-squamous NSCLC. EGFR-tyrosine kinase inhibitors (TKIs) largely improved the survival of EGFR-mutated advanced NSCLC patients [8-10]. Although EGFR-TKIs was reported to pass through BBB and reduce BM in EGFR-mutated

NSCLC patients [11,12], there remain some patients developing metachronous BM during the course of EGFR-TKIs therapy. Lee et al found that 26% of the patients developed central nervous system (CNS) failure and 13% experienced isolated CNS failure among 166 patients with a clinical benefit to first-generation EGFR-TKIs (gefitinib or erlotinib) treatment [13]. Additionally, it was reported that NSCLC patients with EGFR mutation were at a higher incidence of developing BM [14-16]. And patients with longer survival exposed to higher risk of BM [14]. Moreover, EGFR-mutated NSCLC patients with BM had a worse median OS of 25.1 months than the patients without BM (30.2 months) [17]. These results suggested that it was necessary for EGFR-mutated advanced NSCLC patients to reduce risk of developing BM.

How to reduce incidence of developing metachronous BM for EGFR-mutated advanced NSCLC patients? Firstly, prophylactic cranial irradiation (PCI) is a technique that delivers radiation therapy (RT) to the whole brain to prevent BM occurrence. It was reported to significantly reduce incidence of metachronous BM and improve overall survival (OS) in patients with limited-stage small cell lung cancer (SCLC) [18]. In addition, Osimertinib is an oral, irreversible third-generation EGFR-TKIs with higher penetration in CNS [19-21]. FLAURA study showed the lower frequency of CNS progression in the Osimertinib group than in the standard EGFR-TKIs group [21]. However, the results of RTOG-0214 on the effects of PCI in localized NSCLC patients indicated that PCI could reduce BM incidence, whereas failed to improve OS [22] and led to decline in immediate and delayed recall [23]. Interestingly, the 10-years update of RTOG-0214 indicated that only patients non-operatively treated have an improved OS ($P= 0.026$, HR = 1.42, 95% CI: 1.04-1.94) and DFS ($P= 0.014$), suggesting that PCI might just benefit NSCLC patients with higher risk of BM. For Osimertinib, considering the cost of Osimertinib, the first-line Osimertinib treatment for EGFR-mutated advanced NSCLC patients is not widely available in most developing countries. Therefore, it is higher cost-effective to apply the first-line Osimertinib treatment for patients with higher risk of BM. These findings prompted us to identify population subsets with higher risk of BM, who are candidates for PCI or the first-line Osimertinib treatment. Previous studies identified several risk factors of BM in NSCLC, including younger age [24-26], non-squamous cell carcinoma [24], high serum CEA level [27], and disease stages [25,28]. However, they were not specific for EGFR-mutated advanced NSCLC patients, and synchronous BM at initial diagnosis and metachronous BM during their disease course are seldom differentiated in these reports.

Consequently, we established a retrospective single-institutional database including consecutive patients with EGFR-mutated advanced NSCLC between January 2012 and June 2018, to evaluate the impact of BM status on OS, to explore the possible risk factors for developing metachronous BM during the course of first-generation EGFR-TKIs therapy, and to identify the potential candidates for PCI or the first-line Osimertinib treatment.

Methods

Patients

The flow chart of patient enrollment was shown in **Figure 1**. Between January 2012 and June 2018, a total of 157 consecutive EGFR-mutated advanced NSCLC patients without BM at initial diagnosis were reviewed at the Department of Radiation and Medical Oncology, Zhongnan Hospital of Wuhan University. Our inclusion criteria are: (1) NSCLC was confirmed by cytology (14 pts), or histology (143 pts) (World Health Organization, WHO); (2) EGFR mutations were detected by real-time quantitative PCR (ARMS, 126 pts) or Next Generation Sequencing (NGS, 31 pts), using histological or cytological specimens from primary or metastatic lesions; (3) The disease was clinically diagnosed as stage IIIB (10 pts)-IV (147 pts) (American Joint Committee on Cancer, the 7th Edition); (4) Patients had negative results of enhanced magnetic resonance imaging (MRI, 149 pts) or computed tomography (CT, 8 pts) scans of brain before initial treatment; (5) Patients were treatment naive for EGFR-TKIs treatment. All patients received comprehensive assessments within 1 month before treatment, including physical and pathological examination, EGFR mutation test, and TNM stage evaluation.

Treatment and Follow up

Among the 157 patients without BM at initial diagnosis, 24 patients received chemotherapy as their first-line therapy, and the other 133 patients received EGFR-TKIs treatment initially. EGFR-TKIs (gefitinib, erlotinib, or icotinib) were continuously administered until progression of disease (PD) or intolerable side effects. Treatment beyond PD was allowed on the judgement of continuously clinical benefit by the oncologists.

Follow-up examinations were performed every 2 months, including thoracic and abdominal CT scan, brain MRI scan. Progression-free survival (PFS) was defined as the time from EGFR-TKIs treatment to PD (including local, regional, or distant progression) or death from any cause. OS was defined as the time from EGFR-TKIs treatment to death from any cause. Brain-metastasis-free survival (BMFS) was defined as the time from EGFR-TKIs treatment to BM occurrence. Treatment responses were evaluated by the response evaluation criteria in solid tumors as complete response (CR), partial response (PR), stable (SD), and progression (PD).

Statistics

All statistical analyses were conducted using Statistical Package for Social Scientists (SPSS/Windows, Version 22.0, SPSS Inc., Chicago, USA). Descriptive statistics were used for categorical variables (frequency and percentage) and continuous variables (median and range). The cumulative incidence of BM and survival were calculated by the Kaplan-Meier method with 95% confidence intervals (CIs). Univariable and multivariable Cox regression analyses were performed to explore the risk factors of metachronous BM. The multivariable Cox regression analysis simultaneously included factors that had shown associations ($P < 0.100$) in the univariable Cox regression analyses, and variables based on their clinical significance according to previously literature reports. The optimal cut-off values of continuous variables were calculated by X-tile software [29]. All tests were two-sided and $P < 0.05$ were considered statistically significant.

Results

Patient characteristics

The flow chart of patient enrollment was shown in **Figure 1**. Between January 2012 and June 2018, Among the 229 consecutive patients with EGFR-mutated advanced NSCLC, three patients were excluded due to short EGFR-TKI treatment (< 1 month), and 69 patients were excluded due to synchronous BM. A total of 157 patients without BM at initial diagnosis were included: 30 patients (19.1%) developed metachronous BM during EGFR-TKIs treatment and 127 patients (80.9%) didn't. The clinical and treatment characteristics of these patients grouped by BM status are shown in **Table 1**. The median age of the patients without BM and patients developing metachronous BM was 60 and 54 years, respectively. Patients who would develop metachronous BM were more likely to have a more favorable Karnofsky Performance Status (KPS score \geq 80: 100% patients developing metachronous BM vs. 90.5% patients without BM). There was no difference between the two groups with respect to gender, histology, BMI, smoking status, tumor markers level, clinical stages, and extracranial metastatic location.

The incidence of metachronous BM and survival

The median duration of follow-up was 24.1 months (95% CI: 19.6-28.6 months). Thirty patients (19.1%) developed metachronous BM during EGFR-TKIs treatment. Among them, patients with symptomatic and asymptomatic BM were 18 (60%) and 12 (40%) respectively. Fourteen patients (46.7%) received WBRT and 8 patients (26.7%) received stereotactic radiosurgery (SRS) plus continuous EGFR-TKIs treatment, other 2 patients (6.7%) received continuous EGFR-TKIs plus supportive care, and 6 patients (20%) switched to chemotherapy. In addition, 9 patients (9/30, 30%) receiving chemotherapy as the first-line treatment developed metachronous BM during subsequent EGFR-TKIs therapy. The 1-, 2- and 3-year risks of BM were 11.6%, 22.6% and 29.4% respectively (**Figure 2**).

The median OS of these 157 patients was 37.5 months (95% CI: 27.6-47.4 months). The 1-, 2- and 3-year OS rates were 86.9%, 69.8% and 55.9% respectively (**Figure 2**). For PFS, 105 patients (66.9%) progressed during follow-up time. Median PFS was 13.6 months (95% CI:11.2-15.9 months). The 1-, 2- and 3-year PFS rates were 57.8%, 29.4% and 21.3% respectively (**Figure 2**). Our median OS and PFS were longer than those of the clinical trials for patients with EGFR-mutated advanced NSCLC [30].

The overall response rates were partial for 76.4%, stable for 23.0%, and progressive for 0.6% of EGFR-TKIs treatment at the first follow-up examination.

Overall survival of patients grouped by BM status

To evaluate the impact of BM status on OS, the 157 patients were grouped by with metachronous BM and without BM during EGFR-TKIs treatment. Compared with patients without BM, patients developing metachronous BM during the course of EGFR-TKIs treatment were at a higher risk on OS (HR=1.86, 95%CI:1.07-3.26). Our findings confirmed that patients developing metachronous BM during EGFR-TKIs

treatment had poorer outcomes (median OS: 28.6 months) than patients without BM (median OS: 44.8 months, **Figure 3**).

Risk factors of developing metachronous BM

Several clinical factors were associated with metachronous BM in both univariate and multivariate analyses (**Table 2**). In univariate analyses, BM was associated with age, the first-line treatment regimens, types of EGFR mutations, numbers of extracranial metastases, and malignant pleural effusion. Other factors such as gender, KPS score, smoking status, tumor marker levels before treatment, clinical stages, types of EGFR-TKIs, and metastatic locations were not associated with metachronous BM.

The factors showing associations ($P < 0.100$) in the univariable Cox regression analyses, as well as other factors that were reported to be associated with BM in previous studies [27,31] were further examined by multivariable Cox regression analysis. Results of multivariate analysis indicated that age ≤ 49 years ($P = 0.035$), numbers of extracranial metastases ($P = 0.013$), and documented malignant pleural effusion ($P = 0.002$) were independent high-risk factors of developing metachronous BM, while the first-line treatment regimens and types of EGFR mutations were not associated with metachronous BM in multivariate Cox regression analysis.

Furthermore, the 1-year actuarial incidence of developing metachronous BM in patients with no risk factor ($n = 101$), 1 risk factor ($n = 46$), and 2 risk factors ($n = 10$) were 7.01%, 14.61%, and 43.75%, respectively ($P < 0.001$, **Figure 4**). Meanwhile, we performed an internal validation by randomly selecting 52 cases from our patient cohort. The 1-year actuarial incidence of developing metachronous BM in these 52 patients with no risk factor ($n = 37$), 1 risk factor ($n = 11$), and 2 risk factors ($n = 4$) were 5.65%, 22.2%, and 50.0%, respectively ($P = 0.004$, **Figure S1**). These results were consistent with the whole cohort, indicating the credibility of the result to some extent. Therefore, patients with more risk factors had a higher risk of developing metachronous BM. Our studies suggested that the patients with ≥ 1 risk factors were more likely to benefit from PCI or the first-line Osimertinib treatment.

Discussion

During the past two decades, the advances of EGFR-TKIs revolutionarily improved the prognosis of patients with EGFR-mutated advanced NSCLC. The WJTOG3405 trial reported that the median OS of patients with EGFR-mutated advanced NSCLC was up to 30.2 months [32]. Our results of 157 EGFR-mutated advanced NSCLC patients without BM at initial diagnosis showed a median OS of 37.5 months and a median PFS of 13.6 months (**Figure 2**). Although EGFR-TKIs was reported to be more effective than chemotherapy for BM in EGFR-mutated NSCLC patients [11,12], patients with longer survival exposed to a higher risk of BM [14], and patients with EGFR mutations were at a higher incidence of developing BM [14-16]. In our study, 30 patients (30/157, 19.1%) developed metachronous BM during first-generation EGFR-TKIs treatment, and 1-, 2- and 3-year risks of developing BM were 11.6%, 22.6% and 29.4% respectively (**Figure 2**). Therefore, the first-generation EGFR-TKIs therapy resulted in decreased risk of non-BM lesions but had limited impact on BM.

It was well known that BM is a common reason leading to treatment failure [33]. In our study, compared with patients without BM, patients developing metachronous BM during the course of EGFR-TKIs treatment were at a higher risk on OS (HR=1.86, 95%CI:1.07-3.26). Our findings confirmed that patients with metachronous BM had worse outcomes (**Figure 3**) on the condition that there was no difference on clinical and treatment characteristics between the two groups grouped by BM status (**Table 1**). Therefore, reducing incidence of BM in EGFR-mutated advanced NSCLC patients becomes increasingly significant to achieve prolonged survival.

The use of PCI or the first-line Osimertinib treatment could reduce incidence of metachronous BM among EGFR-mutated advanced NSCLC patients. However, existing evidences suggest that PCI might just suitable for patients with high risk of developing BM, and the high cost of Osimertinib led to the limitation of first-line Osimertinib treatment in most developing countries. Therefore, it is important to identify population subsets with higher risk of BM as candidates for PCI or the first-line Osimertinib treatment. But there is a lack of relevant data for EGFR-mutated advanced NSCLC in literature.

In the current study, multivariate analysis indicated that age ≤ 49 years was correlated with higher risk of metachronous BM (**Table. 2**). Despite the difference of age cut-off, our results were consistent with previous studies [25,34]. The underlying mechanism remains unclear. It was partly interpreted that young people may have better performance status, which is associated with longer survival, leading to higher risk of exposure to BM. Moreover, several studies have shown that BM is associated with the angiogenic microenvironment, and the cerebrovascular microenvironment factors of young patients may be better than those of older patients [35]. Further investigations are required to identify the specific reasons that younger patients are more likely to develop BM.

The numbers of extracranial metastases and malignant pleural effusion were also independent risk factors of metachronous BM (**Table. 2**). The underlying mechanism was also unclear. It may be interpreted that both pleural effusion and BM is associated with the angiogenic microenvironment [35]. In addition, the numbers of extracranial metastases are reflection of tumor burden, which was positive correlated with the development of BM.

Furthermore, our results confirmed that the predictive value of gender and KPS score for metachronous BM may remain controversial [36]. Previous studies reported that elevated CEA [27,31,36], NSE [24], and CA125 [24] were independent risk factors of BM. However, there is no correlation between tumor markers levels before treatment (including CEA, NSE, and CA125) and the metachronous BM in our study. And the first-line treatment regimen was also not associated with metachronous BM in our multivariate Cox analysis. In addition, a recent retrospective study [31] showed that point mutations in exon 21 were independent risk factors of BM. However, our results failed to show a statistical difference in the association between types of EGFR mutations and metachronous BM.

Furthermore, the 1-year actuarial risk of developing metachronous BM in patients with no risk factor (n =101), 1 risk factor (n =46), and 2 risk factors (n =10) were 7.01%, 14.61%, and 43.75%, respectively ($P < 0.001$, **Figure 4**). Obviously, patients with more risk factors had a higher risk of developing metachronous

BM. Our studies suggested that the patients with ≥ 1 risk factors were more likely to benefit from PCI or were candidates for the first-line Osimertinib treatment. Certainly, there are several limitations in our study, this was a retrospective study in a single institution, which inevitably resulted in a selection bias. More finely devised prospective and random study is needed to confirm our results, and the mechanisms of the correlation between these risk factors and metachronous BM are to be further explored.

Conclusions

Collectively, the findings of this study were as follows. First, our study confirmed EGFR-mutated advanced NSCLC patients with metachronous BM had worse outcomes. Second, the multivariate Cox analysis indicated that younger age (≤ 49 years), more extracranial metastases, and malignant pleural effusion were independent risk factors of metachronous BM. Third, the patients with more risk factors are more likely to benefit from PCI or the first-line Osimertinib treatment.

List Of Abbreviations

BM, brain metastasis; PCI, prophylactic cranial irradiation; BBB, blood-brain barrier; CI, confidence interval; EGFR, epidermal growth factor receptor; NSCLC, non-small cell lung cancer; SCLC, small cell lung cancer; WBRT, whole brain radiation therapy; SRS, stereotactic radiosurgery; CNS, central nervous system; NGS, Next Generation Sequencing; NSE, neuron-specific enolase; DFS, disease free survival; OS, overall survival; PD, progression of disease; PFS, progression-free survival; BMFS, Brain-metastasis-free survival; CR, complete response; PR, partial response; SD, stable; TKI, tyrosine kinase inhibitor; VEGF, vascular endothelial growth factor.

Declarations

Ethics approval and consent to participate

This study was approved by the Ethics Committee of Zhongnan hospital of Wuhan University. Ethics Committee approved verbal informed consent. Because our study is retrospective, and we collected and analyzed the data of enrolled patients by reviewing their electronic medical records. The information of enrolled patients is anonymous. The collection, analysis and publication of the data will not infringe enrolled patients' health, safety and privacy. Informed consent was obtained orally from the included patients by telephone.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and analyzed in the current study are available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by grants from the National Natural Science Foundation of China (No. 81502308 and 81773236). The funding bodies had no role on the design, data collection, analysis and manuscript writing of this study.

Author Contributions

Conception and design of the work: WO, JY and CX. Acquisition, analysis and interpretation of data: WO, YZ, JY, JH and ZH. Drafting and revising of the article: WO, JZ and CX. Final approval of the manuscript and agreement to be accountable for all aspects of the work: All authors.

Acknowledgements

None

References

1. Govindan R, Page N, Morgensztern D, et al. Changing epidemiology of small-cell lung cancer in the United States over the last 30 years: analysis of the surveillance, epidemiologic, and end results database. *J Clin Oncol.* 2006;24:4539-44. <https://doi.org/10.1200/JCO.2005.04.4859>.
2. Saad AG, Yeap BY, Thunnissen FB, et al. Immunohistochemical markers associated with brain metastases in patients with nonsmall cell lung carcinoma. *Cancer.* 2008;113:2129-38. <https://doi.org/10.1002/cncr.23826>.
3. Lin NU, Wefel JS, Lee EQ, et al. Challenges relating to solid tumour brain metastases in clinical trials, part 2: neurocognitive, neurological, and quality-of-life outcomes. A report from the RANO group. *Lancet Oncol.* 2013;14:e407-16. [https://doi.org/10.1016/S1470-2045\(13\)70308-5](https://doi.org/10.1016/S1470-2045(13)70308-5)
4. Langer CJ, Mehta MP. Current management of brain metastases, with a focus on systemic options. *J Clin Oncol.* 2005;23:6207-19. <https://doi.org/10.1200/JCO.2005.03.145>
5. Kepka L, Cieslak E, Bujko K, et al. Results of the whole-brain radiotherapy for patients with brain metastases from lung cancer: the RTOG RPA intra-classes analysis. *Acta Oncol.* 2005;44:389-98. <https://doi.org/10.1080/02841860510029699>
6. Barlesi F, Mazieres J, Merlio JP, et al. Biomarkers France contributors. Routine molecular profiling of patients with advanced non-small-cell lung cancer: results of a 1-year nationwide programme of the

- French Cooperative Thoracic Intergroup (IFCT). *Lancet*. 2016;387:1415-26. [https://doi.org/10.1016/S0140-6736\(16\)00004-0](https://doi.org/10.1016/S0140-6736(16)00004-0).
7. Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med*. 2009;361:947-57. <https://doi.org/10.1056/NEJMoa0810699>.
 8. Maemondo M, Inoue A, Kobayashi K, et al. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med*. 2010;362:2380-8. <https://doi.org/10.1056/NEJMoa0909530>.
 9. Mok TS, Wu Y-L, Ahn M-J, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med*. 2017;376:629-40. <https://doi.org/10.1056/NEJMoa1612674>.
 10. Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med*. 2018;378:113-25. <https://doi.org/10.1056/NEJMoa1713137>.
 11. Wu YL, Zhou C, Cheng Y, et al. Erlotinib as second-line treatment in patients with advanced non-small-cell lung cancer and asymptomatic brain metastases: a phase II study (CTONG-0803). *Ann Oncol*. 2013;24:993-9. <https://doi.org/10.1093/annonc/mds529>.
 12. Weber B, Winterdahl M, Memon A, et al. Erlotinib accumulation in brain metastases from non-small cell lung cancer: visualization by positron emission tomography in a patient harboring a mutation in the epidermal growth factor receptor. *J Thorac Oncol*. 2011;6:1287-9. <https://doi.org/10.1097/JTO.0b013e318219ab87>.
 13. Lee YJ, Choi HJ, Kim SK, et al. Frequent central nervous system failure after clinical benefit with epidermal growth factor receptor tyrosine kinase inhibitors in Korean patients with non small-cell lung cancer. *Cancer*. 2010;116:1336-43. <https://doi.org/10.1002/cncr.24877>.
 14. Han G, Bi J, Tan W, et al. A retrospective analysis in patients with EGFR-mutant lung adenocarcinoma: is EGFR mutation associated with a higher incidence of brain metastasis?. *Oncotarget*. 2016;7:56998-7010. <https://doi.org/10.18632/oncotarget.10933>.
 15. Bhatt VR, D'Souza SP, Smith LM, Cushman-Vokoun AM, Noronha V, Verma V, et al. Epidermal Growth Factor Receptor Mutational Status and Brain Metastases in Non-Small-Cell Lung Cancer. *J Glob Oncol*. 2016;3:208-17. <https://doi.org/10.1200/JGO.2016.003392>.
 16. Fan Y, Xu X, Xie C. EGFR-TKI therapy for patients with brain metastases from non-small-cell lung cancer: a pooled analysis of published data. *Onco Targets Ther*. 2014;7:2075-84. <https://doi.org/10.2147/OTT.S67586>.
 17. Sperduto PW, Yang TJ, Beal K, et al. The Effect of Gene Alterations and Tyrosine Kinase Inhibition on Survival and Cause of Death in Patients With Adenocarcinoma of the Lung and Brain Metastases. *Int J Radiat Oncol Biol Phys*. 2016;96:406-13. <https://doi.org/10.1016/j.ijrobp.2016.06.006>.
 18. Komaki R, Scott CB, Byhardt R, et al. Failure patterns by prognostic group determined by recursive partitioning analysis (RPA) of 1547 patients on four radiation therapy oncology group (RTOG) studies in inoperable nonsmall-cell lung cancer (NSCLC). *Int J Radiat Oncol Biol Phys*. 1998;42:263-7. [https://doi.org/10.1016/S0360-3016\(98\)00213-2](https://doi.org/10.1016/S0360-3016(98)00213-2).
 19. Cross DA, Ashton SE, Ghiorghiu S, et al. AZD9291, an irreversible EGFR TKI, overcomes T790M-mediated resistance to EGFR inhibitors in lung cancer. *Cancer Discov*. 2014;4:1046-61.

<https://doi.org/10.1158/2159-8290.cd-14-0337>.

20. Ballard P, Yates JW, Yang Z, et al. Preclinical Comparison of Osimertinib with Other EGFR-TKIs in EGFR-Mutant NSCLC Brain Metastases Models, and Early Evidence of Clinical Brain Metastases Activity. *Clin Cancer Res*. 2016;22:5130-40. <https://doi.org/10.1158/1078-0432.ccr-16-0399>.
21. Vansteenkiste J, Reungwetwattana T, Nakagawa K, et al. CNS response to osimertinib vs standard of care (SoC) EGFR-TKI as first-line therapy in patients (pts) with EGFR-TKI sensitising mutation (EGFRm)-positive advanced non-small cell lung cancer (NSCLC): Data from the FLAURA study. *Ann Oncol*. 2017;28. <https://doi.org/10.1093/annonc/mdx729.007>.
22. Gore EM, Bae K, Wong SJ, et al. Phase III comparison of prophylactic cranial irradiation versus observation in patients with locally advanced non-small-cell lung cancer: primary analysis of radiation therapy oncology group study RTOG 0214. *J Clin Oncol*. 2011;29:272-8. <https://doi.org/10.1200/JCO.2010.29.1609>.
23. Sun A, Bae K, Gore EM, et al. Phase III trial of prophylactic cranial irradiation compared with observation in patients with locally advanced non-small-cell lung cancer: neurocognitive and quality-of-life analysis. *J Clin Oncol*. 2011;29:279-86. <https://doi.org/10.1200/JCO.2010.29.6053>
24. Ji Z, Bi N, Wang J, et al. Risk factors for brain metastases in locally advanced non-small cell lung cancer with definitive chest radiation. *Int J Radiat Oncol Biol Phys*. 2014;89:330-7. <https://doi.org/10.1016/j.ijrobp.2014.02.025>
25. Ceresoli GL, Reni M, Chiesa G, et al. Brain metastases in locally advanced nonsmall cell lung carcinoma after multimodality treatment: risk factors analysis. *Cancer*. 2002;95:605-12. <https://doi.org/10.1002/cncr.10687>.
26. Gaspar LE, Chansky K, Albain KS, et al. Time from treatment to subsequent diagnosis of brain metastases in stage III non-small-cell lung cancer: a retrospective review by the Southwest Oncology Group. *J Clin Oncol*. 2005;23:2955-61. <https://doi.org/10.1200/JCO.2005.08.026>.
27. Arrieta O, Saavedra-Perez D, Kuri R, et al. Brain metastasis development and poor survival associated with carcinoembryonic antigen (CEA) level in advanced non-small cell lung cancer: a prospective analysis. *BMC Cancer*. 2009;9:119. <https://doi.org/10.1186/1471-2407-9-119>
28. Robnett TJ, Machtay M, Stevenson JP, et al. Factors affecting the risk of brain metastases after definitive chemoradiation for locally advanced non-small-cell lung carcinoma. *J Clin Oncol*. 2001;19:1344-9. <https://doi.org/10.1200/JCO.2001.19.5.1344>.
29. Camp RL, Dolled-Filhart M, Rimm DL. X-tile: a new bio-informatics tool for biomarker assessment and outcome-based cut-point optimization. *Clin Cancer Res*. 2004;10:7252-9. <https://doi.org/10.1158/1078-0432.CCR-04-0713>.
30. Roeper J, Griesinger F. Epidermal growth factor receptor tyrosine kinase inhibitors in advanced nonsmall cell lung cancer: what is the preferred first-line therapy?. *Curr Opin Oncol*. 2019 ;31:1-7. <https://doi.org/10.1097/CCO.0000000000000495>.
31. Ma X, Zhu H, Guo H, et al. Risk factors of brain metastasis during the course of EGFR-TKIs therapy for patients with EGFR-mutated advanced lung adenocarcinoma. *Oncotarget*. 2016;7:81906-17.

<https://doi.org/10.18632/oncotarget.11918>.

32. Mitsudomi T, Morita S, Yatabe Y, et al. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol*. 2010;11:121-8. [https://doi.org/10.1016/S1470-2045\(09\)70364-X](https://doi.org/10.1016/S1470-2045(09)70364-X).
33. Arrieta O, Villarreal-Garza C, Zamora J, et al. Long-term survival in patients with non-small cell lung cancer and synchronous brain metastasis treated with whole-brain radiotherapy and thoracic chemoradiation. *Radiat Oncol*. 2011;6:166. <https://doi.org/10.1186/1748-717X-6-166>.
34. Dimitropoulos C, Hillas G, Nikolakopoulou S, et al. Prophylactic cranial irradiation in non-small cell lung cancer patients: who might be the candidates?. *Cancer Manag Res*. 2011;3:287-94. <https://doi.org/10.2147/CMR.S22717>.
35. Cedrés S, Nuñez I, Longo M, et al. Serum tumor markers CEA, CYFRA21-1, and CA-125 are associated with worse prognosis in advanced non-small-cell lung cancer (NSCLC). *Clin Lung Cancer*. 2011;12:172-9. <https://doi.org/10.1016/j.clc.2011.03.019>.
36. An N, Jing W, Wang H, et al. Risk factors for brain metastases in patients with non-small-cell lung cancer. *Cancer Med*. 2018;7:6357-64. <https://doi.org/10.1002/cam4.1865>.

Tables

Table 1. Baseline and treatment characteristics of patients grouped by BM status

Characteristic	Patients without BM		Patients developing metachronous BM (n=30)	
	NO.	(n=127) %	NO.	%
Age, years				
≤ 49	24	18.9	12	40.0
> 49	103	81.1	18	60.0
Median (Range)	60 (28-93)		54 (33-75)	
Gender				
Male	63	49.6	14	46.7
Female	64	50.4	16	53.3
KPS score				
≥80	115	90.5	30	100
<80	12	9.5	0	0
Histology				
Adenocarcinoma	122	96.1	28	93.3
Non-adenocarcinoma	5	3.9	2	6.7
BMI				
Mean (95%CI)	21.9 (14.9-28.8)		22.7 (16.3-29.2)	
Smoking status				
Yes	42	33.1	8	26.7
No	85	66.9	22	73.3
CEA (ng/ml)				
Median (Range)	23.5 (0.5-8048)		30.5 (1.5-1819)	
CA125 (ng/ml)				
Median(Range)	52.9 (4.76-3369)		69.4 (11.3-954.5)	
NSE(ng/ml)				
Median (Range)	15.0 (4.4-133.1)		15.2(7.6-55.2)	
First-line treatment regimen				
EGFR-TKI treatment	112	88.2	21	70
Chemotherapy	15	11.8	9	30
Type of EGFR mutations				
L858R	49	38.6	14	46.7
19 deletion	67	52.8	11	36.7
Other	11	8.6	5	16.7
NO. of extracranial metastases				
0	8	6.3	2	6.7
1	65	51.2	14	46.7
2	42	33.1	10	33.3
3 or more	12	9.4	4	13.3
Clinical stages				
Stage IIIB	8	6.3	2	6.7
Stage IV	119	93.7	28	93.3
Location of extracranial metastatic sites				
Pleural effusion	8	6.3	6	20.0
Liver	17	13.4	4	13.3
Adrenal	17	13.4	1	3.3
Bone	73	57.5	18	60
Lung	75	59.1	17	56.6
Other	12	9.4	2	6.7
Types of EGFR-TKIs				
Gefitinib	80	63.0	19	63.3
Erlotinib	31	24.4	7	23.3
Icotinib	16	12.6	4	13.4
Local therapy for BM				

None	NA	8	26.7
WBRT	NA	14	46.7
SRS	NA	8	26.6

Abbreviation: BM, brain metastasis; KPS, Karnofsky Performance Status; CI, confidence interval; EGFR, epidermal growth factor receptor; WBRT, whole brain radiation therapy; SRS, stereotactic radiosurgery; NSE, neuron-specific enolase; TKI, tyrosine kinase inhibitor.

Table 2: Univariate and multivariate analyses for the factors associated with risks for metachronous BM

Factors	Univariate analysis Incidence of BM (%)			Multivariate analysis Incidence of BM (%)		
	HR	95%CI	P	HR	95%CI	P
Gender: female VS male	1.139	0.556-2.337	0.772	1.495	0.506-4.421	0.467
Age, years	0.963	0.931-0.995	0.023			
> 49 VS ≤ 49	0.341	0.162-0.720	0.005	0.396	0.167-0.938	0.035
KPS score: <80 VS ≥80	0.045	0.000-40.173	0.371			
BMI	1.035	0.922-1.161	0.562	1.057	0.919-1.216	0.436
Smoking: yes VS no	0.798	0.353-1.801	0.586	1.302	0.384-4.408	0.672
Tumor markers level before treatment						
CEA (ng/ml)	1.000	0.999-1.000	0.685	1.000	0.999-1.000	0.294
CA125 (ng/ml)	1.000	0.998-1.001	0.498			
NSE (ng/ml)	1.014	0.985-1.043	0.351			
First-line treatment regimen						
Chemotherapy VS EGFR-TKI	2.296	1.050-5.018	0.037	0.504	0.153-1.660	0.260
Type of EGFR mutations			0.071			0.061
19-del VS L858R	0.579	0.263-1.277	0.176	0.490	0.201-1.194	0.116
Other VS L858R	1.968	0.703-5.508	1.968	2.408	0.566-10.246	0.234
Clinical stages: IIIB VS. IV	0.501	0.152-1.653	0.257			
Type of EGFR-TKIs			0.262			
Erlotinib VS Gefitinib	0.422	0.118-1.503	0.183			
Icotinib VS Gefitinib	0.460	0.109-1.946	0.292			
NO. of extracranial metastasis						
0-2 VS 3 or more	0.523	0.181-1.514	0.232	0.200	0.056-0.713	0.013
Location of extracranial metastasis						
Pleural effusion	3.245	1.300-8.098	0.012	5.283	1.854-15.053	0.002
Liver	1.066	0.371-3.062	0.906			
Adrenal	0.242	0.033-1.779	0.163			
Bone	1.161	0.558-2.413	0.690			
Lung	1.543	0.685-3.475	0.295			
Other	1.332	0.317-5.605	0.696			

Abbreviation: BM, brain metastasis; KPS, Karnofsky Performance Status; CI, confidence interval; EGFR, epidermal growth factor receptor; WBRT, whole brain radiation therapy; SRS, stereotactic radiosurgery; NSE, neuron-specific enolase; TKI, tyrosine kinase inhibitor.

Supplementary Figure Legend

Figure S1: Comparison of the actuarial risk of developing metachronous BM among randomly select 52 cases from our patient cohort grouped by different numbers of risk factors.

Figures




Figure 1

Flow chart for patients' enrollment and treatment. *Follow-up examinations were performed every 2 months, including thoracic and abdominal CT scan, brain MRI scan.




Figure 2

Kaplan-Meier plot of OS, PFS, and BMFS in EGFR-mutated advanced NSCLC patients without BM at initial diagnosis. OS, overall survival; PFS, progression-free survival; BMFS, brain-metastasis-free survival; NSCLC, non-small cell lung cancer; BM, brain-metastases; CI, confidence interval.




Figure 3

Kaplan-Meier plot of OS in patients with EGFR-mutated advanced NSCLC grouped on BM status. OS, overall survival; NSCLC, non-small cell lung cancer; BM, brain-metastases; CI, confidence interval.




Figure 4

Comparison of the actuarial risk of developing metachronous BM among patients with different numbers of risk factors.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [STROBEchecklistv4cohort.pdf](#)
- [FigS1.pdf](#)