Role of Human Papillomavirus and Its Association With an Indian OSCC Subjects

Amisha Patel

RK University

Vijaykumar Gupta

: Indian Cancer Society

Sejal Shah (✉ sejal.shah@rku.ac.in)

RK University <https://orcid.org/0000-0002-7664-4086>

Research Article

Keywords: HPV, oral cancer, tissue, India, OSCC

Posted Date: August 26th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-829604/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Oral Cancer; especially Oral Squamous Cell Carcinoma (OSCC) is the eighth most common cancer all over the globe and most common neoplasm in India. Due to the tobacco addiction pattern in Indians specially Gujarati's are well known for high consumption of tobacco in non-smoking form may have the high-risk for developing oral cancer. Apart of this, Human papilloma virus(HPV) might be one of the cause for developing oral cancer hence, the proposed study was carried out for 50 primary OSCC subjects from January 2018 to October 2018. Tumour specimens were HPV-genotyped by type specific PCR method. Statistical analyses were performed by Fisher's exact test. The present study identified 4% HPV infection frequency in advanced staged OSCC Indian subset, where One OSCC stage IV patient observed with HPV 16 infection and one OSCC stage III patient with HPV 18. Our finding differs with rest of the population, as the region have high amount of consumption of tobacco. Current study need to be validated in larger sample size to get the clear view for association of HPV with oral cancer.

Introduction

Oral cancer; the eighth most common neoplasm and blazing dilemma over the globe where 90 % cases are for oral squamous cell carcinoma [1-5]. India has highest incidence and mortality rate throughout the world, mainly males are prominently affected for the oral cancer [6,7]. Average 177,384 mortality occur from 354,864 incidents annually with 913,514 prevalence [8].

High consumption of tobacco in smoking and non-smoking form is the major risk factor for onset of oral cancer end up with 50-fold increase in reactive oxygen species. Gujarat is the hub for tobacco productions leads to high consumption of tobacco in form of Gutkha and mava headed to development of oral cancer [9]. Our previous study has shown most of the patients having genetic alterations found in the patients with the history of tobacco, which support the notion and pointed out the association of tobacco with the development of oral cancer [10]. However, approximately 20% of oral cancers developed in patients lacking these traditional risk factors [11-12]. Apart from tobacco there are several risk factor for developing oral cancer. There is evidence for HPV infected people having the history of tobacco and alcohol have developed oral cancer [13]. Human papilloma virus is characteristic member of papillomaviridae family of small non-enveloped virus that contain double-stranded DNA viral genome of ~8 kb [14-17]. Approximately 120 types are entirely sequenced among the 150 different types of HPV [18]. Previous study suggested that more than 10% of human cancers are developed by HPV infection [19]. There are 24 different types of HPV have been categorised as high risk types that associated with benign lesions of OSCC including 1, 2, 3, 4, 6, 7, 10, 11, 13, 16, 18, 30, 31, 32, 33, 35, 45, 52, 55, 57, 59, 69, 72, and 73, whereas 12 types with malignant OSCC lesions including 2, 3, 6, 11, 13, 16, 18, 31, 33, 35, 52, and 57 [20]. The HPV type 16 if the most leading type that detected in OSCC, followed by HPV 18 [21-23]. The younger patients are affected three times more with the high-risk HPV types than elders, due to viral transmission by direct physical contact [24]. In India, there are no more data which evaluating the prevalence of HPV in OSCC. HPV infection status can become an ordinary element in the diagnosis of

oral cancer. The aim of this study was to determine the association between HPV and OSCC in Saurashtra region of Gujarat.

Materials And Methods

Study design and population

The current study was approved from the Institutional Ethics committee (Human) of P.D.U. Medical College, Rajkot, Gujarat. **All the experiments and methods were performed in accordance with the relevant guidelines and regulations.** Each and every patient provided written informed consent and were personally interviewed in proportion for the structure questionnaire. 50 primary OSCC tumors were obtained from January 2018 to October 2018 during the surgery at Shree Nathalal Parekh Cancer Institute and Shree Gulabchand Talakchand Sheth Cancer Hospital, Rajkot. All the samples were histopathologically confirmed and collected at the time of surgery from the primary site of tumor. All the tissue samples were washed thrice with 1X phosphate-buffered saline (pH 7.2) to remove contamination of blood cells from the samples. The washed samples were stored in RNA later® (Qiagen) at -20 °C for further downstream analysis.

DNA isolation

Each tissue sample were micro dissected, and the genomic DNA was isolated from the tumor tissue using a DNeasy® Blood and tissue mini kit (Qiagen) by following the manufacturer's tissue protocol. The quantity of DNA was analysed by measuring the UV absorption at 260nm and 280nm and quality were analysed by 0.5% agarose gel electrophoresis.

Detection of HPV-16 and HPV-18

The present study was performed by Type-specific PCR (TSPCR) using HPV-16 and 18 primers (Table 1) and amplification was performed on thermal cycler (Biorad T100™). Total volume of 25 µl of PCR mixture containing 1X PCR Master Mix (Thermoscientific #K0171), 100 ng DNA, 10 pmole of each forward and reverse primers and nuclease free water. The PCR condition of HPV-16 amplification was an initial denaturation at 94°C for 3 minutes, followed by 34 cycles each at 94°C for 1 minute, annealing at 52.1°C for 30 seconds and extension at 72°C for 30 seconds which was extended for 5 min at the final cycle. HPV-18 detection was performed with initial denaturation at 94°C for 3 minutes, followed by 34 cycles each at 94°C for 1 minute, annealing at 47.5°C for 1 minute and extension at 72°C for 1 minute. At the final cycle extension was extended for 10 minutes. β-globin gene was used as internal positive control.

Statistical analysis

The presence of reported HPV is indicated as a percentage of the total number of samples screened. Fisher's exact test was used to check correlations between HPV infection and clinicopathological

characteristics of the patients. The p value less than 0.05 were defined as being statistically significant. Two tailed fisher exact test was performed to check association with OSCC risk.

Results

The current study recruited 50 oral cancer tumor tissues for HPV detection. The age of the OSCC patients was ranged from 20 to 80 year (mean age 47.82). Among fifty OSCC patients thirty-eight (76%) were male and twelve (24%) were female. Twenty-four (48%) patients had a history of chewing tobacco and five (10%) patients had a history of alcohol consumption while only two (4%) patient had a history of smoking. Among fifty patients six (12%) patients were habituated with tobacco and smoking while only two (4%) patient was habituated with tobacco and alcohol. Different frequency of OSCC patients as described in (Table 1).

HPV infection was detected by type specific PCR method. The DNA used for the PCR amplification was checked quantitatively and qualitatively by UV spectrophotometer and 0.5% agarose gel electrophoresis, respectively. The DNA samples have 260nm/280nm ratio >1.8 were used for the type specific PCR for HPV 16 and HPV 18. The primers used for the type specific PCR have shown in (table 2). B-globin gene of 268 bp was used as a positive internal control. Bands of 217 bp and 100 bp of HPV-16 and HPV-18 respectively were checked on 2% agarose gel electrophoresis.

The present study identified two sample with HPV infection. One patients identified with HPV-16 history was at advanced stage of oral carcinoma. Another patient was observed with HPV -18 was also at advanced stage of carcinoma. Both the patients neither shown tobacco history nor genetic history in their lifetime, it clearly indicates HPV infection might be the reason for the development of oral carcinogenesis. Both samples were replicated using type specific PCR and confirmed on 2% agarose gel electrophoresis (Fig 1). The current study observed 4% HPV infection frequency in the Gujarati population.

The current study was performed for statistical analysis using fisher exact test to check whether HPV infection has any significant association with any of the clinicopathological characteristics of the OSCC patients or not. The study was not found any statistical significance with Age, Gender, tumor site and disease stage (Table 3).

Discussion

The current study identified 4% HPV infection in the Gujarati oral squamous cell carcinoma subset. We screened total of 50 patients using type specific PCR. Out of fifty patients one patient with advanced stage carcinoma, without any carcinogenic history found infected with HPV-16 and another patient with stage IV, without any personal, genetic or tobacco history observed infected with HPV-18.

The same study has been done in same population recently; they found 2% HPV frequency in oral cancer subset [25]. They have enrolled 200 cancer samples and found 2 HPV infected patients. They did this study in year 2010, so it may be the difference in frequency which support the notion of HPV infection, as

this is the highly tobacco consumed region, the development of OSCC was observed due to tobacco, a rare history of HPV found in both the studies. Another previous study reported in Gujarati population stated absence of HPV infection [12] (Fig 2). It seems, as this is the tobacco growing belt and tobacco consuming region, the frequency of HPV infection is very low compared to another population of India. The main reason of onset of oral cancer is tobacco. The previous study reported 80% of the OSCC patients had history of tobacco in form of chewing and smoking [7]. Hence, it might be the reason of lower frequency of the HPV infection.

Same study in Eastern India [26,27] reported highest (51.3%) prevalence of HPV infection as compared to south India (48.63%) [14,28-32], Followed by North India (35.2%) [33-36]. West India [12,25,37-39] accounted 15.54% frequency of HPV infection and present study in Western India stated only 6.6% prevalence of HPV infection (Fig 3). The western Indian population is showing lower HPV frequency compare to rest of the Indian region, tobacco history might be the main factor for the OSCC carcinogenesis over Gujarati population.

The frequency of HPV infection varies from ethnicity to ethnicity. When we have compared the HPV infection frequency of Western part of the India with other parts of the globe, it is showing contrast result (Fig 4). The difference in results may be due to race, ethnicity, etiological factor, dietary patterns, environmental factors, personal history such as genetic history and mostly tobacco history do affect the results. HPV infection found more frequent in Japan (93.89%) [40-46] then Greece (65.67%) [16,47]. The similar study performed in Venezuela [48-50] and Netherland [51,52] demonstrated 62.93% and 31.27% prevalence of HPV infection respectively. The study in USA [24,53-59], China [60-66] and Brazil [67-69] has shown 22.95%, 22.83% and 23.8% HPV prevalence respectively. Total HPV mutation frequency of Indian population is showing 36.04% [12,14,25-39] which seems similar to Netherland population.

The HPV screening must require before any genetic analysis of the cancer patient. This is the personalized medicine era, based on the patient's genomic profile; the prognosis of the patients can be improved [70]. The patients with HPV infection has better prognosis than any other personal history. Thus, HPV studies must have been carried out in line with all the basic test of the cancer patients. The limitation of the study is lower sample size. The further study with larger cohort should be needed to validate the findings and to establish HPV as a prognostic biomarker.

The present study identified 4% HPV infection frequency in advanced staged OSCC Indian subset. Both the patients were not observed with any personal history. Our finding differs with rest of the population, as the region have high amount of consumption of tobacco, so that might be the reason for developing oral cancer. Another reason race, ethnicity, environmental factor, etiology, dietary factor may affect the results. The future studies with large cohort must be carried out to establish HPV infection has a better prognosis then patients have history of tobacco. The further study would be needed to establish the findings that will open the new vista in cancer biology.

Within the limitation of study, the Saurashtra region of Gujarat is not HPV prone region for onset of OSCC. Tobacco is the prime factor responsible for the development of OSCC. HPV infected patients have shown better prognosis, compared with non-HPV oral cancer patients. Thus, before medication, HPV screening must be done as it shows promising prognostic biomarker.

Declarations

Funding: Not applicable

Competing interest: The authors declare no competing interest.

Availability of data and material: Not applicable

Code availability: Not applicable

Author contributions: The conception and design of study, manuscript editing and polishing by SS, acquisition of data, analysis, interpretation and manuscript drafting by AP and VG has provided oral cancer tumor-control tissues, biopsy reports and all the necessary histopathological analysis.

Ethical approval: The present study is ethically approved by the Institutional Ethics committee (Human) of P.D.U. Medical College, Rajkot, India.

Consent to participate: Written informed consent were taken from each and every patient through personal interview in proportion for the structure questionnaire.

Consent for publication: Authors are giving consent to publish this work in Molecular Biology Reports journal.

Acknowledgement

We thank RK University for financial support and the staff of Shree Nathalal Parekh Cancer Institute, Rajkot for providing tissue samples.

References

1. Warnakulasuriya S (2009) Causes of oral cancer—an appraisal of controversies. Brit Dent J 207:471-475. <https://doi.org/10.1038/sj.bdj.2009.1009>
2. Anantharaman D, Chaubal PM, Kannan S, Bhisey RA and Mahimkar MB (2007) Susceptibility to oral cancer by genetic polymorphisms at CYP1A1, GSTM1 and GSTT1 loci among Indians: tobacco exposure as a risk modulator. Carcinogenesis 28:1455-1462. <https://doi.org/10.1093/carcin/bgm038>
3. Goot-Heah K, Kwai-Lin T, Froemming GRA, Abraham MT, Rosdy NMMNM and Zain RB (2012) Human papilloma virus 18 detection in oral squamous cell carcinoma and potentially malignant lesions

- using saliva samples. *Asian-Pac J Cancer Prev* 13:6109-6113. <https://doi.org/10.7314/APJCP.2012.13.12.6109>
4. Cooper JS, Porter K, Mallin K, Hoffman HT, Weber RS, Ang KK, Gay EG, Langer CJ (2009) National Cancer Database report on cancer of the head and neck: 10-year update. *Head Neck-J Sci Spec* 31:748-758. <https://doi.org/10.1002/hed.21022>
 5. Of the International IP, Maitra A, Biswas NK, Amin K, Kowtal P, Kumar S, Das S, Sarin R, Majumder PP, Bagchi I, Bairagya BB (2013) Mutational landscape of gingivo-buccal oral squamous cell carcinoma reveals new recurrently-mutated genes and molecular subgroups. *Nat Commun* 2:2873. <https://doi.org/10.1038/ncomms3873>
 6. GLOBOCAN 2020. International Agency for research on Cancer. <https://gco.iarc.fr>
 7. Shah S, Shah S, Padh H, Kalia K (2015) Genetic alterations of the PIK3CA oncogene in human oral squamous cell carcinoma in an Indian population. *Or Surg Or Med Or Pa* 120628-635. <https://doi.org/10.1016/j.oooo.2015.08.003>
 8. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A (2018) Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*. 2018 Nov;68(6):394-424. <https://doi.org/10.3322/caac.21492>
 9. Patel A, Gupta V, Pandya M, Shah S. (2021) Metabolomic Profiling of an Indian Oral Squamous Cell Carcinoma Subset. *Analytical Chemistry Letters*, 11(4), 596-606. <https://doi.org/10.1080/22297928.2021.1936168>
 10. Shah S, Jajal D, Mishra G, Kalia K (2017) Genetic profile of PTEN gene in Indian oral squamous cell carcinoma primary tumors. *J Oral Pathol Med* 46:106-111. <https://doi.org/10.1111/jop.12468>
 11. Nair UJ, Nair J, Mathew B, Bartsch H (1999) Glutathione S-transferase M1 and T1 null genotypes as risk factors for oral leukoplakia in ethnic Indian betel quid/tobacco chewers. *Carcinogenesis* 20:743-748. <https://doi.org/10.1093/carcin/20.5.743>
 12. Patel KR, Vajaria BN, Begum R, Desai A, Patel JB, Shah FD, Shukla SN, Patel PS (2014) Prevalence of high-risk human papillomavirus type 16 and 18 in oral and cervical cancers in population from Gujarat, West India. *J Oral Pathol Med* 43:293-297. <https://doi.org/10.1111/jop.12147>
 13. Campisi G, Giovannelli L, Calvino F, Matranga D, Colella G, Di Liberto C, Capra G, Leao JC, Muzio LL, Capogreco M, D'Angelo M (2006) HPV infection in relation to OSCC histological grading and TNM stage. Evaluation by traditional statistics and fuzzy logic model. *Oral Oncol* 42:638-645. <https://doi.org/10.1016/j.oraloncology.2005.11.007>
 14. Bijina BR, Ahmed J, Shenoy N, Ongole R, Shenoy S, Baliga S (2016) Detection of human papilloma virus in potentially malignant and malignant lesions of the oral cavity and a study of associated risk factors. *S Asian J Cancer* 5:179. <https://doi.org/10.4103/2278-330X.195337>
 15. Hagensee ME, Yaegashi N, Galloway DA (1993) Self-assembly of human papillomavirus type 1 capsids by expression of the L1 protein alone or by coexpression of the L1 and L2 capsid proteins. *J Virol* 67:315-322.

16. Bouda M, Gorgoulis VG, Kastrinakis NG, Giannoudis A, Tsoi E, Danassi-Afentaki D, Foukas P, Kyroudi A, Laskaris G, Herrington CS, Kittas C (2000) "High risk" HPV types are frequently detected in potentially malignant and malignant oral lesions, but not in normal oral mucosa. *Modern Pathol* 13:644-653. <https://doi.org/10.1038/modpathol.3880113>
17. Datta S, Agarwal M, Chatterjee S, Gogoi HK, Veer V, Singh L (2015) Detection of human papillomavirus in women attending Pap cervical screening camp at a peripheral hospital of North-Eastern India. *Med J armed forces India* 71:182-185. <https://doi.org/10.1016/j.mjafi.2013.04.014>
18. Rautava J, Syrjänen S (2012) Biology of human papillomavirus infections in head and neck carcinogenesis. *Head neck pathol* 6:3-15. <https://doi.org/10.1007/s12105-012-0367-2>
19. Jain N, Singh V, Hedau S, Kumar S, Daga MK, Dewan R, Murthy NS, Husain SA, Das BC (2005) Infection of human papillomavirus type 18 and p53 codon 72 polymorphism in lung cancer patients from India. *Chest* 128:3999-4007. <https://doi.org/10.1378/chest.128.6.3999>
20. Kojima A, Maeda H, Sugita Y, Tanaka S, Kameyama Y (2002) Human papillomavirus type 38 infection in oral squamous cell carcinomas. *Oral Oncol* 38:591-596. [https://doi.org/10.1016/S1368-8375\(01\)00112-9](https://doi.org/10.1016/S1368-8375(01)00112-9)
21. Smith EM, Ritchie JM, Summersgill KF, Klussmann JP, Lee JH, Wang D, Haugen TH, Turek LP (2004) Age, sexual behavior and human papillomavirus infection in oral cavity and oropharyngeal cancers. *Int J Cancer* 108:766-772. <https://doi.org/10.1002/ijc.11633>
22. Herrero R, Castellsagué X, Pawlita M, Lissowska J, Kee F, Balaram P, Rajkumar T, Sridhar H, Rose B, Pintos J, Fernández L (2003) Human papillomavirus and oral cancer: The International Agency for Research on Cancer multicenter study. *J Natl Cancer Inst* 95:1772-1783. <https://doi.org/10.1093/jnci/djg107>
23. Turakhiya B, Patel A, Shah S (2018). Human Papillomavirus associated burden of Oral Squamous Cell Carcinoma in an Indian inhabitant. *Int J Sci Res rev.* **7**, 81-101 (2018).
24. Schwartz, S. M. Oral cancer risk in relation to sexual history and evidence of human papillomavirus infection. *J Natl Cancer Inst.* **90**(21), 1626-1636 (1998).
25. Parmar MM, Pandhi N, Patel D, Gupta D (2017) Prevalence of High-Risk Human Papilloma Virus (HR-HPV) As a Risk Factor in HNSCC Patients of Saurashtra Region of Gujarat. *Int J Health Science Res* 11:63-69.
26. Nagpal JK, Patnaik S, Das BR (2002) Prevalence of high-risk human papilloma virus types and its association with P53 codon 72 polymorphism in tobacco addicted oral squamous cell carcinoma (oscc) patients of Eastern India. *Int J Cancer* 97:649-653. <https://doi.org/10.1002/ijc.10112>
27. Chatterjee R, Mukhopadhyay D, Chakraborty RN, Mitra RB (1997) Evaluation of argyrophilic nucleolar organizer regions (AgNORs) in oral carcinomas in relation to human papillomavirus infection and cytokinetics. *J oral Pathol Med* 26:310-314. <https://doi.org/10.1111/j.1600-0714.1997.tb00221.x>
28. Balaram P, Nalinakumar KR, Abraham E, Balan A, Hareendran NK, Bernard HU, Chan SY (1995) Human papillomaviruses in 91 oral cancers from Indian betel quid chewers—high

- prevalence and multiplicity of infections. *Int J cancer* 61:450-454.<https://doi.org/10.1002/ijc.2910610403>
29. Pillai MR, Phanidhara A, Kesari AL, Nair P, Nair MK (1999) Cellular manifestations of human papillomavirus infection in the oral mucosa. *J Surg Oncol* 71:10-15.[https://doi.org/10.1002/\(SICI\)1096-9098\(199905\)71:1<10::AID-JSO3>3.0.CO;2-V](https://doi.org/10.1002/(SICI)1096-9098(199905)71:1<10::AID-JSO3>3.0.CO;2-V)
30. Kumar RV, Kadkol SS, Daniel R, Shenoy AM, Shah KV (2003) Human papillomavirus, p53 and cyclin D1 expression in oropharyngeal carcinoma. *Int J Oral Max Surg* 32:539-543.[https://doi.org/10.1016/S0901-5027\(03\)90407-8](https://doi.org/10.1016/S0901-5027(03)90407-8)
31. Kulkarni SS, Kulkarni SS, Vastrad PP, Kulkarni BB, Markande AR, Kadakol GS, Hiremath SV, Kaliwal S, Patil BR, Gai PB (2011) Prevalence and distribution of high risk human papillomavirus (HPV) types 16 and 18 in carcinoma of cervix, saliva of patients with oral squamous cell carcinoma and in the general population in Karnataka, India. *Asian Pac J Cancer Prev* 12:645-648.
32. Elango KJ, Suresh A, Subhadradevi L, Ravindran HK, Iyer SK, Iyer SK, Kuriakose MA (2011) Role of human papilloma virus in oral tongue squamous cell carcinoma. *Asian Pac J Cancer Prev* 12:889-896.
33. Bahl A, Kumar P, Dar L, Mohanti BK, Sharma A, Thakar A, Karthikeyan V, Sikka K, Singh C, Poo K, Lodha J (2014) Prevalence and trends of human papillomavirus in oropharyngeal cancer in a predominantly north Indian population. *Head Neck* 36:505-510.<https://doi.org/10.1002/hed.23317>
34. Khanna R, Rao GR, Tiwary SK, Rai A, Khanna S, Khanna AK (2009) Detection of human papilloma virus 16 and 18 DNA sequences by southern blot hybridization in oral leukoplakia and squamous cell carcinoma. *Indian J Surg* 71:69-72. <https://doi.org/10.1007/s12262-009-0019-2>
35. Katiyar S, Thelma BK, Murthy NS, Hedau S, Jain N, Gopalkrishna V, Husain SA, Das BC (2003) Polymorphism of the p53 codon 72 Arg/Pro and the risk of HPV type 16/18-associated cervical and oral cancer in India. *Mol Cell Biochem* 252:117-124.<https://doi.org/10.1023/A:1025546610920>
36. Jalouli J, Ibrahim SO, Mehrotra R, Jalouli MM, Sapkota D, Larsson PA, Hirsch JM (2010) Prevalence of viral (HPV, EBV, HSV) infections in oral submucous fibrosis and oral cancer from India. *Acta Otolaryngol* 130:1306-1311. <https://doi.org/10.3109/00016481003782041>
37. D'Costa J, Saranath D, Dedhia P, Sanghvi V, Mehta AR (1998) Detection of HPV-16 genome in human oral cancers and potentially malignant lesions from India. *Oral Oncol* 34:413-420.[https://doi.org/10.1016/S1368-8375\(98\)00028-1](https://doi.org/10.1016/S1368-8375(98)00028-1)
38. Koppikar P, deVilliers EM, Mulherkar R (2005) Identification of human papillomaviruses in tumors of the oral cavity in an Indian community. *Int J Cancer* 113:946-950.<https://doi.org/10.1002/ijc.20664>
39. Gheit T, Vaccarella S, Schmitt M, Pawlita M, Franceschi S, Sankaranarayanan R, Sylla BS, Tommasino M, Gangane N (2009) Prevalence of human papillomavirus types in cervical and oral cancers in central India. *Vaccine* 27:636-639. <https://doi.org/10.1016/j.vaccine.2008.11.041>
40. Sugiyama M, Bhawal UK, Kawamura M, Ishioka Y, Shigeishi H, Higashikawa K, Kamata N (2007) Human papillomavirus-16 in oral squamous cell carcinoma: clinical correlates and 5-year survival. *Br*

- J Oral Maxillofac Surg 45:116-122. <https://doi.org/10.1016/j.bjoms.2006.04.012>
41. Koyama K, Uobe K, Tanaka A (2007) Highly sensitive detection of HPV-DNA in paraffin sections of human oral carcinomas. J Oral Pathol Med 36:18-24.<https://doi.org/10.1111/j.1600-0714.2006.00490.x>
42. Zhang ZY, Sdek P, Cao J, Chen WT (2004) Human papillomavirus type 16 and 18 DNA in oral squamous cell carcinoma and normal mucosa. Int J Oral Maxillofac Surg 33:71-74.<https://doi.org/10.1054/ijom.2002.0443>
43. Sugiyama M, Bhawal UK, Dohmen T, Ono S, Miyauchi M, Ishikawa T (2003) Detection of human papillomavirus-16 and HPV-18 DNA in normal, dysplastic, and malignant oral epithelium. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 95:594-600. <https://doi.org/10.1067/moe.2003.36>
44. Chang JY, Lin MC, Chiang CP (2003) High-Risk Human Papillomaviruses May Have an Important Role in Non-Oral Habits-Associated Oral Squamous Cell Carcinomas in Taiwan. Am J Clin Pathol 120:909-916. <https://doi.org/10.1309/C5P6NUQ2NW6LCTBP>
45. Tsuchioka K, Nakazato I, Miyagi J, Iwamasa T, Arasaki A, Hiratsuka H, Sunakawa H, Kohama G, Abo T (2000) Comparative study of oral squamous cell carcinoma in Okinawa, Southern Japan and Sapporo in Hokkaido, Northern Japan; with special reference to human papillomavirus and Epstein-Barr virus infection. J Oral Pathol Med 29:70-79.<https://doi.org/10.1034/j.1600-0714.2000.290204.x>
46. Shindoh M, Chiba I, Yasuda M, Saito T, Funaoka K, Kohgo T, Amemiya A, Sawada Y, Fujinaga K (1995) Detection of human papillomavirus DNA sequences in oral squamous cell carcinomas and their relation to p53 and proliferating cell nuclear antigen expression. Cancer 76:1513-1521.[https://doi.org/10.1002/1097-0142\(19951101\)76:9<1513::AID-CNCR2820760903>3.0.CO;2-4](https://doi.org/10.1002/1097-0142(19951101)76:9<1513::AID-CNCR2820760903>3.0.CO;2-4)
47. Aggelopoulou EP, Skarlos DI, Papadimitriou CO, Kittas CH, TROUNGOS CO (1999) The Human Papilloma Virus DNA Detection in Oral Lesions in the Greek Population. Anticancer Res. 19:1391-1396.
48. Correnti M, Rivera H, Cavazza ME (2004) Detection of human papillomaviruses of high oncogenic potential in oral squamous cell carcinoma in a Venezuelan population. Oral Dis 10:163-166. <https://doi.org/10.1046/j.1601-0825.2003.00989.x>
49. Premoli-De-Percoco G, Ramirez JL (2001) High risk human papillomavirus in oral squamous carcinoma: evidence of risk factors in a Venezuelan rural population. Preliminary report. J Oral Pathol Med 30:355-61. <https://doi.org/10.1034/j.1600-0714.2001.300605.x>
50. Premoli-De-Percoco G, Ramirez JL, Galindo I (1998) Correlation between HPV types associated with oral squamous cell carcinoma and cervicovaginal cytology: An in situ hybridization study. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 86:77-81.[https://doi.org/10.1016/S1079-2104\(98\)90153-6](https://doi.org/10.1016/S1079-2104(98)90153-6)
51. Cruz IB, Snijders PJ, Steenbergen RD, Meijer CJ, Snow GB, Walboomers JM, Van der Waal I (1996) Age-dependence of human papillomavirus DNA presence in oral squamous cell carcinomas. Eur J Cancer B Oral Oncol 32:55-62. [https://doi.org/10.1016/0964-1955\(95\)00060-7](https://doi.org/10.1016/0964-1955(95)00060-7)
52. Melchers LJ, Mastik MF, Cameron BS, Van Dijk BA, De Bock GH, Van Der Laan BF, Van Der Vegt B, Speel EJ, Roodenburg JL, Witjes MJ, Schuurin E (2015) Detection of HPV-associated oropharyngeal

- tumours in a 16-year cohort: more than meets the eye. *Br J Cancer* 112:1349-1357.<https://doi.org/10.1038/bjc.2015.99>
53. Lingen MW, Xiao W, Schmitt A, Jiang B, Pickard R, Kreinbrink P, Perez-Ordóñez B, Jordan RC, Gillison ML (2013) Low etiologic fraction for high-risk human papillomavirus in oral cavity squamous cell carcinomas. *Oral Oncol* 49:1-8. <https://doi.org/10.1016/j.oraloncology.2012.07.002>
 54. El-Mofty SK, Patil S (2006) Human papillomavirus (HPV)-related oropharyngeal nonkeratinizing squamous cell carcinoma: characterization of a distinct phenotype. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 101:339-345. <https://doi.org/10.1016/j.tripleo.2005.08.001>
 55. Ritchie JM, Smith EM, Summersgill KF, Hoffman HT, Wang D, Klusmann JP, Turek LP, Haugen TH (2003) Human papillomavirus infection as a prognostic factor in carcinomas of the oral cavity and oropharynx. *Int J Cancer* 104:336-344. <https://doi.org/10.1002/ijc.10960>
 56. Schwartz SR, Yueh B, McDougall JK, Daling JR, Schwartz SM (2001) Human papillomavirus infection and survival in oral squamous cell cancer: a population-based study. *Otolaryngol Head Neck Surg* 125:1-9 <https://doi.org/10.1067/mhn.2001.116979>
 57. Mao EJ, Schwartz SM, Daling JR, Oda D, Tickman L, Beckmann AM (1996) Human papilloma viruses and p53 mutations in normal, pre-malignant and malignant oral epithelia. *Int J Cancer* 69:152-158. [https://doi.org/10.1002/\(SICI\)1097-0215\(19960422\)69:2<152::AID-IJC15>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1097-0215(19960422)69:2<152::AID-IJC15>3.0.CO;2-B)
 58. Holladay EB, Gerald WL (1993) Viral gene detection in oral neoplasms using the polymerase chain reaction. *Am J Clin Pathol* 100:36-40. <https://doi.org/10.1093/ajcp/100.1.36>
 59. Greer Jr RO, Eversole LR, Crosby LK (1990) Detection of human papillomavirus-genomic DNA in oral epithelial dysplasias, oral smokeless tobacco-associated leukoplakias, and epithelial malignancies. *J Oral Maxillofac Surg* 48:1201-1205. [https://doi.org/10.1016/0278-2391\(90\)90538-D](https://doi.org/10.1016/0278-2391(90)90538-D)
 60. Zhao D, Xu QG, Chen XM, Fan MW (2009) Human papillomavirus as an independent predictor in oral squamous cell cancer. *Int J Oral Sci* 1:119-125. <https://doi.org/10.4248/IJOS.09015>
 61. Chen SF, Yu FS, Chang YC, Fu E, Nieh S, Lin YS (2012) Role of human papillomavirus infection in carcinogenesis of oral squamous cell carcinoma with evidences of prognostic association. *J Oral Pathol Med* 41:9-15. <https://doi.org/10.1111/j.1600-0714.2011.01046.x>
 62. Lee LA, Huang CG, Liao CT, Lee LY, Hsueh C, Chen TC, Lin CY, Fan KH, Wang HM, Huang SF, Chen IH (2012) Human papillomavirus-16 infection in advanced oral cavity cancer patients is related to an increased risk of distant metastases and poor survival. *PloS One* 7:e40767.<https://doi.org/10.1371/journal.pone.0040767>
 63. Huang SF, Li HF, Liao CT, Wang HM, Chen IH, Chang JC, Chen YJ, Cheng AJ (2012) Association of HPV infections with second primary tumors in early-staged oral cavity cancer. *Oral Dis* 18:809-815. <https://doi.org/10.1111/j.1601-0825.2012.01950.x>
 64. Lee LA, Huang CG, Tsao KC, Liao CT, Kang CJ, Chang KP, Huang SF, Chen IH, Fang TJ, Li HY, Yang SL (2013) Increasing rates of low-risk human papillomavirus infections in patients with oral cavity squamous cell carcinoma: association with clinical outcomes. *J Clin Virol* 57:331-337.<https://doi.org/10.1016/j.jcv.2013.04.010>

65. Gan LL, Zhang H, Guo JH, Fan MW (2014) Prevalence of human papillomavirus infection in oral squamous cell carcinoma: a case-control study in Wuhan, China. *Asian Pac J Cancer Prev* 15:5861-5865. <https://doi.org/10.7314/APJCP.2014.15.14.5861>
66. Huang CG, Lee LA, Tsao KC, Liao CT, Yang LY, Kang CJ, Chang KP, Huang SF, Chen IH, Yang SL, Lee LY (2014) Human papillomavirus 16/18 E7 viral loads predict distant metastasis in oral cavity squamous cell carcinoma. *J Clin Virol* 61:230-236. <https://doi.org/10.1016/j.jcv.2014.07.007>
67. Kaminagakura E, Villa LL, Andreoli MA, Sobrinho JS, Vartanian JG, Soares FA, Nishimoto IN, Rocha R, Kowalski LP (2012) High-risk human papillomavirus in oral squamous cell carcinoma of young patients. *Int J Cancer* 130:1726-1732. <https://doi.org/10.1002/ijc.26185>
68. Simonato LE, Garcia JF, Sundefeld ML, Mattar NJ, Veronese LA, Miyahara GI (2008) Detection of HPV in mouth floor squamous cell carcinoma and its correlation with clinicopathologic variables, risk factors and survival. *J Oral Pathol Med* 37:593-598. <https://doi.org/10.1111/j.1600-0714.2008.00704.x>
69. Gabrielli Fregonesi PA, Teresa DB, Duarte RA, Neto CB, de Oliveira MR, Soares CP (2003) p16INK4A immunohistochemical overexpression in premalignant and malignant oral lesions infected with human papillomavirus. *J Histochem Cytochem* 51:1291-1297. <https://doi.org/10.1177/002215540305101006>
70. Shah S, Mishra G, Kalia K (2018) Single nucleotide polymorphism rs17849071 G/T in the PIK3CA gene is inversely associated with oral cancer. *Oral Cancer*, 2:83-89. <https://doi.org/10.1007/s41548-018-0012-1>

Tables

Table 1. Clinicopathological parameters of the OSCC patients

Characteristics	Oral Cancer Patients (n=50), no (%)		
	Male (n=38) (76%)	Female (n=12) (24%)	Both (n=50) (100%)
Age Range (years)			
Mean age – 47.82			
20-39	10 (26.3%)	1 (8.3%)	11 (22%)
40-59	18 (47.3%)	9 (75%)	27 (54%)
60-79	10 (26.3%)	2 (16.6%)	12 (24%)
Site of primary tumor			
Buccal mucosa	28 (73.6%)	6 (50.0%)	34 (68%)
Tongue	3 (7.8%)	4 (33.3%)	7 (14%)
Lip	1 (2.6%)	1 (8.3%)	2 (4%)
GB complex	3 (7.8%)	0 (0%)	3 (6%)
Alveolus	1 (2.6%)	1 (8.3%)	2 (4%)
Hard palate	2 (5.2%)	0 (0%)	2 (4%)
Risk habit			
Tobacco chewing	19 (50.0%)	5 (41.6%)	24 (48%)
Alcohol consumption	4 (10.5%)	1 (8.3%)	5 (10%)
Smoking	2 (5.2%)	0 (0%)	2 (4%)
Tobacco + Smoking	6 (15.7%)	0 (0%)	6 (12%)
Tobacco + Alcohol	2 (5.2%)	0 (0%)	2 (4%)
Non-habited	5 (13.1%)	6 (50.0%)	11 (22%)
HPV genotype			
HPV-16	1 (2.6%)	0 (0%)	1 (2%)
HPV-18	0 (0%)	1 (8.3%)	1 (2%)

Table 2. Co-relation of HPV and clinicopathological characteristics of the OSCC patients

Characteristics		HPV infection	NO HPV infection	p value
Total number (%)		2 (4%)	48 (96%)	
Age (years) [mean]		50 (40-60)	47.82 (27-70)	
Gender	Male	1	37	p<0.5
	Female	1	11	
Tumor site	BM	1	33	p<0.5
	Others	1	15	
Disease stage	I-III	1	28	p<0.5
	IV	1	22	

Table 3. Sequences of Primers used for PCR amplification

HPV genotype	Primers	Tm [°C]
HPV-16	Forward primer – 5'-AAGGCCAACTAAATGTCAC-3'	52.4
	Reverse primer – 5'-CTGCTTTTATACTAACCGG-3'	
HPV-18	Forward primer – 5'-ACCTTAATGAAAAACCACGA-3'	54.3
	Reverse primer – 5'-CGTCGTTTAGAGTCGTTCTTG-3'	
β-globin	Forward primer – 5'-CAACTTCATCCACGTTCCACC-3'	48-58
	Reverse primer – 5'-GAAGAGCCAAGGACAGGTAC-3'	

Figures

Figure 1

The confirmation of observed HPV 16, and HPV 18 infection using type specific PCR

Figure 2

HPV infection frequency in Gujarati population

Figure 3

Frequency of HPV infection in an Indian population

Figure 4

Global HPV infection frequency