

Effects of training podiatrists to use imagery-based motivational interviewing to improve self-care for people with diabetes-related foot disease: A mixed-methods pilot study

Tracey Kaczmarek (✉ lyontracey241@gmail.com)

Queensland University of Technology <https://orcid.org/0000-0002-9661-153X>

Jaap Van Netten

Amsterdam University Medical Centres: Amsterdam Universitair Medische Centra

Peter Lazzarini

Queensland University of Technology IHBI: Queensland University of Technology Institute of Health and Biomedical Innovation

David Kavanagh

Queensland University of Technology Institute of Health and Biomedical Innovation

Research

Keywords: Diabetic foot, Diabetes, Treatment, Motivational Interviewing, Imagery, Communication, Training, Podiatry

Posted Date: October 1st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-84108/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on February 10th, 2021. See the published version at <https://doi.org/10.1186/s13047-021-00451-1>.

Abstract

Background Self-care in diabetic foot disease (DFD) is challenging and can contribute to poor outcomes. Motivational Interviewing (MI) engages people in self-care and integrating imagery may further improve its outcomes. No previous studies have trained podiatrists in using MI to address DFD self-care. This was the first study on training podiatrists to conduct imagery-based motivational interviewing (MI) when treating people with diabetes-related foot disease (DFD), and to examine impacts on MI related skills, job satisfaction and subjective experiences in a mixed-methods pilot study. Methods Eleven recruited podiatrists (Median age 35, 9 female) received two 4-hour training sessions and three received later mentoring. MI and imagery skills were rated using validated tools during two clinical sessions per participant at baseline, and 2- and 12-weeks post-training. Job satisfaction was assessed at baseline and 12 weeks. Semi-structured interviews at 12 weeks were analysed using the framework approach. Results Significant improvements over time ($p = .006-.044$) with substantial effect sizes ($\eta^2 = .50-.67$) were found in three of four global MI related communication skills and two of four MI behaviours. However, effects on these indices were not sustained to 12 weeks, and imagery was rarely used. Job satisfaction was high at baseline and unchanged at follow-up ($p = 0.34$, $\eta^2 = .100$). In qualitative interviews, MI training and skills were valued, but significant challenges in using MI when treating people with DFD were reported. Conclusion Training podiatrists in MI may have potential but more training, observation and mentoring appear needed to obtain sustained communication changes in practice.

Background

An estimated 451 million people worldwide were diagnosed with Type 2 Diabetes Mellitus (DM) in 2017, a number that is expected to become 693 million by 2045 (1). A major complication of diabetes is foot disease (DFD), which includes foot ulceration (2). Multiple pathophysiological factors including peripheral artery disease and peripheral neuropathy lead to foot ulceration (3). Around half of these ulcers become complicated by infection, often leading to hospitalisation, amputation and increased mortality (2). This all contributes to DFD being a leading cause of the global burden of disability (4, 5), yet, up to 90% of this burden may be prevented when evidence-based clinical care is optimised (6, 7).

Optimal evidence-based care of DFD often relies upon sustained engagement to recommended self-care (8), such as regularly changing ulcer dressings, monitoring for signs of infection and wearing offloading devices (9). Maintaining optimal self-care is extremely difficult for people with DFD to achieve (9–12). While education improves self-care knowledge and behaviour in the short-term, it has yet to show longer-term benefits (13). A traditional “physician-directed” style of education and communication by clinicians, where people are told to adhere to recommended behaviours (14, 15), has limited positive effects on longer-term outcomes and may undermine commitment to effective self-management if it elicits arguments in favour of the status quo (14). However, practitioners may help people with DFD engage with self-care and facilitate ulcer prevention and healing when they build collaborative relationships (10, 16, 17). One approach to establishing these relationships is motivational interviewing (MI) (18).

MI is an evidence-based approach that enhances motivation for functional behavioural change by helping people resolve ambivalence (16, 19). Its empathic and accepting manner provides a safe setting to consider behaviour change, while its agenda encourages exploration of the advantages and possibility of improved self-care (20). MI has been successfully used in behaviour change in healthcare, including prevention of Human Immunodeficiency Virus, modification of substance abuse and improved outcomes in diabetes (19, 21). However, not all studies report substantial and sustained success (e.g.22).

A recent modification and extension of MI involves incorporating mental imagery throughout sessions (23, 24). Mental imagery is more closely linked to emotional reactions than is verbal discussion, and the affect is central to the experience of being motivated (25). In Functional Imagery Training (FIT), participants are encouraged to use individually tailored mental imagery when considering the benefits of improved self-care, and when remembering past successes (23, 24). If they become committed to changing their behaviour, they create mental images about their plans, and rehearse those at home to

motivate and guide their self-management. An early form of this approach was tested in people with type 2 diabetes and showed substantial acceptability (26). A multi-session version has shown stronger outcomes than standard MI in dieting, exercise and weight reduction (27, 28).

Effective application of imagery-based MI requires that practitioners are sufficiently trained to apply it routinely with high fidelity in their work (19). MI skills can be acquired by nurses or general practitioners who care for people with diabetes, albeit to varying success (22, 29–33), depending in part on the length of MI training and availability of supervision (34). However, training podiatrists in using MI to address DFD self-care has not been investigated yet. Podiatrists are in a prime position to engage people with DFD in carrying out self-care, and training podiatrists in imagery-based MI could be a valuable addition to their clinical skills (35). An observational study of podiatrists working with people with DFD identified that 90% perceived their role in promoting self-care as vital, but only 25% reported use of “MI type” skills (36). Increasing this level represents an acute need.

Our aim was to train podiatrists to conduct imagery-based MI when treating people with DFD, and to evaluate effects on related skills, job satisfaction, training experiences and implementation attempts.

Methods

This mixed-methods pilot study involved podiatrists who primarily treated people with DFD. Its quantitative component comprised a single-group pre/post design, testing the effects of a training program on MI and FIT-related skills of participants at baseline, 2- and 12-weeks post-training, and self-reported job satisfaction at baseline and 12 weeks. The qualitative component comprised a semi-structured interview at 12-weeks post-training. The study received human ethical approval from the health service and university (HREC/2018/QPCH/45318).

Participants

Participants for this study were consenting podiatrists who were registered by the Australian Health Practitioner Regulation Agency and worked within government funded DFD Services in Brisbane, Australia; they primarily treated people with DFD (i.e. with a diagnosis of diabetes and a history of foot ulceration). Exclusion criteria included prior training in MI. Potential participants were informed about the study by recruitment emails and posters displayed in clinical rooms.

Participating podiatrists recruited people with DFD to assist with outcome assessment. These people also provided voluntary written consent to participate in recorded clinical sessions during which podiatrists’ skills in imagery-based MI were assessed.

Intervention

The training was planned in collaboration with the manager of the podiatry team, who gave the project strong support, linked training sessions to team meetings and actively participated in the training sessions. The intervention training program was jointly developed and delivered by two authors: a clinical psychology researcher (DK) and a senior podiatrist in the trained team (TK). DK is a co-developer of FIT and has led multiple controlled trials using MI or FIT (27, 37, 38). TK was trained in MI and FIT before study commencement and had experience applying these skills in her own clinical sessions, allowing her to incorporate detailed descriptions of her experience in training.

The program comprised two 4-hour training sessions in imagery-based MI, separated by 2 weeks. Training was delivered in groups of 5–6 participants. The first session included group discussion of issues participants had experienced in attempting to facilitate DFD-related self-care, and of advantages for morale of defining success in terms of delivering a favourable context for change, rather than reaching behavioural or clinical targets. It outlined core concepts of MI and provided video examples of MI, followed by group discussion of potential application in podiatry sessions. A written manual gave a script

for use in consultations. To maximise compatibility with existing sessions, trainers suggested incorporating elements of MI when assessing self-management, conducting physical treatment and collaboratively planning future self-care.

The second session described the use of imagery and home-practice in podiatry sessions, with demonstrations, roleplayed practice and feedback to build skills and confidence. Following feedback from participants, an abbreviated version of the manual was provided. Key elements of the manual were later converted into a flowchart placed in the clinic rooms, to prompt the application of skills in sessions.

All participants were offered follow-up peer support opportunities from TK to observe and feedback on consultation sessions or provide peer-mentoring via email, phone or face-to-face. Such feedback and advice were positive and practical, recognising incremental improvements and helping participants solve challenges in their application of MI related skills. Emails reminding participants about key elements of the intervention were sent 4 and 10 weeks after the workshop training.

Outcomes of interest

The outcomes of interest for the quantitative component included MI related skills, use of imagery and job satisfaction. MI related skills and imagery were assessed using audio recordings from two clinical consultation sessions by each podiatrist at baseline, and 2- and 12-weeks post-training. Sessions were selected opportunistically, based on availability of the participants and researcher. No data on characteristics of the people with DFD or outcomes of sessions were collected, since the focus was on podiatrists' conduct of sessions. If a friend or relative of the person with DFD were also present, they were asked to remain silent during the consultation. If they became involved, the session was omitted from data collection and another was used.

MI related skills were assessed using the Motivational Interviewing Treatment Integrity Tool (MITI) 4.2.1 (39). MITI has two components: global scores and behaviour counts. Four global characteristics (Cultivating Change Talk, Soften Sustain Talk, Partnership, Empathy) are each scored from 1 (low) to 5 (high). Three core MI adherent behaviours (Affirm, Seeking Collaboration, Emphasizing Autonomy) and two non-MI adherent behaviours (Persuade, Confront) are each counted. Counts of other relevant behaviours (Giving Information, Persuasion with Permission, Questions, Simple and Complex reflections) were used to guide the global scores. Some Questions and Information Giving were expected, since consultations had to include clinical assessment and instructions for foot care, and the frequency of these behaviours were downplayed in global scores. Consistent with MITI instructions, only the first 20 minutes of each session were assessed. Assessment of clinical sessions were undertaken independently by two authors (TK and DK), who reached consensus after discussion and replay of audio segments where required.

FIT-related skills during the same session segments were assessed using the Functional Imagery Training – Quality Coding (FIT-QC) 1.0 (40) which includes introduction to imagery, instruction, delivery, use and adaption of imagery, refinement of imagery quality and focus, reflection and promotion of home practice. Each component was scored from 1 (poor) to 5 (high). FIT skills were assessed by the same authors (TK and DK).

Job satisfaction at baseline and 12-weeks post-training was assessed using the Hoppock Job Satisfaction Scale (41), which has 4 items rated 1 to 7, with higher total scores denoting higher satisfaction. (41)

Individual face-to-face semi-structured interviews at 12 weeks post-training were undertaken by the primary author (TK), audio recorded and transcribed verbatim. The interview was designed to gain insight into the podiatry participant's experiences in training and their application of MI skills into clinical practice. Each primary question in Table A1 was augmented by non-specific questions to obtain additional responses.

Table 1
 Questions used in the Semi-Structured Interviews

Interviewer Question
Before the training, what aspects of working with your patients did you find most challenging?
Have you noticed any differences in your patient's responses since doing the training?
What was your experience of the training like?
What aspects of training seemed most useful for your practice?
What aspects of the training seemed least useful for your practice?
What aspects of the training were easiest to apply into clinical practice?
What particular aspects were more difficult to apply?
What challenges did you face in applying what you learned in the training?
What could we do to help you keep using your skills from the workshop routinely with your patients?

Analysis

Quantitative data were analysed using repeated measures general linear models on IBM SPSS v26, with *F* values using Pillai's Trace. The primary focus was the overall effect for time, but nonorthogonal contrasts examining changes from baseline to 2- and 12-weeks post-training were also examined. Because this was a small pilot, η^2 (the proportion of total variance accounted for by an effect) was the primary focus rather than statistical significance, although both are reported.

Qualitative interviews were analysed using the framework approach (42). After transcription of interviews, participants' responses were analysed using: 1) familiarisation (re-reading transcripts and field notes), 2) identification of the thematic framework (identifying key themes from data), 3) indexing (arranging transcripts by themes), 4) charting (creating a matrix with responses collated), and 5) mapping and interpretation (interpreting the data). Themes were identified by one author (TK) and checked and discussed by another (JvN) until consensus was reached.

Results

Of 14 eligible podiatrists who were approached, 11 consented to participate. Of the remaining three, one was on vacation, one did not express interest and one expressed interest but did not consent. Participants were aged between 29–43 years (median 35), nine (82%) were female, and years of DFD-treatment experience ranged from 2–17 years (median 11). Two training sessions were delivered to all participants. Five participants requested peer support following training and three of these received feedback about a clinical consultation session.

Three of the four global score ratings (Change Talk, Partnership and Empathy) and two of the four core behaviour counts (Affirm and Persuade) showed statistically significant improvements and substantial effect sizes ($\eta^2 = .50-.67$) across the three time points (Table A2). Contrasts comparing scores with those at baseline showed significant improvements on these six indices 2 weeks but not 12 weeks post-training.

None of the five other behaviour counts showed statistically significant improvements, but Simple Reflections, Complex Reflections and Giving Information each showed moderate effect sizes ($\eta^2 = .43-.48$). Only Giving Information showed a statistically significant improvement from baseline to 12 weeks. As anticipated, both Giving Information and Questions were frequent throughout the study.

No substantial use of FIT Skills was evident in recordings, with only one podiatrist attempting to use motivational imagery after the training. Accordingly, detailed results on the FIT-QC are not reported.

Ten podiatrists completed the Hoppock scale. Job satisfaction was high at both baseline (Mean = 19.4, SD = 3.2) and 12 weeks (Mean = 20.4, SD = 3.9), with no significant change over time ($F(1, 9) = 1.00$; $p = 0.34$, $\eta^2 = .100$).

Table 2
MI related skills at baseline, 2- and 12-weeks post-training

Variable	Overall Time Effect						Time Contrasts					
	Baseline	2 weeks post-training	12 weeks post-training	from Pillai's Trace			Baseline – 2 weeks			Baseline – 12 weeks		
				<i>F</i> (2,9)	<i>p</i>	η^2	<i>F</i> (1,10)	<i>p</i>	η^2	<i>F</i> (1,10)	<i>p</i>	η^2
M (SD)	M (SD)	M (SD)										
Global scores												
Change Talk	1.77 (0.79)	3.08 (1.28)	2.08 (1.08)	5.60	.026	.554	10.92	.008	.522	0.79	.395	.073
Soften Sustain	1.09 (1.46)	1.36 (1.64)	1.00 (1.32)	0.14	.874	.030	0.20	.665	.019	0.02	.883	.002
Partnership	2.21 (0.74)	3.08 (0.96)	2.36 (0.98)	9.30	.006	.674	6.78	.026	.404	0.19	.671	.019
Empathy	2.34 (1.18)	3.09 (1.04)	2.27 (1.03)	6.91	.015	.605	3.31	.099	.249	0.03	.859	.003
Focal MI behaviour counts												
Affirm	0.68 (0.93)	1.23 (1.08)	0.64 (0.81)	6.12	.021	.576	2.68	.133	.211	0.02	.905	.001
Seeking Collaboration	1.65 (1.68)	2.91 (2.33)	2.56 (2.12)	1.66	.243	.270	3.69	.084	.270	1.74	.216	.148
Emphasising Autonomy	0.05 (0.15)	0.36 (0.55)	0.36 (0.50)	2.00	.192	.307	3.06	.111	.234	4.22	.067	.297
Persuade*	1.36 (1.12)	0.41 (0.66)	2.36 (2.60)	4.52	.044	.501	4.51	.060	.311	1.43	.260	.125
Other behaviour counts												
Giving Information*	19.30 (8.55)	13.41 (5.13)	11.05 (6.04)	3.36	.081	.428	3.50	.091	.259	7.12	.024	.416
Persuade with Permission	0.00 (0.00)	0.14 (0.32)	0.14 (0.24)	2.43	.143	.351	1.96	.192	.164	3.75	.082	.273
Question	19.55 (7.92)	18.89 (5.25)	18.64 (5.80)	0.58	.944	.013	0.07	.798	.007	0.13	.731	.012
Simple Reflection	1.36 (1.23)	2.38 (1.70)	0.76 (0.50)	4.14	.053	.479	3.13	.107	.239	2.05	.183	.170
Complex Reflection	0.18 (0.34)	1.14 (1.19)	1.05 (1.65)	3.99	.057	.470	6.58	.028	.397	2.87	.121	.223
Note: *Lower Persuade or Giving Information scores indicate better MI adherence. The non-adherent behaviour 'Confront' was not seen in any recordings and therefore omitted from the table. M: Mean. SD: Standard Deviation.												

Semi-structured interviews were conducted with all 11 participants and lasted a median 5.5 minutes (Range = 5.1–10.4). Three main themes with 10 subthemes were identified from the interviews (Table A3).

Table 3
Main themes identified in semi-structured interviews

Main Theme	Subtheme 1	Subtheme 2	Subtheme 3	Subtheme 4
1. Clinical Issues	1.1 Challenging situations	1.2 Communication challenges		
2. Training Content	2.1 Overall training experience	2.2 Role-play	2.3 Imagery	2.4 Ongoing training and support
3. Training Outcomes	3.1 New communication skills	3.2 Increased patient engagement	3.3 Long-term application of MI skills	3.4 Appropriateness of MI

Main theme 1. Clinical issues

Subtheme 1.1 Challenging situations:

Before training, all podiatrists identified struggles with fostering self-care by people with DFD. These struggles included situations where podiatrists tried to achieve commitment to self-care.

“... a bit like that feeling like your hitting your head against a brick wall. That you know they know what they should be doing and they’re just not doing it and it’s frustrating to be able to have the solution but them not being invested enough in their own care to do it” (Pod B)

Subtheme 1.2 Communication challenges:

Podiatrists also reported challenges with communication, especially when the person had multiple comorbidities or limited education, and when relatives or friends joined consultations.

“The relatives and the family, because it becomes a joint problem a joint concern...some of the carers have sort of pushed their needs onto the patient and you have...to steer the consult back to the patient sometimes.” (Pod F).

Main theme 2. Training Content

Subtheme 2.1 Overall training experience:

Most podiatrists reported that MI training was enjoyable, beneficial, interesting and informative. They liked the small group and felt engaged.

“I found the training days very beneficial and informative, and I think that we were very engaged. It was quite good because it was interactive.” (Pod G).

“It was good....I definitely did learn better communication skills to a certain extent” (Pod K).

Some felt:

“There was a lot of information in a short period of time” (Pod C)

“Hard to relate it sometimes to a clinical setting” (Pod D)

Sub theme 2.2 Role-play:

Role-play practice elicited mixed feelings. Some found it useful, despite some negative feelings.

"Nice to see how different clinicians worked." (Pod A).

"As much as I hate role-play, it did help." (Pod I).

Others said:

"[Role-play is] tricky as we think differently than the patients." (Pod J).

"[I] hated it." (Pod B).

Sub theme 2.3 Imagery:

Participants struggled fitting imagery into their clinical practice:

"I found imagery most difficult to implement in a clinical environment. I thought we were reasonably well trained in it, but when I tried to execute it, I found it difficult to approach that kind of a thing with the patients. And when I did, they weren't particularly receptive." (Pod C)

Subtheme 2.4 Ongoing training and support:

More training was requested, via ongoing support in the form of peer support and via hands-on training where actual difficult patient situations can be discussed:

"If we were able to maybe isolate some of the difficult patients and have a group discussion about how you would apply that training to that particular patient. Then everyone can come away with a new way to deal with similar patients." (Pod C)

Theme 3. Training outcomes

Subtheme 3.1 New communication skills:

Applying new strategies when communicating with people with DFD, such as reflections and open-ended questions, was seen as useful.

"...having those open-ended questions other than short-ended questions, so you could really just learn to shut up and let the patient speak." (Pod E)

"Strategies of getting the patient to commit themselves to something as opposed to us dictating to them what they needed to do." (Pod E)

Participants liked the fact that training allowed them to ask people with DFD to reflect on what they had previously done and wanted to do:

"I think the open-ended questions and the reflections back were really useful for developing a rapport with the patients." (Pod D)

Other useful skills were asking for goals, allowing the patient to talk more, and reducing podiatrists' talk time:

"I have allowed the patients to speak more which has given me an insight that I did not have before" (Pod F).

"To be quiet as well, that's a hard one. It's a very practical thing to let them just speak, to fill the silence rather than us." (Pod B).

"Getting the patient to tell you what their plan is and what their goals are, because at the end of the day it's their health care." (Pod A).

Subtheme 3.2 Increased patient engagement:

The new communication skills appeared to result in increased patient engagement:

"They've seemed to be more engaged with their self-care. And instead of telling them what to do I feel that they respond better because they are thinking what they can do better. They're taking responsibility more." (Pod A)

"Initially getting people to open up and talk about experiences with foot ulcers and how they can change things, its changing their mentality from them thinking that its actually our problem and our responsibility to them actually having to think about it as their wound and their responsibility." (Pod G)

Subtheme 3.3 Long-term implementation of MI skills:

Participants reported difficulty with changing their habitual behaviour in sessions. While anticipating that applying MI over the longer-term would be hard, some were optimistic about achieving this.

"...it was definitely still challenging. Hard not problem-solving" (Pod D)

Participants suggested ways to support them to sustain their use of MI related skills over the longer term:

"[We need] more training within motivational interviewing to keep your skills up and keep it fresh. The more you do it, the more it will embed into your practice, but at the beginning it's easy to go back to how you used to do." (Pod A)

"We slowly slip back into our old ways because we don't reinforce it and we're really time poor." (Pod F)

Subtheme 3.4 Appropriateness of MI:

MI was not seen as an appropriate approach for every patient:

"Some of them still do the same thing regardless." (Pod J).

"I guess I use it where I see the need." (Pod I).

Discussion

This was the first study to investigate the impact of delivering MI training to podiatrists who treat people with DFD. The training resulted in some moderate short-term improvements in MI related skills after 2 weeks, but no improvements remained at the 12-week follow-up, and only one podiatrist attempted to use imagery in the assessed sessions. Improvements in job satisfaction were not observed, although satisfaction was already high at baseline and the training did not encompass broader issues related to the work setting.

The qualitative responses were largely consistent with other studies. Participants saw the interactive nature of workshops as beneficial, although some expressed discomfort about role-play (43). Participants had positive experiences with engaging and empowering patients after training, as found in another study (44), but expressed a need for ongoing supervision and support and noted the risk of reverting to old habits (45).

The short-term improvements in MI related skills of Empathy (30, 31, 46, 47), Change Talk (22, 30, 48) and Partnership (29) were consistent with changes in previous studies on MI training. Reductions in Persuasion are less often reported explicitly

(32); in the current study, this behaviour appeared to return to baseline levels at 12 weeks. The skill Soften Sustain did not change during the study, but statements requiring this type of response by podiatrists were surprisingly rare in the assessed audios. Rises in Simple and Complex Reflections were small in number, and effects for time fell short of statistical significance, suggesting that they require further attention in training. While both Questions and Giving Information were very frequent, these were required elements in the podiatry sessions, and it is not clear what would be an ideal frequency. The 43% reduction in Giving Information was encouraging, if it reflected a reduction in the repetition of information within and across consultations, since such repetition runs the risk of undermining rapport and collaborative self-management.

Participants' positive experiences and short-term MI related skill improvements provide a solid basis for further changes in practice. However, the lack of maintained changes in core MI skills and reported difficulties with their routine use suggest a need for additional training, including follow-up peer support session observation and feedback. The qualitative responses suggested this would be highly valued. With 8 hours of training provided in the current study, half of the median 16 hours found in a systematic review (34), more training may indeed be needed for lasting positive effects on practice. In particular, further group discussion of difficulties experienced in applying MI is recommended. Difficulties in maintaining the use of new skills has often been observed in other studies on practitioner training (22), including training on MI. A reliance on workshops alone typically gives limited skill improvement (49) and little sustained benefit (45, 50, 51). Monitoring accompanied by encouraging and corrective feedback appears critical (44, 45, 50–52), at least until use of the skills becomes habitual. The positive short-term training outcomes and positive evaluation in this pilot study suggest that podiatrists are receptive of more intensive training, can be seen as an encouragement for further research and implementation.

A strength of this study was our incorporation of several principles of implementation science to facilitate training uptake (53, 54). The project had strong management support, and one of the authors (TK) was a senior member of the team and modelled early adoption of the approach in her own work. Further, the training highlighted incentives for use of MI, including the potential for MI to reduce podiatrists' guilt or frustration by defining success in terms of their conduct of sessions rather than clinical or behavioural outcomes. Finally, we minimised change from existing practice by integrating MI with assessment, physical treatment and planning within the existing session length.

Despite these facilitators, podiatrists had difficulties integrating MI into clinical sessions if these required substantial clinical assessment and physical treatment. This integration probably needed more training than we were able to deliver, and for some attempting to use a new skill while conducting another task may have been too difficult. Podiatrists were also discouraged when people with DFD did not respond positively, and additional encouragement to persist in the face of this is required. The complexity of people's co-occurring problems also presented significant challenges for use of MI, as did addressing the needs of carers.

A limitation of this novel pilot study was the small sample size. However, we recruited sufficient participants to provide estimates of potential effect sizes for the planning of future randomised controlled trials, as well as important findings on training content and implementation. A further potential limitation was involvement of the trainers in the conduct and rating of assessments, which may result in assessment reactivity and rating biases. We minimized this by using validated tools, and having two assessors working independently. Qualitative interviews were short, but derived rich data on participants' experiences. Use of mental imagery may have also been a limitation, as it proved a step too far for the participants. They found it difficult to introduce and use in sessions, which may have blocked additional uptake of MI related skills. In retrospect, improvements in MI related skills should have been consolidated before introducing mental imagery, and more training on introducing and using imagery was needed.

Conclusion

We provided eight hours of valued training in MI to podiatrists who treat people with DFD. This resulted in some uptake of MI related skills, although maintenance of these changes was short lived. Findings of this study provide a solid basis for

refinement of the training and support for implementation and future research of its impact on both podiatry practice and outcomes for people with DFD.

Declarations

Ethics Approval and Consent to Participate

Approvals were obtained from human ethics committees from the Research Governance Unit The Prince Charles Hospital and Queensland University of Technology (# 45318), and all participating podiatrists and people with DFD provided informed consent before participation began.

Consent for publication

Not applicable

Data availability

Deidentified data sets generated/or analysed during this current study are stored at <https://github.com/dkavanagh212/DFDPodiatry.git>.

Competing interests

The authors declare they have no competing interests.

Funding

No funding was provided for the study.

Authors' contributions

All authors contributed to the design of the study. Training and ratings of the clinical interviews were conducted by the TK and DK, rating of the qualitative interviews was by TK and JvN . Data entry was by the TK, and analysed by DK. The first draft of the paper was written by TK and DK, JvN and PL critically reviewed the manuscript. All authors contributed to the manuscript and agree with the final version.

References

1. Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract.* 2018;138:271–81.
2. Armstrong DG, Boulton AJM, Bus SA. Diabetic foot ulcers and their recurrence. *N Engl J Med.* 2017;376(24):2367–75.
3. Perrin BM, Swerissen H, Payne C. The association between foot-care self efficacy beliefs and actual foot-care behaviour in people with peripheral neuropathy: a cross-sectional study. *J Foot Ankle Res.* 2009;2:3.
4. Lazzarini PA, Pacella RE, Armstrong DG, Van Netten JJ. Diabetes-related lower-extremity complications are a leading cause of the global burden of disability. *Diabet Med.* 2018;35(9):1297–9.
5. Zhang Y, McPhail SM, Netten JJv, Armstrong DG, Pacella RE, Lazzarini PA. Global disability burdens of diabetes-related lower-extremity complications in 1990 and 2016. *Diabetes Care.* 2020.

6. Lazzarini PA, Gurr JM, Rogers JR, Schox A, Bergin SM. Diabetes foot disease: the Cinderella of Australian diabetes management? *J Foot Ankle Res.* 2012;5(1):24.
7. Heidemann C, Du Y, Paprott R, Haftenberger M, Rathmann W, Scheidt-Nave C. Temporal changes in the prevalence of diagnosed diabetes, undiagnosed diabetes and prediabetes: findings from the German Health Interview and Examination Surveys in 1997–1999 and 2008–2011. *Diabet Med.* 2016;33(10):1406–14.
8. Schaper NC, Van Netten JJ, Apelqvist J, Lipsky BA, Bakker K. Prevention and management of foot problems in diabetes: A Summary Guidance for Daily Practice 2015, based on the IWGDF guidance documents. *Diabetes Res Clin Pract.* 2017;124:84–92.
9. Van Netten JJ, Seng L, Lazzarini PA, Warnock J, Ploderer B. Reasons for (non-)adherence to self-care in people with a diabetic foot ulcer. *Wound Repair Regen.* 2019;27(5):530–9.
10. Vluggen S, Hoving C, Schaper NC, de Vries H. Exploring beliefs on diabetes treatment adherence among Dutch type 2 diabetes patients and healthcare providers. *Patient Educ Couns.* 2018;101(1):92–8.
11. Barg FK, Cronholm PF, Easley EE, Davis T, Hampton M, Malay DS, et al. A qualitative study of the experience of lower extremity wounds and amputations among people with diabetes in Philadelphia. *Wound Repair Regen.* 2017;25(5):864–70.
12. Coffey L, Mahon C, Gallagher P. Perceptions and experiences of diabetic foot ulceration and foot care in people with diabetes: A qualitative meta-synthesis. *Int Wound J.* 2019;16(1):183–210.
13. Dorresteijn JA, Kriegsman DM, Assendelft WJ, Valk GD. Patient education for preventing diabetic foot ulceration. *Cochrane Database Syst Rev.* 2014;12(12):CD001488.
14. Funnell M, Anderson RM. The problem with compliance in diabetes. *JAMA.* 2000;284(13):1709-.
15. Gabbay RA, Kaul S, Ulbrecht J, Scheffler NM, Armstrong DG. Motivational interviewing by podiatric physicians. *J Am Podiatr Med Assoc.* 2011;101(1):78–84.
16. Emmons KM, Rollnick S. Motivational interviewing in health care settings: Opportunities and limitations. *Am J Prev Med.* 2001;20(1):68–74.
17. Price P. How can we improve adherence? *Diabetes Metab Res Rev.* 2016;32(S1):201–5.
18. Moser C. Toward a theory of motivational interviewing. *J Dev Behav Pediatr.* 2010;31(1):79.
19. Soderlund P. Effectiveness of motivational interviewing for improving physical activity self-management for adults with type 2 diabetes: A review. *Chronic Illn.* 2018;14(1):54–68.
20. Gabbay RA, Dellasega C, Martinez-King N., ; Orlando. Content of a motivational interviewing intervention for high risk patients with type 2DM. 70th Scientific Sessions. Florida. Arlington: American Diabetes Association; American Diabetes Association; 2010.
21. Rollnick S. Motivational interviewing in health care helping patients change behavior. In: Miller WR, Butler C, editors. *Applications of motivational interviewing.* New York: Guilford Press; 2008.
22. Jansink R, Braspenning J, Laurant M, Keizer E, Elwyn G, Weijden T, et al. Minimal improvement of nurses' motivational interviewing skills in routine diabetes care one year after training: a cluster randomized trial. *BMC Fam Pract.* 2013;14:44.
23. Kavanagh DJ, Andrade J, May J, Connor JP. Motivational interventions may have greater sustained impact if they trained imagery-based self-management. *Addiction.* 2014;109(7):1062–3.
24. Jon E, Jackie E, David JK. An imagery-based road map to tackle maladaptive motivation in clinical disorders. *Front Psychiatry.* 2015;6.
25. Kavanagh D, Andrade J, May J. Imaginary relish and exquisite torture: the elaborated intrusion theory of desire. *Psychol Rev.* 2005;112(2):446–67.
26. Parham SC, Kavanagh DJ, Shimada M, May J, Andrade J. Qualitative analysis of feedback on functional imagery training: A novel motivational intervention for type 2 diabetes. *Psychol Health.* 2018;33(3):416–29.

27. Solbrig L, Whalley B, Kavanagh DJ, May J, Parkin T, Jones R, et al. Functional imagery training versus motivational interviewing for weight loss: a randomised controlled trial of brief individual interventions for overweight and obesity. *Int J Obes (Lond)*. 2019;43(4):883.
28. Andrade J, Khalil M, Dickson J, May J, Kavanagh DJ. Functional Imagery Training to reduce snacking: Testing a novel motivational intervention based on Elaborated Intrusion theory. *Appetite*. 2016;100:256–62.
29. Rubak S, Sandbaek A, Lauritzen T, Borch-Johnsen K, Christensen B. An education and training course in motivational interviewing influence: GPs' professional behaviour–ADDITION Denmark. *Br J Gen Pract*. 2006;56(527):429–36.
30. Brug J, Spikmans F, Aartsen C, Breedveld B, Bes R, Fereira I. Training dietitians in basic motivational interviewing skills results in changes in their counseling style and in lower saturated fat intakes in their patients. *J Nutr Educ Behav*. 2007;39:8–12.
31. Magill N, Graves H, de Zoysa N, Winkley K, Amiel S, Shuttlewood E, et al. Assessing treatment fidelity and contamination in a cluster randomised controlled trial of motivational interviewing and cognitive behavioural therapy skills in type 2 diabetes. *BMC Fam Pract*. 2018;19(1):60.
32. Nightingale B, Gopalan P, Azzam P, Douaihy A, Conti T. Teaching brief motivational interventions for diabetes to family medicine residents. *Fam Med*. 2016;48(3):187–93.
33. Welch G, Zagarins SE, Feinberg RG, Garb JL. Motivational interviewing delivered by diabetes educators: Does it improve blood glucose control among poorly controlled type 2 diabetes patients? *Diabetes Res Clin Pract*. 2011;91(1):54–60.
34. Van Netten TKaczmarek, D.Kavanagh JJ, Warnock J, P.Lazzarini. A systematic review of motivational interviewing training outcomes for health practitioners treating persons with diabetes. 8th International Symposium on the Diabetic Foot; The Hague, The Netherlands. Harmelen, The Netherlands: Diabetic Foot; May 2019.
35. Knight KM, McGowan L, Dickens C, Bundy C. A systematic review of motivational interviewing in physical health care settings. *Br J Health Psychol*. 2006;11(2):319–32.
36. Gabbay R, Tinloy J, Kaul S, Ulbrecht J, Schaefer E. Podiatrists' perspectives on their role in promoting self-care in high-risk patients with diabetes. *J Am Podiatr Med Assoc*. 2014.
37. Kavanagh DJ, Young R, White A, Saunders JB, Wallis J, Shockley N, et al. A brief motivational intervention for substance misuse in recent-onset psychosis. *Drug Alcohol Rev*. 2004;23(2):151–5.
38. Baker AL, Kavanagh DJ, Kay-Lambkin FJ, Hunt SA, Lewin TJ, Carr VJ, et al. Randomized controlled trial of MICBT for co-existing alcohol misuse and depression: Outcomes to 36-months. *J Subst Abuse Treat*. 2014;46(3):281–90.
39. Moyers TB, Manuel JK, Ernst D. Motivational interviewing treatment integrity coding manual 4.2.1. 2015 (Unpublished manual).
40. Connolly JM, Kavanagh DJ, Andrade J, Solbrig L. Functional Imagery Training – Quality Coding (FIT-QC) V1.0. 2017 (Unpublished manual).
41. Hoppock R. Job satisfaction. New York: Harper & Row; 1935.
42. Pope C, Mays N. Qualitative research in health care. 3rd ed. Malden: Blackwell; 2006.
43. Doherty Y, Hall D, James PT, Roberts SH, Simpson J. Change counselling in diabetes: The development of a training programme for the diabetes team. *Patient Educ Couns*. 2000;40(3):263–78.
44. Szczekala K, Wiktor K, Kanadys K, Wiktor H. Benefits of motivational interviewing application for patients and healthcare professionals. *Pol J Public Health*. 2018;128:170–3.
45. van Eijk-Hustings YJL, Daemen L, Schaper NC, Vrijhoef HJM. Implementation of motivational interviewing in a diabetes care management initiative in the Netherlands. *Patient Educ Couns*. 2011;84(1):10–5.
46. Ismail K, Maissi E, Thomas S, Chalder T, Schmidt U, Bartlett J, et al. A randomised controlled trial of cognitive behaviour therapy and motivational interviewing for people with type 1 diabetes mellitus with persistent sub-optimal glycaemic control: A diabetes and psychological therapies (ADaPT) study. *Health Technol Assess*. 2010;14(22):1–127.

47. Keukenkamp R, Merx MJ, Busch-Westbroek TE, Bus SA. An explorative study on the efficacy and feasibility of the use of motivational interviewing to improve footwear adherence in persons with diabetes at high-risk of foot ulceration. *J Am Podiatr Med Assoc.* 2017;108:90–9.
48. Welch G, Zagarins S, Shayne R, Garb J. Does a motivational interviewing intervention delivered by diabetes educators improve HbA1c for poorly controlled type 2 diabetes patients? *Diabetes.* 2010.
49. Chilton R, Pires-Yfantouda R, Wylie M. A systematic review of motivational interviewing within musculoskeletal health. *Psychol Health Med.* 2012;17(4):392–407.
50. Martins RK, McNeil DW. Review of Motivational Interviewing in promoting health behaviors. *Clin Psychol Rev.* 2009;29(4):283–93.
51. Britt E, Blampied NM. Motivational interviewing training: A pilot study of the effects on practitioner and patient behaviour. *Behav Cogn Psychother.* 2010;38(2):239–44.
52. Söderlund LL, Madson MB, Rubak S, Nilsen P. A systematic review of motivational interviewing training for general health care practitioners. *Patient Educ Couns.* 2011;84(1):16–26.
53. Rogers EM. *Diffusion of innovations.* 3rd ed. New York: Free Press; 1983.
54. Greenhalgh T, Robert G, Bate P, Macfarlane F, Kyriakidou O. *Diffusion of innovations in health service organisations a systematic literature review.* 1st ed. Malden: Blackwell; 2005.