

Pulmonary mucoepidermoid carcinoma: Clinicopathological characteristics of 20 cases and a literature review

Jing Zhang

East Hospital

Yan Xiao Chen

Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital

Jun Zi Qian

Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital

Ping Yue

Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital

Jialei Wang

Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital

Yi Xiang (✉ xiangyiht@163.com)

<https://orcid.org/0000-0002-0724-6359>

Research article

Keywords: pulmonary mucoepidermoid carcinoma clinicopathology

Posted Date: November 26th, 2019

DOI: <https://doi.org/10.21203/rs.2.17790/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Pulmonary mucoepidermoid carcinoma is a rare tumor of the lung. The clinicopathological characteristics of pulmonary mucoepidermoid carcinoma are not well defined due to the low incidence. This study was performed to provide more supplementary clues for the identification and understanding of pulmonary mucoepidermoid carcinoma. **Methods:** We reviewed the medical records since January 1, 2000 to December 31, 2018. The patients' medical records—including age at the time of diagnosis, gender, smoking history, preoperative evaluations, operative procedures, tumor location, tumor size, tumor stage, lymph node metastasis, pathological markers, prognosis and survival information were extracted and reviewed. Categorical variables were presented as parameters and percentages. A comparison was performed between patients with high and low grade of pulmonary mucoepidermoid carcinoma. **Results:** 20 patients were identified and the age span is from 18 to 67 year-old with the average age is 45. Mucoepidermoid carcinomas were commonly found in men (60%). 80% patients had clinical presentations and the positive rate of tumor markers was 78%, although no specific tumor markers were found. TTF-1 were negative in all cases. ALK rearrangement was identified in a non-smoking woman with high grade pulmonary mucoepidermoid carcinoma. Surgery is the main procedure. 3-year survival rate is 72% and 80% patients achieved disease-free alive. High-grade patients tend to harbor older age ($p=0.035$), larger tumor volume ($p=0.026$) and higher index of ki-67 ($p=0.0005$). **Conclusions:** Pulmonary mucoepidermoid carcinoma could occur in a wide age span. Early diagnosis and complete surgical resection may promise a good prognosis. Grading is a key factor to predict the overall survival time. Combined TTF-1 and MAML2 will benefit the identification of pulmonary mucoepidermoid carcinoma from other lung tumors. Future prospective randomized controlled trials and larger, multi-centric series are needed.

Background

Pulmonary mucoepidermoid carcinoma (PMEC) is a rare tumor that accounts for 0.1% to 0.2% of all pulmonary tumors. Most pulmonary MECs arise in the proximal bronchi [1]. Histologically, the classification of the PMEC is mainly based on the degree of mitosis–necrosis and abnormalities of the nucleus in the tumor. Low-grade mucoepidermoid carcinoma consists mainly of glandular-like structures and mucin-secreting cells, whereas high-grade PMECs include a large number of squamous cells [2]. However, in a few previous studies, intermediated grade had been suggested because of different grade systems described by Brandwein[3]. So far, most researches on PMEC belonged to either a series of cases report or small sample retrospective studies, especially in China [4-6]. The characteristics of PMEC are not well defined due to the low incidence of PMEC and there are a lot of controversies in the diagnosis and treatment of PMEC. In this paper, we analyzed the clinicopathological features, treatment and prognostic information of 20 PMEC patients who were exposed to 2000-2018 in Shanghai Ruijin Hospital and Shanghai Cancer center. This study was performed to provide more information for the identification and understanding of pulmonary mucoepidermoid carcinoma.

Methods

Patients and tissue specimens

The study was conducted under the approval by the Institutional Review Board. Ethical approval was granted by the clinical Research Ethics Committee of Shanghai Ruijin Hospital, Jiaotong University and Shanghai Cancer center, Fudan university. We reviewed the medical records, and outpatient pathology visits since January 1, 2000 to December 31, 2018 and collected 20 patients with PMEC, including 12 inpatients and 8 cases of pathology consultation patients whose data were partly missing. Signed informed consent was obtained from all induced patients for the acquisition and use of tissue samples and anonymized clinical data. Tissue specimens were collected from surgical resection or biopsy. The bronchoscopy was performed in 8 inpatients. In the preoperative period, two inpatients had a diagnosis of PMEC by bronchoscopy, whereas the other patients were finally diagnosed through resected specimens. Hematoxylin and eosin(H&E) stained slides in all patients were reviewed by two experienced pathologists for histopathological confirmation. PMEC was divided into low or high grade according to the World Health Organization(WHO) criteria of lung cancer [7].

Data collection and extraction

The patients' medical records—including age at the time of diagnosis , gender, smoking history, preoperative evaluations, operative procedures, tumor location, tumor size, tumor stage, lymph node metastasis, pathological markers, prognosis and survival information were extracted and reviewed. Tumor stages were classified according to the American Joint Committee on Cancer Staging System, 7th edition. We acquired follow-up data from outpatient follow-ups or telephone and the last follow up time is December 31, 2018.

Statistical analysis

Categorical variables were presented as parameters and percentages. Nominal variables included gender, smoker, lymph node metastasis and TNM stage were conducted using Fisher's exact test. Age, tumor size and Ki-67 labeling index were compared by way of an independent-samples t-test. All tests were two-sided, and p-values less than 0.05 were considered to be statistically significant.

Results

Clinical characteristics

Table.1 summarized the main clinical features of the 20 patients. Twenty cases of PMEC included 8 females and 12 males ranged from 18 to 67 years old with a mean age of 45. Five of 13 patients were smokers. Among ten patients, eight had symptoms which were presenting as cough(7/8) fever(1/8) hemoptysis(1/8) and bone pain(1/8). Two patients were asymptomatic. All 12 inpatients underwent CT examination. CT imaging showed a solitary nodule located at the periphery or the hilum of lung (Figure

1). Almost half of patients (42%) had complications, presented as atelectasis or obstructive pneumonia. 8 patients performed bronchoscopy and the neoplasm grew into the lumen in 7(88%) patients. 13(65%) tumors were located in right lung, including right upper lobe(n=2), right middle lobe(n=5), right lower lobe(n=6). 7(35%) tumors were located in left lung, including left upper lobe(n=4), left lower lobe(n=3). The levels of serum tumor markers were detected in 9 patients. 7(78%) cases had higher levels of tumor markers, including CA125(n=3), NSE□CYFRA21-1(n=2), CEA□AFP□CA153□SCCA(n=1), while two(22%) patients were normal.

Histopathology

Twenty patients were divided into low-grade (85%) and high-grade (15%) according to pathological diagnosis (Figure 2). Some patients' pathological characteristics were available and summarized in Table2. The results showed that resection margins were negative in all 14 (100%) patients and no perineural invasion were found in 6(100%) cases who had been recorded. Intravascular embolus were found in 2(33%) cases. Pleural involvement were positive in 4(36%) patients among 11 patients. Bronchus invasion existed in 2 (21%)patients. TTF-1 were negative in all patients and P63 were detected in 67% patients.

Treatment

The main treatment was surgery (Table1). Operation procedures included 12 (63%)lobectomies, 4 (21%)bilobectomies, 1(5%) sleeve resection and 2(11%) pneumonectomies. However, there was one patient lost the opportunity of surgery because of bone metastasis. Chemotherapies were performed in 3(25%)patients after operation among 12 cases and the patient accompanied with bone metastasis received local radiotherapy. 11 patients had follow-up records and the follow-up time ranged from 8 to 126 months (mean 52m). Two patients died of tumor at 8 and 18 months after diagnosis. Among these two patients, one did not undergo surgery because of bone metastasis and the other one suffered recurrence in ipsilateral lung after surgery. The remaining 9 patients were alive without evidence of recurrence or metastasis. A comparison was performed between patients with high and low grade of pulmonary mucoepidermoid carcinoma. There were statistically significant differences in the age(p=0.035), tumor size(p=0.026) and the index of ki-67(p=0.0005).

Discussion

Pulmonary mucoepidermoid carcinoma is the most common type of salivary gland tumor [8]. Previous studies indicated that P MEC had a broad spectrum in age and often occurred in 40s [5,9]. The age span of our series is from 18 to 67 year-old and the average age is 45. Interestingly, in accordance with recent reports, we also found that P MECs were commonly found in men [10]. 80% patients had clinical presentations and the positive rate of tumor markers was 78%, although no specific markers were found. Chest CT had an important value in the diagnosis of P MEC despite rare tumors exhibited an endobronchial pattern. If these patients were accompanied with persistent respiratory symptoms or elevated tumor markers, bronchoscopy examination and biopsy should be conducted to avoid

misdiagnosis. TTF-1 was negative in all PMECs, conforming that it was of great value in distinguishing PMEC from lung adenocarcinoma, adenosquamous carcinoma and PMEC-like tumors [11]. Surgery is the main procedure of PMEC. Most patients were alive without evidence of disease after completely resection [12]. Our study showed that 3-year survival rate of PMEC is 72%, and 80 percent of those patients achieved disease-free alive. Histological grade was widely recognized as a prognostic factor. Our series also demonstrated that high-grade PMEC patients tend to harbor older age, larger tumor volume and higher index of ki-67. Due to the small size of samples, we did not find significant difference in TNM stage and metastasis between low and high grade of PMEC. However, some large series had revealed the worse outcomes of patients who accompanied with advanced TNM staging and earlier metastasis [13-14].

At present, there were few studies focused on driven genes of PMEC. Evidences indicated that the most frequent gene mutation in PMEC is MAML2, which belongs to MasterMind-like gene family on chromosome 11q21. In salivary gland tumors, MAML2 rearrangement has been identified in 40–75% of mucoepidermoid carcinoma [15-16]. PMEC has a low preoperative diagnosis rate through bronchoscopy and CT-guide percutaneous lung puncture biopsy. Most of our cases had been treated as non small cell lung cancer (NSCLC) before and during surgery. This may suggest that the application of molecular techniques to detect the MAML2 rearrangement and TTF-1 expression may be helpful in improving the preoperative diagnosis. Targeted therapy has brought survival benefit to NSCLC patients. So far there is no effective treatment for PMEC patients who cannot undergo surgery. Targeted therapy for PMEC-driven genes may be a wise choose in future. However, the association between MAML2 rearrangement and prognosis is still in suspense. The current view suggests that MALML2 is more prone to occur in lower-grade PMEC and has no value for prognosis [17-19]. Few studies had paid attention to other driven genes such as EGFR and ALK. It was noted that the mutational frequency of EGFR in PMEC was 40% and mutations were all L858R mutation. A case of ALK-rearranged PMEC was reported by Wong et.al in 2009 [20-22]. In our retrospective cases, we identified ALK rearrangement in a non-smoking woman with high grade PMEC.

Conclusions

Our study has provided supplementary clues for the identification of PMEC and updated the understanding of PMEC. However, as a retrospective study, it had inherent limitations due to the small size of sample. Meanwhile, some clinical data and follow-up information were not available. Future prospective randomized controlled trials and larger, multi-centric series are needed to overcome the challenges related to the accurate diagnosis and prediction of PMEC.

Abbreviations

PMEC: Pulmonary mucoepidermoid carcinoma

WHO: World Health Organization

CT: Computed tomography

CA125: Carbohydrate antigen 125

NSE: Neuron-specific enolase

CYFRA21: Cytokeratin 19 fragments

CA153: Carbohydrate antigen 153

SCCA: Squamous cell carcinoma antigen

CEA: Carcinoembryonic antigen

AFP: Alpha fetoprotein

H&E: Hematoxylin and eosin

TTF-1: Thyroid transcription factor-1

NSCLC: Non small cell lung cancer

ALK: Anaplastic lymphoma kinase

EGFR: Epidermal growth factor receptor

Declarations

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgements: Not applicable.

Ethics declarations:

Authors' contributions: Yi Xiang and Jialei Wang contributed the central idea. Jing Zhang and Xiaoyan Chen analyzed most of the data, and wrote the initial draft of the paper. Zijun Qian and Ping Yue contributed to refining the ideas, carrying out additional analyses and finalizing this paper.

Funding: This study was supported by National Natural Science Foundation of China (Grant number 81672271), Shanghai Jiao Tong University Project and Shanghai Key Discipline for Respiratory Disease (Grant number 2017ZZ02014). The funders have played an important role in the design of the study and collection of data.

Ethics approval and consent to participate: The study was conducted under the approval by the Institutional Review Board. Ethical approval was granted by the clinical Research Ethics Committee of Shanghai Ruijin Hospital, Jiaotong University and Shanghai Cancer center, Fudan university. Signed informed consent was obtained from all induced patients for the acquisition.

Patient consent for publication: All patients signed consent for publication.

Competing interests: The authors declared that they have no conflict of interest.

References

1. Hsieh CC, Sun YH, Lin SW, Yeh YC, Chan ML. Surgical outcomes of pulmonary mucoepidermoid carcinoma: A review of 41 cases. *PLoS One*. 2017; 12: e0176918.
2. Yamamoto T, Nakajima T, Suzuki H, Tagawa T, Iwata T, Mizobuchi T, Yoshida S, Nakatani Y, Yoshino I. Surgical treatment of mucoepidermoid carcinoma of the lung: 20 years' experience. *Asian Cardiovasc Thorac Ann*. 2016; 24: 257-261.
3. Lee GD, Kang DK, Kim HR, Jang SJ, Kim YH, Kim DK, Park SI. Surgical outcomes of pulmonary mucoepidermoid carcinoma: a review of 23 cases. *Thorac Cardiovasc Surg*. 2014; 62: 140-146.
4. Jiang L, Li P, Xiao Z, Qiu H, Zhang X, Xiao Y, Zhang B. Prognostic factors of primary pulmonary mucoepidermoid carcinoma: a clinical and pathological analysis of 34 cases. *Int J Clin Exp Pathol*. 2014; 7: 6792-6799.
5. Wang M, Ouyang S, Sun P, Li D, Huang G. Pulmonary mucoepidermoid carcinoma in Chinese population: a clinicopathological and radiological analysis. *Int J Clin Exp Pathol*. 2015; 8: 3001-3007.
6. [Chen H, Zhang J, Qiu XJ, Wang J, Pei YH, Wang YL](#). Interventional Bronchoscopic Therapy in Adult Patients with Tracheobronchial Mucoepidermoid Carcinoma. *Chin Med J (Engl)*. 2017 Oct 20;130(20):2453-2458.
7. Ishikawa Y, Alvarez-Fernandez E, Aubry MC, Dacic S, Nicholson AG. Mucoepidermoid carcinoma. In: Travis WD, Brambilla E, Burke AP, Marx A, Nicholson AG, editors. WHO classification of tumours of the lung, pleura, thymus and heart. Lyon: IARC Press; 2015. pp. 99–100.
8. [Kumar V1, Soni P2, Garg M2, Goyal A1, Meghal T3, Kamholz S2, Chandra AB](#) A Comparative Study of Primary Adenoid Cystic and Mucoepidermoid Carcinoma of Lung. *Front Oncol*. 2018 May 15;8:153.
9. PhD JRMM, Aubry MC, Lewis JE, Wampfler JA, Ms BAW, Midthun DE, PhD PYM, Cassivi SD. Primary salivary gland-type lung cancer. *Cancer*. 2007; 110: 2253-2259.
10. [Cheng DL1, Hu YX1, Hu PQ1, Wen G2, Liu K](#) Clinicopathological and multisection CT features of primary pulmonary mucoepidermoid carcinoma. *Clin Radiol*. 2017 Jul;72(7):610.e1-610.e7.
11. Huo Z, Wu H, Li J, Li S, Wu S, Liu Y, Luo Y, Cao J, Zeng X, Liang Z. Primary Pulmonary Mucoepidermoid Carcinoma: Histopathological and Molecular Genetic Studies of 26 Cases. *PLoS*

- One. 2015; 10: e0143169.
12. Yamamoto T, Nakajima T, Suzuki H, Tagawa T, Iwata T, Mizobuchi T, Yoshida S, Nakatani Y, Yoshino I. Surgical treatment of mucoepidermoid carcinoma of the lung: 20 years' experience. *Asian Cardiovasc Thorac Ann.* 2016 Mar;24(3):257-61.
 13. Komiya T, Perez RP, Yamamoto S, Neupane P. Primary lung mucoepidermoid carcinoma: analysis of prognostic factors using surveillance, epidemiology and end results program. *Clin Respir J.* 2017 Nov;11(6):847-853.
 14. Qin BD, Jiao XD, Liu K, Wu Y, He X, Liu J, Zang YS. Clinical, pathological and treatment factors associated with the survival of patients with primary pulmonary salivary gland-type tumors. *Lung Cancer.* 2018 Dec;126:174-181.
 15. Jee KJ, Persson M, Heikinheimo K, Passador-Santos F, Aro K, Knuutila S, Odell EW, Makitie A, Sundelin K, Stenman G, Leivo I. Genomic profiles and CRTC1-MAML2 fusion distinguish different subtypes of mucoepidermoid carcinoma. *Mod Pathol.* 2013; 26: 213-222.
 16. Saade RE, Bell D, Garcia J, Roberts D, Weber R. Role of CRTC1/MAML2 Translocation in the Prognosis and Clinical Outcomes of Mucoepidermoid Carcinoma. *JAMA Otolaryngol Head Neck Surg.* 2016; 142: 234-240.
 17. Zhu F, Wang W, Hou Y, Shi J, Liu Z, He D, Bai C, Li S, Jiang L. MAML2 rearrangement in primary pulmonary mucoepidermoid carcinoma and the correlation with FLT1 expression. *PLoS One.* 2014; 9: e94399.
 18. Roden AC, Garcia JJ, Wehrs RN, Colby TV, Khor A, Leslie KO, Chen L. Histopathologic, immunophenotypic and cytogenetic features of pulmonary mucoepidermoid carcinoma. *Mod Pathol.* 2014; 27: 1479-1488.
 19. Kalhor N1, Moran CA Pulmonary mucoepidermoid carcinoma: diagnosis and treatment. *Expert Rev Respir Med.* 2018 Mar;12(3):249-255.
 20. Han SW, Kim HP, Jeon YK, Oh DY, Lee SH, Kim DW, Im SA, Chung DH, Heo DS, Bang YJ, Kim TY. Mucoepidermoid carcinoma of lung: potential target of EGFR-directed treatment. *Lung Cancer.* 2008; 61: 30-34.
 21. Yu Y, Song Z, Gao H, Zhu L, Lu S, Zhang J, Luo Q. EGFR L861Q mutation is a frequent feature of pulmonary mucoepidermoid carcinoma. *J Cancer Res Clin Oncol.* 2012; 138: 1421-1425.
 22. Wong DW1, Leung EL, So KK, Tam IY, Sihoe AD, Cheng LC, Ho KK, Au JS, Chung LP, Pik Wong M; University of Hong Kong Lung Cancer Study Group. The EML4-ALK fusion gene is involved in various histologic types of lung cancers from nonsmokers with wild-type EGFR and KRAS. 2009 Apr 23.15;115(8):1723-33.

Tables

Table1. Clinical data of 20 patients with pulmonary mucoepidermoid carcinoma

Parameters	Percentage	Parameters	Percentage
Age,yr(n=20)		Left lower lobe	3
Mean	45	Tumor markers(n=9)	
Range	18-67	Present	7(78%)
Gender(n=20)		CA125	3
Female	8(40%)	NSE	2
Male	12(60%)	CYFRA21-1	2
Smoking(n=13)		CA153	1
Never	8(62%)	SCCA	1
Have smoked	5(38%)	CEA	1
Symptoms(n=10)		AFP	1
Present	8(80%)	Absent	2(22%)
Fever	1	Operation method(n=19)	
Cough	7	Lobectomy	12(63%)
Hemoptysis	1	Sleeve resection	1(5%)
Bone pain	1	Pneumonectomy	2(11%)
Absent	2(20%)	BiLobectomy	4(21%)
Chest CT findings(n=12)		Adjuvant therapy(n=12)	
Atelectasis or obstructive pneumonia	5(42%)	Present	3(25%)
A solitary nodule	7(58%)	Radiotherapy only	0
Bronchoscopy (n = 8)		Chemotherapy only	2
Neoplasm in lumen	7(88%)	Both	1
Normal	1(12%)	Absent	9(75%)
Tumor Location (n = 20)		Follow-up time.mo(n=11)	
Right Lung	13(65%)	Mean	52
Right upper lobe	2	Rang	8-126
Right middle lobe	5	Prognosis(n=11)	
Right lower lobe	6	3-year survival rate	72%
Left lung	7(35%)	Disease free	8(80%)
Left upper lobe	4	Death	2(20%)

Table2. Histopathological information of patients with pulmonary mucoepidermoid carcinoma

Parameters	Percentage	Parameters	Percentage
Tumor grade(n=20)		Pleural involvement(n=11)	
Low grade	17(85%)	Positive	4(36%)
High grade	3(15%)	Negative	7(64%)
TNM stage(n=12)		Bronchus invasion (n=13)	
I-IIA	9(75%)	positive	2(15%)
IIB-V	3(25%)	negative	11(85%)
Resection margin (n=14)		TTF-1(n=10)	
Positive	0(0%)	Positive	0
Negative	14(100%)	Negative	10(100%)
Perineural invasion(n=6)		P63(n=9)	
Positive	0(0%)	Positive	6(67%)
Negative	6(100%)	Negative	3(33%)
Intravascular embolus(n=6)		ALK(n=3)	
Positive	2(33%)	Positive	1(33%)
Negative	4(67%)	Negative	2(67%)

Table3. Comparison between patients with high and low grade of pulmonary mucoepidermoid.

Parameters	High grade	Low grade	P-value
Age(n=20)			0.035
Mean, range	61(56-67)	42(18-64)	
Ki-67(n=11)			0.0005
Mean, range	32%(15%-50%)	3.6%(3%-10%)	
Tumor size.cm(n=12)			0.026
Mean, range	4.7(3-6)	2.6(1-4)	
Gender(n=20)			1.000
Female	1	7	
male	2	10	
TNM stage (n=12)			0.127
I-IIA	1	7	
IIB-IV	2	1	
Local invasion(n=19)			0.546
Positive	2	6	
Negative	1	10	
Metastasis(n=17)			0.465
Positive	1	2	
Negative	2	12	

Local invasion included perineural invasion, blood vessel invasion, pleural involvement and bronchus invasion; Metastasis included lymph node, bone, abdominal and brain.

Figures




Figure 1

CT features of Pulmonary mucoepidermoid carcinoma. (A-B) Pulmonary mucoepidermoid carcinoma in a 64-year-old woman. CT shows a peripheral pulmonary nodule with slight lobulated and pleural involvement. (C-D) Pulmonary mucoepidermoid carcinoma in a 21-year-old man with a well-circumscribed mass at central site.




Figure 2

Hematoxylin-eosin staining images of low and high grade mucoepidermoid carcinoma. (A-B) Low-grade mucoepidermoid carcinoma showing glandular-like structures and mucin-secreting cells (red arrows); (C-D) High-grade mucoepidermoid carcinoma exhibiting evidently cellular atypia, islands of squamous cells (black arrows), with only scattered mucin