

Analysis of the Early Clinical Outcomes of Arthroscopic Debridement in the Treatment of Shoulder Tuberculosis

Yan Wei He

The Affiliated Hospital of Southwest Medical University

Jun-Cai Liu

The Affiliated Hospital of Southwest Medical University

Zhi Wang

The Affiliated Hospital of Southwest Medical University

Peng Zhou

The Affiliated Hospital of Southwest Medical University

Xiang tian Deng

Nankai University

Li Yang

The Affiliated Hospital of Southwest Medical University

Zan Chen

The Affiliated Hospital of Southwest Medical University

Zhong Li (✉ 3201996908@qq.com)

The Affiliated Hospital of Southwest Medical University

Research article

Keywords: shoulder, tuberculosis, arthroscopy, treatment

Posted Date: October 7th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-84451/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on November 20th, 2020. See the published version at <https://doi.org/10.1186/s13018-020-02086-7>.

Abstract

Background Due to atypical clinical symptoms, it is difficult to diagnose joint tuberculosis infection, which often results in misdiagnosis and missed diagnosis. It is easy to cause joint disability, and there are few reports of using arthroscopy to diagnose and treat shoulder tuberculosis. This case series aims to introduce the clinical outcomes of arthroscopic treatment of shoulder tuberculosis.

Methods 29 patients with shoulder tuberculosis from September 2013 to February 2019, 10 males, 19 females. Age range from 22-69, The average age is 37.6 years. All patients underwent arthroscopic lesion debridement, preoperative and postoperative regular use of isoniazid, rifampicin, pyrazinamide, streptomycin quadruple anti-tuberculosis drugs. The erythrocyte sedimentation rate (ESR) and C reactive protein (CRP) were recorded before and at the last follow-up. The shoulder function was evaluated according to the visual analogue scoring method (visual analogue scale, VAS) pain score and Constant score.

Results 29 patients were followed up for 12-24 months, the average follow-up time was 15.7 months. The pathological diagnosis of all patients after surgery was shoulder tuberculosis. No serious complications were found at the last follow-up, and the incision healed well. VAS pain score, Constant score, ESR and CRP at the last follow-up were significantly improved compared with those before treatment ($p < 0.05$).

Conclusion On the basis of the standard use of anti-tuberculosis drugs before and after surgery, shoulder arthroscopy is used to treat early and mid-term shoulder tuberculosis, which can be diagnosed by direct observation under the arthroscope and postoperative pathological examination. It has the advantages of thorough lesion removal, minimally invasive, rapid recovery, and reliable clinical effect.

Background

Tuberculosis (TB) is a chronic, granulomatous, and necrotizing disease caused by infection with *Mycobacterium tuberculosis* (MTB). Tuberculosis has always been one of the world's public health problems. According to statistics from the World Health Organization, there were approximately 6.4 million new tuberculosis cases worldwide in 2018. 866000 new cases in China in 2018^[1]. Osteoarticular tuberculosis accounts for about 10% of extrapulmonary tuberculosis^[2]. The shoulder joint is a rare part of joint tuberculosis^[3]. With the increase of drug-resistant tuberculosis strains and patients with autoimmune diseases, the incidence of shoulder tuberculosis has also increased^[2,4-6]. Shoulder tuberculosis can cause bone and joint damage, seriously affecting the ability to move and quality of life. Because of its atypical symptoms, shoulder tuberculosis is difficult to obtain etiological evidence and it is difficult to distinguish from malignant tumors, chronic infections and other diseases in imaging, which makes the diagnosis difficult, so it is often missed and misdiagnosed, and the disability rate is very high^[7]. After years of popularization and development, shoulder arthroscopy technology has become one of the important ways to solve shoulder diseases with its advantages of small impact on anatomical structures, minimally invasive, rapid recovery. This article retrospectively analyzed 29 cases of minimally

invasive treatment of tuberculosis under shoulder arthroscopy performed in our unit. The clinical outcomes are satisfactory. The report is as follows.

Method

Inclusion criteria: (1) Patients with regular anti-tuberculosis treatment for more than 3 weeks before surgery. (2) Patients with joint tuberculosis diagnosed pathologically after surgery. (3) Patients treated with arthroscopy. (4) Patients who take anti-tuberculosis drugs regularly after surgery.

Exclusion criteria: (1) Patients who received conservative treatment and did not receive arthroscopy. (2) In addition to arthroscopy, patients who need other surgical procedures. (3) Patients who fail to take anti-tuberculosis drugs on time and regularly as prescribed by a doctor. (4) Patients with infectious active tuberculosis.

From September 2013 to February 2019, a total of 29 patients met the inclusion criteria, 10 males and 19 females, age range from 22-69 years, with an average age of 37.6 years. All patients had single joint involvement of the shoulder joint. 11 patients had a history of tuberculosis. The average course of disease is 13 months (7-24 months). All patients undergo detailed clinical, laboratory and imaging examinations before surgery. Including blood routine examination, liver and kidney function, X-ray, CT, (SPE/CT) MRI, tuberculosis T-SPOT examination and other routine examinations. All patients had elevated ESR and CRP before surgery. No sinus formation was seen in all patients.

Preoperative management

All patients should be prescribed anti-tuberculosis drugs before surgery, and be treated for at least 3 weeks according to the principles of regularity, whole course, combination, and appropriate amount. The drugs included isoniazid, rifampicin, pyrazinamide, and ethambutanol in a quadruple anti tuberculosis treatment, with supportive treatment. When the symptoms of tuberculosis poisoning have basically disappeared, the systemic symptoms and nutritional status have basically improved, and other basic diseases had stabilized, the preoperative erythrocyte sedimentation rate ESR is adjusted to below 40mm/h, and surgical treatment can be arranged.

Surgical purpose and technique

All operations were performed by the same physician who was proficient in arthroscopic techniques. After the patient underwent tracheal intubation and general anesthesia. Those without contraindications use controlled blood pressure to control blood pressure at 100mmHg/60mmHg. A modified lateral position was taken, with the trunk tilted back 20-30 degrees, shoulder abduction 30-70 degrees, and flexion 15-20 degrees. The surgical area was sterilized and draped, the sterile arm was pulled by 3-5 kilograms, the soft cushion under the armpit assisted in abduction, and the anatomical position and approach were marked. First, the conventional shoulder arthroscopy approach was adopted to explore the joint cavity, and the synovial hyperplasia in the joint cavity and the damage of various anatomical structures were observed.

An anterior approach was established through the rotator cuff space under arthroscopic monitoring, and typical diseased synovial tissues were selected for pathological examination with nucleus pulposus clamp. The synovium, articular cartilage denudation and part of the diseased subchondral bone were removed, and the adhesion tissue in the shoulder joint was released. The shoulder joint has been moved in all directions, which can suck up the tissue debris in the joint cavity. When necessary, the anterior and posterior approach was used to enter the subacromial space for acromial formation and drainage of pus in the bursa. After the operation, the bleeding was completely stopped, the wound was sutured, and cotton pads were used to protect the underarms and then bandaged appropriately. Usually under arthroscopy, the synovial fluid is yellow and turbid, with varying degrees of synovial hyperplasia, hyperemia, edema, hypertrophy, and paleness. Granuloma and pannus. Cartilage degeneration defect, cartilage exfoliation, subchondral bone exposure, subchondral bone destruction.

Postoperative management

Nutritional support treatment is given after surgery, such as a high-protein diet. And it is necessary to take isoniazid, rifampicin, ethambutol, pyrazinamide quadruple anti-tuberculosis treatment for 12 months. The shoulder joint abduction brace has been worn and fixed in the abduction and external rotation position for 6-8 weeks. After awakening from anesthesia, the patient was instructed to start wrist flexion and dorsiflexion activities, and elbow joint flexion and extension activities. On the second day after the operation, the patient was instructed to perform mild shoulder joint passive abduction and external rotation function training, and the amount of activity was gradually increased 6 weeks after the operation until the maximum tolerance or nearly returned to normal. Regularly review blood routine, ESR, CRP, liver and kidney function and other indicators after surgery, and pay attention to prevent postoperative joint stiffness and muscle atrophy.

Observation indicators and Statistical treatment

Follow-up was 1 week, 4 weeks, 3, 6, 12, 18, and 24 months after surgery. The VAS pain score, Constant score, ESR value and CRP value were recorded before treatment and at the last follow-up. The statistical data were analyzed by SPSS 17.0 statistical software. The enumeration data was expressed with mean \pm standard deviation. Before and after treatment VAS pain score, Constant score, ESR value and CRP value were conducted by paired t test. The test level was 0.05. $P < 0.05$ indicated statistical significance.

Results

All the patients had a smooth operation, the incision was healed in stage I, and no serious complications such as postoperative infection and nerve injury occurred. All 29 cases were followed up. The follow-up time was 12-24 months, with an average follow-up time of 15.7 months. The postoperative pathological diagnosis of 29 patients was shoulder tuberculosis. At the last follow-up evaluation, the VAS pain score, Constant score, ESR and CRP were significantly improved compared to before treatment.

Case

A female, 36 years old, had left shoulder pain and limited mobility for two years. Physical examination revealed pain in the front and back of his left shoulder. The left shoulder joint is limited in abduction, adduction, internal rotation, and extension. The X-ray showed that the bone of the left shoulder joint was slightly damaged. MRI showed narrowing of the glenohumeral joint space, thickening of the glenohumeral joint synovium, effusion in the joint cavity, and swelling of the soft tissue around the joint muscles. The diagnosis of tuberculosis of the left shoulder joint has been considered. The symptoms were improved after three weeks of four anti-tuberculosis treatments. Shoulder arthroscopic debridement was performed 3 weeks later. During the operation, a large number of synovial hyperplasia, congestion and swelling in the joint cavity were seen, and the articular cartilage of the humeral head was extensively destroyed, and the exfoliated cartilage was removed. Postoperative pathological examination confirmed shoulder tuberculosis. Regular anti-tuberculosis treatment was continued for 12 months after the operation. After 24 months of follow-up, the symptoms of the shoulder joint improved significantly.

Discussion

Tuberculosis caused by mycobacterium tuberculosis has been accompanied by the development of human civilization. With the advent of anti-tuberculosis drugs, tuberculosis has gradually been effectively controlled. Bone and joint tuberculosis accounts for about 10% of extrapulmonary tuberculosis^[2,8,9]. Spine tuberculosis, hip tuberculosis and knee tuberculosis account for the majority of bone and joint tuberculosis, while shoulder tuberculosis is rarely reported. In recent years, due to the increase in world population movement and the increase in drug-resistant strains. The incidence of tuberculosis is showing an upward trend. Especially in underdeveloped areas in western China, tuberculosis, especially bone and joint tuberculosis, is still relatively common. This study uses arthroscopy to diagnose and treat patients and achieve significant results. It can not only completely clear the lesion, but also reduce the effect of inflammatory factors. It can also help undiagnosed patients before the operation to confirm the diagnosis through intraoperative observation and postoperative pathological examination. Shoulder arthroscopic surgery has minimally invasive, a wide field of vision under the arthroscope, thorough removal of lesions, and rapid functional recovery. Continuing to cooperate with anti-tuberculosis drug treatment can effectively control the development of the disease and protect joint function.

Diagnosis of shoulder tuberculosis

The clinical symptoms of shoulder tuberculosis are atypical. Most patients do not have the characteristic symptoms of tuberculosis such as hot flashes, night sweats, and cough. Moreover, most patients failed to find a history of tuberculosis. Shoulder tuberculosis often has no specific manifestations on imaging and laboratory indicators, and often needs to be distinguished from shoulder joint purulent infection, shoulder joint malignant tumors, rheumatoid arthritis, gouty arthritis and pigmented villonodular synovitis^[10-14]. Diagnostic puncture and extraction of joint fluid applied to shoulder tuberculosis often results in negative results and needs to be performed multiple times. Not only does it not help diagnose, but also increases the risk of joint cavity infection and tuberculosis dissemination. Furthermore, clinical orthopedic doctors do not have a deep understanding of shoulder tuberculosis, and they often fail to

consider the possibility of tuberculosis. The onset of shoulder tuberculosis is insidious, and many patients do not seek medical attention until they have joint dysfunction or joint swelling and pain. Therefore, early diagnosis and treatment are often not available. Causes a prolonged course of shoulder tuberculosis patients, and the disease continues to develop and spread. From the early stage simple synovial tuberculosis or subchondral bone destruction develops into total joint tuberculosis. It even involves surrounding muscles, tendons, nerves and lymph nodes. The disability rate is very high, which seriously affects the quality of life of patients^[15].

The early diagnosis and early treatment of shoulder tuberculosis are for the recovery of the patient's limb function and the improvement of symptoms. In daily diagnosis and treatment activities, clinicians should pay attention to inquiring about the patient's medical history and grasp every information that is helpful for diagnosis. The patient's past history, with or without tuberculosis poisoning symptoms or tuberculosis-specific signs, such as cough, night sweats, low fever, fatigue, weight loss, etc. In some cases, recurrent shoulder swelling and pain with limited mobility are present. In imaging examinations, MRI is the most important auxiliary diagnostic method. MRI of shoulder tuberculosis generally manifests as roughness and narrowing of the joint space, thickening of synovial edema, joint capsule effusion, discontinuous bone cortex, increased bone signal, and bone quality Irregular destruction, etc. When a highly suspicious case is found, diagnostic anti-tuberculosis treatment can be performed. The diagnosis of shoulder tuberculosis should be based on the patient's clinical symptoms, medical history and signs, imaging findings, and laboratory indicators to make a comprehensive consideration. Pathological diagnosis is the gold standard of shoulder joint diagnosis. In this study, the minimally invasive method of shoulder arthroscopy can obtain pathological examination specimens as soon as possible, which has high utility value.

Arthroscopy for the treatment of shoulder tuberculosis

In recent years, with the continuous development and maturity of shoulder arthroscopy technology, shoulder arthroscopy technology has been widely developed in orthopedics clinics. Compared with open operation, arthroscopy has the advantages of minimally invasive, wider, clearer vision and larger cleaning area under the arthroscope, and thorough removal of the lesion. However, there have been few reports on the treatment of shoulder tuberculosis by arthroscopy. In other joint tuberculosis, the treatment with arthroscopy has achieved good results^[16-18]. This is similar to the clinical results achieved in this study. Shoulder arthroscopy can effectively control the development of tuberculosis through thorough and effective removal of necrotic tissues and drug treatment, so as to achieve good therapeutic effects.

The early stage of shoulder tuberculosis usually manifests as synovial tuberculosis and or subchondral tuberculosis. Synovial tuberculosis spreads from the blood source and infects the synovium. Synovial congestion and swelling into villonodula, and synovial tissue hyperplasia. Sometimes granuloma or pannus can be seen. Due to the poor blood supply to cartilage, Mycobacterium tuberculosis invades the joints and spreads along the subchondral area, causing subchondral bone destruction and cartilage denudation.

arthroscopic treatment

Early diagnosis or high suspicion of shoulder tuberculosis can be roughly diagnosed by observing synovial tissue and cartilage denudation under a arthroscope after a period of anti-tuberculosis treatment. After obtaining the synovial tissue and other specimens, perform pathological examination to achieve the purpose of confirming the diagnosis. At the same time, the hyperplastic synovium, fibrous necrotic tissue, cartilage denudation and tuberculosis foci are removed, and the adhesions are loosened under the arthroscope, and the dead space is eliminated. Continuous lavage of the joint cavity under the arthroscope can not only completely remove the necrotic tissue and inflammatory factors, but also improve the internal environment of the joint and effectively prevent the recurrence and spread of tuberculosis^[19,20]. For end-stage shoulder tuberculosis, since it has developed into total joint tuberculosis, a large amount of bone has been destroyed by tuberculosis, arthroscopy can still remove dead bones, synovial tissue and tuberculosis foci, reduce inflammatory mediators and slow the progression of the disease. Creating opportunities for other operations and treatments for the future, increase the success rate of operations and the cure rate of diseases. Due to the small sample size in this study, it is impossible to compare drug therapy with arthroscopy diagnosis and treatment, and a larger number of patients will be needed for research in the future.

Conclusion

Arthroscopic debridement is a minimally invasive treatment for shoulder tuberculosis, with complete debridement and short functional recovery time. Combined regular drug therapy can effectively treat shoulder tuberculosis with satisfactory results and is worthy of clinical promotion.

Abbreviations

TB Tuberculosis MTB Mycobacterium tuberculosis CRP C reactive protein

ESR: Erythrocyte sedimentation rate; MRI: Magnetic resonance imaging; VAS: Visual analogue scale; WHO: World Health Organization

Declarations

Acknowledgements

Not applicable

Funding

This research is supported by Cooperation program between Sichuan university and Luzhou municipal people's government (No. 2019cdlz-17)

Availability of data and materials

The datasets analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YWH contributed to the data analysis and interpretation, and manuscript draft.; ZW, ZL, JCL study design; PZ,XTD,LY,ZC enrolled the patients in the study,data collection and participated in the interpretation of the data.

Both authors read and approved the final manuscript.

Corresponding author

Correspondence to Zhong Li.

Ethics approval and consent to participate

This study was approved by the Ethics Committee of The Affiliate Hospital of Southwest University.

Consent for publication

Informed consent was obtained from all individual participants included in the study.

Competing interests

The authors declare that they have no competing interests

References

1. World Health Organization. Global Tuberculosis Report 2018. Geneva: World Health Organization; WHO/CDS/TB/2018.20. Available at: https://www.who.int/tb/publications/global_report/en/. Accessed October 2, 2018
2. Chen ST, Zhao LP, Dong WJ, *et al*. The Clinical Features and Bacteriological Characterizations of Bone and Joint Tuberculosis in China. *Sci Rep*. 2015; 5: 11084.
3. Agathangelidis F, Boutsiadis A, Fouka E, *et al*. Concomitant acromioclavicular and miliary tuberculosis. *BMJ Case Rep*. 2013; 2013:
4. Davies PD. The world-wide increase in tuberculosis: how demographic changes, HIV infection and increasing numbers in poverty are increasing tuberculosis. *Ann Med*. 2003; 35(4): 235-43.
5. Gardam M, Lim S. Mycobacterial osteomyelitis and arthritis. *Infect Dis Clin North Am*. 2005; 19(4): 819-30.
6. Jellis JE. Human immunodeficiency virus and osteoarticular tuberculosis. *Clin Orthop Relat Res*. 2002; 10.1097/00003086-200205000-00005(398): 27-31.

7. Dhillon MS, Aggarwal S, Prabhakar S, *et al.* Tuberculosis of the foot: An osteolytic variety. *Indian J Orthop.* 2012; 46(2): 206-11.
8. Chen ST, Zhao LP, Dong WJ, *et al.* The Clinical Features and Bacteriological Characterizations of Bone and Joint Tuberculosis in China. *Sci Rep.* 2015; 5: 11084.
9. Qian Y, Han Q, Liu W, *et al.* Characteristics and management of bone and joint tuberculosis in native and migrant population in Shanghai during 2011 to 2015. *BMC Infect Dis.* 2018; 18(1): 543.
10. Rodriguez-Takeuchi SY, Renjifo ME, Medina FJ. Extrapulmonary Tuberculosis: Pathophysiology and Imaging Findings. *Radiographics.* 2019; 39(7): 2023-37.
11. Parmar H, Shah J, Patkar D, *et al.* Tuberculous arthritis of the appendicular skeleton: MR imaging appearances. *Eur J Radiol.* 2004; 52(3): 300-9.
12. De Backer AI, Mortelé KJ, Vanhoenacker FM, *et al.* Imaging of extraspinal musculoskeletal tuberculosis. *Eur J Radiol.* 2006; 57(1): 119-30.
13. Sawlani V, Chandra T, Mishra RN, *et al.* MRI features of tuberculosis of peripheral joints. *Clin Radiol.* 2003; 58(10): 755-62.
14. Sanghvi DA, Iyer VR, Deshmukh T, *et al.* MRI features of tuberculosis of the knee. *Skeletal Radiol.* 2009; 38(3): 267-73.
15. Leonard MK, Blumberg HM. Musculoskeletal Tuberculosis. *Microbiol Spectr.* 2017; 5(2):
16. Duan X, Yang L. Arthroscopic management for early-stage tuberculosis of the ankle. *J Orthop Surg Res.* 2019; 14(1): 25.
17. Shen HL, Xia Y, Li P, *et al.* Arthroscopic operations in knee joint with early-stage tuberculosis. *Arch Orthop Trauma Surg.* 2010; 130(3): 357-61.
18. Tiwari V, Khan SA, Kumar A, *et al.* Functional improvement after hip arthroscopy in cases of active paediatric hip joint tuberculosis: a retrospective comparative study vis-à-vis conservative management. *J Child Orthop.* 2015; 9(6): 495-503.
19. Arora S, Sabat D, Maini L, *et al.* The results of nonoperative treatment of craniovertebral junction tuberculosis: a review of twenty-six cases. *J Bone Joint Surg Am.* 2011; 93(6): 540-7.
20. Teklali Y, El Alami ZF, El Madhi T, *et al.* Peripheral osteoarticular tuberculosis in children: 106 case-reports. *Joint Bone Spine.* 2003; 70(4): 282-6.

Tables


Due to technical limitations, table 1 is only available as a download in the Supplemental Files section.

Figures




1.

2.



3.

4.



5.

6.

Figure 1

A 38-year-old female patient with joint tuberculosis of left shoulder 1.Preoperative anteroposterior X - ray 2.Preoperative coronal MRI 3,4. Arthroscopic images; 5. postoperative pathological sections; 6.Coronal MRI 2 months after surgery.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table1.JPG](#)