Point-of-Care Ultrasound May Be Useful for Detecting Pediatric Intussusception at an Early Stage

Jeong-Yong Lee (✉ pedkorea@gmail.com)

Asan Medical Center <https://orcid.org/0000-0001-8471-901X>

Jung Heon Kim

Ajou University School of Medicine

Seung Jun Choi

Asan Medical Center

Jong Seung Lee

Asan Medical Center

Jeong-Min Ryu

Asan Medical Center

Research article

Keywords: Child; Diagnosis; Emergency Service, Hospital; Intussusception; Point-of-Care Systems; Ultrasonography

Posted Date: November 27th, 2019

DOI: <https://doi.org/10.21203/rs.2.17852/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Pediatrics on April 13th, 2020. See the published version at <https://doi.org/10.1186/s12887-020-02060-6>.

Abstract

Background: This study aimed to verify the usefulness of point-of-care ultrasound (POCUS) performed by pediatric emergency physicians for detecting intussusception at an early stage. **Methods:** This retrospective study included 1-month- to 6-year-old children who underwent POCUS for the detection of intussusception in the pediatric emergency department between December 2016 and February 2018. The criteria for performing POCUS were set to lower standards wherein POCUS was performed in the presence of any one of the following symptoms: intermittent abdominal pain, bloody stool, ≥ 2 causes of nonspecific abdominal pain, abdominal mass, vomiting, or lethargy. POCUS results were interpreted and categorized as “negative” or “suspicious,” and a radiologist performed confirmatory ultrasound in “suspicious” cases. **Results:** We analyzed 575 POCUS scans from 549 patients (mean age, 25.5 months). Among the 92 “suspicious” cases (16.0%), 70 (12.2%) were confirmed to have intussusception. POCUS showed 100% sensitivity, 95.6% specificity, and 97.8% accuracy. Patients with confirmed intussusception were mainly diagnosed in the early stages, with a mean symptom duration of 11.7 hours, and most patients (97.1%) were treated successfully via air enema reduction. Compared to the non-intussusception group, the intussusception group had more intermittent abdominal pain ($P < 0.001$), but less vomiting ($P = 0.001$); the other clinical features showed no intergroup differences. **Conclusion:** POCUS performed using the criteria set to lower standards by pediatric emergency physicians may be useful for detecting intussusception at an early stage, which may present with obscure clinical symptoms.

Introduction

Intussusception is a significant cause of intestinal obstruction in the pediatric population, and it necessitates a visit to the emergency department (ED) [1]. Intussusception is often alleviated by therapeutic air or liquid enema, but a delay in diagnosis may lead to intestinal gangrene, perforation, and peritonitis, which may require unexpected surgery [2]. Therefore, screening suspected cases of intussusception and detecting it at an early stage in the ED are important.

Detecting intussusception by evaluating the clinical features or plain abdominal radiographs may be challenging. The classic triad of abdominal pain/irritability, a palpable sausage-shaped mass, and bloody stool occurs in less than 40% of cases of intussusception, and these are often indistinguishable from the symptoms of acute gastroenteritis [3]. The use of plain abdominal radiographs interpreted by pediatric emergency physicians has a low sensitivity of 48% in the diagnosis of intussusception [4]. In contrast, intussusception can be accurately diagnosed using ultrasound [5]. However, requesting radiologist-performed ultrasound (RADUS) in all suspected cases may not only be time-consuming and inefficient for the ED workflow, but also exhausting for the on-duty radiologist.

Several studies have suggested that point-of-care ultrasound (POCUS) performed by pediatric emergency physicians could be a practical measure [6-8]. POCUS for detecting intussusception is relatively easy to learn and readily available in the ED [8-10]. The aim of this study was to verify the usefulness of POCUS performed by pediatric emergency physicians for detecting intussusception at an early stage.

Methods

This retrospective study included 1-month- to 6-year-old children with clinically suspected ileocolic intussusception, who underwent POCUS performed by pediatric emergency physicians in the pediatric ED of a tertiary-care university-affiliated hospital between December 2016 and February 2018. Patient data were collected by reviewing the electronic medical records. The institutional review board approved this present study and waived the requirement for informed consent.

POCUS

The criteria for performing POCUS were set to lower standards to detect intussusception at an early stage wherein POCUS was performed in the presence of any one of the following symptoms: intermittent abdominal pain/irritability, bloody stool, two or more causes of nonspecific abdominal pain/irritability, abdominal mass/distension, vomiting, or lethargy. The criteria were modified from the diagnostic criteria (a proposal) in the Japanese guidelines for the management of intussusception in children given in 2011 [11]. Patients transferred from another hospital with a confirmed diagnosis of intussusception were excluded. POCUS was performed by one of seven pediatric emergency physicians. They completed a 4-hour pediatric ultrasound course certified by the Korean Society of Pediatric Emergency Medicine, and had a mean experience of approximately 3 years (range, 0–6 years) of performing POCUS in the ED. All POCUS procedures were performed in a separate dedicated room by using the iE33 (Philips Ultrasound, Bothell, Washington) with a 3- to 11-MHz linear or 5- to 8-MHz curvilinear transducer. POCUS scans were interpreted and categorized as “negative” or “suspicious” for intussusception. A “negative” result on POCUS was strictly defined as absolutely no observable target or pseudokidney sign.

Management of patients

Patients with “suspicious” POCUS results were treated using fluid replacement and were transferred for RADUS. In addition, the pediatric emergency physicians requested RADUS in a few cases of “negative” POCUS results for evaluating conditions other than ileocolic intussusception. Intussusceptions confirmed using RADUS were evaluated for reducibility, and the radiologists subsequently attempted air enema reduction. Patients with successfully reduced intussusceptions were discharged after approximately 6 hours of observation in the ED for possible recurrence or other complications, whereas patients in whom more than two attempts of air enema reduction resulted in failure were considered for surgery. Patients in whom intussusceptions recurred after their discharge from the ED were counted separately. Patients requiring surgery or showing recurrent intussusception within 48 hours were admitted to the hospital. Most of the patients with “negative” POCUS results were discharged safely after ensuring they or their parents had a full understanding of the symptoms that would necessitate a revisit to the ED.

Data analysis

Patient data were categorized according to clinical features (age, sex, time of ED visit, previous ED visit within 24 hours, duration of symptoms, and clinical symptoms), POCUS and RADUS (interpretations and time from ED arrival to performing POCUS or RADUS), and management outcomes (air enema reduction, surgery, recurrence within 48 hours, admission, and ED observation time). The duration of symptoms was defined on the basis of the time of onset of abdominal pain/irritability (otherwise, lethargy or bloody stool) as determined by the parents; the time of onset was double-checked by a nurse as well as a physician with the time interval in all cases.

The POCUS results were analyzed using MedCalc version 18.11.6 (MedCalc software, Ostend, Belgium) and presented as 95% confidence intervals. Comparisons of clinical features between the intussusception and non-intussusception groups were analyzed using the χ^2 test for categorical variables and t test for continuous variables by using IBM SPSS Statistics for Windows, Version 20.0 (IBM Corp., Armonk, New York). $P < 0.05$ was considered statistically significant.

Results

Patient characteristics

We analyzed 575 POCUS scans from 549 patients. The mean age of the patients was 25.5 ± 15.9 months (range, 1–81 months) and 297 patients (51.7%) were male. The mean duration of time from arrival at the ED to undergoing POCUS was 54.7 ± 74.7 minutes. Among 92 patients (16.0%) with “suspicious” POCUS results, 70 (12.2%) were confirmed to have intussusception using RADUS. These patients mainly had intussusception in the early stages, showing symptoms for a mean duration of 11.7 ± 15.6 hours. Treatment of all patients with intussusception was attempted via air enema reduction, which was successful in 68 patients (97.1%). More than two attempts of air enema reduction resulted in failure in only 2 patients (one with suspected appendiceal intussusception requiring appendectomy and another with a delayed ED visit after 48 hours of abdominal pain), and they underwent surgical reduction. Table 1 shows the patient characteristics, including time management in the ED.

POCUS performance

The performance characteristics of POCUS are presented as a flowchart in Figure 1. Among the 92 patients with “suspicious” POCUS results, 22 were considered false positives, including 6 with an edematous ileocecal valve and 2 with small bowel intussusception. The remaining 483 patients with “negative” POCUS results were considered true negatives. Among them, RADUS was performed in 9 patients for evaluating conditions other than ileocolic intussusception (suspicious small bowel intussusception, unexplained ileus, or prominent intra-abdominal fluid collection), including 2 with small bowel intussusception and one with an intestinal duplication cyst, but no ileocolic intussusception. The

remaining 474 patients with “negative” POCUS results, who did not undergo RADUS, were discharged safely from the ED, except for 10 patients who were admitted to the hospital for supportive care; none of these patients were diagnosed with intussusception within 48 hours. The performance results of POCUS are summarized in Table 2.

Clinical features

The common clinical symptoms of the enrolled patients were abdominal pain/irritability (87.5%), vomiting (52.7%), loose stool (20.2%), fever (19.8%), and bloody stool (15.3%). The intussusception group demonstrated abdominal pain/irritability (92.9%), vomiting (37.1%), and bloody stool (21.4%). Compared to the non-intussusception group, the intussusception group showed more intermittent abdominal pain/irritability (58.6% vs. 33.9%, $P < 0.001$), but significantly less vomiting (37.1% vs. 58.2%, $P = 0.001$). Other clinical features, including bloody stool, lethargy, ED visit during night-time (7 pm to 8 am), or duration of symptoms, were not significantly different between the two groups (Table 3).

Discussion

This study verified the usefulness of POCUS performed by pediatric emergency physicians by applying criteria set to lower standards for detecting intussusception at an early stage. Our results demonstrated a relatively short symptom duration of 11.7 hours in intussusception cases, excellent performance outcome of POCUS, and the limitations of clinical features for distinguishing the intussusception and non-intussusception groups.

In this study, POCUS performed by pediatric emergency physicians seemed highly reliable (sensitivity, 100%; specificity, 95.6%; and accuracy, 97.8%) and useful (positive likelihood ratio, 23.0; and negative likelihood ratio, 0) in detecting intussusception. A previous study also showed a high degree of accuracy with 100% sensitivity and 94% specificity, even though it enrolled only 49 patients, which was a relatively small number [7]. Among the 92 patients with “suspicious” POCUS results in this study, 22 were confirmed as false positives, with a relatively low positive predictive value of 76.1%. This might be because the intussusception reduced spontaneously in 6 patients with edematous ileocecal valves, given the time interval between performing POCUS and RADUS. Moreover, the uncertain cases, wherein the patients were not definitely concluded as having no intussusception, were interpreted as indicating “suspicious” POCUS results; thus, the low positive predictive value was expected. In the ED setting, a missed diagnosis of intussusception possibly leads to serious consequences; therefore, the interpretation criteria of POCUS should be strict regarding the exclusion of intussusception as the diagnosis. In this study, RADUS was preferably requested in cases of “suspicious” POCUS results for confirming the diagnosis, as well as for evaluating reducibility or the possible presence of a pathologic lead point in cases of definite intussusception. We intended to use POCUS primarily as a utility for screening suspected intussusception rather than for confirming an exact diagnosis in the ED.

Proactively performing POCUS by applying criteria set to lower standards seems to facilitate the detection of intussusception at an early stage. Accordingly, this study aimed to perform POCUS by applying criteria set to lower standards in patients showing any one of the following symptoms: intermittent abdominal pain/irritability, bloody stool, two or more causes of nonspecific abdominal pain/irritability, abdominal mass/distension, vomiting, or lethargy. Consequently, the enrolled patients with intussusception were considered to be in the early stage, which presented symptoms for a much shorter mean duration of 11.7 hours instead of a duration of over 18.5 hours reported in previous studies [12-14]. The favorable treatment outcome in 97.1% of patients with successful air enema reduction also indirectly indicates that the patients were in early stages of intussusception; only 2 patients required surgical reduction. Moreover, the intussusception group in this study presented a lower prevalence of vomiting (37.1%) and bloody stool (21.4%) than did those in previous studies, which reported vomiting in 85% and bloody stool in up to 65% of patients [15]. According to the clinical course of intussusception, as intestinal obstruction progresses, abdominal pain appears first, followed by vomiting and bloody stool [3, 11]. Thus, our findings indicated that most patients with intussusception were in the early stage and had not yet developed vomiting.

Compared with the non-intussusception group, the intussusception group presented more intermittent abdominal pain ($P < 0.001$), but less vomiting ($P = 0.001$); however, the other clinical features were not significantly different. Only intermittent abdominal pain/irritability (58.6%) seems helpful in distinguishing intussusception in the early stages in a clinical setting, and this may suggest that detecting intussusception would still be challenging without performing POCUS.

This study has several limitations owing to its retrospective design. Most of the patients with “negative” POCUS results were not confirmed to have intussusception using RADUS; thus, the possibility of false-negative results exists. However, we strictly ruled out patients without intussusception, and none of them were proven to have intussusception within 48 hours of ED discharge. Moreover, defining the onset of intussusception on the basis of the duration of symptoms determined by the parents might be incorrect. However, we double-checked the presumed time in order to reduce any recall bias. We also did not consider the individual POCUS experience of pediatric emergency physicians, cost-effectiveness of POCUS, and satisfaction of the children or parents. Further prospective studies are required to address these issues.

In conclusion, POCUS may be performed by pediatric emergency physicians to detect intussusception. Furthermore, performing POCUS by applying criteria set to lower standards in the ED could help detect intussusception at an early stage, which may present with obscure clinical symptoms.

Declarations

Funding sources: The authors received no financial support for the research, authorship, and/or publication of this article.

Ethical approval: The institutional review board approval was obtained for this study.

Conflicts of interests: The authors have no potential conflicts of interest to disclosure.

Disclosure of previous presentation

Portions of this study were presented as an e-poster discussion at the 7th Congress of the European Academy of Pediatric Societies in Paris, France, on November 2, 2018.

References

1. Fischer TK, Bihrmann K, Perch M, Koch A, Wohlfahrt J, Kare M, et al. Intussusception in early childhood: a cohort study of 1.7 million children. *Pediatrics*. 2004;114:782-5.
2. Justice FA, Auldist AW, Bines JE. Intussusception: trends in clinical presentation and management. *J Gastroenterol Hepatol*. 2006;21:842-6.
3. Waseem M, Rosenberg HK. Intussusception. *Pediatr Emerg Care*. 2008;24:793-800.
4. Morrison J, Lucas N, Gravel J. The role of abdominal radiography in the diagnosis of intussusception when interpreted by pediatric emergency physicians. *J Pediatr*. 2009;155:556-9.
5. Carroll AG, Kavanagh RG, Ni Leidhin C, Cullinan NM, Lavelle LP, Malone DE. Comparative effectiveness of imaging modalities for the diagnosis and treatment of intussusception: a critically appraised topic. *Acad Radiol*. 2017;24:521-9.
6. Kim JH, Lee JY, Kwon JH, Cho HR, Lee JS, Ryu JM. Point-of-care ultrasound could streamline the emergency department workflow of clinically nonspecific intussusception. *Pediatr Emerg Care*. 2017; doi:10.1097/PEC.0000000000001283.
7. Lam SH, Wise A, Yenter C. Emergency bedside ultrasound for the diagnosis of pediatric intussusception: a retrospective review. *World J Emerg Med*. 2014;5:255-8.
8. Riera A, Hsiao AL, Langan ML, Goodman TR, Chen L. Diagnosis of intussusception by physician novice sonographers in the emergency department. *Ann Emerg Med*. 2012;60:264-8.
9. Eshed I, Gorenstein A, Serour F, Witzling M. Intussusception in children: can we rely on screening sonography performed by junior residents? *Pediatr Radiol*. 2004;34:134-7.
10. Gallagher RA, Levy JA. Advances in point-of-care ultrasound in pediatric emergency medicine. *Curr Opin Pediatr*. 2014;26:265-71.
11. Ito Y, Kusakawa I, Murata Y, Ukiyama E, Kawase H, Kamagata S, et al. Japanese guidelines for the management of intussusception in children, 2011. *Pediatr Int*. 2012;54:948-58.
12. Tareen F, Ryan S, Avanzini S, Pena V, Mc Laughlin D, Puri P. Does the length of the history influence the outcome of pneumatic reduction of intussusception in children? *Pediatr Surg Int*. 2011;27:587-9.
13. Herwig K, Brenkert T, Losek JD. Enema-reduced intussusception management: is hospitalization necessary? *Pediatr Emerg Care*. 2009;25:74-7.
14. Flaum V, Schneider A, Gomes Ferreira C, Philippe P, Sebastia Sancho C, Lacreuse I, et al. Twenty years' experience for reduction of ileocolic intussusceptions by saline enema under sonography control. *J Pediatr Surg*. 2016;51:179-82.

15. Mandeville K, Chien M, Willyerd FA, Mandell G, Hostetler MA, Bulloch B. Intussusception: clinical presentations and imaging characteristics. *Pediatr Emerg Care.* 2012;28:842-4.

Tables

Table 1. Patient characteristics

Characteristic	Value
Age, months, mean ± SD	25.5 ± 15.9
Sex, male, n (%)	297/575 (51.7%)
Duration of symptoms, hours, mean ± SD	9.9 ± 13.2
Door-to-POCUS time, minutes, mean ± SD	54.7 ± 74.7
ED length of stay, hours, mean ± SD	3.5 ± 3.0
Ileocolic intussusception, n (%)	70/575 (12.2%)
Successful air enema reduction, n (%)	68/70 (97.1%)
Surgical reduction, n (%)	2/70 (2.9%)
Recurrence within 48 hours, n (%)	12/70 (17.1%)
Admission, n (%)	14/70 (20%)
Door-to-RADUS time, minutes, mean ± SD	72.2 ± 56.7
ED observation time after reduction, hours, mean ± SD	6.9 ± 2.9

SD, standard deviation; POCUS, point-of-care ultrasound; ED, emergency department; RADUS, radiologist-performed ultrasound

Table 2. Performance results of point-of-care ultrasound

Indicator	Value (95% CI)
Sensitivity	100% (94.9%–100%)
Specificity	95.6% (93.5%–97.3%)
Accuracy	97.8% (96.3%–98.9%)
Positive predictive value	76.1% (67.9%–82.7%)
Negative predictive value	100% (100%–100%)
Positive likelihood ratio	23.0 (15.3–34.5)
Negative likelihood ratio	0 (0–0)

CI, confidence interval

Table 3. Comparison of clinical features between the intussusception and non-intussusception groups

	Intussusception group (n = 70)	Non-intussusception group (n = 505)	<i>P</i>
Age, mean $\pm$ SD, months	25.6 $\pm$ 16.8	25.5 $\pm$ 15.8	0.966
Sex, male, n (%)	41/70 (58.6%)	256/505 (50.7%)	0.216
Time of ED visit			
Night-time (7 pm–8 am)	36/70 (51.4%)	311/505 (61.6%)	0.104
Day-time (8 am–7 pm)	34/70 (48.6%)	194/505 (38.4%)	
Previous ED visit within 24 hours	7/70 (10%)	25/505 (5%)	0.084
Duration of symptoms, mean $\pm$ SD, hours	11.7 $\pm$ 15.6	9.7 $\pm$ 12.9	0.238
Abdominal pain/irritability	65/70 (92.9%)	438/505 (86.7%)	0.147
Intermittent	41/70 (58.6%)	171/505 (33.9%)	<0.001
Nonspecific	24/70 (34.3%)	267/505 (52.9%)	0.004
Bloody stool	15/70 (21.4%)	73/505 (14.5%)	0.129
Abdominal mass/distension	10/70 (14.3%)	64/505 (12.7%)	0.706
Vomiting	26/70 (37.1%)	294/505 (58.2%)	0.001
Lethargy	5/70 (7.1%)	51/505 (10.1%)	0.434
Fever	13/70 (18.6%)	101/505 (20.0%)	0.779
Loose stool	10/70 (14.3%)	106/505 (21.0%)	0.190

SD, standard deviation; ED, emergency department

Figures

Figure 1

Flowchart showing the performance characteristics of point-of-care ultrasound (POCUS) *Radiologist-performed ultrasound (RADUS) was performed for evaluating conditions other than ileocolic intussusception.