The Epidermal Microbiome Within An Aggregation of Leopard Sharks (*Triakis semifasciata*) Has Taxonomic Flexibility With Gene Functional Stability Across Three Time-Points

Michael Doane (✉ michael.doane@flinders.edu.au)

Sydney Institute of Marine Science <https://orcid.org/0000-0001-9820-2193>

Colton Johnson

San Diego State University

Shaili Johri

Stanford University Hopkins Marine Station

Emma N. Kerr

San Diego State University

Megan M. Morris

Lawrence Livermore National Laboratory

Ric Desantiago

San Diego State University

Abigail C. Turnlund

University of Queensland - Saint Lucia Campus: The University of Queensland

Asha Goodman

San Diego State University

Maria Mora

San Diego State University

Laís Farias Oliveira Lima

San Diego State University

Andrew P. Nosal

University of California San Diego Scripps Institution of Oceanography

Elizabeth A. Dinsdale

Flinders University

Research

Keywords: Microbiome, leopard shark, *Triakis semifasciata*, shark skin, next-generation sequencing

Posted Date: September 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-850465/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

**The epidermal microbiome within an aggregation of leopard sharks (*Triakis semifasciata*)
has taxonomic flexibility with gene functional stability across three time-points**

Michael P. Doane^{1#}, Colton Johnson^{2#}, Shaili Johri³, Emma N. Kerr¹, Megan M. Morris⁴, Ric
Desantiago², Abigail C. Turnlund⁵, Asha Goodman², Maria Mora², Laís Farias Oliveira Lima²,
Andrew P. Nosal^{6,7}, Elizabeth A. Dinsdale²

¹College of Science and Engineering, Flinders University, Bedford Park, South Australia,
Australia

²Department of Biology, San Diego State University, San Diego, California, USA

³Hopkins Marine Station, Stanford University, Pacific Grove, California, USA

⁴Laurence Livermore National Labs, Livermore, California, USA

⁵Australian Centre for Ecogenomics, University of Queensland, St Lucia, Queensland, Australia

⁶Department of Environmental and Ocean Sciences, University of San Diego, San Diego,
California, USA

⁷Scripps Institution of Oceanography, University of California – San Diego, La Jolla, California,
USA

#co-first authors

Michael P. Doane: 0000-0001-9820-2193

michael.doane@flinders.edu.au

Colton Johnson: 0000-0001-8895-6813

cjohnson1412@sdsu.edu

Shaili Johri: 0000-0001-7481-7548

shailij@stanford.edu

Emma N. Kerr

Emma.n.kerr@gmail.com

Megan M. Morris: 0000-0002-7024-8234

morris81@llnl.gov

Ricardo Desantiago

rdesantiago@sdsu.edu

31 Abigail C. Turnlund
32 **acturnlund@gmail.com**
33 Asha Z. Goodman
34 **azgoodman@sdsu.edu**
35 Maria F. Mora
36 **moramariaf21@gmail.com**
37 Laís Farias Oliveira Lima: 0000-0002-7616-3637
38 **laisfolima@gmail.com**
39 Andrew P. Nosal: 0000-0002-1815-9269
40 **anosal@ucsd.edu**
41 Elizabeth A. Dinsdale: 0000-0002-2177-203X
42 **elizabeth.dinsdale@flinders.edu.au**
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58

The epidermal microbiome within an aggregation of *Triakis semifasciata* (leopard shark) has taxonomic flexibility with gene functional stability across three time-points

Michael P. Doane^{1#}, Colton Johnson^{2#}, Shaili Johri³, Emma N. Kerr¹, Megan M. Morris⁴, Ric Desantiago², Abigail C. Turnlund⁵, Asha Goodman², Maria Mora², Laís Farias Oliveira Lima², Andrew P. Nosal^{6,7}, Elizabeth A. Dinsdale²

¹College of Science and Engineering, Flinders University, Bedford Park, South Australia, Australia

²Department of Biology, San Diego State University, San Diego, California, USA

³Hopkins Marine Station, Stanford University, Pacific Grove, California, USA

⁴Laurence Livermore National Labs, Livermore, California, USA

⁵Australian Centre for Ecogenomics, University of Queensland, St Lucia, Queensland, Australia

⁶Department of Environmental and Ocean Sciences, University of San Diego, San Diego, California, USA

⁷Scripps Institution of Oceanography, University of California – San Diego, La Jolla, California, USA

Abstract

Background: The epidermis of Chondrichthyan fishes consists of dermal denticles with production of minimal but protein rich mucus that influence the attachment and biofilm development of microbes, facilitating a unique epidermal microbiome. Here, we use metagenomics to provide the taxonomic and functional characterization of the epidermal microbiome of the *Triakis semifasciata* (leopard shark) across three time-points to identify links between microbial groups and host metabolism. Our aims include 1) describing the variation of microbiome taxa over time and identify those members which are recurrent (present across all time-points, 2) investigating the relationship between the recurrent and flexible taxa (those which are not found consistently across time-points, 3) describing the functional compositions of the microbiome which may suggest links with the host metabolism; and 4) identifying whether the metabolisms are share across microbial genera or found in specific taxa.

Results: Microbial members of the microbiome showed high similarity between all individuals (average similarity: 82.74) with relative abundance of those members varying across years, suggesting flexibility of taxa in the microbiome. One hundred and eighty-eight genera were

identified as recurrent, including *Pseudomonas*, *Erythrobacter*, *Alcanivorax*, *Marinobacter* and *Sphingopxis* being consistently abundance across time-points, while *Limnobacter* and *Xyella* exhibited switching patterns with high relative abundance in 2013, *Sphingobium* and *Sphingomona* in 2015, and *Altermonas*, *Leeuwenhoekiella*, *Gramella* and *Maribacter* in 2017. Of the 188 genera identified as recurrent, the top 19 relative abundant genera forming three recurrent groups. The microbiome also displayed high functional similarity between individuals (average similarity: 97.65) with gene function composition being consistent across time-points.

Conclusion: These results show that while presence of microbial genera exhibit consistency across time-points, their abundances do fluctuate. Functions however remain stable across time points; thus, we suggest the leopard shark microbiomes exhibit functional redundancy. We hypothesize this may be the result of the host's epidermal attributes structuring the microbiome. In addition, we show the co-existence of many microbial genera that carry genes which may enable the microbes to use the nutrients provided by the elasmobranch's metabolism.

Keywords: Microbiome, leopard shark, *Triakis semifasciata*, shark skin, next-generation sequencing

Introduction

The epidermis of marine organisms is biologically active (i.e. not keratinised like most terrestrial organisms)(Meyer and Seegers 2012), serving as the first line of defence against environmental influences. Along with providing broad protections, the skin of elasmobranchs interacts with the microbes from the surrounding environment, forming a specific microbiome (Doane et al., 2017). The epidermal microbiome of six elasmobranchs, including thresher sharks (*Alopias vulpinus*), whale sharks (*Rhincodon typus*), leopard sharks (*Triakis semifasciata*), blacktip reef sharks (*Carcharhinus melanopterus*), and round rays (*Urobatis halleri*) confirmed that the elasmobranch epidermal microbiomes are species-specific (Doane et al. 2017; Doane et al. 2020; Pogoreutz et al. 2019). However, the selection and maintenance of the microbiome on sharks remains an outstanding question, and may be the result of several processes, including 1) biophysical properties of the epidermis, 2) metabolism of the shark, and 3) interspecific interactions between the microbes.

The biophysical properties of the chondrichthyan fish epidermis are uniquely covered with dermal denticles that improve hydro-dynamism (Lang 2020). Dermal denticles have ridges and troughs arranged in overlapping patterns that alter the hydrodynamic properties of water close to the epidermis (Sullivan and Regan 2011b; Zhang et al. 2011; Wen et al. 2015) and this biophysical property potentially affects microbial growth. In synthetic models mimicking shark skin, two bacteria, *Escherichia coli* and *Staphylococcus aureus*, showed differential growth patterns compared to a smooth surface. *E. coli* had high attachment rates to the structure surface, whereas *S. aureus* was not able to attach (Chien et al. 2020), therefore, suggesting that only specific types of microbes are able to attach to the shark skin. Further, modeling of microbial community dynamics shows that high surface structural complexity can reduce competition between microbes that result in chaotic spatial species distribution pattern on the surface (Lowery and Ursell 2019). Therefore, the high micro-structured nature of the shark skin surface, may produce similar dynamics in the microbiome. Further, the epidermis of Chondrichthyan fish is characterized by a lower level of secretory cells compared with Actinopterygian fishes (Meyer and Seeger 2012) and the product is a thin heterogeneous outer mucous layer (5–8 μm thickness; pH between 6 and 7). The mucus contains high levels of proteins, with only one protein being identified, which is lectin pentraxin (Tsutsui et al. 2014) and has moderate to high disulphide concentrations that is suspected to provide mechanical stabilization of the mucus coating (Meyer, Seegers, and Stelzer 2007).

The metabolism of Elasmobranchs may also affect the functions of the microbiomes in the form of secretions being transported by the skin mucus. Elasmobranchs rely on ketone bodies, fatty acids, carbohydrates, and amino acids for energy production (Speers-Roesch and Treberg 2010). Amino acids are ketogenic precursors, oxidative fuel, and nitrogen suppliers (Speers-Roesch and Treberg 2010). Nitrogen metabolism is also important due to the high amount of urea synthesised for osmoregulation, along with trimethylamine N-oxide (TMAO), which could also be transported to the skin surface via mucus production (Ballantyne 1997; Speers-Roesch and Treberg 2010). Sharks bioaccumulate toxins and heavy metals from lower trophic levels (Escobar-Sanchez, Galvan-Magana, and Rosiles-Martinez 2010; Maz-Courrau et al. 2012) and have been shown to exhibit maternal off-loading to their offspring in mucus surrounding

developing embryo (Lyons and Lowe, 2013). Therefore, metabolic and harmful contaminants are being leached from elasmobranchs in their mucus and if this occurs at the skin surface may structure the skin microbiome. The microbiome signature from thresher sharks (*Alopias vulpinus*) suggests potential heavy metal concentration at the skin interface signified by high abundance of heavy metal resistance genes (Doane et al., 2017).

Interactions between the microbes on the surface and those from the surrounding water are also expected to affect the characteristics of the microbiome. The low levels of mucus secreted by the sharks and minimal biofilm development that occurred on the synthetic models of the shark skin, compared with the pronounced biofilm development on the smooth surface (Chien et al. 2020), suggests that microbial growth on the epidermis is suppressed, and the turnover of microbial communities may be high. Conversely, stability of a microbiome is important for the host's health and functioning, and rapid or unexpected variations in microbiome taxonomy leads to a state of dysbiosis, which can result in a decline in organismal health (Costello et al. 2012; Egan and Gardiner 2016). Therefore, an understanding of how the microbiome varies through time is important, but there are limited investigations of microbiome stability over time in the marine environment. A few examples include the epidermis microbiome of humpback whales (*Megaptera novaeangliae*) and Atlantic cod (*Gadus morhua*) which varied seasonally (Bierlich et al. 2018; Wilson, Danilowicz, and Meijer 2008), and similar seasonal changes occurred in the microbiome of the stony coral, *Montastrea faveolata* (Kimes et al. 2013). While the respiratory microbiome of bottlenose dolphin species (*Tursiops truncatus* and *Tursiops aduncus*) maintained taxonomic stability over two months, but individual dolphins had significantly different microbiome taxonomic structures (Lima et al. 2012). In an analysis of 8 months, captive dolphins had an individual signature of their lung microbiome and shared between 17 – 41 % of the microbiome with other individuals (Vendl et al. 2021). In contrast, 77.6 % of the microbes in the epidermal microbiome of thresher sharks (*A. vulpinus*) were shared across individual sharks (Doane et al. 2017), however, this study was only conducted at one timepoint. Therefore, whether the epidermal microbiomes of marine vertebrate hosts are temporal stable or flexible remains an outstanding question in science.

Here we analyse both the taxonomic make-up and functional capabilities of the leopard shark (*Triakis semifasciata*) epidermal microbiome over five years (three sampling time-points) using the following objectives: 1) to describe the variation of microbiome taxa over time and identify those members which are recurrent (present across time-points, 2) to investigate the relationship between the recurrent and flexible taxa (those which are not found consistently across time-points, 3) to describe the functional compositions of the microbiome which may suggest links with the host metabolism; and 4) to identify whether the metabolisms are share across microbial genera or found in specific taxa.

Materials and Methods

Epidermal Microbiome Sampling

Leopard sharks (*Triakis Semifasciata*) sharks form consistent aggregations in late summer, early fall at near-shore locations (Nosal et al. 2014; Nosal et al. 2013) along the coast of California making this species a good candidate to investigate temporal microbiome dynamics of a wild animal. Sharks in this study were sampled from La Jolla Cove, California, USA (32°51'12" N, 117°15'48" W). We targeted at least three sharks per year, and captured eight sharks in 2013, three in 2015, and seven in 2017, thus the study was conducted over a five-year period. The sharks were caught with a handline on baited barbless circle hooks and brought onboard a small skiff for processing using a scoop net; all sharks were released after processing. Microbiome samples from the sharks were collected using a microbial collection tool, which flushes the epidermis with ~250 ml of sterile seawater and dislodges the microbes. The microbial slurry is collected into the back end of the microbial tool, as we have conducted previously (Doane et al. 2017; Doane et al. 2018; Doane et al. 2020; Johri et al. 2019; Lima et al. 2020). Each epidermal microbiome sample was taken from the flank region below the dorsal fin and above the lateral line. The collected microbiome samples were filtered through a 0.22 µm Sterivex filter (Millipore, Inc.) and stored in a -20 °C freezer until processing.

DNA Extraction and Sequencing

Microbiome DNA was extracted using a cell lysis and spin-column filtration method by Macherey Nagel Tissue Kit. The purified DNA was prepared for random shot-gun sequencing using the standard protocol from Swift 2S Plus Kit (Swift Biosciences). The DNA library was

sequenced using a Mi-Seq Illumina sequencer with Mi-Seq v3 Reagent Kit. Samples were bar-coded and the *T. semifasciata* microbiomes were mixed in with a range of microbiome samples and run on several sequencing runs.

Metagenome Annotation

The sequenced metagenomes were processed through PRINSEQ $\pm\pm$ software for quality control, removing sequences less than 70 bp, those that had a single N, or a quality score below 25. MID tags were also removed with PRINSEQ $\pm\pm$ (Cantu, Sadural, and Edwards 2019; Schmieder and Edwards 2011). The two fasta files were paired using FLASH software (Magoc and Salzberg 2011). After pairing, metagenomes were uploaded into MG-RAST Version 4.0.3, which was used to provide taxonomic and functional gene annotations of the metagenomes using standard cut-offs (Aziz et al. 2008; Meyer et al. 2008). The sequences with the highest bit score were reported. The taxonomic annotation outputs were filtered in MG-RAST to only include those from Bacteria and Archaea domains. Viruses and Eukaryotes were not used in the analysis because they were underrepresented in the metagenome. The functional gene characterization was obtained from the SEED platform (Overbeek et al. 2005), similar to previous analysis (Dinsdale, Edwards, et al. 2008; Dinsdale, Pantos, et al. 2008).

Quantitative evaluation of leopard shark microbiomes

To identify whether the taxonomy of the microbiome remained consistent or varied over time, a Permanova on the relative abundance of each taxon level was conducted across years (Anderson 2017). All data was fourth root transformed to reduce abundance variation among samples. A simpler analysis was used to identify the taxa contributing to the dissimilarity between years (Clarke, Somerfield, and Gorley 2008). A principal coordinate analysis was performed on the taxonomic data to visualize the variation in the structure of the microbiomes over the years. β -diversity was identified using a Bray-Curtis analysis with 100 - similarity index, where 0 = high overlap, 100 = no overlap. Last, a PERMDISP analysis compared the distribution of microbial taxa across each year around the year's group centroid (Clarke, Somerfield, and Gorley 2016; Anderson 2006).

We identified a recurrent group of microbes (or core) as those genera that were present on every individual shark, on a presence/absence basis and non-recurrent microbes, as all other genera that were an occasional member of the microbiome. The 19 most abundant recurrent and non-recurrent microbial taxa (abundances calculated within the recurrent/non-recurrent datasets, separately) were compared with a Bray-Curtis similarity analysis to identified groups of genera that had a similar proportional abundance across years. A Pearson correlation was conducted to identify positive or negative correlations between these taxonomic groupings. The output of the Pearson correlation was organized into a heatmap and combined with the Bray-Curtis dendrogram. The analysis identified negative and positive correlations between the microbial groups that live on the shark epidermis.

To identify whether the functional potential of the shark epidermis microbiome remains stable or varies temporally, a PERMANOVA was conducted to compare variation in functional components across years. The SEED provides a hierarchal description including major metabolisms, metabolic pathways, gene clusters and gene functions (Overbeek et al. 2005). The SIMPER analysis identified potential functions that contributed to the dissimilarity between years. A principal coordinate analysis was performed to visualize the variation in the functional potential of the microbiomes over the years and a PERMDISP analysis compared the distribution of functional potential of *T. semifasciata* epidermal microbiome across years. All analyses were performed in PRIMER Version 6.1.15 & PERMANOVA± Version 1.0.5 from PRIMER-E (2012). GraphPad Prism Version 8.3.1 was used to visualize the data.

We identified the functional potential of the 17 most proportionally abundant genera using the Krona plug-in (Ondov, Bergman, and Phillippy 2011) in MG-RAST (Aziz et al. 2008; Meyer et al. 2008). Only 17 genera were use because they occurred in sufficient abundance to explore the functions of the genera. The sequences were further investigated using Kyoto Encyclopedia of Genes and Genomes (KEGG) Orthology (KO) database's KEGG mapper plug-in on MG-RAST (Kanehisa et al. 2012; Kanehisa et al. 2016), which provided identification of metabolic and biochemical pathways. The proportion of sequences within each genus for each functional group was calculated as the number of sequences within a genus divided by the total number of sequences within the function across the 17 genera. The proportion of sequences associated with

each function in each genus was categorized into one of four quartiles: < 75 %, 50 – 75 %, 25 – 50 %, 1 – 25 %, and 0 % and visualized using a heatmap. The heatmap was used to visualize the connection between the microbes and the sharks over time and identified whether there was switching of specific functional roles between genera in the microbiome.

Results

Temporal Taxonomy of the Epidermal Microbiome

The epidermal microbiomes were constructed from 18 *T. semifasciata*, including seven individuals in 2013, three in 2015 and eight in 2017 caught in La Jolla, California. Metagenomes ranged from 312,135 – 2,347,886 sequences that had 89,080,935 – 605,042,370 base pairs (Table 1). Of those sequences identified as protein coding, 24.01 – 83.74 % were classified as unknown sequences, which could not be matched to database (Table 1). The abundant phyla represented in the *T. semifasciata* epidermal microbiome were Proteobacteria (78.50 ± 2.02 S.E. %), Bacteroidetes (15.83 ± 2.34 %), and Actinobacteria (2.09 ± 0.30 %). Within the Proteobacteria the classes Gammaproteobacteria (38.99 ± 2.28 %) and Alphaproteobacteria (31.32 ± 2.20 %) were evenly represented, with Betaproteobacteria (7.25 ± 0.81 %) and Deltaproteobacteria (0.81 ± 0.05 %) in lower relative abundance (SI Figure 1). Within the Bacteroidetes, the Flavobacteria class made up 12.08 % (± 2.15) of the metagenome and Actinobacteria class accounted for 2.40 % (± 0.30) (SI Figure 2).

Across the 18 leopard sharks, 597 genera made up the epidermal microbiome. Highly represented genera present within the microbiomes included *Pseudomonas* (9.85 ± 1.81 %), *Erythrobacter* (5.88 ± 0.69 %), *Leeuwenhoekiella* (2.52 ± 0.72 %), and *Limnobacter* (1.51 ± 0.42 %) (Figure 1). The *T. semifasciata* microbiomes collected across the three time-points showed high taxonomic similarity between all individuals, with average Bray-Curtis similarity of 82.74 (0 = no overlap, 100 = total overlap) among all pairs of samples. The similarity of the genera on the sharks across time-points suggests a selective process is occurring and that the shark epidermis has a specialized microbiome (Figure 1). The 19 most relative abundant genera (except for *Providencia* 2.29 ± 0.94 %) were recurrent over sampling periods, but the proportional abundance varied throughout the years (Figure 3). The PERMANOVA analysis identified significant differences in the proportional abundances of all taxonomic levels between years (PERMANOVA: Phylum, Pseudo-F_{df = 2} = 3.696, P(permanova) = 0.001; Class, Pseudo-F_{df = 2} =

3.628, $P(\text{perm}) = 0.001$; Order, Pseudo- $F_{df=2} = 3.222$, $P(\text{perm}) < 0.01$; Family, Pseudo- $F_{df=2} = 2.762$, $P(\text{perm}) < 0.01$; Genus, Pseudo- $F_{df=2} = 2.967$, $P(\text{perm}) = 0.049$). The microbial genera of the *T. semifasciata* epidermal microbiomes formed distinct clusters for each year in the principal coordinate analysis, which explained 64.7 % of the variation in the first two axes (Figure 2). A few outliers identified in 2013 and 2017 may be associated with variation in an individual shark's physiology, i.e., such as pregnancy.

The SIMPER analysis identified a similarity coefficient of 83.05 between 2013 & 2015; 80.29 for 2013 & 2017; and 84.64 for 2015 & 2017 (Bray-Curtis similarity index, 100 = similar, 0 = no overlap). Genera of the Gammaproteobacteria class contributed the most to the dissimilarity between the years, followed by genera in the Alphaproteobacteria and Betaproteobacteria, and the Flavobacteria class from the Bacteroidetes phylum (Table 2). The variation in the relative abundance of *Alcanivorax* and *Alteromonas* contributed the most to the dissimilarity across years.

Last, we explored the variance within years of microbial taxonomy with PERMDISP analysis. Each year the microbiome showed a similar amount of variation within the microbiomes at each taxonomic level (PERMDISP: Phylum, $F_{df=2, 15} = 2.120$, $P(\text{perm}) > 0.1$; Class, $F_{df=2, 15} = 2.439$, $P(\text{perm}) > 0.1$; Order, $F_{df=2, 15} = 1.513$, $P(\text{perm}) > 0.5$; Family, $F_{df=2, 15} = 2.019$, $P(\text{perm}) > 0.1$; Genus, $F_{df=2, 15} = 2.142$, $P(\text{perm}) > 0.05$). The microbiomes from 2013, 2015, and 2017 showed a dispersion of 11.7 (± 1.8), 6.0 (± 0.8), and 10.1 (± 1.3) respectively. In summary, the individual sharks shared many microbial genera suggesting a consistent taxon across the years, with flexibility in the proportional abundance.

Temporally Recurrent Microbes

Microbial genera present in all 18 metagenomes, were categorized as 'recurrent' which included 188 of 597 genera. The stability was considered further by comparing the proportional abundance of sequences within the 188 genera across sampling periods and identified a significant difference (PERMANOVA: Pseudo- $F_{df=2, 15} = 10.462$, $P(\text{perm}) = 0.001$). The variability in the recurrent microbial genera, measured by PERMDISP (PERMDISP: $F_{df=2, 15} = 0.029$, $P(\text{perm}) = 0.983$) indicated no difference across years with 2013, 2015, and 2017 showing a dispersion of 4.09 (± 0.47), 4.07 (± 0.57), 3.974 (± 0.22), respectively. Collectively the

fluctuations in the dominate genera along with similar variability across time suggest a flexible but regulated recurrent groups of microbes in the *T. semifasciata* microbiome.

Relationship between recurrent and non-recurrent genera

In the clustering analysis, the 19 dominant recurrent microbes formed three groups and the 19 non-recurrent microbes formed five separate groups (Figure 4). The recurrent group A consisted of *Pseudomonas*, *Erythrobacter*, *Alcanivorax*, and *Marinobacter*, which showed an even proportional abundance across the five years and with *Erythrobacter* and *Marinobacter* having higher proportional abundance in 2013. The recurrent group B consisted of seven genera which showed higher proportional abundance in 2017 and included *Leeuwenhoekiella*, *Altermonas*, *Gramella*, *Zunongwangia*, *Pseudoaltermonas*, *Flabacterium*, and *Maribacter*. The recurrent group C consisted of six genera with lower proportional abundance, including *Burkholderia*, *Novoshingobium* and *Sphingopyxis*, which showed even distribution across the years, *Rugeria*, which showed proportional higher abundance in 2013 and 2017 and *Sphingobium* and *Sphingomonas*, which showed higher proportional abundance in 2015. The non-recurrent (top 19 most relative abundant genera not occurring across all sampling points) microbial group A consisted of three genera with lower proportional abundance in 2017, including *Propionibacterium*, *Pirellula*, and *Plesiocystis*. The non-recurrent microbial group B consisted of two genera that showed low but even proportional abundance across all years and included *Aromatoleum* and *Chromobacterium*. The non-recurrent microbial group C only consisted of *Dokdonia*, which displayed higher proportional abundance in 2017. The non-recurrent group D included *Acidiphillum*, *Azorhizobium*, *Gluconobacter*, and *Sphingobacterium* and the first two genera displayed low even proportional abundances throughout all years *Gluconobacter* and *Sphingobium* showed higher proportional abundance in 2017. The non-recurrent microbial group E, including *Leadbetterella*, *Klebsiella*, *Neisseria*, *Oligotropha*, *Thauara*, and *Frankia*. *Leadbetterella* displayed its highest proportional abundance in 2015. *Klebsiella*, *Neisseria*, and *Thauara* showed even proportional abundance across all years. *Oligotropha* and *Frankia* displayed lower proportional abundances at all years (Figure 4). There also occurred a clustering group termed ‘mixed’ because genera included those which were classified as both recurrent and non-recurrent including *Endoritia*, *Limnobacter*, *Xylella*, and *Providencia* genera.

We then investigated the correlative relationship of recurrent and non-recurrent groups (Figure 4). The heatmap showed that *Erythrobacter* and *Marinobacter* (recurrent group A) were negatively correlated with all microbes in recurrent group B. Whereas *Pseudomonas* and *Alcanivorax* (recurrent group A) showed positive correlations with recurrent group B. Recurrent group A, however, was positively correlated with non-recurrent group A and B. In contrast, recurrent group B was negatively correlated with non-recurrent group A and B but strongly associated with non-recurrent group C, *Dokdonia*. Recurrent group C was positively correlated with non-recurrent groups A, B, D, E, and the mixed group (Figure 4). These associations between taxa will be investigated further in the functional analysis to identify metabolisms that may drive the microbiome composition.

Functional potential of the shark epidermis microbiome

Annotated microbial sequences were categorized into 26 major metabolic groups, 153 metabolic pathways, 949 functional gene clusters, and 5,358 functional genes. The gene function profiles were highly stable across the three time-points as evidenced by the high functional similarity between individual sharks (mean 97.65, Bray-Curtis similarity index). There was not a significant difference between time-points in the functional potential of the microbiome (PERMANOVA: major metabolic groups, Psuedo-F_{df = 2, 15} = 1.8, P(perm) > 0.05; metabolic pathways, Psuedo-F_{df = 2, 15} = 0.525, P(perm) > 0.5; functional gene clusters, Pseudo-F_{df = 2, 15} = 1.182, P(perm) > 0.1). The similarity of the functional potential was shown by the single cluster on the principal coordinate analysis that explained 64.4 % of the variation in the first two axes (Figure 5).

The major metabolic groups included carbohydrates (mean 10.96 ± 0.32 %); amino acids & derivatives (9.17 ± 0.19 %); protein metabolism (7.88 ± 0.11 %); and fatty acids, lipids, & isoprenoids (2.61 ± 0.04 %) (SI Figure 3). In nitrogen metabolism (1.23 ± 0.11 %), there were 11 functional gene clusters in the microbiome including ammonia assimilation (0.51 ± 0.01 %), nitrate and nitrite ammonification (0.38 ± 0.001 %), and denitrification (0.10 ± 0.02 %), but few sequences were associated with nitrogen fixing genes (0.02 ± 0.007 %) suggesting high levels of organic nitrogen that are present on the shark. Urea decomposition (0.27 ± 0.02 %) was present and there were a few sequences associated with trimethylamine N-oxide (TMAO) reductase (0.004 ± 0.001 %). The functional gene clusters that perform fatty acid biosynthesis and

metabolism are fatty acid biosynthesis FASII (0.72 ± 0.02 %), fatty acid degradation regulons (0.22 ± 0.02 %), and fatty acid metabolism (0.13 ± 0.01 %). The functional gene clusters that complete protein biosynthesis and degradation are ATP dependent proteolysis in bacteria (0.37 ± 0.01 %), bacterial proteasome (0.30 ± 0.02 %), protein degradation (0.20 ± 0.01 %), nucleolar protein complex (1.01 ± 0.05 %), aminopeptidase (0.18 ± 0.01 %), metallocarboxypeptidase (0.04 ± 0.006 %), and bacterial ribosome LSU (0.38 ± 0.02 %). A high proportional abundance of flagellar motility (1.27 ± 0.06 %) and lower proportional abundance of bacterial chemotaxis (0.36 ± 0.02 %) is consistent with low biofilm formation predicted on the shark epidermis. There was a high proportional abundance of functions associated with resistance to antibiotics and toxic compounds (5.08 ± 0.13 %) and cobalt-zinc-cadmium resistance had a mean of 1.58 ± 0.12 %, multidrug resistance efflux pumps (0.85 ± 0.05 %), copper homeostasis (0.69 ± 0.05 %), and the production of exopolysaccharide biosynthesis (EPS) (0.058 ± 0.008 %) and Rhamnolipids produced by *Pseudomonas* (0.005 ± 0.002 %) that are used in heavy metal detoxification, consistent with bioaccumulation of heavy metals and toxins by the large-bodied sharks.

Microbial gene function across genera

The 17 dominant recurrent genera were used to investigate whether functional genes were equally dispersed across the genera or confined to a small subset of microbes (Figure 6). *Pseudomonas*, had the highest proportion of sequences annotated to all metabolisms, including those involved in nitrate and nitrite ammonification and denitrification, and in association with *Alcanivorax*, also conducted ammonia assimilation. These two genera were positively correlated (Figure 6) with most other microbes, suggesting these were key functions in microbiome growth. In comparison, *Marinobacter* also specialized in these two functions plus urea decomposition and was negatively correlated *Pseudomonas* and *Alcanivorax*. *Ruegeria* and *Novosphingobium* were the only other genera that had a high proportion abundance of genes associated with urea decomposition, which may have caused the negatively correlated with recurrent group B. The genera within recurrent group B had a high proportion of genes associated with carbon metabolism, with *Gramella* having a high proportion of genes within TCA cycle, Glycolysis and Serine cycle. *Marinobacter* has a high proportion of genes in Glycolysis, Denitrification, and fatty acid biosynthesis. Many genera, including *Pseudomonas*, *Alcanivorax*, *Erythrobacter*, *Leeuwenhoekiella*, *Altermonas*, *Gramella* and *Flavobacterium*, had many sequences associated

with protein biosynthesis and degradation, and fatty acid biosynthesis, which are associated with the shark ketosis dominated metabolism. Genera within recurrent group C, including *Ruegeria*, *Sphingopyxis* and *Novosphingobium* had a high proportion of genes associated with Denitrification and fatty acid biosynthesis. All 17 genera, except 1, had sequences that matched the copper, cobalt and cambium resistance and copper homoeostasis genes. The abundance of these functions across most genera in the metagenomes suggests the important heavy metal tolerance or manipulation for microbes living on the surface of sharks.

Discussion

The *T. semifasciata* epidermal microbiome was consistent between individual sharks with many genera shared across the three time-points spanning 5 years, one of the longest studies of marine vertebrate microbiomes to date. A recurrent set of 188 microbial genera were present on all individuals. However, there was flexibility in the proportional makeup of the microbiome across years, but the microbiome variability was consistent across years, suggesting similar ecological dynamics are occurring from year to year, despite some turnover in microbial genera. Regardless of the proportional abundance of each microbial genera, the functional potential was constant, suggesting that the epidermal microbes were responding to the unique environment of the *T. semifasciata* epidermis, including the dermal denticles that reduce biofilm build-up and potentially elements secreted via the minimal mucus associated with shark's metabolism.

Leopard shark microbiome exhibits specificity and flexibility

The *T. semifasciata* epidermal microbiome has a high relative abundance of *Pseudomonas*, *Erythrobacter*, *Alcanivorax*, and *Marinobacter*; whereas the thresher shark (*Alopias vulpinus*) epidermal microbiome had a high relative abundance of *Pseudoalteromonas*, *Erythrobacter*, *Idiomarina*, and *Limnobacter* (Doane et al. 2017); and the *Carcharhinus melanopterus* epidermal microbiome a high relative abundance of *Psychrobacter*, *Pseudoalteromonas*, *Rhodobacter*, and *Alteromonas* (Pogoreutz et al. 2019). While some genera were shared across shark species, the proportional abundance differed, suggesting that each elasmobranch microbiome has distinct characteristics which select for differing microbiome members.

The highly shared microbiome across individual sharks within a timepoint and flexible communities across timepoints coupled with evidence that species microbiomes are distinct may be caused by the dermal denticles (Chien et al. 2020; Dillon, Norris, and O'Dea 2017) and metabolites secreted in the skin mucus as a result of the sharks metabolism. Dermal denticles alter the hydrodynamic properties of water close to the epidermis, reducing drag on the shark (Sullivan and Regan 2011a) and micro-organism growth (Wen et al. 2015; Zhang et al. 2011). A study that tested the recruitment of two lab microbial species found that only one species was able to successfully recruit, and biofilms did not develop (Chien et al. 2020; Kim et al. 2012). Biofilm development is dependent on quorum sensing, which is interrupted on rough surfaces (Kim et al. 2012). We showed that specific microbial taxa recruit to the epidermis of *T. semifasciata* and the proportion of motility genes are relatively high compared to the chemotaxis genes which are proportionally low in abundance in the leopard shark microbiomes. These features are consistent across the microbiome of four shark species (Doane et al. 2017; Doane et al. 2020).

Modelling of microbial community dynamics of surfaces with structural complexity, like the conditions on a sharkskin, potentially suggests there is lower interspecific competition, which results in a chaotic spatial distribution pattern of microbial species (Lowery and Ursell 2019). The *T. semifasciata* epidermal microbiome supports the modelling where 188 genera coexisted on all sharks across the three years, with flexibility in the relative abundance of those genera, suggesting that the epidermis may provide multiple niches for microbial taxa to survive. An emergent feature of the modelling is that competing genera do not go extinct (Lowery and Ursell 2019), and we show that many microbial species co-occurred on the sharkskin across the three time-points. The co-occurrence or positive correlations within the microbiome suggest niche partitioning and sharing of resources with several genera exhibiting abundance relationships (Figure 4). For example, *Pseudomonas* and *Alcanivorax* showed positive correlations with recurrent group B members, including *Leeuwenhoekiella*, *Altermonas*, *Pseudoaltermonas*, *Flabacterium*, and *Maribacter*, suggesting that these two genera could share resources. *Erythrobacter* and *Marinobacter* were negatively correlated with all microbes in recurrent group B, suggesting strong competition; this group was however positively correlated

with non-recurrent group A and B, suggesting transient microbes add to the microbiome's growth at different years.

Microbiome functional groups

We then investigated the gene functions present in the groups of microbial genera that may explain the positive or negative relationships between genera in Figure 4. In addition, because host metabolism and microbial functions are linked (Lynch and Hsiao 2019), we identified functional genes in each genus that may be linked to the host metabolism and classify three broad functional groups, generalists (recurrent group A), opportunists (recurrent group B) and specialists (recurrent group C) based on the presence of potential genes in these microbial genera. The group A recurrent microbes are typical marine species that have highly flexible genomes and growth strategies, including heterotrophic and photo-heterotrophic growth. These organisms can utilize many of the nutrients that may be present on the shark epidermis or excreted in the mucus, including proteins, fatty acids and disulphides (Meyer and Seegers 2012). *Pseudomonas*, for example, had the highest relative abundance and are capable of mixotrophy and simultaneous use of sulphur, nitrogen, and carbon from the surrounds (Guo, Chen, and Lee 2019) and sequences similar to these genes were highly represented in the *Pseudomonas* in leopard shark microbiomes. *Pseudomonas* were positively correlation with many microbes (Figure 4), which may be enhanced by traits, such as phosphate solubilization, and ammonia production (Yang et al. 2019; Goswami et al. 2013). Several, *Pseudomonas* species poses urease activity (Goswami et al. 2015) and tolerate, transition metals via the production of extracellular polysaccharides (EPS), such as Rhamnolipids, that form complexes with heavy metals (Meliani 2015).

Alcanivorax species, a highly abundant recurrent group A microbe, reduced alkane hydrocarbons, and possessed many enzymes for β -oxidation of fatty acids, the glyoxylate bypass, and the gluconeogenesis. These gene functions carried by *Alcanivorax* suggest they break down the complex hydrocarbons and fatty acid rich metabolites and may facilitate other member of the microbiome by providing simpler metabolic products that can be utilized. They also include enzymes for synthesis of riboflavin and unsaturated fatty acids and cardiolipin (Sabirova et al. 2020). Cardiolipin is a diphosphatidylglycerol lipid (Schlame, Brody, and Hostetler 1993) and is

516 more highly saturated in sharks than other vertebrates (Shadwick, Farrell, and Brauner 2015) and
517 sequences similar to Cardiolipin were identified in the metagenomes.

518 *Erythrobacter* species, a highly abundance group A microbe that was negatively correlated with
519 many other microbes are mostly aerobic anoxygenic phototrophic bacteria, containing
520 bacteriochlorophyll a, but lack the genes of autotrophic CO₂ fixation pathway, thus
521 photoheterotrophic metabolism requiring a supply of organic substrates. Several *Erythrobacter*
522 species have genes encoding enzymes for glycolysis and the tricarboxylic acid cycle but lack
523 genes for nitrogenase or nitrate reductase thus are reliant on other forms of organic nitrogen
524 (Koblizek et al. 2003), which matches the lower proportion of sequences identified in the
525 *Erythrobacter* in the *T. semifasciata* microbiome. *Erythrobacter* carry heavy metal resistance
526 genes, including resistance to lead, cadmium, zinc, mercury, nickel, cobalt, and arsenicals
527 (Zheng et al. 2016) and these genes were highly present in the metagenomic profiles. The
528 behaviour of the leopard sharks, being in shallow water with higher light conditions may have
529 provided the *Erythrobacter* with a competitive advantage over other microbial groups.

530 The last group A recurrent microbe was *Marinobacter* and these organisms are opportunistic
531 generalists that switch rapidly between lithoheterotrophy to heterotrophy, in both anaerobic and
532 aerobic conditions, in response to nutrient pulses. They have fast growth, outgrowing other
533 species in culture (Handley and Lloyd 2013). *Marinobacter* are able to respire inorganic
534 compounds that are usually found in metal rich environments (Handley and Lloyd 2013), and we
535 show they have many genes associated with metal resistance and transport. *Marinobacter* use
536 many hydrocarbons as a sole energy source, and produce large quantities of bioemulsifier, which
537 aid in bacterial adhesion to hydrophobic surfaces.

538 The recurrent group B microbes, the opportunists, were negatively correlated with most other
539 groups and had a high relative abundance of genes associated with carbon metabolism. *Gramella*
540 species degrades high molecular weight organic matter and encoding hydrolytic enzymes
541 predicting a preference for polymeric carbon sources and have range of gliding motility that
542 provide for surface adhesion (Bauer et al. 2006), features that enable the organism to live on the
543 surface of elasmobranchs. However, these microbes had fewer genes in ammonia assimilation,
544 promoting their coexistence with the more abundant species. *Leeuwenhoekiella*, a marine
545 Flavobacteria, has guiding motility, heterotrophic growth, is pigmented and often found in

associated with photosynthetic microbes and converts nitroalkanes into nitrite (Tahon et al. 2020). *Alteromonas* are non-phototrophic, heterotrophic, with flexible genome for taking advantage of influx of nutrients, particularly polymers, and some degrade polycyclic aromatic hydrocarbons (Math et al. 2012). *Alteromonas* have high flexibility in the metal resistant genes and efflux pumps and show strain differences in the copper-zinc-cobalt gene (*czc*) cassettes by successive insertions at a tRNA gene, with the numbers of *czc* cassettes varying from zero to three (Lopez-Perez et al. 2012) and these highly abundant in this genus in the *T. semifasciata* microbiome. *Pseudoalteromonas* and *Alteromonas* utilized extracellular hydrolysis as the major decomposition pathway of peptides and released fragments of amino acids into the surrounding environment (Liu and Liu 2020) and may have enable them to outcompete *Marinobacter* which is unable to conduct extracellular hydrolysis processes. However, these free amino acids may be used by other microbes, such as the non-recurrent group C and D microbes.

The group C recurrent microbes, the specialists, had less relative abundance overall but showed high relative abundance of a few metabolic pathways (Figure 5). *Ruegeria* had a high proportion of sequences associated with urea metabolism, compared with its relatively low proportional abundance. *Ruegeria* are able to use Trimethylamine (TMA) and trimethylamine N-oxide (TMAO) as an energy source to product intracellular ATP (Lidbury, Murrell, and Chen 2015). TMAO and TMA are used by elasmobranchs for osmoregulation.

Sphingopyxis also had low relative abundance, utilizing carbon and nitrogen compounds as substrates for growth but are generally slow growing (Moran et al. 2007). Serine, a pathway that was overrepresented in *Sphingopyxis*, can serve as a nitrogen source for the growth of some species, but not as a carbon source (Williams et al. 2009). Ammonia and nitrate are used as nitrogen sources, but there is a high energetic cost associated with nitrate utilization (Verma et al. 2020). *Sphingomonas* are closely related to *Sphingopyxis* and both produces exopolysaccharides (EPS) (Chang et al. 2021). The production of these compounds may provide biofilm development on the shark skin. Therefore, each recurrent microbial group was providing a different set of functional genes that enhanced coexistence and reflected the products that may be associated with shark metabolism and excreted via the mucus.

Gene functions linked to host metabolic processes

The stability of the functional potential of the epidermal microbiome suggests the elasmobranch physiology and dermal denticle topography are strong drivers of microbiome function, and we have developed a model to link the microbiome function with shark metabolism (Figure 7). Elasmobranchs rely on protein as the primary food source and use ketogenic pathways and produce high ketone bodies and hydrocarbons (Speers-Roesch and Treberg 2010). The functions that were present in metagenomes suggests that the microbes were responding to the presence of these compounds in the minimal mucus that covers the sharks. For example, *Alxavorax* and *Marinobacter* carried the genes to degrade hydrocarbons. There are few descriptions of the compounds within the mucus of Elasmobranchs, and these included high levels of proteins (Tsutsui et al., 2009) with moderate to high disulphide concentrations (Meyer et al., 2007a). A recent transcriptome analysis of the integument of cookie cutter sharks identified Glyceraldehyde-3-phosphate dehydrogenase and Fructose-bisphosphate aldolase A transcripts, both play key roles in the glycolysis and gluconeogenesis (Delroisse et al. 2021) suggesting these products are being produced in the skin and may be available to the microbial community. Protein metabolism and nitrate and nitrite ammonification, urea decomposition with genes such as trimethylamine N-oxide (TMAO) reductase were significant functions in the microbiomes. The low levels of nitrogen fixation in the microbiome and the high proportional abundance of nitrogen metabolism and urea decomposition and the presences of within the microbiome suggest the elasmobranch is a nitrogen-rich environment (Hazon et al. 2003; Dowd et al. 2010). Proteasomes, metallocarboxypeptidase, aminopeptidase genes, protein degradation and biosynthesis functional genes suggest a propensity of the microbes for protein degradation. There was no evidence of dentine, hydrolysate, or collagen degradation in the microbiome, suggesting that the microbes are not breaking down the dermal denticles, which signifies a lack of negative impact on the sharks. Therefore, a commensal connection between *T. semifasciata* and the epidermis microbiome is established through the passive resource subsidization of these microbiome functions.

Cobalt-zinc-cadmium resistance and ton/to transport genes are highly prevalent functional genes shared across epidermis microbiomes of *T. semifasciata* (this study), *A. vulpinus* (Doane et al. 2017), *Rhincodon typus* and *Urobatis halleri* (Doane et al. 2020). Elasmobranchs bioaccumulate heavy metals and shunt these compounds to the epidermis (Escobar-Sanchez, Galvan-Magana,

and Rosiles-Martinez 2010; Maz-Courrau et al. 2012), and the epidermal microbiomes are responding to the presence of these compounds. The prevalence of heavy metal resistance genes may provide a biomarker of elasmobranchs health and identify those exposed to anthropogenic sources of heavy metals in the environment providing an early warning sign of declining health.

Conclusions

The highly structured epidermis of elasmobranchs, in this case, *T. semifasciata*, promotes a diverse microbiome that has a flexible taxonomic makeup where many genera coexist. The microbial community was maintained by functional redundancy driven by the taxonomic flexibility and various microbes "flip-flopping" in abundance with one another, which kept the microbial niches filled across the years. The high relative abundance of genes involved in nitrification and denitrification, urea decomposition, and heavy metal resistance genes in the epidermis microbiome across all *T. semifasciata* individuals suggests a connection with the host metabolism potentially through passive subsidization of organic nitrogen, proteins, hydrocarbons, and heavy metals.

622 **Abbreviations**

623 Not applicable

624

625 **Ethical Approval and Consent to participate**

626 Animal handling and ethics were reviewed at San Diego State University through IACUC under
627 permit APF #14-05-011D, APF #17-11-010D, APF # 18-05-007D. Sampling was conducting
628 under state permit SCP #12847 and SCP #9893 from the California Department of Fish and
629 Wildlife.

630

631 **Consent for publication**

632 Not applicable

633

634 **Availability of data and materials**

635 All data will be publicly released upon publication.

636

637 **Competing interests**

638 The authors declare no conflict of interest.

639

640 **Funding**

641 We acknowledge support from S. Lo and B. Billings. We acknowledge funding support from
642 NSF Division of Undergraduate Education # 1323809 and NSF Division of Molecular and
643 Cellular Science # 1330800.

644

645 **Author's contribution**

646 MD collected all field samples, constructed the ideas and help write the paper, CJ analyzed the
647 data and wrote the manuscript, EK, AT, AG, SJ, captured and prepared samples and provide
648 feedback on the manuscript, MMM and LL provided sequencing and analysis, AN facilitated the
649 catching of leopard sharks, RD provided the illustration, ED conceptualized the idea, analyzed,
650 wrote and lead the project.

651

652 **Acknowledgements**

We would like to thank the students in the SDSU undergraduate Ecological Metagenomic class assisting with the processing and sequencing of the leopard shark metagenomes.

Authors' information

¹College of Science and Engineering, Flinders University, Bedford Park, South Australia, Australia; ²Department of Biology, San Diego State University, San Diego, California, USA; ³Hopkins Marine Station, Stanford University, Pacific Grove, California, USA; ⁴Laurence Livermore National Labs, Livermore, California, USA; ⁵Australian Centre for Ecogenomics, University of Queensland, St Lucia, Queensland, Australia; ⁶Department of Environmental and Ocean Sciences, University of San Diego, San Diego, California, USA; ⁷Scripps Institution of Oceanography, University of California – San Diego, La Jolla, California, USA

Figure 1. The relative proportions of the most abundant microbial genera represented in the *Triakis semifasciata* microbiome across four years.

673

674

675

676 Figure 2 The microbial genera of the *Triakis semifasciata* epidermal microbiome clustered by year, with a
 677 few outliers on the principal component analysis conducted on proportional sequence distribution.

678

Figure 3. The mean relative abundance ($\pm$ S.E.) of abundant recurrent genera per year on the *Triakis semifasciata* epidermal microbiome, showing flexibility with no abundant genera becoming extinct over the four years.

Figure 4. Positive and negative correlations occurred between most abundant 19 recurrent and non-recurrent microbes in the *Triakis semifasciata* epidermal microbiome. Bray Curtis similarity was used to cluster the genera with similar proportional abundance across years and the relationship with other genera was compared using a Pearson's correlation, displayed as a heat map where red is a negative correlation and blue is a positive correlation.

Figure 5. The functional potential of the *Triakis semifasciata* epidermal microbiome was stable across the four-year period, shown using a principal component analysis.

Function Group	Recurrent Genera	Carbon Metabolism			Nitrogen Metabolism				Protein Metabolism		Fatty Acids		Heavy Metals		Motility	
		TCA Cycle	Glycolysis	Serine Cycle	Nitrate & Nitrite Ammonification	Ammonia Assimilation	Denitrification	Urea Decomposition	Protein Biosynthesis	Protein Degradation	Fatty Acid Biosynthesis	Fatty Acid Metabolism	CoZnCd Resistance	Copper Homeostasis	Flagellar Motility	Bacterial Chemotaxis
Generalist	Pseudomonas															
	Alcanivorax															
	Erythrobacter															
	Marinobacter															
Opportunist	Leeuwenhoekeiella															
	Alteromonas															
	Gramella															
	Zunongwangia															
	Pseudoalteromonas															
	Flavobacterium															
Specialist	Maribacter															
	Ruegeria															
	Sphingopyxis															
	Burkholderia															
	Novosphingobium															
	Sphingobium															
	Sphingomonas															

Figure 6. Heat map of the mean proportion of genes associated with each metabolism in the 17 most abundant recurrent taxon in group A (aqua), group B(grey) and group c (green), shows genes were mostly shared across taxon, but some specialization occurred. The proportion of sequences were divided into quartiles, where darker colors represent higher percentile (>75– dark red, 50-75 orange/red, 25-50 - orange, and 0-25 - pale orange). No color represented no sequences for that metabolism held by that genus.

Figure 7. The links between the functional potential of the *Triakis semifasciata* epidermal microbiome and the host metabolism. The proportion of each gene in each taxon were identified and classed into quartiles. Most taxa had most functions suggesting the microbiome displays weak competition and is responding to the metabolic processes of the host.

Table 1. The *Triakis semifasciata* metagenome statistics, including collection date, total sequences and proportion of the metagenome that was annotated by the database.

Metagenome Name	Date	Location	SRA#	Number of sequences	Annotated proteins (%)
LS 1 2013	2013	La Jolla		1188249	20.69
LS 2 2013	2013	La Jolla		1060680	24.36
LS 3 2013	2013	La Jolla		1024296	20.16
LS 4 2013	2013	La Jolla		1154296	21.33
LS 5 2013	2013	La Jolla		2347886	19.79
LS 6 2013	2013	La Jolla		3346424	18.84
LS 7 2013	2013	La Jolla		2330953	20.36
LS 1 2015	2015	La Jolla		1143301	22.68
LS 2 2015	2015	La Jolla		2012275	15.87
LS 3 2015	2015	La Jolla		1635251	20.4
LS 1 2017	2017	La Jolla		701418	75.63
LS 2 2017	2017	La Jolla		898767	74.86
LS 3 2017	2017	La Jolla		636435	46.71
LS 4 2017	2017	La Jolla		324102	68.57
LS 5 2017	2017	La Jolla		312135	61.76
LS 6 2017	2017	La Jolla		693318	43.43
Ls 7 2017	2017	La Jolla		613946	62.52
LS 8 2017	2017	La Jolla		1189453	43.69

Table 2. The genera that contributed to the flexibility in the *Triakis semifasciata* epidermal microbiome across four years, calculated via a Simper analysis, with dissimilarity coefficient of 16.95 between 2013 & 2015; 19.71 for 2013 & 2017; and 15.32 for 2015 & 2017

Years compared	Phyla	Class	Genus	Percent Contribution
2013 vs 2015	Proteobacteria	Gammaproteobacteria	<i>Providencia</i>	0.9
			<i>Endorifia</i>	0.86
			<i>Marinobacter</i>	0.72
			<i>Pseudomonas</i>	0.61
			<i>Xylella</i>	0.58
			<i>Alcanivorax</i>	0.58
		Betaproteobacteria	<i>Limnibacter</i>	0.87
		Alphaproteobacteria	<i>Erythrobacter</i>	0.7
			<i>Hyphomonas</i>	0.56
			<i>Oceanibulbus</i>	0.66
2013 vs 2017	Bacteroidetes	Flavobacteria	<i>Leeuwenkoekiella</i>	0.9
			<i>Zuonwangia</i>	0.7
			<i>Gramella</i>	0.58
	Proteobacteria	Betaproteobacteria	<i>Limnibacter</i>	0.89
		Gammaproteobacteria	<i>Providencia</i>	0.89
			<i>Alteromonas</i>	0.73
			<i>Xylella</i>	0.67
			<i>Endorifia</i>	0.54
		Alphaproteobacteria	<i>Hyphomonas</i>	0.48
		Planctomycetia	<i>Planctomyces</i>	0.47
2015 vs 2017	Proteobacteria	Gammaproteobacteria	<i>Providencia</i>	1.41
			<i>Endorifia</i>	1.11
			<i>Psychrobacter</i>	0.91
			<i>Xylella</i>	0.81
			<i>Alcanivorax</i>	0.75
			<i>Alteromonas</i>	0.69
			<i>Pseudomonas</i>	0.52
		Alphaproteobacteria	<i>Ruegeria</i>	0.59
		Betaproteobacteria	<i>Achromobacter</i>	0.56
	Bacteroidetes	Flavobacteria	<i>Leeuwenkoekiella</i>	0.95

752
753
754   Supplementary Figure 1. The phylum profile of the *Triakis semifasciata* epidermal microbiome had
755   stable across four years.

Supplementary Figure 2.

The relative distribution of microbial classes in the *Triakis semifasciata* microbiome were similar over a four-years.

Supplementary Figure 3. The major metabolisms identifying within the *Triakis semifasciata* microbiomes.

References

- Anderson, M. J. 2017. 'Permutational multivariate analysis of variance (PERMANOVA).' in T. Colton N. Balakrishnan, B. Everitt, W. Piegorisch, F. Ruggeri, and J. Teugels (ed.), *Wiley statsref: statistics reference online* (John Wiley and Sons: Hoboken, NJ, USA.).
- Anderson, M.J. 2006. 'Distance-based tests for homogeneity of multivariate dispersions. ', *Biometrics*, 62: 245–53.
- Aziz, R. K., D. Bartels, A. A. Best, M. DeJongh, T. Disz, R. A. Edwards, K. Formsma, S. Gerdes, E. M. Glass, M. Kubal, F. Meyer, G. J. Olsen, R. Olson, A. L. Osterman, R. A. Overbeek, L. K. McNeil, D. Paarmann, T. Paczian, B. Parrello, G. D. Pusch, C. Reich, R. Stevens, O. Vassieva, V. Vonstein, A. Wilke, and O. Zagnitko. 2008. 'The RAST server: Rapid annotations using subsystems technology', *Bmc Genomics*, 9.
- Ballantyne, J. S. 1997. ' Jaws: the inside story. The metabolism of elasmobranch fishes. ', *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* . 118: 703-42.
- Bauer, M., M. Kube, H. Teeling, M. Richter, T. Lombardot, E. Allers, C. A. Wurdemann, C. Quast, H. Kuhl, F. Knaust, D. Woebken, K. Bischof, M. Mussmann, J. V. Choudhuri, F. Meyer, R. Reinhardt, R. I. Amann, and F. O. Glockner. 2006. 'Whole genome analysis of the marine Bacteroidetes'Gramella forsetii' reveals adaptations to degradation of polymeric organic matter', *Environmental Microbiology*, 8: 2201-13.
- Bierlich, K. C., C. Miller, E. DeForce, A. S. Friedlaender, D. W. Johnston, and A. Apprill. 2018. 'Temporal and Regional Variability in the Skin Microbiome of Humpback Whales along the Western Antarctic Peninsula', *Appl Environ Microbiol*, 84.

- Cantu, V. A., J. Sadural, and R. A. Edwards. 2019. 'PRINSEQ ++, a multi-threaded tool for fast and efficient quality control and preprocessing of sequencing datasets.', *PeerJ Preprints*, 7: 43–45.
- Chang, A. P., J. Qian, H. Li, Y. L. Wang, J. Y. Lin, Q. M. He, Y. L. Shen, and H. Zhu. 2021. 'Characterization and Function of a Novel Welan Gum Lyase From Marine *Sphingomonas* sp. WG', *Frontiers in Microbiology*, 12: 638355.
- Chien, H. W., X. Y. Chen, W. P. Tsai, and M. Lee. 2020. 'Inhibition of biofilm formation by rough shark skin-patterned surfaces', *Colloids and Surfaces B-Biointerfaces*, 186.
- Clarke, K. R., P. J. Somerfield, and R. N. Gorley. 2008. 'Testing of null hypotheses in exploratory community analyses: similarity profiles and biota-environment linkage', *Journal of Experimental Marine Biology and Ecology*, 366: 56-69.
- . 2016. 'Clustering in non-parametric multivariate analyses', *Journal of Experimental Marine Biology and Ecology*, 483: 147-55.
- Costello, E. K., K. Stagaman, L. Dethlefsen, B. J. Bohannan, and D. A. Relman. 2012. 'The application of ecological theory toward an understanding of the human microbiome', *Science*, 336: 1255-62.
- Delroisse, J., L. Duchatelet, P. Flammang, and J. Mallefet. 2021. 'Photophore Distribution and Enzymatic Diversity Within the Photogenic Integument of the Cookie-Cutter Shark *Isistius brasiliensis* (Chondrichthyes: Dalatiidae)', *Frontiers in Marine Science*, 8.
- Dillon, E. M., R. D. Norris, and A. O'Dea. 2017. 'Dermal denticles as a tool to reconstruct shark communities', *Marine Ecology Progress Series*, 566: 117-34.
- Dinsdale, E. A., R. A. Edwards, D. Hall, F. Angly, M. Breitbart, J. M. Brulc, M. Furlan, C. Desnues, M. Haynes, L. L. Li, L. McDaniel, M. A. Moran, K. E. Nelson, C. Nilsson, R. Olson, J. Paul, B. R. Brito, Y. J. Ruan, B. K. Swan, R. Stevens, D. L. Valentine, R. V. Thurber, L. Wegley, B. A. White, and F. Rohwer. 2008. 'Functional metagenomic profiling of nine biomes', *Nature*, 452: 629-U8.
- Dinsdale, E. A., O. Pantos, S. Smriga, R. A. Edwards, F. Angly, L. Wegley, M. Hatay, D. Hall, E. Brown, M. Haynes, L. Krause, E. Sala, S. A. Sandin, R. V. Thurber, B. L. Willis, F. Azam, N. Knowlton, and F. Rohwer. 2008. 'Microbial ecology of four coral atolls in the Northern Line Islands', *Plos One*, 3: e1584.
- Doane, M. P., J. M. Haggerty, D. Kacev, B. Papudeshi, and E. A. Dinsdale. 2017. 'The skin microbiome of the common thresher shark (*Alopias vulpinus*) has low taxonomic and gene function -diversity', *Environmental Microbiology Reports*, 9: 357-73.
- Doane, M. P., D. Kacev, S. Harrington, K. Levi, D. Pande, A. Vega, and E. A. Dinsdale. 2018. 'Mitochondrial recovery from shotgun metagenome sequencing enabling phylogenetic analysis of the common thresher shark (*Alopias vulpinus*)', *Meta Gene*, 15: 10-15.
- Doane, M. P., M. M. Morris, B. Papudeshi, L. Allen, D. Pande, J. M. Haggerty, S. Johri, A. C. Turnlund, M. Peterson, D. Kacev, A. Nosal, D. Ramirez, K. Hovel, J. Ledbetter, A. Alker, J. Avalos, K. Baker, S. Bhide, E. Billings, S. Byrum, M. Clemens, A. J. Demery, L. F. O. Lima, O. Gomez, O. Gutierrez, S. Hinton, D. Kieu, A. Kim, R. Loaiza, A. Martinez, J. McGhee, K. Nguyen, S. Parlan, A. Pham, R. Price-Waldman, R. A. Edwards, and E. A. Dinsdale. 2020. 'The skin microbiome of elasmobranchs follows phylosymbiosis, but in teleost fishes, the microbiomes converge', *Microbiome*, 8.
- Dowd, W. W., B. N. Harris, J. J. Cech, Jr., and D. Kultz. 2010. 'Proteomic and physiological responses of leopard sharks (*Triakis semifasciata*) to salinity change', *Journal of Experimental Biology*, 213: 210-24.
- Egan, S., and M. Gardiner. 2016. 'Microbial Dysbiosis: Rethinking Disease in Marine Ecosystems', *Frontiers in Microbiology*, 7: 991.
- Escobar-Sanchez, O., F. Galvan-Magana, and R. Rosiles-Martinez. 2010. 'Mercury and selenium bioaccumulation in the smooth hammerhead shark, *Sphyrna zygaena* Linnaeus, from the Mexican Pacific Ocean', *Bull Environ Contam Toxicol*, 84: 488-91.

- Goswami, D., K. Patel, S. Parmar, H. Vaghela, N Muley, P Dhandhukia, and J.N. Thakker. 2015. 'Elucidating multifaceted urease producing marine *Pseudomonas aeruginosa* BG as a cogent PGPR and bio-control agent.', *Plant Growth Regulation*, 75: 253–63.
- Goswami, D., H. Vaghela, S. Parmar, P Dhandhukia, and J. Thakker. 2013. 'Plant growth promoting potentials of *Pseudomonas* spp. strain OG isolated from marine water.', *J Plant Interact*, 8: 281–90.
- Guo, H., C. Chen, and D. J. Lee. 2019. 'Nitrogen and sulfur metabolisms of *Pseudomonas* sp. C27 under mixotrophic growth condition', *Bioresour Technol*, 293: 122169.
- Handley, K. M., and J. R. Lloyd. 2013. 'Biogeochemical implications of the ubiquitous colonization of marine habitats and redox gradients by *Marinobacter* species', *Frontiers in Microbiology*, 4.
- Hazon, N., A. Wells, R. D. Pillans, J. P. Good, W. Gary Anderson, and C. E. Franklin. 2003. 'Urea based osmoregulation and endocrine control in elasmobranch fish with special reference to euryhalinity', *Comp Biochem Physiol B Biochem Mol Biol*, 136: 685-700.
- Johri, S., M. P. Doane, L. Allen, and E. A. Dinsdale. 2019. 'Taking Advantage of the Genomics Revolution for Monitoring and Conservation of Chondrichthyan Populations', *Diversity-Basel*, 11.
- Kanehisa, M., S. Goto, Y. Sato, M. Furumichi, and M. Tanabe. 2012. 'KEGG for integration and interpretation of large-scale molecular data sets', *Nucleic Acids Research*, 40: D109-D14.
- Kanehisa, M., Y. Sato, M. Kawashima, M. Furumichi, and M. Tanabe. 2016. 'KEGG as a reference resource for gene and protein annotation', *Nucleic Acids Research*, 44: D457-D62.
- Kim, M., S. Lee, H. D. Park, S. I. Choi, and S. Hong. 2012. 'Biofouling control by quorum sensing inhibition and its dependence on membrane surface', *Water Science and Technology*, 66: 1424-30.
- Kimes, N. E., W. R. Johnson, M. Torralba, K. E. Nelson, E. Weil, and P. J. Morris. 2013. 'The *Montastraea faveolata* microbiome: ecological and temporal influences on a Caribbean reef-building coral in decline', *Environmental Microbiology*, 15: 2082-94.
- Koblizek, M., O. Beja, R. R. Bidigare, S. Christensen, B. Benitez-Nelson, C. Vetriani, M. K. Kolber, P. G. Falkowski, and Z. S. Kolber. 2003. 'Isolation and characterization of *Erythrobacter* sp strains from the upper ocean', *Archives of Microbiology*, 180: 327-38.
- Lang, Amy. 2020. 'The speedy secret of shark skin', *Physics Today*, 73: 58-59.
- Lidbury, I. D., J. C. Murrell, and Y. Chen. 2015. 'Trimethylamine and trimethylamine N-oxide are supplementary energy sources for a marine heterotrophic bacterium: implications for marine carbon and nitrogen cycling', *Isme Journal*, 9: 760-9.
- Lima, L. F. O., M. Weissman, M. Reed, B. Papudeshi, A. T. Alker, M. M. Morris, R. A. Edwards, S. J. de Putron, N. K. Vaidya, and E. A. Dinsdale. 2020. 'Modeling of the Coral Microbiome: the Influence of Temperature and Microbial Network', *Mbio*, 11.
- Lima, N., T. Rogers, K. Acevedo-Whitehouse, and MV. Brown. 2012. 'Temporal stability and species specificity in bacteria associated with the bottlenose dolphins respiratory system.', *Environ Microbiol Rep.*, 4: 89–96.
- Liu, S., and Z. Liu. 2020. 'Distinct capabilities of different Gammaproteobacterial strains on utilizing small peptides in seawater', *Sci Rep*, 10: 464.
- Lopez-Perez, M., A. Gonzaga, A. B. Martin-Cuadrado, O. Onyshchenko, A. Ghavidel, R. Ghai, and F. Rodriguez-Valera. 2012. 'Genomes of surface isolates of *Alteromonas macleodii*: the life of a widespread marine opportunistic copiotroph', *Sci Rep*, 2: 696.
- Lowery, N. V., and T. Ursell. 2019. 'Structured environments fundamentally alter dynamics and stability of ecological communities', *Proceedings of the National Academy of Sciences of the United States of America*, 116: 379-88.
- Lynch, J. B., and E. Y. Hsiao. 2019. 'Microbiomes as sources of emergent host phenotypes', *Science*, 365: 1405-08.

- Magoc, T., and S. L. Salzberg. 2011. 'FLASH: fast length adjustment of short reads to improve genome assemblies', *Bioinformatics*, 27: 2957-63.
- Math, R. K., H. M. Jin, J. M. Kim, Y. Hahn, W. Park, E. L. Madsen, and C. O. Jeon. 2012. 'Comparative genomics reveals adaptation by *Alteromonas* sp. SN2 to marine tidal-flat conditions: cold tolerance and aromatic hydrocarbon metabolism', *Plos One*, 7: e35784.
- Maz-Courrau, A., C. Lopez-Vera, F. Galvan-Magana, O. Escobar-Sanchez, R. Rosiles-Martinez, and A. Sanjuan-Munoz. 2012. 'Bioaccumulation and biomagnification of total mercury in four exploited shark species in the Baja California Peninsula, Mexico', *Bull Environ Contam Toxicol*, 88: 129-34.
- Meliani, A. 2015. 'Bioremediation Strategies Employed by *Pseudomonas* Species. .' in Maheshwari D. (ed.), *Bacterial Metabolites in Sustainable Agroecosystem. Sustainable Development and Biodiversity* (Springer,).
- Meyer, F., D. Paarmann, M. D'Souza, R. Olson, E. M. Glass, M. Kubal, T. Paczian, A. Rodriguez, R. Stevens, A. Wilke, J. Wilkening, and R. A. Edwards. 2008. 'The metagenomics RAST server - a public resource for the automatic phylogenetic and functional analysis of metagenomes', *BMC Bioinformatics*, 9.
- Meyer, W., and U. Seegers. 2012. 'Basics of skin structure and function in elasmobranchs: a review', *Journal of Fish Biology*, 80: 1940-67.
- Meyer, W., U. Seegers, and R. Stelzer. 2007. 'Sulphur, thiols, and disulphides in the fish epidermis, with remarks on keratinization', *Journal of Fish Biology*, 71: 1135-44.
- Moran, M. A., R. Belas, M. A. Schell, J. M. Gonzalez, F. Sun, S. Sun, B. J. Binder, J. Edmonds, W. Ye, B. Orcutt, E. C. Howard, C. Meile, W. Palefsky, A. Goesmann, Q. Ren, I. Paulsen, L. E. Ulrich, L. S. Thompson, E. Saunders, and A. Buchan. 2007. 'Ecological genomics of marine Roseobacters', *Appl Environ Microbiol*, 73: 4559-69.
- Nosal, A. P., A. Caillat, E. K. Kisfaludy, M. A. Royer, and N. C. Wegner. 2014. 'Aggregation behavior and seasonal philopatry in male and female leopard sharks *Triakis semifasciata* along the open coast of southern California, USA', *Marine Ecology Progress Series*, 499: 157-75.
- Nosal, A. P., D. C. Cartamil, J. W. Long, M. Luhrmann, N. C. Wegner, and J. B. Graham. 2013. 'Demography and movement patterns of leopard sharks (*Triakis semifasciata*) aggregating near the head of a submarine canyon along the open coast of southern California, USA', *Environmental Biology of Fishes*, 96: 865-78.
- Ondov, B. D., N. H. Bergman, and A. M. Phillippy. 2011. 'Interactive metagenomic visualization in a Web browser', *BMC Bioinformatics*, 12: 385.
- Overbeek, R., T. Begley, R. M. Butler, J. V. Choudhuri, H. Y. Chuang, M. Cohoon, V. de Crecy-Lagard, N. Diaz, T. Disz, R. Edwards, M. Fonstein, E. D. Frank, S. Gerdes, E. M. Glass, A. Goesmann, A. Hanson, D. Iwata-Reuyl, R. Jensen, N. Jamshidi, L. Krause, M. Kubal, N. Larsen, B. Linke, A. C. McHardy, F. Meyer, H. Neuweeger, G. Olsen, R. Olson, A. Osterman, V. Portnoy, G. D. Pusch, D. A. Rodionov, C. Ruckert, J. Steiner, R. Stevens, I. Thiele, O. Vassieva, Y. Ye, O. Zagnitko, and V. Vonstein. 2005. 'The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes', *Nucleic Acids Research*, 33: 5691-702.
- Pogoreutz, C., M. A. Gore, G. Perna, C. Millar, R. Nestler, R. F. Ormond, C. R. Clarke, and C. R. Voolstra. 2019. 'Similar bacterial communities on healthy and injured skin of black tip reef sharks', *Anim Microbiome*, 1: 9.
- Sabirova, JS, M Ferrer, D Regenhart, KN Timmis, and PN Golyshin. 2020. 'Proteomic Insights into Metabolic Adaptations in *Alcanivorax borkumensis* Induced by Alkane Utilization', *Journal of Bacteriology*, 188: 3763-73.
- Schlame, M., S. Brody, and K. Y. Hostetler. 1993. 'Mitochondrial Cardiolipin in Diverse Eukaryotes - Comparison of Biosynthetic Reactions and Molecular Acyl Species', *European Journal of Biochemistry*, 212: 727-35.

- Schmieder, R., and R. Edwards. 2011. 'Fast Identification and Removal of Sequence Contamination from Genomic and Metagenomic Datasets', *Plos One*, 6.
- Shadwick, R.E. , A.P. Farrell, and C.J. Brauner. 2015. *Physiology of Elasmobranch Fishes: Internal Processes* (Academic Press).
- Speers-Roesch, B, and JR Treberg. 2010. 'The unusual energy metabolism of elasmobranch fishes.', *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 155: 417-34.
- Sullivan, T., and F. Regan. 2011a. 'The characterization, replication and testing of dermal denticles of *Scyliorhinus canicula* for physical mechanisms of biofouling prevention', *Bioinspiration & Biomimetics*, 6.
- . 2011b. 'The characterization, replication and testing of dermal denticles of *Scyliorhinus canicula* for physical mechanisms of biofouling prevention', *Bioinspiration & Biomimetics*, 6: 046001.
- Tahon, G., L. Lebbe, M. De Troch, K. Sabbe, and A. Willems. 2020. 'Leeuwenhoekiella aestuarii sp. nov., isolated from salt-water sediment and first insights in the genomes of Leeuwenhoekiella species', *Int J Syst Evol Microbiol*, 70: 1706-19.
- Tsutsui, S., Y. Dotsuta, A. Ono, M. Suzuki, H. Tateno, J. Hirabayashi, and O. Nakamura. 2014. 'A C-type lectin isolated from the skin of Japanese bullhead shark (*Heterodontus japonicus*) binds a remarkably broad range of sugars and induces blood coagulation.', *Journal of Biochemistry*, 157: 345–56.
- Vendl, C., T. Nelson, B. Ferrari, T. Thomas, and T. Rogers. 2021. 'Highly abundant core taxa in the blow within and across captive bottlenose dolphins provide evidence for a temporally stable airway microbiota', *BMC Microbiol*, 21: 20.
- Verma, H., G. G. Dhingra, M. Sharma, V. Gupta, R. K. Negi, Y. Singh, and R. Lal. 2020. 'Comparative genomics of *Sphingopyxis* spp. unravelled functional attributes', *Genomics*, 112: 1956-69.
- Wen, L., P. J. M. Thornycroft, J. C. Weaver, and G. V. Lauder. 2015. 'Hydrodynamic function of biomimetic shark skin: effect of denticle pattern and spacing', *Integrative and Comparative Biology*, 55: E196-E96.
- Williams, T. J., H. Ertan, L. Ting, and R. Cavicchioli. 2009. 'Carbon and nitrogen substrate utilization in the marine bacterium *Sphingopyxis alaskensis* strain RB2256', *Isme Journal*, 3: 1036-52.
- Wilson, B., B. S. Danilowicz, and W. G. Meijer. 2008. 'The diversity of bacterial communities associated with Atlantic Cod *Gadus morhua*', *Microbial Ecology*, 55: 425-34.
- Yang, L., X. H. Wang, S. Cui, Y. X. Ren, J. Yu, N. Chen, Q. Xiao, L. K. Guo, and R. H. Wang. 2019. 'Simultaneous removal of nitrogen and phosphorous by heterotrophic nitrification-aerobic denitrification of a metal resistant bacterium *Pseudomonas putida* strain NP5', *Bioresour Technol*, 285: 121360.
- Zhang, D. Y., Y. Y. Li, X. Han, X. A. Li, and H. W. Chen. 2011. 'High-precision bio-replication of synthetic drag reduction shark skin', *Chinese Science Bulletin*, 56: 938-44.
- Zheng, Q., W. X. Lin, Y. T. Liu, C. Chen, and N. Z. Jiao. 2016. 'A Comparison of 14 *Erythrobacter* Genomes Provides Insights into the Genomic Divergence and Scattered Distribution of Phototrophs', *Frontiers in Microbiology*, 7.