Phosphate starvation enhances phagocytosis of *Mycobacterium bovis*/BCG by macrophages.

Patricia Espinosa-Cueto

Universidad Nacional Autonoma de Mexico Instituto de Investigaciones Biomedicas

Alejandro Magallanes-Puebla

Universidad Nacional Autonoma de Mexico Instituto de Investigaciones Biomedicas

Raul Mancilla (✉ mancilla@biomedicas.unam.mx)

Universidad Nacional Autonoma de Mexico Instituto de Investigaciones Biomedicas

<https://orcid.org/0000-0002-5752-8995>

Research article

Keywords: Phosphate, deprivation, phagocytosis

Posted Date: December 1st, 2019

DOI: <https://doi.org/10.21203/rs.2.17903/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Immunology on June 9th, 2020. See the published version at <https://doi.org/10.1186/s12865-020-00364-x>.

Abstract

Tuberculosis (TB) is an important health problem worldwide. The only available TB vaccine is *Mycobacterium bovis* bacillus Calmette-Guerin, a saprophytic mycobacterium that activates the innate and the acquired immune system after being phagocytosed by macrophages and dendritic cells. BCG vaccination fails to prevent adult pulmonary TB although it may have a protective effect in childhood TB. Understanding how BCG interacts with macrophages and other immunocompetent cells is crucial to develop new vaccines. In this study we showed that macrophages phagocytose *Mycobacterium bovis*/BCG bacilli with higher efficiency when they are cultured without phosphate. The increased phagocytosis is associated with diminished phagosome acidification. We isolated mycobacterial cell wall and membranes to search for mycobacterial molecules that could be involved in these processes; by immunoblot, it was found that the plasma membranes of phosphate-deprived bacilli express the adhesins PstS-1, LpqH, LprG, and the APA antigen. These proteins are not detected in membranes of bacilli grown with usual amounts of phosphate. The interest of our observations is to show that under the metabolic stress implied in phosphate deprivation, mycobacteria respond upregulating adhesins that could improve their capacity to infect macrophages. These observations are relevant to understand how *M. bovis*/BCG induces protective immunity.

Introduction

Tuberculosis (TB) remains an important health problem worldwide. In 2017, there were 10.4 million new cases and 1.7 million deaths [1]. The main etiologic agent of TB is *Mycobacterium tuberculosis* (Mtb), a facultative intracellular bacterium that infects host cells, primarily macrophages (MO) which generally eliminate the bacilli. However, highly virulent Mtb strains may inhibit the fusion of the phagosome with the lysosome, thus allowing its survival and intracellular replication [2]. In pulmonary TB, in an initial step Mtb binds alveolar MOs through mycobacterial adhesins or opsonins that interact with a variety of receptors, including CR1, CR3, and CR4, IgGfC, surfactant A protein and C-type lectins [3]. Information on the repertoire of Mtb adhesins is scanty; a few have been characterized including lipoarabinomannan, a cell wall glycolipid [4], LpqH, a 19-kDa surface located lipoprotein [5], the heparin-binding hemagglutinin [6], PE_PGRS33, a member of the PE-PGRS family [7], LprG a 25-kDa glycolipoprotein [8], APA a 45-47-kDa mannose-containing glycoprotein [9] and PstS-1, a phosphate transporter [10]. It is important to know better the repertoire of mycobacterial adhesins since it could be helpful to design novel strategies for TB prevention and treatment. [11].

Recently, it has been reported that PstS-1, the 38-kDa glycoprotein, binds the MO mannose receptor triggering the phagocytosis of mycobacteria [10]. PstS-1 has focused much interest as a phosphate (Pi) transporter whose expression is upregulated under Pi starvation conditions [12, 13, 14]. Pi deprivation can activate the expression of a variety of genes not involved in phosphate metabolism that participate in the pathogenesis of TB [15]. Herein, we show that *M. bovis*/BCG bacilli grown without phosphate are engulfed with increased efficiency by MO, an event that was associated with decreased phagosome acidification. Considering that Pi starvation increases the surface expression of PstS-1, we searched its

expression in isolated plasma membranes; we found that together with PstS-1, the membranes of Pi-deprived mycobacteria express the mycobacterial adhesins LpqH, LprG, and the APA antigen, glycoproteins that are not directly involved in phosphate regulation.

Results

Mycobacteria cultured under Pi deprivation conditions are phagocytosed with higher efficiency by macrophages.

The phagocytosis assays were carried out incubating J774A.1 macrophage-like cells with mycobacteria labeled with the lipophilic red fluorescent dye PKH-26 at 1:2, 1:5 and 1:20 MOI. Immunofluorescence microscopy revealed that the number of MOs with engulfed or bound bacilli was significantly higher with Pi-deprived mycobacteria (Fig. 1). Phagocytosis was quantitated by FACS with PKH-26 stained bacilli using a Calibur cytometer (Beckton Dickinson, San Diego, CA, USA). Phagocytosis expressed in mean fluorescence intensity (MFI) and percent was dose-dependent and considerably higher with Pi-deprived bacilli (Fig. 1). At a 1:20 MOI, the percent of phagocytosis of bacilli grown without Pi was 80.2% and 38.7% of bacilli grown with usual amounts of Pi.

Figure. 1. Phosphate deprivation increases *M. bovis*/BCG phagocytosis. J774A.1 MOs (5×10^5) were incubated with *M. bovis*/BCG bacilli labeled red with the PKH-26; nuclei were stained blue with DAPI. Immunofluorescence microscopy shows a marked difference in the degree of phagocytosis between bacilli grown with and without Pi (a, b). FACS analysis considering the percent and the mean fluorescence index (MFI), showed dose-dependent phagocytosis that was significantly higher with Pi-deprived mycobacteria (c, d, e, f). At a 1:20 MOI, 80.2% of MOs phagocytosed bacilli cultured without Pi while the percent phagocytosis of bacilli cultured with Pi was 38.7. $p < 0.05$; unpaired t Student's test. Results of 5 independent experiments are shown.

Assays to verify the viability of phagocytosed mycobacteria grown with and without phosphate.

After 4 h of MO phagocytosis at 1:10 and 1:20 MOI the viability of ingested bacilli was assessed using the LIVE/DEAD BacLight bacterial viability Kit (Molecular Probes, Invitrogen, Carlsbad, CA EUA) following manufacturer's instructions. Bacilli were fixed in 4 % paraformaldehyde for 20 min and rinsed in TBS. Viability was analyzed by flow cytometry separating bacilli in two regions of the log-integrated red fluorescence propidium iodide versus Syto 9; the number of bacilli found within these regions were used to estimate the percentage of viability (Fig. 2). Viability is defined by the equation: measured % live bacilli = (# of bacilli in live region/# of bacilli in dead region) x 100. The results showed that at 1:10 MOI the viability of the Pi-deprived bacilli was significantly increased (Fig. 2).

Figure 2. Phosphate deprivation increases the viability of mycobacteria within macrophages. After 4 h phagocytosis, mycobacteria viability was analyzed by FACS using the LIVE/DEAD viability assay. A dot plot was set to differentiate living from dead cells (a). The survival rate was calculated using the formula: measured % live bacteria = (# of bacteria in live region / # of bacteria in dead region) x 100 (b). $p < 0.05$ Mann Whitney test. Results of 3 experiments are presented.

High phagocytosis of microbeads coated with plasma membrane proteins of phosphate-deprived mycobacteria

To study the role of mycobacteria proteins in the phagocytosis of bacilli grown under Pi starvation conditions, we carried out assays with fluorescent microbeads coated with proteins obtained from plasma membranes isolated from whole mycobacteria grown with and without Pi. For phagocytosis assays, MOs (5×10^6) cultured in RPMI 1640 with 10% heat-inactivated fetal bovine serum (FBS) were placed on acid-washed glass coverslips. Microbeads were added to cells at 1:5, 1:20 and 1:100 rate for 4 h at 37°C in 5% CO₂. To quantitate phagocytosis of microbeads, the slides were analyzed with a Nikon epifluorescence microscope examining at least 400 cells in random fields of 3 slides at each experimental time (Fig. 2). MOs were scored positive regardless of the number of bound or engulfed microbeads. Binding/phagocytosis of microbeads was dose-dependent and significantly higher with microbeads coated with membrane proteins obtained from bacilli grown without Pi (Fig. 2); at 1:100 rate, 84.44% of MOs bound or engulfed one or more microbeads. With microbeads coated with proteins from bacilli grown with Pi, binding/phagocytosis of microbeads was 45.4%.

Figure 3. Phagocytosis of membrane protein-coated microbeads is higher with phosphate deprived bacilli proteins. Assays were carried out with J774A.1 cells and green fluorescent microbeads coated with plasma membrane proteins obtained from mycobacteria grown with and without Pi. MOs (5×10^5) were placed on acid-washed glass coverslips. Microbeads were added to cells at a 1:5, 1:20 and 1:100 rate for 4 h at 37°C in 5% CO₂. The percentage of cells with ingested or bound microbeads was estimated counting at least 400 MOs (a, b). At 1:100 rate the phagocytosis of microbeads coated with proteins of Pi-deprived bacilli was 84.44% and of bacilli grown with Pi it was 45.4% (a, c). $p < 0.05$; unpaired t Student's test. Results of 5 experiments are presented.

Phosphate-deprived mycobacteria alter phagosome acidification

An outstanding ability of virulent mycobacteria is to inhibit phagosome acidification and phagolysosome fusion [2]. We studied the phagosomal acidification process by immunofluorescence microscopy using *M.*

bovis/BCG grown with or without Pi labelled green with FITC and MOs labelled with LysoTracker, a marker that turns red at low pH (Fig.4). Bacilli located in LysoTracker positive vacuoles were yellow fluorescent due to overlapping of FITC and the red fluorescence of LysoTracker. The engulfed bacilli emitted a FITC green fluorescence when were located in LysoTracker negative vacuoles (Fig.4). When infected with bacilli grown with usual amounts of Pi, 80.75 % of cells were LysoTracker positive and with Pi-deprived bacilli the percent of LysoTracker labeled MOs was reduced to 73.06% FACS analysis of the phagocytosis assays confirmed the immunofluorescence findings showing that Pi-deprived bacilli phagocytosis is associated with decreased LysoTracker labeling (Fig. 4).

Figure. 4. Decreased acidification in MOs that engulfed mycobacteria grown without phosphate.

Phagocytosis assays were performed with MOs labelled with LysoTracker red and FITC-labelled bacilli cultured with or without Pi. Immunofluorescence microscopy was performed counting randomly at least 400 MO in 3 different slides. MO emitting red fluorescence were scored as LysoTracker positive (a). It was found that 80.75% of MO incubated with mycobacteria grown with usual amount of Pi were LysoTracker labelled while incubation with Pi-deprived bacilli reduced the number of LysoTracker marked MO (76.06%, b). The degree of LysoTracker labelling was also analyzed by FACS which showed that the MIF was less in MO that phagocytosed Pi-deprived mycobacteria (c). The immunofluorescence patterns are shown; bacilli were located in non-acidified compartments (LysoTracker negative) are green fluorescent (arrows). In acidified vacuoles (LysoTracker positive) bacilli emitted yellow fluorescence due to overlapping of FITC-labelled bacilli and LysoTracker red fluorescence (asterisks). Four independent experiments were performed.

Phosphate-starved mycobacteria increment the expression of adhesins in isolated plasma membranes.

We have previously shown by immunoelectron microscopy that PstS-1 is a cell surface located adhesin that triggers the phagocytosis of mycobacteria [10]. To study its location in Pi- starved mycobacteria, we isolated cell walls and plasma membranes to carry out 1-D electrophoresis and immunoblot (Fig. 5). By Coomassie blue staining and immunoblot with a polyclonal anti-*M.bovis*/BCg antiserum the number of protein bands was higher in cell-walls of Pi-deprived mycobacteria. In the isolated plasma membranes immunoblotting with mAb available in our laboratory that recognizes adhesins that participate in the phagocytosis of mycobacteria 5 reactive bands of 38, 19, 23, and 47 kDa that correspond to PstS-1, LpqH, LprG, and the APA antigen, respectively (Fig. 5). In the membranes of bacilli grown with Pi band mild APA reactivity was observed.

Figure 5. Cell membrane located PstS-1, LpqH, LprG, and APA are virulence factors expressed by Pi-deprived mycobacteria. *M.bovis*/BCG bacilli were cultured with and without Pi for 24 h. The mycobacteria

were sonicated and by centrifugation the cell walls and the plasma membranes were obtained. The cell walls were resolved in 12% SDS-PAGE and transferred to PVDF for immunoblot. Coomassie blue stain and immunoblotting with an anti-*M. bovis*/BCG antiserum detected many protein bands more numerous in cell walls of Pi-deprived bacilli (a,b). Using mAbs available in our laboratory the expression of PstS-1, LpqH, LprG, and APA were demonstrated in isolated plasma membranes of Pi-deprived mycobacteria (c-f). These proteins are recognized virulence factors that participate as adhesins in the infection of host cells.

Discussion

In this study, we observed that *M. bovis*/BCG cultured without Pi acquire properties that could indicate increased virulence. First, bacilli were phagocytosed by MOs with increased efficiency, a property suggesting increased infectivity. Additionally, reduced phagosome acidification was detected in MOs that had phagocytosed Pi-deprived bacilli. These changes were associated with the expression in isolated plasma membranes of glycoproteins that have been characterized as adhesins that play a critical role in the infection of host cells.

Pulmonary TB is initiated when inhaled bacilli reach the alveoli, the terminal part of the respiratory tract, where they encounter host cells, especially MO, the best-equipped cell to contend with intracellular microbes [3]. After phagocytosis, mycobacteria can be lysed within the phagolysosome, although virulent bacilli have developed the ability to avoid destruction impeding the fusion of the phagosome with the lysosome, thus transforming the MO in a friendly niche where they proliferate and persist [2, 4].

It is noteworthy that mycobacteria subjected to the metabolic stress implied in Pi deprivation increment the expression of adhesins which are virulence molecules that bind to complementary molecules present in the surface of MOs and other host cells [11]. Information on the repertoire of Mtb adhesins is needed since it could help to design prevention and treatment strategies aimed to block the crucial step that is the infection of host cells by the mycobacteria. A few mycobacterial adhesins have been characterized including PstS-1, the 38-kDa antigen which is high-affinity protein that transports Pi [10,13]. It has been recognized its high antigenicity at the humoral level that allowed the design of diagnostic serologic tests [16, 17, 18, 19]. PstS-1 was characterized as a glycoprotein with mannose residues and more recently its role of adhesin in the phagocytosis of mycobacteria was recognized [10]. PstS-1 has focused much interest as a high-affinity Pi transporter that carries Pi from the periplasm to the cytoplasm. It has been reported that Pi-starved mycobacteria upregulates PstS-1 which is 30% similar in amino acid sequence to PstS-1 from *Escherichia coli* [12].

Our current findings suggesting increased virulence in mycobacteria subjected to Pi deprivation are in keeping with previous studies [20]. Pst genes are overexpressed in mycobacteria grown under Pi starvation conditions; mutations in these genes affect the import of Pi and reduces the proliferation of mycobacteria inside macrophages and in experimentally infected mice [21, 22]. On the other hand It has been observed that Mtb Pst system neutralizes IFN-gamma-dependent host immunity [22]

There is much information about the role membrane proteins of Mtb and other microbes play in the pathogenesis of infection [23]; their strategic location at the microbe surface allow their efficient interaction with host cells to promote phagocytosis and when they are antigenically active an immune response that can be cell or antibody-mediated [23]. Mycobacterial membrane proteins may also exert inhibitory effects on host cell activity. In this study we found that in addition to PstS-1, Pi-deprived mycobacteria express membrane proteins that are considered adhesins involved in the pathogenesis of the mycobacterial infection. LpqH, the 19 kDa glycolipoprotein, participates in MO infection as an adhesin that binds the mannose receptor triggering the phagocytosis of bacilli and downregulating MHC-II antigen presentation interfering with TLR2 and MAPK signaling [22]. The APA antigen plays a prominent role in Mtb virulence; it functions as a mycoyl-transferase that catalyzes the attachment of mycolic acids to arabinogalactan and the biogenesis of cord factor, a very active virulence factor [24]. APA is expressed at the bacterial surface, which explains its interaction with fibronectin, elastin and the surfactant protein A. LprG, the 25-kDa glycolipoprotein transports ManLAM to the mycobacterial cell-wall and as an adhesin it may bind the MO mannose receptor facilitating phagocytosis and inhibition of the phagosome-lysosome fusion. Isolated LprG was found to inhibit MHC-II Ag processing by THP-1 cells and primary human MO through prolonged TLR2 activation [25, 26].

Conclusion

We have found that the capacity of *M. bovis*/BCG to infect and survive inside MOs can be increased when mycobacteria are cultivated in the absence of Pi. MO phagocytosis by Pi-deprived mycobacteria was increased and phagosomal acidification was impaired. This phenotype was associated with an increase in the expression of PstS-1, the Pi transporter protein that is recognized as an adhesin that promotes the phagocytosis of mycobacteria. In addition, Pi-deprived mycobacteria upregulate the expression of the membrane located adhesins LpqH, LprG, and APA, which do not participate directly in Pi regulation.

Materials And Methods

Culture of *Mycobacterium bovis*/BCG under phosphate deprivation conditions

The method of Lefevre et al. was followed [16]. Briefly, *M. bovis*/BCG was cultured in Sauton medium with K_2HPO_4 (0.5 g/L) at 37°C, until optic density of 460 was reached. Thereafter, aliquots were taken and inoculated in Sauton medium that was supplemented with an excess of K_2HPO_4 to a final concentration of 3 g/L; the mycobacteria were cultured at 37°C until optic density of 460 was reached. After that, two aliquots were rinsed, centrifuged at 6,000g, and inoculated in Sauton medium. One aliquot was cultured without K_2HPO_4 and the other with 0.5 g/L K_2HPO_4 . After 24 h of culture, bacilli were collected, rinsed with TBS and frozen until use.

Phagocytosis assays with phosphate-deprived mycobacteria

We carried assays with mycobacteria that were grown with and without Pi as described above. The Balb/c-derived murine macrophage-like tumor cell line J774A.1 was obtained from the American Type Culture Collection (Manassas, VA, USA). Cells were cultured in RPMI 1640 supplemented with 10% fetal calf serum. MO (5×10^5) were incubated with PKH26 labeled bacilli at 1:2, 1:5, and 1:20 MOI, at 37°C for 4 h. After extensive rinsing with PBS, the cells were fixed in 4% paraformaldehyde in PBS for 20 min. For immunofluorescence microscopy, cytospin slides were mounted with ProLong Antifade with DAPI (Invitrogen, Eugene, OR, USA) and examined with a Nikon epifluorescence microscope. Phagocytosis was also analyzed by FACS.

Analysis of the mycobacterial viability after phagocytosis

Assays to verify the viability of phagocytosed mycobacteria were carried out. After 4 h phagocytosis of mycobacteria at 1:10 and 1:20 MOIs, MOs were recovered in TBS, lysed with 0.1 % sodium deoxycholate for 5 min, and centrifuged twice at 12,000 rpm for 10 min. Thereafter, to assess viability the LIVE/DEAD BacLight bacterial viability Kit (Molecular Probes) was used following the manufacturer instructions. Afterward, bacilli were fixed in 4 % paraformaldehyde for 20 min, rinsed in TBS and analyzed by flow cytometry. Because both live and dead cells exhibit green fluorescence, a threshold was set to eliminate debris. Populations of bacteria were discriminated as two regions of the log-integrated red fluorescence propidium iodide versus Syto 9, and the numbers of bacteria found within these regions were used to estimate the percentage of viable bacteria.

Binding/phagocytosis assays with microbeads coated with mycobacterial membrane proteins

We assessed a possible role of mycobacterial membrane proteins in the phagocytosis of mycobacteria employing green fluorescent Polystyrene 1 μ m microbeads (Molecular Probes, Eugene, OR, USA) coated with plasma membrane proteins obtained from mycobacteria grown with and without Pi. Microbeads and 500 μ g membrane proteins were mixed in 1 mL MES Buffer 2-[N-morpholino] ethane sulfonic acid (pH 6). After that, 1-ethyl-3-(3-dimethylamine npropyl)-carbodiimide was added to a 2.5/ml final concentration. After overnight incubation, 100 mM glycine was added to stop the reaction. To block remaining reactive sites, microbeads were rinsed with PBS and kept at 4°C in PBS containing 1% BSA. For phagocytosis assays, 5×10^5 MOs were cultured in RPMI 1640 with 10% heat-inactivated FBS and placed on acid-washed glass coverslips. Microbeads were added to cells at a 1:20 rate and incubated for 4 h at 37°C in 5% CO₂. The slides were rinsed in PBS to remove non-adherent microbeads and fixed in 1% paraformaldehyde for 10 min.

Phagosome maturation after phagocytosis of mycobacteria

For these analyses bacilli grown with and without Pi were labeled green with fluorescein isothiocyanate (FITC). Briefly, 50 µl of a bacilli suspension was suspended in 950 µl carbonate buffer 0.1 M, pH 9 and 0.5 mg FITC were added for 2 h. To finish labeling the mycobacteria were centrifuged at 10,500g for 10 min and rinsed 3 times in PBS. To examine whether Pi deprived mycobacteria acquired the ability to alter phagosome maturation, we use the acidotropic dye LysoTracker Red DND-99 (Molecular Probes, Eugene, OR). After rinsing, FITC labeled mycobacteria were added for 4 h at a 1:10 MOI. In a final step, LysoTracker was added in RPMI (1:20,000) for 20 min before phagocytosis was finished. MOs were rinsed, fixed with 4% paraformaldehyde and placed in glass slides. To verify LysoTracker labeling slides were examined by immunofluorescence microscopy. Besides, we quantitated by FACS the extent of LysoTracker phagosome labeling and the phagocytosis of FITC labeled bacilli.

Isolation of mycobacterial cell walls and plasma membranes to identify proteins expressed during phosphate deprivation

Mycobacteria were grown with and without Pi as described above. To obtain mycobacterial cell walls, bacilli were sonicated at 60 kHz in iced water (20 cycles 1 min each) and centrifuged at 10,500g to obtain the cell wall fraction in the precipitate and the plasma membrane in the supernatant. Protein concentration was measured by the Lowry protein assay (Bio-Rad Laboratories, Hercules, CA, USA). To identify cell wall proteins, membranes proteins they were separated by 12% SDS-PAGE, stained with Coomassie blue, transferred to PVDF membranes and incubated with a rabbit antiserum to M.bovis/BCG. Isolated plasma membranes were with mAb to LpqH, PstS-1 and APA and with Polyclonal antibodies to LprG (donated by PhD Clara Espitia); bound antibodies were detected with goat anti-mouse and a goat anti-rabbit IgG antibodies labeled with horseradish peroxidase. The reactive bands were visualized by chemiluminescence with SuperSignal West Dura kit (Pierce, Rockford, IL, USA).

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data are included in the manuscript

Competing interests

The authors declare that they have no competing interests

Funding

This work was funded by Programa de Apoyos a Proyectos de Investigación e Innovación Tecnológica No. IN210816 (URL: <http://dgapa.unam.mx/index.php/impulso-a-la-investigacion/papiit>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

PEC acquired, analyzed data and revised the manuscript; AMP acquired, analyzed data and revised the manuscript; RM designed the work, data interpretation and wrote the original manuscript.

References

1. Global Tuberculosis Report 2017. Geneva, Switzerland: WHO Press. 2017.
2. Clemens DL, Horwitz MA. Characterization of the *Mycobacterium tuberculosis* phagosome and evidence that phagosomal maturation is inhibited. *J Exp Med*. 1995;181:257-70.
3. Ernst JD. Macrophage receptors for *Mycobacterium tuberculosis*. *Infect Immun*. 1998; 66:1277–81.
4. Schlesinger LS, Hull SR, Kaufman TM. Binding of the terminal mannosyl units of lipoarabinomannan from a virulent strain of *Mycobacterium tuberculosis* to human macrophages. *J Immunol*. 1994;152:4070–9.
5. Diaz-Silvestre H, Espinosa-Cueto P, Sanchez-Gonzalez A, Esparza-Ceron MA, Pereira-Suarez AL, Bernal-Fernandez G, et al. The 19-kDa antigen of *Mycobacterium tuberculosis* is a major adhesin that binds the mannose receptor of THP-1 monocytic cells and promotes phagocytosis of mycobacteria. *Microbial Pathogenesis*. 2005; 39:97–107.
6. Menozzi FD, Bischoff R, Fort E, Brennan MJ, Locht C. Molecular characterization of heparin-binding hemagglutinin, a mycobacterial adhesin. *Proc Natl Acad Sci USA*. 1998; 95:12625–1263.
7. Palucci I, Camassa S, Cascioferro A, Sali M, Anoosheh S, Zumbo A, et al. PE_PGRS33 Contributes to *Mycobacterium tuberculosis* entry in macrophages through interaction with TLR2. *PLoS One*. 2016. doi:10.1371/journal.Pine.0150800.
8. Drage MG, Tsai HC, Pecora ND, Cheng TY, Arida AR, Shukla S, et al. *Mycobacterium tuberculosis* lipoprotein LprG (Rv1411c) binds triacylated glycolipid agonists of toll-like receptor 2. *Nat Struct Mol Biol*. 2010; 17:1088–1095.
9. Ragas A, Roussel L, Puzo G, Riviere M. The *Mycobacterium tuberculosis* cell surface glycoprotein apa as a potential adhesin to colonize target cells via the innate immune system pulmonary C-type lectin surfactant protein A. *J Biol Chem* 2007; 282:5133-5142.
10. Esparza M, Palomares B, García T, Espinosa P, Zenteno E, Mancilla R. PstS-1, the 38-kDa *Mycobacterium tuberculosis* glycoprotein, is an adhesin, which binds the macrophage mannose receptor and promotes phagocytosis. *Scand J Immunol*. 2015; 81:46-55.

11. Govender VS, Ramsugit S, Pillay M. *Mycobacterium tuberculosis* adhesins: Potential biomarkers as anti-tuberculosis therapeutic and diagnostic targets. *Microbiology* 2014; 160:1821-1831.
12. Andersen AB, Ljungqvist L, Olsen M. Evidence that protein antigen b of *Mycobacterium tuberculosis* is involved in phosphate metabolism. *J. Gen. Microbiol.* 1990; 136:477–480.
13. Espitia C, Elinos M, Hernández-Pando R, Mancilla R. Phosphate starvation enhances expression of the immunodominant 38-kilodalton protein antigen of *Mycobacterium tuberculosis*: demonstration by immunogold electron microscopy. *Infect Immun.* 1992; 60:2998-3001.
14. Braibant M, Lefèvre P, De Wit L, Peirs P, Ooms J, Huygen K, et al, Content J. A *Mycobacterium tuberculosis* gene cluster encoding proteins of a phosphate transporter homologous to the *Escherichia coli* Pst system. *Gene.* 1996;176:171–176.
15. Santo-Beneit F. The Pho regulon: a huge regulatory network in bacteria. *Front Microbiol.* 2015; 6:402. doi: 10.3389/fmicb.2015.00402
16. Harboe M, Wiker HG. The 38-kDa protein of *Mycobacterium tuberculosis*: a review. *J Infect Dis.* 1992; 166:874-84.
17. Wilkinson RJ, Hasløv K, Rappuoli R, Giovannoni F, Narayanan PR, Desai CR, et al. Evaluation of the recombinant 38-kilodalton antigen of *Mycobacterium tuberculosis* as a potential immunodiagnostic reagent. *J Clin Microbiol.* 1997; 35:553-7.
18. Espitia C, Cervera I, Gonzalez R, Mancilla R. A 38-kD *Mycobacterium tuberculosis* antigen associated with infection. Its isolation and serologic evaluation. *Clin. Exp. Immunol.* 1989; 77:373-377.
19. Uma Devi KR, Ramalingam B, Brennan PJ, Narayanan PR, Raja A. Specific and early detection of IgG, IgA and IgM antibodies to *Mycobacterium tuberculosis* 38-kDa antigen in pulmonary tuberculosis. 2001; 81:249 – 53.
20. Peirs P, Lefevre P, Boarbi S, Wang XM, Denis O, Braibant M, et al. *Mycobacterium tuberculosis* with disruption in genes encoding the phosphate binding proteins PstS1 and PstS2 is deficient in phosphate uptake and demonstrates reduced in vivo virulence. *Infect Immun.* 2005;73:1898–1902.
21. Lamarche MG, Wanner BL, Crépin S, Harel J. The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. *FEMS Microbiol.* 2008;32:461–473.
22. Pennini ME, Pai RK, Schultz DC, Boom WH, Harding CV. *Mycobacterium tuberculosis* 19-kDa Lipoprotein Inhibits IFN- γ -Induced Chromatin Remodeling of MHC2TA by TLR2 and MAPK Signaling. *J Immunol.* 2006;176:4323–4330.
23. Målen H, Berven FS, Sjøfteland T, Arntzen MØ, D'Santos CS, De Souza GA, et al. Membrane and membrane-associated proteins in Triton X-114 extracts of *bovis*/BCG identified using a combination of gel-based and gel-free fractionation strategies. *Proteomics.* 2008;8(9):1859-70.
24. Karbalaee Zadeh Babaki Z, SoleimanPur S, Rezaee SA. Antigen 85 complex as a powerful *Mycobacterium tuberculosis* immunogen: Biology, immune-pathogenicity, applications in diagnosis, and vaccine design. *Microbial Pathogenesis.* 2017;112:20-29.

25. Gaur RL, Ren K, Blumenthal A, Bhamidi S, Gonzáles-Nilo FD, Jackson M, Zare RN, et al. LprG-mediated surface expression of lipoarabinomannan is essential for virulence of *Mycobacterium tuberculosis*. PLoS Pathog. 2014;10(9):e1004376 doi:10.1371/journal.ppat.1004376.
26. Gehring AJ, Dobos KM, Belisle JT, Harding CV, Boom WH. *Mycobacterium tuberculosis* LprG (Rv1411c): A Novel TLR-2 ligand that inhibits human macrophage Class II MHC antigen processing. J Immunol. 2004;173:2660–2668.

Figures

Figure 1

Phosphate deprivation increases *M. bovis*/BCG phagocytosis. J774A.1 MOs (5×10^5) were incubated with *M. bovis*/BCG bacilli labeled red with the PKH-26; nuclei were stained blue with DAPI. Immunofluorescence microscopy shows a marked difference in the degree of phagocytosis between bacilli grown with and without Pi (a, b). FACS analysis considering the percent and the mean fluorescence index (MFI), showed dose-dependent phagocytosis that was significantly higher with Pi-deprived

mycobacteria (c, d, e, f). At a 1:20 MOI, 80.2% of MOs phagocytosed bacilli cultured without Pi while the percent phagocytosis of bacilli cultured with Pi was 38.7. $p < 0.05$; unpaired t Student's test. Results of 5 independent experiments are shown.

Figure 2

Phosphate deprivation increases the viability of mycobacteria within macrophages. After 4 h phagocytosis, mycobacteria viability was analyzed by FACS using the LIVE/DEAD viability assay. A dot plot was set to differentiate living from dead cells (a). The survival rate was calculated using the formula: measured % live bacteria = ($\#$ of bacteria in live region / $\#$ of bacteria in dead region) $\times$ 100 (b). $p < 0.05$ Mann Whitney test. Results of 3 experiments are presented.

Figure 3

Phagocytosis of membrane protein-coated microbeads is higher with phosphate deprived bacilli proteins. Assays were carried out with J774A.1 cells and green fluorescent microbeads coated with plasma membrane proteins obtained from mycobacteria grown with and without Pi. MOs (5×10^5) were placed on acid-washed glass coverslips. Microbeads were added to cells at a 1:5, 1:20 and 1:100 rate for 4 h at 37°C in 5% CO₂. The percentage of cells with ingested or bound microbeads was estimated counting at

least 400 MOs (a, b). At 1:100 rate the phagocytosis of microbeads coated with proteins of Pi-deprived bacilli was 84.44% and of bacilli grown with Pi it was 45.4% (a, c). $p < 0.05$; unpaired t Student's test. Results of 5 experiments are presented.

Figure 4

Decreased acidification in MOs that engulfed mycobacteria grown without phosphate. Phagocytosis assays were performed with MOs labelled with LysoTracker red and FITC-labelled bacilli cultured with or without Pi. Immunofluorescence microscopy was performed counting randomly at least 400 MO in 3 different slides. MO emitting red fluorescence were scored as LysoTracker positive (a). It was found that 80.75% of MO incubated with mycobacteria grown with usual amount of Pi were LysoTracker labelled while incubation with Pi-deprived bacilli reduced the number of LysoTracker marked MO (76.06%, b). The degree of LysoTracker labelling was also analyzed by FACS which showed that the MIF was less in MO that phagocytosed Pi-deprived mycobacteria (c). The immunofluorescence patterns are shown; bacilli were located in non-acidified compartments (LysoTracker negative) are green fluorescent (arrows). In acidified vacuoles (LysoTracker positive) bacilli emitted yellow fluorescence due to overlapping of FITC - labelled bacilli and LysoTracker red fluorescence (asterisks). Four independent experiments were performed.

Figure 5

Cell membrane located PstS-1, LpqH, LprG, and APA are virulence factors expressed by Pi-deprived mycobacteria. *M. bovis*/BCG bacilli were cultured with and without Pi for 24 h. The mycobacteria were sonicated and by centrifugation the cell walls and the plasma membranes were obtained. The cell walls were resolved in 12% SDS-PAGE and transferred to PVDF for immunoblot. Coomassie blue stain and immunoblotting with an anti-*M. bovis*/BCG antiserum detected many protein bands more numerous in cell walls of Pi-deprived bacilli (a,b). Using mAbs available in our laboratory the expression of PstS-1, LpqH, LprG, and APA were demonstrated in isolated plasma membranes of Pi-deprived mycobacteria (c-f). These proteins are recognized virulence factors that participate as adhesins in the infection of host cells.