

Mechanism of Quercetin Therapeutic Targets for Alzheimer Disease and type 2 Diabetes Mellitus

Guoxiu Zu

Shandong University of Traditional Chinese Medicine

Keyun Sun

Shandong University of Traditional Chinese Medicine

Ling Li

Shandong University of Traditional Chinese Medicine

Xiuli Zu

Shandong University of Chinese Medicine

Tao Han

Shandong University of Traditional Chinese Medicine

Hailiang Huang (✉ 06000031@sducm.edu.cn)

Shandong University of Traditional Chinese Medicine

Research Article

Keywords: quercetin, type 2 diabetes mellitus, Alzheimer's disease, associated diseases, comorbid diseases, drugs targeting multiple diseases, network pharmacology

Posted Date: August 31st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-853709/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on November 25th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-02248-5>.

Abstract

Quercetin has demonstrated antioxidant, anti-inflammatory, hypoglycemic, and hypolipidemic activities, suggesting therapeutic potential against type 2 diabetes mellitus (T2DM) and Alzheimer's disease (AD). In this study, potential molecular targets of quercetin were first identified using the Swiss Target Prediction platform and pathogenic targets of T2DM and AD were identified using Online Mendelian Inheritance in Man (OMIM), DisGeNET, TTD, DrugBank, and GeneCards databases. The 95 targets shared among quercetin, T2DM, and AD were used to establish a protein–protein interaction (PPI) network, top 25 core genes, and protein functional modules using MCODE. Metascape was then used for Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis. A protein functional module with best score was obtained from the PPI network using CytoHubba, and 6 high-probability quercetin targets (AKT1, JUN, MAPK, TNF, VEGFA, and EGFR) were confirmed by docking simulations. KEGG pathway enrichment analysis suggested that the major shared mechanisms for T2DM and AD include “AGE-RAGE signaling pathway in diabetic complications,” “pathways in cancer,” and “MAPK signaling pathway” (the key pathway). We speculate that quercetin may have therapeutic applications in T2DM and AD by targeting MAPK signaling, providing a theoretical foundation for future clinical research.

1. Introduction

According to the 2017 global burden of disease study, type 2 diabetes mellitus (T2DM) and Alzheimer's disease (AD) are the sixth and tenth leading causes of premature death worldwide,¹ and many studies have found strong epidemiological, pathological, and clinical associations between T2DM and AD.²⁻⁷ The incidence of AD is about twofold greater in T2DM patients than in non-diabetic individuals, while a disproportionate number of AD patients have a history of early or clinical diabetes. Diabetes and AD also share numerous common risk factors, including metabolic disease (obesity), cardiovascular disease, older age, and progressive insulin resistance,⁴ as well as common pathogenic processes such as chronic hyperglycemia, tissue amyloid β (A β) deposition and toxicity, cellular oxidative stress, and chronic inflammation.

Indeed, the term “type 3 diabetes” has been proposed to describe AD based on the hypothesis that AD-associated dementia is caused in part by insulin resistance and insufficient insulin-like growth factor signaling in the brain.³ It is believed that endocrine and metabolic diseases characterized by elevated blood glucose due to insulin resistance and death of insulin-producing pancreatic islet β -cells can lead to chronic cardiovascular diseases and ultimately to AD. Alzheimer's disease is characterized by the accumulation of A β in the brain parenchyma (termed senile plaques, SPs) and abnormal hyperphosphorylation of the intermediate filament protein tau within neurons, forming neurofibrillary tangles (NFTs). The presence of these inclusions results in synaptic loss, particularly in neocortex and hippocampus, oxidative stress, and neuroinflammation, ultimately leading to neuronal dysfunction and death.⁵ Tau protein and human islet amyloid polypeptide (hIAPP) deposits are also found in the

pancreatic islets of T2DM patients. Moreover, pharmacological studies have shown that some traditional hypoglycemic drugs can slow the progression of AD.⁷

While there is strong evidence for an association between T2DM and AD at multiple levels, the seminal molecular mechanisms linking these disorders remain unknown. One potential strategy for identifying these molecular linkage pathways is to screen for common drug targets, which would also provide new possibilities for the treatment of T2DM and AD.

Quercetin (3, 3', 4', 5, 7-pentahydroxyflavone) is a natural polyhydroxyflavonoid found in the flowers, leaves, and fruits of edible plants such as onion, apple, lettuce, and cabbage.^{2,8} In addition, many Chinese herbal medicines such as *Sophora japonica*, *Ginkgo biloba* leaves, *Hypericum perforatum*, and cortex moutan are enriched in quercetin.⁹ Numerous studies have documented bioactivities of quercetin relevant to T2DM and AD pathology, such as antioxidant, anti-inflammatory, hypoglycemic, and hypolipidemic effects. Quercetin has also been shown to reduce blood glucose concentration and preserve islet function, insulin sensitivity, and β -cell numbers in diabetic model rats and mice.^{10,11}

At present, there is no broadly effective drug treatment to delay AD onset, slow progression, or improve outcome,¹² and the few approved medications can only transiently improve symptoms.¹³ Also, many synthetic drugs with antioxidant, anti-inflammatory, hypoglycemic, and (or) hypolipidemic activities induce adverse events that preclude clinical use. The use of natural substances to treat neurodegenerative diseases such as AD and metabolic diseases such as T2DM is a promising alternative as such agents are often easily and inexpensively isolated and have well documented biomechanisms and safety profiles. Recent epidemiological and experimental studies have determined that Chinese herbal medicines as well as fruit and vegetable juices containing flavonoids can reduce the probability of cognitive decline. Quercetin has also shown anti-dementia and neuroprotective efficacies in models of dementia/AD and ischemia–reperfusion injury by mitigating neuronal oxidative stress and neuroinflammation, and by improving calcium homeostasis, growth factor signaling, and neuroplasticity,¹⁴⁻¹⁸ ultimately preventing neuronal apoptosis.¹⁸ Further, a number of large double-blind clinical studies have demonstrated that quercetin can reduce the composite end-point of death in patients with T2DM and AD. In the present study, we used a systematic bioinformatics approach (network pharmacology) and molecular docking simulations to identify quercetin-binding targets that may also be potential therapeutic targets for AD and T2DM.

2. Materials And Methods

2.1 Screening of potential quercetin targets

The Traditional Chinese Medicine Systems Pharmacology and Analysis Platform (*TCMSP*) was used to search for preparations with “quercetin” as a key word, and results were then combined with the Chinese Medicine Encyclopedia database (ETCM) to find Chinese herbal medicines enriched in quercetin. Quercetin targets were then identified using the PubChem database with “quercetin” as the key word.

Based on the notion that drugs of similar structure will have common (overlapping) targets, five potential targets of quercetin were predicted using Swiss Target Prediction (<http://www.swisstargetprediction.ch/>)¹⁹ with species “Homo sapiens” and a probability >0.1 as the selection criteria. Target names were standardized according to Uniprot protein database terminology (<https://www.uniprot.org/>).²⁰

2.2 Screening of quercetin targets for T2DM and AD treatment

Pathogenic molecules contributing to T2DM and AD was screened using Online Mendelian Inheritance in Man (OMIM, <https://omim.org/>)²¹ TTD (<http://db.idrblab.net/ttd/>), DisGeNET (<https://www.disgenet.org/>), DrugBank (<https://www.drugbank.ca/>),²² and GeneCards (<https://www.genecards.org/>)²³ using the terms “Type 2 diabetes mellitus” and “Alzheimer’s Disease.” By intersecting quercetin targets with disease-associated molecules targets using a Venn diagram, we obtained targets potentially relevant to T2DM and AD treatment. Cytoscape software²⁴ was then used to map the component-target-disease network as described in detail below.

2.3 Construction of protein–protein interaction (PPI) network

To identify interaction targets of quercetin potentially relevant to T2DM and AD pathology and treatment, we used the online drawing tool Interactive Venn (<http://www.interactivenn.net/>)²⁵ to construct a Venn diagram. The overlapping targets were then uploaded to the STRING11.0 platform (<https://string-db.org/>)²⁶ to obtain a protein–protein interaction (PPI) network. The maximal clique centrality (MMC) algorithm in the CytoHubba plug-in was used for comprehensive analysis of network topology to obtain the top 25 hub genes.²⁷ Then MCODE,²⁸ a cluster analysis plug-in for Cytoscape3.6.0 (<https://cytoscape.org/>), was utilized to further analyze the PPI network. The targets in the protein functional module with best MCODE score were selected as core targets for further molecular docking simulations.

2.4 Enrichment analysis

Gene Ontology (GO) functional analysis and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis were performed using Metascape (<https://metascape.org/>),²⁹ with $p < 0.01$ as the cutoff criterion. The top 10 GO items and top 20 KEGG pathways that meet this criterion were visualized using OmicShare (<http://www.omicshare.com/>) and microscopic letter (<http://www.bioinformatics.com.cn>).

2.5 Construction of a component-target-pathway network

Cytoscape3.6.0 was used to construct a component-target-pathway network based on KEGG enrichment analysis, and the topological parameters of the network were analyzed using the built-in Network Analyzer tool.

2.6 Molecular docking verification

The PDB formats of the core target proteins were downloaded from the RCSB PDB database (<http://www.rcsb.org/>). We used PyMol 2.4 software to remove water molecules and separate the original ligand from the core target protein, and then imported the processed protein targets into AutoDock Tools 1.5.6 software³⁰ for hydrogenation, calculation of total charge, and setting the atomic type. Results were saved in “pdbqt” format. The mol2 structures of the core targets were downloaded from the TCMSP database, and AutoDock Tools was used to set the rotatable bonds. Files were again saved in “pdbqt” format and then imported to AutoDock Vina to perform molecular docking. Finally, PyMol software was used to visualize the docking results and establish the docking interaction pattern. A flowchart of the network pharmacology study is shown in **Figure 1**.

3. Results

3.1 Candidate quercetin targets for T2DM and AD treatment

After screening the literature and online databases, we identified 263 candidate gene products potentially targeted by quercetin. Screening for T2DM pathogenic targets retrieved 30 genes from OMIM, 175 from TTD, one from DisGeNET (with relevance score >0.1), 266 from DrugBank, and 1108 from GeneCards (relevance score >1). For AD, we retrieved 9 gene products from OMIM, 143 from TTD, 218 from DisGeNET, 67 from DrugBank, and 2712 from GeneCards. After removing duplicates, 1294 T2DM-related and 2865 AD-related genes were included in subsequent analyses. To obtain quercetin targets with relevance to T2DM and AD pathology and treatment, we constructed a Venn diagram, and a total of 95 overlapping targets were obtained (**Figure 2**).

3.2 PPI network of quercetin targets for T2DM and AD treatment.

To estimate the contributions of these candidate common targets to disease and reveal the interactive effects, all 95 were imported into the STRING database with highest confidence set at 0.9 to obtain a PPI network. The PPI network results were saved in TSV text format and imported into Cytoscape3.6.0 for network topology analysis to identify the top 25 targets (arranged in descending order of relevance as a concentric circle in **Figure 3-A**). The PPI network (**Figure 3-B**) includes 273 edges among the 95 nodes

(circles), where edge thickness represents the interaction strength. To further refine the list of quercetin targets relevant to T2DM and AD treatment, we conduct cluster analysis on the PPI network based on shared mechanisms using MCODE. This analysis yielded a function module of five key nodal proteins (**Figure 3-C**). It is generally believed that proteins in such functional modules are more closely related and may interact to perform specific biological functions. Therefore, these proteins may have important regulatory effects on T2DM and AD.

3.3 Possible molecular targets of quercetin and associated regulatory signaling pathways relevant to T2DM and AD treatment.

We further conducted GO functional analysis and KEGG pathway enrichment analysis to elucidate the biological effects of quercetin targets and related signaling pathways relevant to T2DM and AD treatment. The top 10 GO items was selected based on p-values (**Figure 4**). Similarly, the top 20 KEGG pathway items were selected based on p-values (**Figure 5-A**) and arranged according to category (**Figure 5-B**).

The common targets were enriched in molecules with GO molecular function terms “cytokine receptor binding,” “serine hydrolase activity,” “oxidoreductase activity,” “protein kinase binding,” and “antioxidant activity,” biological processes terms “positive regulation of cell apoptosis” (GO:0097190, GO:0010942, GO:0008285), “regulation of inflammatory response” (GO:0050727), “regulation of reactive oxygen species metabolic process” (GO:0072593), “cellular response to external stimulus” (GO:0071496), and “regulation of protein serine/threonine kinase activity” (GO:0071900), and cellular components terms “membrane raft,” “external side of plasma membrane,” “extracellular matrix,” “vesicle lumen,” “endoplasmic reticulum lumen,” “cell body,” “receptor,” “perinuclear region of cytoplasm,” “actin-based cell projection,” and “apical part of cell.” According to KEGG pathway enrichment analysis, most of these common targets are involved in “MAPK signaling pathway,” “TNF signaling pathway,” “AGE-RAGE signaling pathway in diabetic complications,” “pathways in cancer,” “proteoglycans in cancer,” “interleukin (IL)-17 signaling pathway,” “hypoxia-inducible factor (HIF)-1 signaling pathway,” “PI3K-Akt signaling pathway,” “EGFR tyrosine kinase inhibitor resistance,” “Toll-like receptor signaling pathway,” “NF-kappa B signaling pathway,” “cytokine-cytokine receptor interaction,” “VEGF signaling pathway,” and (or) “Alzheimer’s disease.” Thus, the common targets of quercetin, T2DM, and AD are strongly enriched in “MAPK signaling pathway” components as indicated by four of the MCODE nodes (Fig. 3-C) and two of the top 10 GO molecular function items. In addition, most of the biological processes items are associated with MAPK activity, such as “kinase activity regulation,” “cell proliferation,” and “apoptosis,” suggesting that the MAPK signaling pathway is the core quercetin target relevant to T2DM and AD treatment.

3.4 Component-target-pathway network construction

A component-target-pathway network was constructed based on KEGG pathway enrichment analysis using Cytoscape3.6.0 (**Figure 6**) in which the importance of a given node is indicated by the number of connections to other nodes. The average degree of importance (average number of connection to a given node) was 8.64, and there were 11 targets with degrees higher than 8.64: MAPK1 (degree = 18), AKT1 (degree = 15), PIK3R1 (degree = 14), IL6 (degree = 13), PRKCB (degree = 12), TNF (degree = 12), VEGFA (degree = 11), EGFR (degree = 11), CASP3 (degree = 10), BCL2 (degree = 9), and IL1B (degree = 9). Since MAPK1, AKT1, and EGFR were also included in the best scoring protein functional module, we speculate that these may be the key targets of quercetin for T2DM and AD treatment.

3.5 Molecular docking results analysis

Finally, the interactions with core proteins in the PPI, MCODE, and component-target-pathway networks were verified by molecular docking simulations. The most likely and stable molecular interactions are generally those with lower energy after binding. Six core molecules were tested in docking simulations (**Table 1** and **Figure 7**), and all showed good binding affinity with an average free energy change of -6.47 kcal/mol. For the quercetin–MAPK1 docking model, the small molecule ligand dapagliflozin potentially fit into the interface pocket formed by interacting amino acid residues in the protein. As shown in Figure 7, a hydrogen bond was formed between quercetin and VAL302, HIS297, HIS139, and LYS205 near the active site of MAPK1.

PDB ID: Protein Data Bank Identification Code

Table 1

Binding energy of quercetin with MAPK1, AKT1, VEGFA, EGFR, JUN, and TNF

Bioactive component	Target name	PDB ID	Binding energy (kcal/mol)
	MAPK1	5K4I	-6.28
	AKT1	2UZR	-5.94
quercetin	VEGFA	6D30	-5.81
	EGFR	3POZ	-5.76
	JUN	5T01	-6.64
	TNF	7ATB	-8.4

4. Discussion

Type 2 diabetes is the third most common non-infectious disease in the world. It is characterized by poor blood glucose control due to insulin resistance and progression apoptotic death of insulin-producing islet

β -cells.^{31,32} As T2DM progresses, the loss of insulin production creates an absolute insulin deficiency that can only be addressed by exogenous replacement. Therefore, enhancing insulin sensitivity and reducing β -cell apoptosis are the ultimate strategies for the prevention and treatment of T2DM.³³ Several studies have found that quercetin can reduce blood glucose, peripheral insulin resistance, and indicators of oxidative stress induced by high glucose in rats, a response profile similar to the widely used antidiabetic drug metformin.

Alzheimer's disease is the most common neurodegenerative disorder. Clinically, it is characterized by progressive cognitive impairment, memory loss, and marked changes in personality and behavior. The two primary hypotheses proposed to explain the progressive neurodegeneration and neurological impairments of AD are cytotoxic β -amyloid deposition (plaque formation) due to abnormal processing and hyperphosphorylation of tau as these are the pathological hallmarks of the AD brain. In turn, these plaques and NFTs may cause oxidative stress and inflammation, leading to synaptic failure, neuronal apoptosis (especially basal forebrain cholinergic neurons and various hippocampal and cortical neurons), and brain atrophy.^{34,35} These pathomechanisms are not mutually exclusive and indeed may interact, creating pathogenic cascades. Thus, many protein signaling pathways and gene regulation pathways may contribute to the pathological changes in AD.

Many natural components found in traditional herbal preparations have documented antioxidant, anti-inflammatory, antidiabetic, and anti-apoptotic activities, suggesting potential utility for AD and T2DM treatment. For instance, quercetin can improve insulin production by rat insulinoma INS-1 cells in response to high glucose, as well as downregulate the relative expression levels of pro-apoptotic Bax and PDX-1 mRNAs, inhibit oxidative stress caused by H_2O_2 , and reduce INS-1 cell apoptosis.³⁶ Quercetin can also clear reactive oxygen species (ROS) and reactive nitrogen species and inhibit the level of low density lipoprotein peroxidase, thereby reducing the incidence of cardiovascular disease.³⁷ Further, quercetin was shown to reduce ischemic injury and peroxynitrate injury by inhibiting nitric oxide synthase and xanthine dehydrogenase activities.³⁸

The current study revealed several potential molecular mechanisms for AD and T2DM as well as possible treatment targets by combining network pharmacology with molecular docking. By integrating and collating information from several databases, we identified 95 potential quercetin targets involved in T2DM and AD pathology, and further defined a potential protein functional module of five proteins and 25 core genes from the PPI network. The biological processes mediated by these functional module proteins include protein kinase signaling and cell proliferation, consistent with pathogenic pathways of T2DM and AD identified in previous studies.

CytoHubba and MCODE identified AKT1, JUN, MAPK, TNF, VEGFA, and EGFR as core targets of quercetin that may contribute to AD and T2DM. The PI3K/Akt signaling pathway is anti-apoptotic, and AKT1 inhibits Akt phosphorylation (activation), which in turn upregulates the activity of GSK3 β , a kinase implicated in the pathogenesis of both T2DM and AD. JUN is a member of the mitogen-activated protein kinase (MAPK) family implicated in the onset, progression, and reversal of vascular diseases through

modulation of vascular cell protein expression, proliferation, oxidative stress, and apoptosis. Downregulation of JUN can significantly reduce the expression of inflammatory factors and inhibit endothelial cell apoptosis.³⁹ Tumor necrosis factor (TNF) is a multifunctional cytokine that can directly damage islet β -cells and induce insulin resistance by inhibiting the transduction of insulin signals. MAPK is a serine/threonine protein kinase involved in cell proliferation, differentiation, apoptosis, and inflammatory responses.⁴⁰ Activation of MAPK can also increase tau phosphorylation in neurons,⁴¹ while inhibition of MAPK expression in hippocampus can significantly improve memory, cognitive function, synaptic plasticity, and neuronal metabolism.⁴² Suppression of MAPK can also improve insulin resistance by upregulating the expression of the insulin-dependent glucose transporter GLUT-4 in striatal muscle and adipose tissue, and by enhancing the sensitivity of insulin receptors, thereby promoting the absorption and utilization of glucose. Therefore, MAPK may be a core target for the treatment of T2DM and AD.⁴³

To better understand the interactions among these target genes, we conducted GO and KEGG pathway analyses. The target genes were mainly related to cell apoptosis, regulation of inflammatory response, regulation of ROS, metallic process, and other biological processes. KEGG pathway enrichment analysis revealed that target proteins are involved in the MAPK, TNF, HIF-1, IL-17, PI3K, NF-kappa B, Toll-like receptor, and VEGF signaling pathways, AGE-RAGE signaling pathway in diabetic complications, pathways in cancer, proteoglycans in cancer, EGFR tyrosine kinase inhibitor resistance, cytokine-cytokine receptor interaction, and Alzheimer's disease. We hypothesize that quercetin may play a therapeutic role in T2DM and AD through the regulation of these pathways, but especially through modulation of MAPK signaling (**Figure 8**).

A variety of extracellular signaling molecules (growth factors, neurotransmitters, cytokines, hormones, etc.) stimulate MAPK signaling via the MAPKKK-MAPKK cascade, ultimately regulating both cytoplasmic processes and gene expression pathways underlying cell proliferation, differentiation, apoptosis, and stress responses. Three parallel MAPK signaling pathways are found in mammalian cells, extracellular signal regulated kinase (ERK), c-Jun N-terminal kinase (JNK), and p38 mitogen-activated protein kinase (p38 MAPK). Inhibiting MAPK signaling can reduce oxidative stress and ensure the normal proliferation, differentiation, and insulin secretion capacity of islet cells.⁴⁴ Further, inhibition of the MAPK pathway can improve learning and memory⁴⁵ and suppress apoptosis of hippocampal CA1 neurons by downregulating the transcription factor c-Fos. Activation of the MAPK/ERK pathway may be an important mechanism underlying early and medium-term cognitive impairment in rats, while inhibition can reduce damage and promote the repair of damaged neurons. Quercetin acts on membrane EGFRs and TNFR1 by promoting the release of VEGFA and TNF, thereby affecting the activation of Ras, Raf, MEK, and phosphorylation of MAPK1 and MAPK3, leading to changes in angiogenesis, cell apoptosis, and cell proliferation. More in-depth studies on these mechanisms and preclinical studies in T2DM and AD animal models are warranted to provide a foundation for future clinical studies targeting the MAPK signaling pathway as a novel therapeutic strategy for T2DM and AD.

Conclusion

In conclusion, we have identified several potential molecular mechanisms by which quercetin can simultaneously interfere with T2DM and AD progression. Among these, suppression of MAPK signaling appears to be the most promoting strategy for the treatment of T2DM and AD.

Declarations

Author Contribution Statement

Hai-Liang Huang and Tao Han, as corresponding authors with equal contributions, conceived and designed the research. Guo-Xiu Zu performed the research and wrote the paper. Ke-Yun Sun arranged the data and revised the manuscript. Guo-Xiu Zu, Xiu-Li Zu, Ke-Yun Sun, Ling Li, and Hai-Liang Huang analyzed the data and drew figures. Guo-Xiu Zu and Ke-Yun Sun are equal contributors.

Acknowledgments

This work was supported by a grant from the Key Program of Shandong Province, China (No. 2016CYJS08A01-6).

This work was supported by the first batch of outstanding research and innovation team of Shandong University of Traditional Chinese Medicine. Mechanism and effect evaluation of prevention of major diseases (220316).

Guoxiu Zu received a graduate studentship award from the University of "Shandong university of Traditional Chinese Medicine"

The authors would like to thank Yinglun Pavilion (www.enago.cn) for providing English touch up.

Conflicts of Interest Statement

The authors declare that they have no competing interests.

Data Statement

The data that supports the findings of this study are available in the supplementary material of this article.

References

1. GBD 2017 Causes of Death Collaborators. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*, **392** (10159), 1736–1788 (2018).
2. van Sloten, T. T., Sedaghat, S., Carnethon, M. R., Launer, L. J. & Stehouwer, C. D. A. Cerebral microvascular complications of type 2 diabetes: stroke, cognitive dysfunction, and depression. *Lancet Diabetes Endocrinol*, **8** (4), 325–336 (2020).
3. Neth, B. J. & Craft, S. Insulin resistance and Alzheimer's disease: bioenergetic linkages. *Front Aging Neurosci*, **9**, 345 (2017).
4. de la Monte, S. M. Contributions of brain insulin resistance and deficiency in amyloid-related neurodegeneration in Alzheimer's disease. *Drugs*, **72** (1), 49–66 (2012).
5. Rabinovici, G. D. *et al.* Testing and disclosures related to amyloid imaging and Alzheimer's disease: common questions and fact sheet summary. *Alzheimers Dement*, **12** (4), 510–515 (2016).
6. Huang, C. C. *et al.* Diabetes mellitus and the risk of Alzheimer's disease: a nationwide population-based study. *PLOS ONE*, **9** (1), e87095 (2014).
7. Rabinovici, G. D. *et al.* Testing and disclosures related to amyloid imaging and Alzheimer's disease: common questions and fact sheet summary. *Alzheimers Dement*, **12** (4), 510–515 (2016).
8. Manach, C., Scalbert, A., Morand, C., Rémésy, C. & Jiménez, L. Polyphenols: food sources and bioavailability. *Am J Clin Nutr*, **79** (5), 727–747 (2004).
9. Li, Y. *et al.* Quercetin, inflammation and immunity. *Nutrients*, **8** (3), 167 (2016).
10. Henagan, T. M., Lenard, N. R., Gettys, T. W. & Stewart, L. K. Dietary quercetin supplementation in mice increases skeletal muscle PGC1 α expression, improves mitochondrial function and attenuates insulin resistance in a time-specific manner. *PLOS ONE*, **9** (2), e89365 (2014).
11. Vessal, M., Hemmati, M. & Vasei, M. Antidiabetic effects of quercetin in streptozocin-induced diabetic rats. *Comp Biochem Physiol C Toxicol Pharmacol*, **135C** (3), 357–364 (2003).
12. Atri, A. Current and future treatments in Alzheimer's disease. *Semin Neurol*, **39** (2), 227–240 (2019).
13. Thompson, T. L. *et al.* Lack of efficacy of Hydergine in patients with Alzheimer's disease. *N Engl J Med*, **323** (7), 445–448 (1990).
14. Denny, J. K. & Muralidhara Enhanced neuroprotective effect of fish oil in combination with quercetin against 3-nitropropionic acid induced oxidative stress in rat brain. *Prog Neuropsychopharmacol Biol Psychiatry*, **40**, 83–92 (2013).
15. Xu, S. L. *et al.* Flavonoids induce the expression of synaptic proteins, synaptotagmin, and postsynaptic density protein-95 in cultured rat cortical neuron. *Planta Med*, **79** (18), 1710–1714 (2013).
16. Yao, Y., Han, D. D., Zhang, T. & Yang, Z. Quercetin improves cognitive deficits in rats with chronic cerebral ischemia and inhibits voltage-dependent sodium channels in hippocampal CA1 pyramidal neurons. *Phytother Res*, **24** (1), 136–140 (2010).

17. Rattanajarasroj, S. & Unchern, S. Comparable attenuation of Abeta(25–35)-induced neurotoxicity by quercitrin and 17beta-estradiol in cultured rat hippocampal neurons. *Neurochem Res*, **35** (8), 1196–1205 (2010).
18. Kanter, M., Unsal, C., Aktas, C. & Erboga, M. Neuroprotective effect of quercetin against oxidative damage and neuronal apoptosis caused by cadmium in hippocampus. *Toxicol Ind Health*, **32** (3), 541–550 (2016).
19. Daina, A., Michielin, O. & Zoete, V. SwissTargetPrediction: updated data and new features for efficient prediction of protein targets of small molecules. *Nucleic Acids Res*, **47** (W1), W357–W364 (2019).
20. UniProt Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res*, **47** (D1), D506–D515 (2019).
21. Amberger, J. S., Bocchini, C. A., Schiettecatte, F., Scott, A. F. & Hamosh, A. OMIM.org: Online Mendelian Inheritance in Man (OMIM®), an online catalog of human genes and genetic disorders. *Nucleic Acids Res*, **43** (Database issue), D789–D798 (2015).
22. Wishart, D. S. *et al.* DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res*, **46** (D1), D1074–D1082 (2018).
23. Stelzer, G. *et al.* The GeneCards suite: from gene data mining to disease genome sequence analyses. *Curr Protoc Bioinformatics*. 2016;54:1.30.1–1.30.33.
24. Smoot, M. E., Ono, K., Ruscheinski, J., Wang, P. L. & Ideker, T. Cytoscape 2.8: new features for data integration and network visualization., **27** (3), 431–432 (2011).
25. Heberle, H., Meirelles, G. V., da Silva, F. R., Telles, G. P. & Minghim, R. InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics*, **16** (1), 169 (2015).
26. Szklarczyk, D. *et al.* STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res*, **47** (D1), D607–D613 (2019).
27. Chin, C. H. *et al.* cytoHubba: identifying hub objects and sub-networks from complex interactome. *BMC Syst Biol*. 2014;8(suppl 4) [suppl:S11]:S11.
28. Bader, G. D. & Hogue, C. W. An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics*, **4**, 2 (2003).
29. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun*, **10** (1), 1523 (2019).
30. Trott, O. & Olson, A. J. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem*, **31** (2), 455–461 (2010).
31. Kahn, S. E., Cooper, M. E. & Del Prato, S. Pathophysiology and treatment of type 2 diabetes: perspectives on the past, present, and future., **383** (9922), 1068–1083 (2014).
32. Saisho, Y. Importance of beta cell function for the treatment of Type 2 diabetes. *J Clin Med*, **3** (3), 923–943 (2014).

33. White, S. A., Kimber, R., Veitch, P. S. & Nicholson, M. L. Surgical treatment of diabetes mellitus by islet cell and pancreas transplantation. *Postgrad Med J*, **77** (908), 383–387 (2001).
34. Soria, L. J., González, H. M. & Léger, G. C. Alzheimer's disease. *Handb Clin Neurol*, **167**, 231–255 (2019).
35. Ramirez, M. J., Lai, M. K., Tordera, R. M. & Francis, P. T. Serotonergic therapies for cognitive symptoms in Alzheimer's disease: rationale and current status. *Drugs*, **74** (7), 729–736 (2014).
36. Tixi-Verdugo, W., Contreras-Ramos, J., Sicilia-Argumedo, G., German, M. S. & Fernandez-Mejia, C. Effects of biotin supplementation during the first week postweaning increases pancreatic islet area, beta-cell proportion, islets number, and beta-cell proliferation. *J Med Food*, **21** (3), 274–281 (2018).
37. Bischoff, S. C. Quercetin: potentials in the prevention and therapy of disease. *Curr Opin Clin Nutr Metab Care*, **11** (6), 733–740 (2008).
38. Forman, H. J. *et al.* The chemistry of cell signaling by reactive oxygen and nitrogen species and 4-hydroxynonenal. *Arch Biochem Biophys*, **477** (2), 183–195 (2008).
39. Wu, L. *et al.* Exendin-4 protects HUVECs from tunicamycin-induced apoptosis via inhibiting the IRE1a/JNK/caspase-3 pathway., **55** (3), 764–772 (2017).
40. Liao, Y. *et al.* Cordycepin induces cell cycle arrest and apoptosis by inducing DNA damage and up-regulation of p53 in leukemia cells., **14** (5), 761–771 (2015).
41. Giraldo, E., Lloret, A., Fuchsberger, T. & Viña, J. A β and tau toxicities in Alzheimer's are linked via oxidative stress-induced p38 activation: protective role of vitamin E. *Redox Biol*, **2**, 873–877 (2014).
42. von Holstein-Rathlou, S. *et al.* FGF21 mediates endocrine control of simple sugar intake and sweet taste preference by the liver. *Cell Metab*, **23** (2), 335–343 (2016).
43. Dai, H. L. *et al.* p38 MAPK Inhibition Improves Synaptic Plasticity and Memory in Angiotensin II-Dependent Hypertensive Mice. *Sci. Rep*, **6**, 27600 (2016).
44. Salminen, A., Ojala, J., Kauppinen, A., Kaarniranta, K. & Suuronen, T. Inflammation in Alzheimer's disease: amyloid-beta oligomers trigger innate immunity defence via pattern recognition receptors. *Prog Neurobiol*, **87** (3), 181–194 (2009).
45. Samuels, I. S. *et al.* Deletion of ERK2 mitogen-activated protein kinase identifies its key roles in cortical neurogenesis and cognitive function. *J Neurosci*, **28** (27), 6983–6995 (2008).

Figures




Figure 1

Flowchart of the network pharmacology study investigating quercetin targets for T2DM and AD treatment.




Figure 2

Venn diagram showing the overlap among quercetin targets (blue), T2DM-related gene products (red), and AD-related gene productions (green). A total of 95 gene products were identified as common.




Figure 3

PPI network





Figure 4

GO functional analysis of quercetin targets relevant to T2DM and AD treatment.



A KEGG pathway dot bubble plot

B KEGG pathway classification plot

Figure 5

KEGG pathway enrichment analysis of quercetin targets relevant to T2DM and AD treatment. (A) KEGG pathway dot bubble plot. (B) KEGG pathway classification plot.




Figure 6

Component-target-pathway network for quercetin targets relevant to T2DM and AD treatment.




Figure 7

Molecular docking diagrams of quercetin binding to (from top to bottom) MAPK1, AKT1, VEGFA, EGFR, JUN, and TNF.




Figure 8

Components of MAPK signaling pathway influenced by quercetin.