Microscopic and submicroscopic Plasmodium infections in indigenous and non-indigenous communities in Colombia.

Jehidys Montiel

Universidad de Antioquia

Lina M. Zuluaga

Universidad de Antioquia

Daniel C. Aguirre

Universidad de Antioquia

Cesar Segura

Universidad de Antioquia

Alberto Tobon-Castaño

Universidad de Antioquia

Ana M Vasquez (✉ amaria.vasquez@udea.edu.co)

Universidad de Antioquia <https://orcid.org/0000-0001-9418-3298>

Research

Keywords: malaria, indigenous communities, asymptomatic infections, submicroscopic infections.

Posted Date: March 25th, 2020

DOI: <https://doi.org/10.21203/rs.2.17952/v3>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Malaria Journal on April 16th, 2020. See the published version at <https://doi.org/10.1186/s12936-020-03226-4>.

Abstract

Background: The indigenous population is considered a highly susceptible group to malaria because they usually live in areas with high exposure to *Anopheles* and poverty, and have limited access to health services. There is a great diversity of indigenous communities in Colombia living in malaria-endemic areas; however, the burden of infection in these populations has not been studied extensively. This study aimed to determine the prevalence of *Plasmodium* infections in indigenous and non-indigenous communities in two malaria-endemic areas in Colombia. **Methods:** A Community-based cross-sectional survey was conducted in seven villages of Turbo and El Bagre municipalities; three of these villages were indigenous communities. Inhabitants of all ages willing to participate were included. Sociodemographic and clinical data were recorded as well as household information. The parasitological diagnosis was performed by microscopy and nested PCR. The prevalence of microscopy and submicroscopic infection was estimated. An adjusted GEE model was used to explore risk factors associated with the infection. **Results:** Among 713 participants, 60.7% were from indigenous communities. *Plasmodium* spp. was detected in 30 subjects (4.2%, CI 95% 2.9-5.9); from those, 29 were in the indigenous population, 47% of infections were afebrile, and most of them submicroscopic (10/14). Microscopic and submicroscopic prevalence was 2.5% (CI 95% 1.6-3.9) and 1.7% (CI 95% 0.9-2.9) respectively. In El Bagre, all infections occurred in indigenous participants (3.9%, CI 95% 2.2-7.1), and 81% were submicroscopic. By contrast, in Turbo, the highest prevalence occurred in indigenous people (11.5%; CI 95%: 7.3-17.5), but 88.8% were microscopic. Living in an indigenous population increased the prevalence of infection compared with a non-indigenous population (PR 19.4; CI 95% 2.3-166.7). **Conclusion:** There is a high proportion of *Plasmodium* infection in indigenous communities. A substantial proportion of asymptomatic and submicroscopic carriers were detected. The identification of these infections, not only in indigenous but also in the non-indigenous population, as well as their associated factors, could help to implement specific malaria strategies for each context.

Background

Although significant advances have been made towards malaria elimination in several endemic countries, malaria remains a significant public health problem [1]; the World Malaria Report 2018 estimated that the global malaria burden was around 219 million reported cases and 43,5000 deaths worldwide [2]. Besides, the situation in the Americas comes up with further challenges for control and malaria elimination, given the high proportion of cases by *Plasmodium vivax* [3]. Particularly in Colombia, the number of malaria cases officially reported in 2018 was 63,143; with Chocó, Nariño, Córdoba, and Antioquia, the departments with the highest number of malaria cases (27.0%, 20.6%, 15.6%, and 8.8% respectively) [4].

In the Americas, the indigenous population is considered one of the most vulnerable groups to suffer from malaria. The elevated vulnerability is not solely explained by the fact that they live in areas with a high *Anopheles* bite exposure, but also because they have high poverty rates and little to no access to diagnostic and treatment services [5]. Information about health conditions of these populations is not always collected, so their risk is not well understood, but in general, it is known that indigenous communities have poor health indicators as compared to non-indigenous populations, including the morbidity and mortality due to transmissible diseases, child undernutrition, infant mortality rates, and years of potential life lost [6].

In Colombia, 3.4% of the population is indigenous, and there are around 710 indigenous communities located in 27 departments [7], many of them living in malaria-endemic regions. Between 2009 and 2014, 75.8% of the Colombian indigenous population was at risk of being infected by any microorganism, where *Plasmodium* spp caused 46.7% of the total infections [8]. Unfortunately, there are few studies in the indigenous population, so that the risk is not well understood. Only eight of the 21 malaria-endemic countries of the Americas Region reported cases of ethnic groups and indigenous peoples in 2014 [5]. Without adequate data, it is difficult to follow up on malaria trends, recognize the risk factors in these communities, and establish malaria control strategies. In Colombia, the majority of malaria cases occur at the pacific coast and amazon region and affects mainly Afro-Colombian and indigenous populations [1, 9]. In 2018, 62,141 malaria cases were reported, of these, 14,714 (23.7%) were in the indigenous population [10]. Autochthonous malaria transmission has also been reported mainly among indigenous communities in Chocó (Pacific coast) [11].

Antioquia was one of the departments with the highest malaria prevalence in Colombia for many years. However, the cases have decreased markedly from 20,511 in 2008 to 4,971 in 2017 [12]. This could be explained by the several malaria control strategies implemented between 2007 and 2010, such as vector control activities, strengthening of diagnosis network, distribution of insecticide-treated bed nets (ITNs), and chemoprophylaxis. Despite this reduction, a significant proportion of malaria cases is related to gold-mining activities, which play an important role in the maintenance of malaria transmission and are considered to be an important barrier to malaria elimination [13]. Like miners, indigenous populations are also considered a significant reservoir of malaria transmission.

Unfortunately, these populations have been scarcely studied, and there is not enough information about the *Plasmodium* prevalence in them.

One of the main challenges of malaria control programs is the early diagnosis and treatment, not only for symptomatic but also for asymptomatic infections, which represent a silent reservoir of parasites [14]. Compared to patients with acute malaria disease, who generally seek treatment in health facilities, people with low-density infections that often are asymptomatic, do not seek medical attention or anti-malarial treatment [15, 16]. These infections can contribute to local transmission in an endemic region [17]. It has been reported that in Peru, 50% of *P. falciparum* and 22% of *P. vivax* asymptomatic infections can harbor gametocytes [18]; similarly, in Colombia, 57% of the samples from asymptomatic volunteers were infective to mosquitoes [19].

In Colombia, the prevalence of low-density infections by *Plasmodium* has been previously explored, finding frequencies from 2% to 15% in general population, with most of the infections being submicroscopic [20-22]. In pregnant women frequencies reached from 1.1% in peripheral to a 2.1% in placental blood [23]. In the Urabá region located in Antioquia-Colombia, the prevalence of asymptomatic infections detected by PCR was 2.6% [24]. Together these studies suggest that in low endemic settings such as Colombia, molecular tests are more useful than microscopy to detect this kind of infection [25-27].

Most of the studies about low-density infections in Colombia have been conducted in the general population and pregnant women, but there are not reports in special populations such as indigenous people. We aimed to determine the prevalence of microscopic and submicroscopic *Plasmodium* infections in indigenous and non-indigenous communities in two malaria-endemic areas in Antioquia-Colombia and to explore the associated factors to the *Plasmodium* infections.

Methods

Study design

A cross-sectional study was conducted in seven malaria-endemic villages of Turbo and El Bagre municipalities between November 2016 and June 2017. Villages were selected based on the historical records of malaria cases, and the accessibility for field staff. Three of these villages were indigenous communities (Los Aguacates and Los Almendros in El Bagre and Arqua in Turbo).

Study sites description

The two towns selected for this study are located in the Antioquia department (figure 1). Turbo (8° 05' 42" N, 76° 44' 123") is located in the Urabá region of Colombia, has an area of 3055 m², an altitude of 2 meters above sea level and an average temperature of 28°C. The main economic activity is banana production [28, 29]. El Bagre (7° 35'25" N, 74° 48'27"), is located in the Bajo Cauca region of Colombia, has an area of 1563 m², an average temperature of 32°C and an altitude of 50 meters above sea level [29, 30]. The main economic activity is gold mining. The annual parasite index (parasite incidence per 1,000 population) reported for 2017 was 0.77 in Turbo and 21.29 in El Bagre, in both, *P. vivax* was the predominant species [12].

Study population and sample collection

A census was carried out in each village to know the number of houses and people; all individuals were invited to participate in the study. Individual and household data regarding sociodemographic, self-report of previous malaria episodes, and characteristics of the household were collected. Axillary temperature was measured, and 6mL of whole blood was collected by venipuncture in tubes with anticoagulant EDTA (BD Vacutainer, BD Franklin Lakes, USA). Samples were stored at 4°C until processing at the laboratory of the local hospital in each municipality. The sample was used to perform diagnosis by microscopy using thick and thin blood smears and to measure hemoglobin levels using by a HemoCue (Hb 201+, Lake Forest, California). A blood sample was dropped on to Whatman filter paper #3 (Fisher, Ref 1003-917) and later used for molecular diagnosis. Mild anemia was defined as a hemoglobin level between 10 g/dL and 10.9 g/dL, moderate anemia between 7 g/dL and 9.9 g/dL and severe anemia below 7 g/dL following the WHO criteria [2].

Laboratory procedures

Conventional microscopy

Field-stained thick and thin blood smears were read in the field by an expert microscopist according to national guidelines [31] with a limit of detection (LOD) around 50 parasites/uL [32]. Parasitaemia was estimated against 200 leukocytes (assuming a standard value

of 8,000 leukocytes/ μ L of blood) and was expressed as parasites/ μ L (p/ μ L). A sample was considered negative if, after the examination of 200 microscopic fields at 100x magnification, no parasites were observed. As a quality control, a second reading was performed in all PCR and/or microscopy positives samples and 10% of negative ones. Discordant results were solved by a third reading.

All participants with positive thick smear received anti-malarial treatment according to national guidelines; *P. vivax* cases were treated with chloroquine for 3 days and primaquine (PQ) for 14 days, *P. falciparum* cases were treated with artemether plus lumefantrine for 3 days [31].

Molecular diagnostic

DNA was extracted from half a blood spot of the filter cards (approximately 30 μ L of blood) using the QIAamp DNA Mini Kit following the manufacturer's recommendations (QIAGEN, Germany). Nested PCR was performed as a two-step procedure following the protocol described by Singh et al. with a LOD of 1-6 parasites/uL [33]. Amplification products were resolved in a 1.5% agarose gel stained with GelRed™ (Biotium, United States) and visualized under UV light. This protocol consists of a universal PCR that detects and amplifies a region of the 18S ribosomal gene from the *Plasmodium* genus. Positive samples for *Plasmodium* spp. were further processed using species-specific primers for *P. falciparum* and *P. vivax*.

Ethical considerations

This study was reviewed and approved by the Ethics Committee of the Medicine school, Universidad de Antioquia, Colombia (Record 011 dated 28 July 2016). Before to start the fieldwork, permission from community leaders was approved in each village. Signed informed consent was obtained before the interview and blood sampling of all participants. In the case of individuals <18 years old, an additional informed consent from parents or legal guardians was also obtained.

Statistical analysis

All data from questionnaires and forms were entered into a Microsoft Access database, and statistical analyses were conducted in STATA 14 (StataCorp. 2015. Stata Statistical Software: Release 14. College Station, TX: StataCorp LP). A description of the population was carried out for both, individual (anemia, sex, age group, occupation, residence time in endemic region, number of malaria episodes, malaria in the previous year, municipality, use of mosquito nets) and housing factors (inhabitants per household, draining of stagnant water, access to electricity, water and sewer system, mosquito mesh on windows, roof and wall materials). Frequencies were expressed as numbers and percentages for qualitative variables and medians with the interquartile range for the quantitative ones. The prevalence of *Plasmodium* infection was estimated by diagnostic tests (microscopy and nPCR), locality (Turbo and El Bagre), and type of community (indigenous and non-indigenous communities). 95% confidence intervals (CI) are shown for each estimation. A generalized estimating equation (GEE) model [34] was used to handle the nested structure of sampled data (713 individuals nested within 212 households), GEE model was fitted for a Poisson family with a logarithmic link and an exchangeable correlation. Crude (PR) and adjusted (aPR) Prevalence ratio (by occupation and malaria in the previous year) were calculated for the *Plasmodium* infection (outcome) with 95% confidence intervals for each factor.

Results

Socio-demographic and household characteristics

In this study, 713 subjects from 212 houses were enrolled in seven villages of two municipalities in the Antioquia department. El Bagre town had more indigenous people (n=276; 38.7%) than Turbo (n=157; 22.0%). The proportion of *Plasmodium* infection in both municipalities was higher in indigenous communities than non-indigenous communities (6.69% vs. 0.36%) (**Figure 2**).

The characteristics of participants, according to the residence in the indigenous communities, are shown in **Table 1**. There was a slightly higher proportion of anemia in non-indigenous people (15.4%) compared to indigenous (12.7%). Additionally, the proportion of moderate anemia was higher in indigenous communities than non-indigenous communities (11.6% and 32.7%, respectively). The same pattern was observed in severe anemia (2.4% and 3.6% respectively).

The gender and age distribution was similar in both groups, and the majority of people have lived in an endemic malaria region for more than four years (72.5% and 75% in non-indigenous and indigenous communities respectively). Most of the indigenous people lived in

houses with more than five people (59.1%) and had animals in their homes (97%). Additionally, a high proportion of them did not drain standing water in their homes (93%), and 34% did not have access to electricity.

***Plasmodium* infections prevalence**

The overall prevalence of *Plasmodium* spp infection was 4.21% (95% CI 2.95%-5.96%), 60% of these infections were detected by microscopy, and *P. vivax* was the predominant species (3.09%; 95% CI 2.04%-4.65%). The prevalence of infection was higher in Turbo compared to El Bagre (5.67%, 95% CI 3.64%-8.73% versus 2.91%, 95% CI 1.62%-5.19% respectively); most of the infections in Turbo were detected by microscopy (16/19, 84.2%) and 93.75% (15/16) had malaria symptoms (**Table 2**), while most of the infections in El Bagre were asymptomatic (temperature <37.5°C) and submicroscopic (9/11). In both towns, a considerably higher prevalence of *Plasmodium* was found in indigenous communities (**Figure 3**), resulting in a 12-fold increase in the prevalence of malaria as compared to non-indigenous communities.

Association between *Plasmodium* infections and risk factors

The analysis of the association of the subjects and household characteristics with *Plasmodium* infections is shown in **Table 3**. The prevalence of *Plasmodium* infections was higher in males (5.7%) than females (2.9%), with an aPR for the male sex of 2.38 (95% CI: 1.07-5.31). People who have lived in the endemic region for more than four years had a higher prevalence of infection than those living in the endemic region for less time (4.2% and 2.3%, respectively). The infection was also higher in people with previous self-reported malaria as compared to people with no malaria history (4.8% and 3.2%, respectively). Furthermore, living in an indigenous community increased the prevalence of infection as compared to the non-indigenous population (aPR 17.86, 95% CI: 2.12-150.19). Regarding household factors, it was found that participants with no electricity services in their households have a higher prevalence of *Plasmodium* infection (aPR 4.62, 95% CI: 1.89-11.32). The PR could not be calculated for the association with water and sewage services (the convergence was not achieved in the GEE model); because none of the infected people had access to these services.

Living in an indigenous community as well as some characteristics of the houses were associated with a higher incidence rate of *Plasmodium* infections. A household description of indigenous and non-indigenous communities is detailed in **Table 4**. A total of 125 from 212 houses belong to indigenous communities (59%), 30.4% of these did not have electricity service, and 91.2% did not drain the standing water; these proportions were substantially higher in indigenous compared to non-indigenous communities.

Discussion

This study evaluated the prevalence of microscopic and submicroscopic *Plasmodium* infections in indigenous and non-indigenous communities from Antioquia, Colombia, and its associated factors, to describe the distribution of disease prevalence among heterogeneous populations; this knowledge is necessary to implement proper control strategies for each context [35]. We found that the prevalence of *Plasmodium* infections it was 12-fold increased in indigenous communities as compared to non-indigenous communities in both municipalities. Even more, all infections in El Bagre were detected in indigenous communities (11/11), and most of them were asymptomatic and submicroscopic (9/11). On the contrary, most of the infections in indigenous communities in Turbo were symptomatic and microscopic (84.2%).

It is known that malaria transmission in Colombia varies among the endemic regions [35]; in this way, our findings could be explained by differences in malaria profiles in each municipality. Although the general prevalence of malaria in Antioquia has decreased in recent years, the number of cases in El Bagre has been higher than in Turbo (from 190.45 cases/1000 people in 2007 to 21.29 cases/1000 people in 2017 in El Bagre and from 61.53 cases/1000 in 2007 to 0.77cases/1000 people in 2017 in Turbo).

Most of the malaria cases were caused by *P. vivax* and it is well known that the PQ used for the *P. vivax* treatment can induce haemolytic crises in individuals with Glucose 6-phosphate dehydrogenase (G6PD) deficiency. The G6PD deficiency is distributed worldwide, however, its frequency varies among regions and ethnic groups. A previous report in malaria-endemic areas of Colombia located on the Pacific coast, found a frequency of G6PD deficiency of 6.56% [36], studies in indigenous populations (Amerindian) are lacking. Considering the high number of *P. vivax* infections in these populations found in this study, there is a need for further evaluation of the frequency of G6PD deficiency in malaria-endemic areas in view that the primaquine treatment (14 days) is required for the radical cure.

As previously reported, malaria immunity is determined by previous *Plasmodium* exposure, where an anti-disease immunity is first achieved, resulting in a reduction of severe malaria and mortality. Then, an anti-parasitic immunity is slowly acquired and confers

protection against high parasitic densities, which in turn protect against the severe disease [37]; this could explain the highest prevalence of submicroscopic infections in El Bagre, where 50.3% of individuals had had more than one malaria episode over life compared to 35.8% in Turbo (**Supplementary Table 1**). Nevertheless, we could not find an association of this variable with the *Plasmodium* infections using a GEE analysis. However, a previous study in Nariño- Colombia showed that having suffered from more than one malaria episode was associated with an increased risk of having asymptomatic infections (aOR 2.4, 95% CI 1.1–5.4) [22]. These differences could be explained because our model included not only asymptomatic but also symptomatic infections.

Household factors are also associated with malaria risk [38]. We found that having no access to electricity was associated with an increase in the malaria rate. These findings are in agreement with previous studies that reported that the poorest households had a 29% greater risk of microscopic parasitemia compared to the poorest houses (aRR 1.29; 95% CI 1.07–1.55) [39]. Additionally, lack of household electricity increased the childhood mortality in Rwanda, including malaria mortality (aOR: 1.4, 95% CI: 1.0–1.8) [40]. The above is important because housing quality can affect malaria risk through its effect on house entry of the malaria vector [39].

Taken together, the individual and housing characteristics could help to understand why the indigenous population has a higher prevalence than its counterpart, the non-indigenous population does. Ethnicity is an important determinant of health conditions, influencing the morbidity and mortality rates in different ethnic groups and interfering with access to health services for some populations [41]. In Colombia, the exclusion of indigenous people is reflected in poverty rates, lack of land and employment, school desertion, unsatisfied basic needs, a higher prevalence of transmissible diseases and limited access to health services compared to the general Colombian population [42]. Regarding this last point, we noticed that the indigenous villages were farther from the health services (1 to 2 hours by motorcycle) compared to the non-indigenous villages, and the road conditions were far worse. Furthermore, the indigenous population frequently lives close to rainforests or wetlands where they have more vector exposure, resulting in an increased risk of getting sick with vector-borne diseases such as malaria [8].

It is possible to suggest that the diversity of epidemiologic characteristics of malarial infection among the Colombian subpopulations account for an ideal environment for parasite evolution. In this environment, the parasite can interact with susceptible populations from different ethnicities and under different public health interventions [35]. We suggest that prevention efforts should be population-specific and vary according to the individual, housing, and environmental characteristics. Given the heterogeneity of the prevalence of malaria in Colombia, it has become necessary to adjust malaria control activities according to each population and context.

Further studies are needed to evaluate the potential integration of molecular tests into the surveillance programs to promptly detect malaria infection in the community in order to contribute to the control and future elimination strategies. However, due the low prevalence of infection in this region of Colombia, there is also a need to evaluate the costs per assay comparing to conventional test, including equipment, reagents, staff, training, and maintenance in order to evaluate the cost-effectiveness of molecular test for their potential integration into surveillance strategies and explore alternatives as serological surveillance. A previous study in a low transmission setting in Indonesia, suggest that reactive-active detection of cases in the community using molecular test had high costs per individual screened, however, compared to microscopy, molecular test was most cost-effective for the detection of infections [43]. Also, another study in a low transmission setting in Africa concludes that standards test as rapid diagnosis test (RDTs) are not useful to detect infections in the community and suggest that the achievement of malaria elimination may require active case detection using more sensitive point-of-care diagnostics, especially in high-risk groups [44], that in our cases are the indigenous communities among others.

This study has some limitations. First, due to cross-sectional design, the association with malaria status should be interpreted with caution, as they do not imply causality. Second, it was no possible to analyze the risk factors for asymptomatic infections exclusively due to the low number of this type of infection. This last could, in turn, affect the accuracy of confidence intervals for some of the factors analyzed due to the sample size. Third, as we mentioned before, the villages in this study were selected based on the historical records of malaria cases, the distance to the urban area, and the accessibility for field staff, and we did not make a random selection of the villages included in the study. The results of this study cannot be extrapolated to the general population; nevertheless, they are useful to exhibit the problems around the asymptomatic infections in the indigenous and non-indigenous people. Fourth, considering that nowadays there are ultra-sensitive molecular tests for the detection of low-density infections, the prevalence in this study could be underestimated due to the limit of detection of the nPCR used, nevertheless, the nPCR used in this study was able to detect 1.6 times more infections than microscopy showing the presence of a significant *Plasmodium* reservoir in the region. At last, it is possible that other variables which we did not consider in the GEE model could explain the associated factors to *Plasmodium* infections. Future studies are required to improve the knowledge that we have about the risk factors of the *Plasmodium* infections in indigenous

communities. Despite these limitations, these results are useful to understand malaria transmission in studied places and to suggest prevention efforts according to the individual, housing, and environmental characteristics.

Conclusion

This study reveals that in both municipalities, most of the *Plasmodium* infections were in indigenous communities. Nevertheless, the infection profile was different for each town. A substantial proportion of asymptomatic and submicroscopic carriers were detected in El Bagre, while most of the symptomatic and microscopic infections were identified in Turbo. Our findings provide an understanding of the key characteristics of asymptomatic, submicroscopic, and microscopic infections in the study population: to live in an indigenous community, having had previous malaria episodes and not having access to electricity, sewage system, and water services. The current malaria control efforts could benefit from the implementation of targeted interventions in indigenous villages using molecular tests to identify submicroscopic reservoirs that could be contributing to malaria transmission, although further studies are needed to evaluate the potential integration of molecular test into the surveillance programs to promptly detect malaria infection in the community, especially in high-risk groups. Additionally, the identification of these infections not only in indigenous but also in the non-indigenous populations, as well as demographic, social and household factors related to them, could help to implement specific malaria strategies for each context.

List Of Abbreviations

aPR: adjusted Prevalence ratio; aOR: adjusted odds ratio; CI: confidence interval; DNA: deoxyribonucleic acid; GEE: generalized estimating equation; G6PD: Glucose 6-phosphate dehydrogenase; PR: Prevalence ratio; IQR: inter quartile range; ITNs: insecticide-treated bed nets; LOD: limit of detection; p/μL: parasite/μL *P. falciparum*: *Plasmodium falciparum*; *Plasmodium spp.*: *Plasmodium* species; *P. vivax*: *Plasmodium vivax*; PCR: polymerase chain reaction; nPCR: nested polymerase chain reaction; PQ: Primaquine; *spp.*: species; UV: ultraviolet.

Declarations

Ethical approval and consent to participate

The study was reviewed and approved by the Ethics Committee of the Medical School at the Universidad de Antioquia, Medellín, Colombia (Record 011 dated 28 July 2016). Before starting any study procedure, written informed consent or an informed assent in the case of participants under 18 years old were obtained from each participant. For participants <18 years old additional informed consent from her parents or legal guardian was also obtained.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study has been conducted with funding from Universidad de Antioquia (Code 2015-7765) and COLCIENCIAS (RC-075-2016, Code 111571249820).

Authors' contributions

JM, LZ, AT, CS, AV conceived the design of this study. JM and LZ supervised data collection and work field. JM, LZ and DA analysed the data. JM drafted the manuscript. All authors contributed to the manuscript edit, review and revising, and approved the final version of the manuscript.

Acknowledgements

The authors thank Hospital Francisco Valderrama in Turbo and Hospital Nuestra Señora del Carmen in El Bagre for their support in this research. A special thanks to the people from the communities for allowing and participating in this study, and to Deiber Toro, Rafael Mercado, Luisa Carbal Reyes y Veronica Sierra for their commitment in the fieldwork and to Juan Camilo Pérez and Gabriel Velez for the molecular diagnostic.

References

1. Recht J, Siqueira AM, Monteiro WM, Herrera SM, Herrera S, Lacerda MVG: **Malaria in Brazil, Colombia, Peru and Venezuela: current challenges in malaria control and elimination.** *Malar J* 2017, **16**:273.
2. **World Health Organization. (2018). World malaria report 2018.** World Health Organization. <http://www.who.int/iris/handle/10665/275867>. Licencia: CC BY-NC-SA 3.0 IGO.
3. Lindblade KA, Steinhart L, Samuels A, Kachur SP, Slutsker L: **The silent threat: asymptomatic parasitemia and malaria transmission.** *Expert Rev Anti Infect Ther* 2013, **11**:623-639.
4. **Instituto Nacional de Salud. (2018). Informe de Evento Malaria, Colombia, 2018.** Instituto Nacional de Salud [https://www.ins.gov.co/buscador-eventos/Informesdeevento/MALARIA_2018.pdf]
5. **Informe de la situación de la Malaria en las Américas, 2014.** Washington, D.C. : PAHO, 2017.
6. Coimbra CE, Santos RV, Welch JR, Cardoso AM, de Souza MC, Garnelo L, Rassi E, Follér ML, Horta BL: **The First National Survey of Indigenous People's Health and Nutrition in Brazil: rationale, methodology, and overview of results.** *BMC Public Health* 2013, **13**:52.
7. Monje-Carvajal J: **Colombian indigenous peoples' life plan, a structure of ethno-eco development.** *Luna Azul* 2015, **41**:29-56.
8. **Perfil de Salud de la Población Indígena, y medición de desigualdades en salud.** Colombia 2016. Colombia, Ministerio de Salud y Protección Social, 2016.
9. Coura JR, Suárez-Mutis M, Ladeia-Andrade S: **A new challenge for malaria control in Brazil: asymptomatic Plasmodium infection—a review.** *Mem Inst Oswaldo Cruz* 2006, **101**:229-237.
10. **Comportamiento de la notificación malaria 2018.** Boletín Epidemiológico semanal. Bogotá: Instituto Nacional de Salud; 2019.
11. Molina Gómez K, Caicedo MA, Gaitán A, Herrera-Varela M, Arce MI, Vallejo AF, Padilla J, Chaparro P, Pacheco MA, Escalante AA, et al: **Characterizing the malaria rural-to-urban transmission interface: The importance of reactive case detection.** *PLoS Negl Trop Dis* 2017, **11**:e0005780.
12. **Dirección Seccional de Salud de Antioquia. Eventos de interés en salud pública por subregiones y municipios.** Antioquia 2007-2017 [<http://www.dssa.gov.co/index.php/estadisticas/eventos-en-salud-publica/item/71-eventos-de-interes-en-salud-publica-por-subregiones-y-municipios-antioquia-2007-2012>]
13. Castellanos A, Chaparro-Narváez P, Morales-Plaza CD, Alzate A, Padilla J, Arévalo M, Herrera S: **Malaria in gold-mining areas in Colombia.** *Mem Inst Oswaldo Cruz* 2016, **111**:59-66.
14. Cohen JM, Moonen B, Snow RW, Smith DL: **How absolute is zero? An evaluation of historical and current definitions of malaria elimination.** *Malar J* 2010, **9**:213.
15. Okell LC, Ghani AC, Lyons E, Drakeley CJ: **Submicroscopic infection in Plasmodium falciparum-endemic populations: a systematic review and meta-analysis.** *J Infect Dis* 2009, **200**:1509-1517.
16. Bousema T, Okell L, Felger I, Drakeley C: **Asymptomatic malaria infections: detectability, transmissibility and public health relevance.** *Nat Rev Microbiol* 2014, **12**:833-840.
17. Carrasco-Escobar G, Miranda-Alban J, Fernandez-Miño C, Brouwer KC, Torres K, Calderon M, Gamboa D, Llanos-Cuentas A, Vinetz JM: **High prevalence of very-low Plasmodium falciparum and Plasmodium vivax parasitaemia carriers in the Peruvian Amazon: insights into local and occupational mobility-related transmission.** *Malar J* 2017, **16**:415.
18. Branch O, Casapia WM, Gamboa DV, Hernandez JN, Alava FF, Roncal N, Alvarez E, Perez EJ, Gotuzzo E: **Clustered local transmission and asymptomatic Plasmodium falciparum and Plasmodium vivax malaria infections in a recently emerged, hypoendemic Peruvian Amazon community.** *Malar J* 2005, **4**:27.
19. Vallejo AF, García J, Amado-Garavito AB, Arévalo-Herrera M, Herrera S: **Plasmodium vivax gametocyte infectivity in sub-microscopic infections.** *Malar J* 2016, **15**:48.

20. Vallejo AF, Chaparro PE, Benavides Y, Álvarez Á, Quintero JP, Padilla J, Arévalo-Herrera M, Herrera S: **High prevalence of sub-microscopic infections in Colombia.** *Malar J* 2015, **14**:201.
21. Vásquez-Jiménez JM, Arévalo-Herrera M, Henao-Giraldo J, Molina-Gómez K, Arce-Plata M, Vallejo AF, Herrera S: **Consistent prevalence of asymptomatic infections in malaria endemic populations in Colombia over time.** *Malar J* 2016, **15**:70.
22. Cucunubá ZM, Guerra Á, Rivera JA, Nicholls RS: **Comparison of asymptomatic Plasmodium spp. infection in two malaria-endemic Colombian locations.** *Trans R Soc Trop Med Hyg* 2013, **107**:129-136.
23. Vásquez AM, Medina AC, Tobón-Castaño A, Posada M, Vélez GJ, Campillo A, González IJ, Ding X: **Performance of a highly sensitive rapid diagnostic test (HS-RDT) for detecting malaria in peripheral and placental blood samples from pregnant women in Colombia.** *PLoS One* 2018, **13**:e0201769.
24. Rodríguez Vásquez C, Barrera Escobar S, Tobón-Castaño A: **Low Frequency of Asymptomatic and Submicroscopic Plasmodial Infections in Urabá Region in Colombia.** *J Trop Med* 2018, **2018**:8506534.
25. Okell LC, Bousema T, Griffin JT, Ouedraogo AL, Ghani AC, Drakeley CJ: **Factors determining the occurrence of submicroscopic malaria infections and their relevance for control.** *Nat Commun* 2012, **3**:1237.
26. Lin JT, Saunders DL, Meshnick SR: **The role of submicroscopic parasitemia in malaria transmission: what is the evidence?** *Trends Parasitol* 2014, **30**:183-190.
27. Hofmann NE, Gruenberg M, Nate E, Ura A, Rodriguez-Rodriguez D, Salib M, Mueller I, Smith TA, Laman M, Robinson LJ, Felger I: **Assessment of ultra-sensitive malaria diagnosis versus standard molecular diagnostics for malaria elimination: an in-depth molecular community cross-sectional study.** *Lancet Infect Dis* 2018, **18**:1108-1116.
28. Alcaldía de Turbo - Antioquia. Indicadores del Municipio de Turbo. [<http://www.turbo-antioquia.gov.co/indicadores.shtml#poblacion>]
29. Carmona-Fonseca J: **La malaria en Colombia, Antioquia y las zonas de Urabá y Bajo Cauca: panorama para interpretar la falla terapéutica antimalárica. Parte 1.** *Iatreia* 2003, **16**:299-318.
30. Alcaldía de El Bagre - Antioquia. Información del Municipio [<http://www.elbagre-antioquia.gov.co/Paginas/default.aspx>]
31. Instituto Nacional de Salud (INS). Ministerio de Salud y Protección Social. Manual para el diagnóstico de malaria no complicada en puestos de diagnóstico y tratamiento. Milenio Editores, editor. Social Impresión 2015.
32. Policy brief on malaria diagnostics in low-transmission settings [<https://www.who.int/malaria/publications/atoz/policy-brief-diagnosis-low-transmission-settings/en/>]
33. Singh B, Bobogare A, Cox-Singh J, Snounou G, Abdullah MS, Rahman HA: **A genus- and species-specific nested polymerase chain reaction malaria detection assay for epidemiologic studies.** *Am J Trop Med Hyg* 1999, **60**:687-692.
34. Twisk J. Generalized estimating equations. In: Press CU, editor. Applied Longitudinal Data Analysis for Epidemiology. United States of America: Cambridge University Press; 2003.
35. Feged-Rivadeneira A, Ángel A, González-Casabianca F, Rivera C: **Malaria intensity in Colombia by regions and populations.** *PLoS One* 2018, **13**:e0203673.
36. Valencia SH, Ocampo ID, Arce-Plata MI, Recht J, Arévalo-Herrera M: **Glucose-6-phosphate dehydrogenase deficiency prevalence and genetic variants in malaria endemic areas of Colombia.** *Malar J* 2016, **15**:291.
37. Marsh K, Kinyanjui S: **Immune effector mechanisms in malaria.** *Parasite Immunol* 2006, **28**:51-60.
38. Deressa W: **Individual and household factors associated with ownership of long-lasting insecticidal nets and malaria infection in south-central Ethiopia: a case-control study.** *Malar J* 2017, **16**:402.
39. Okiring J, Olwoch P, Kakuru A, Okou J, Ochokoru H, Ochieng TA, Kajubi R, Kamya MR, Dorsey G, Tusting LS: **Household and maternal risk factors for malaria in pregnancy in a highly endemic area of Uganda: a prospective cohort study.** *Malar J* 2019, **18**:144.
40. Gupta N, Hirschhorn LR, Rwabukwisi FC, Drobac P, Sayinzoga F, Mugeni C, Nkikabahizi F, Bucyana T, Magge H, Kagabo DM, et al: **Causes of death and predictors of childhood mortality in Rwanda: a matched case-control study using verbal social autopsy.** *BMC Public Health* 2018, **18**:1378.
41. Sandes LFF, Freitas DA, de Souza MFNS, Leite KBS: **[Primary health care for South-American indigenous peoples: an integrative review of the literatureAtención primaria en salud a indígenas de América del Sur: revisión integrativa de la bibliografía].** *Rev Panam Salud Publica* 2018, **42**:e163.

42. Hernández Sarmiento JM, Dávila Osorio VL, Martínez Sánchez LM, Restrepo Serna L, Grajales Ospina DC, Toro Montoya AE, Arango Urrea V, Vargas Grisales N, Estrada Gómez M, Lopera Valle JS, et al: **Tuberculosis in indigenous communities of Antioquia, Colombia: epidemiology and beliefs.** *J Immigr Minor Health* 2013, **15**:10-16.
43. Zelman BW, Baral R, Zarlinda I, Coutrier FN, Sanders KC, Cotter C, Herdiana H, Greenhouse B, Shretta R, Gosling RD, Hsiang MS: **Costs and cost-effectiveness of malaria reactive case detection using loop-mediated isothermal amplification compared to microscopy in the low transmission setting of Aceh Province, Indonesia.** *Malar J* 2018, **17**:220.
44. McCreesh P, Mumbengegwi D, Roberts K, Tambo M, Smith J, Whittemore B, Kelly G, Moe C, Murphy M, Chisenga M, et al: **Subpatent malaria in a low transmission African setting: a cross-sectional study using rapid diagnostic testing (RDT) and loop-mediated isothermal amplification (LAMP) from Zambezi region, Namibia.** *Malar J* 2018, **17**:480.

Tables

Table 1. Clinical and demographic characteristics and malaria history in the study population.

Characteristic	Non-Indigenous community		Indigenous community		Total	
	n= 280		n= 433		n=713	
	n	%	n	%	n	%
<i>Individual characteristics</i>						
Hemoglobin <11mg/dL	43	15.4	55	12.7	98	13.7
Sex						
Female	145	51.8	234	54.0	379	53.2
Age						
<5	27	9.6	43	9.9	70	9.8
5 -15	87	31.1	133	30.7	220	30.9
> 15	166	59.3	255	58.9	421	59.0
Occupation						
Outdoor	152	54.3	207	47.8	359	50.4
Residence time in endemic region						
≥5 years	210	75.0	314	72.5	524	73.5
Self-report of number of symptomatic malaria episodes						
0	106	37.9	173	40.0	279	39.1
1	58	20.7	66	15.2	124	17.4
>1	116	41.4	194	44.8	310	43.5
Self-report of malaria last year						
	12	4.3	51	11.8	63	8.8
Municipality						
Turbo	102	36.4	157	36.3	259	36.3
El Bagre	178	63.6	276	63.7	454	63.7
Use of bed net						
	262	93.6	405	93.5	667	93.5
<i>Household characteristics</i>						
Number of inhabitants per household						
1 - 4	127	45.4	162	37.4	289	40.5
>5	146	52.1	256	59.1	402	56.4
Participants who have animals in their households						
	259	92.5	420	97.0	679	95.2
Participants who drain standing water in their households						
	78	27.9	20	4.6	98	13.7
Participants who have no access to electricity in their households						
	23	8.2	149	34.4	172	24.1
Participants who have no access to water in their households						
	266	95.0	382	88.2	648	90.9
Participants who have no access to sewage system in						
	271	96.8	415	95.8	686	96.2

Table 2. Overall *Plasmodium* prevalence in the study population by diagnostic test.

Municipality	Overall prevalence			Prevalence detected by microscopy and PCR			Prevalence detected only by PCR		
	n	%	95% CI	n	%	95% CI	n	%	95% CI
TOTAL n=713	30	4.21	2.95 - 5.96	18	2.52	1.59 - 3.98	12	1.68	0.96 - 2.94
<i>P. falciparum</i>	6	0.84	0.38 - 1.86	3*	0.42	0.14 - 1.30	3	0.42	0.14 - 1.30
<i>P. vivax</i>	22	3.09	2.04 - 4.65	15**	2.10	1.27 - 3.46	7	0.98	0.47 - 2.05
<i>Mixed infection</i>	2	0.28	0.07 - 1.12	0	0.00	0.00 - 0.00	2	0.28	0.07 - 1.12
El Bagre n=378	11	2.91	1.62 - 5.19	2	0.53	0.13 - 2.10	9	2.38	1.24 - 4.52
Indigenous community n=276	11	3.99	2.22 - 7.07	2	0.72	0.18 - 2.86	9	3.26	1.70 - 6.16
Non-indigenous community n=102	0	0.00	0.00 - 0.00	0	0.00	0.00 - 0.00	0	0.00	0.00 - 0.00
Turbo n=335	19	5.67	3.64 - 8.73	16	4.78	2.94 - 7.66	3	0.90	0.29 - 2.75
Indigenous community n=157	18	11.46	7.33 - 17.50	16	10.19	6.32 - 16.02	2	1.27	0.32 - 4.98
Non-indigenous community n=178	1	0.56	0.08 - 3.91	0	0.00	0.00 - 0.00	1	0.56	0.08 - 3.91

*Median parasitemia (IQR: inter quartile range) [parasites/ μ l]) 360 (40-28186), ** Median parasitemia (IQR) [parasites/ μ l]) 3147 (2805-15615)

Table 3. Association between individual and household characteristics with *Plasmodium* infections

Characteristic	Infected by <i>Plasmodium</i>				Crude IRR*				Adjusted IRR**			
	No		Yes		IRR	IC 95%			IRR	IC 95%		
	n	%	n	%								
<i>Individual factors</i>												
Sex												
Female	368	97.1	11	2.9	1				1			
Male	315	94.3	19	5.7	1.95	0.95	-	4.02	2.38	1.07	-	5.31
Age												
median (IQR)	20	(10 -37)	17	(11 -29)	0.99	0.97	-	1.01	0.99	0.97	-	1.01
Age												
<5	67	95.7	3	4.3	1				1			
5 -15	209	95.0	11	5.0	1.29	0.36	-	4.61	0.93	0.20	-	4.48
> 15	405	96.2	16	3.8	0.94	0.27	-	3.22	0.68	0.15	-	3.07
Residence time in endemic region												
median (IQR)	10	(4 -20)	12	(6 -20)	1.00	0.97	-	1.03	1	0.97	-	1.03
Residence time in endemic region												
<5 years	172	97.7	4	2.3	1				1			
≥5 years	502	95.8	22	4.2	1.64	0.57	-	4.70	1.83	0.55	-	6.08
Number of episodes of symptomatic malaria												
Median (IQR)	1	(0 -3)	2	(0 -4)	1.02	0.99	-	1.06	1.01	0.98	-	1.05
0	270	96.8	9	3.2	1				1			
1	120	96.8	4	3.2	0.98	0.30	-	3.21	0.75	0.19	-	2.92
>1	293	94.5	17	5.5	1.77	0.78	-	4.03	1.55	0.59	-	4.10
Self-report of malaria history												
0	270	96.8	9	3.2	1				1			
≥1	413	95.2	21	4.8	1.53	0.70	-	3.36	1.26	0.50	-	3.19
Participants living in the indigenous community												
No	279	99.6	1	0.4	1				1			
Yes	404	93.3	29	6.7	20.77	2.14	-	201.12	17.86	2.12	-	150.19
Use of bed net												
Yes	638	95.7	29	4.3	1				1			
No	42	97.7	1	2.3	0.44	0.05	-	4.10	0.51	0.06	-	4.14
<i>Household factors</i>												
Number of inhabitants per												

household										
Median (IQR)	5	(4 - 6)	4	(4 - 7)	1.08	0.90 -	1.31	1.05	0.88 -	1.27
1 - 4	280	96.9	9	3.1	1			1		
≥5	381	94.8	21	5.2	1.47	0.62 -	3.44	1.38	0.59 -	3.20
Participants who have animals in their households										
No	24	96.0	1	4.0	1			1		
Yes	650	95.7	29	4.3	0.88	0.12	6.51	0.79	0.10 -	5.97
Participants who drain standing water in their households										
Yes	97	99.0	1	1.0	1			1		
No	577	95.2	29	4.8	3.99	0.48 -	32.88	3.90	0.49	30.78
Participants who have access to electricity in their households										
Yes	518	97.7	12	2.3	1.00			1		
No	154	89.5	18	10.5	4.94	2.17	11.28	4.62	1.89 -	11.32
Participants who have access to water in their households										
Yes	56	100.0	0	0.0	1			1		
No	618	95.4	30	4.6	CNA***			CNA		
Participants who have access to sewage system in their households										
Yes	18	100.0	0	0.0	1			1		
No	656	95.6	30	4.4	CNA			CNA		

*Incidence Rate Ratio, **Adjusted by occupation and malaria last year, ***Convergence not achieved

Table 4. Household characteristics by indigenous and non-indigenous communities

Variable	Non-Indigenous community		Indigenous community		Total	
	n= 87		n= 125		n=212	
	n	%	n	%	n	%
Number of inhabitants						
1 - 4	55	63.2	70	56.0	125	59.0
>5	31	35.6	51	40.8	82	38.7
Ownership						
Own	79	90.8	113	90.4	192	90.6
Leased	8	9.2	9	7.2	17	8.0
Electric service						
Yes	77	88.5	84	67.2	161	75.9
No	9	10.3	38	30.4	47	22.2
Water service						
Yes	5	5.7	12	9.6	17	8.0
No	82	94.3	110	88.0	192	90.6
Sewage system						
Yes	3	3.4	4	3.2	7	3.3
No	84	96.6	118	94.4	202	95.3
Kind of water bodies						
Lake	3	3.4	2	1.6	5	2.4
Stagnant rain water	6	6.9	3	2.4	9	4.2
River	77	88.5	114	91.2	191	90.1
Fumigation (vector control program)						
Yes	78	89.7	115	92.0	193	91.0
No	9	10.3	7	5.6	16	7.5
Use of insecticides						
Yes	5	5.7	10	8.0	15	7.1
No	82	94.3	112	89.6	194	91.5
Draining standing water						
Yes	23	26.4	8	6.4	31	14.6
No	64	73.6	114	91.2	178	84.0
Mosquito mesh for windows						
Yes	0	0.0	7	5.6	7	3.3
No	87	100.0	115	92.2	202	95.3
Wall material						
Wood	74	85.1	103	82.4	177	83.5
Cement blocks	10	11.5	10	8.0	20	9.4

Others	3	3.4	9	7.2	12	5.7
Roof material						
Palm leaf	29	33.3	19	15.2	48	22.6
Mud	7	8.0	13	10.4	20	9.4
Tin	49	56.3	90	72.0	139	65.6
Plastic	2	2.3	0	0.0	2	0.9

Figures

Figure 1

Study sites in Antioquia department, Colombia

Figure 2

Flowchart of enrolled participants and proportion of infected people by villages. * Plasmodium infections detected by either microscopy or PCR

Figure 3

Proportion of Plasmodium infection in indigenous and non-indigenous communities. A. Proportion of Plasmodium infection by species B. Proportion of Plasmodium infection by diagnostic test (Microscopic infection were detected by microscopy and PCR and submicroscopic infection were only detected by PCR)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarytable1.docx](#)