

Long noncoding RNA Gm44593 attenuates oxidative stress from age-related hearing loss by regulating miR-29b/WNK1

Qian Li

People's Hospital of Zhengzhou University

Yanzi Zang

People's Hospital of Zhengzhou University

Zhanwei Sun

People's Hospital of Zhengzhou University

Wenqi Zhang

People's Hospital of Zhengzhou University

Hongjian Liu (✉ liuhjhn78@yeah.net)

Otolaryngology Head and Neck Surgery, People's Hospital of Zhengzhou University

<https://orcid.org/0000-0001-6561-3949>

Research article

Keywords: lncRNA Gm44593, oxidative stress, AHL

Posted Date: October 8th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-86114/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Bioengineered on December 30th, 2021.
See the published version at <https://doi.org/10.1080/21655979.2021.2012062>.

Abstract

Amis

Long noncoding RNA has been reported to play important role in various disease. However, the function of lncRNA in age-related hearing loss still unclear. The aim of our study is to investigate the function and mechanism of lncRNA Gm44593 in AHL

Materials and methods

ATP content, JC-1 assay, mitochondrial content, cell death rates and dual-luciferase reporter assay were performed to assess the function of lncRNA Gm44593 in HEI-OC1 cells and the mouse cochlea.

Results

The expression of lncRNA Gm44593 was significantly upregulated upon H_2O_2 and starvation treatment. Overexpression of lncRNA Gm44593 manifestly reduced the cell death rates. The ATP content, mtDNA content and mitochondrial membrane potential were alleviated upon overexpression of lncRNA Gm44593. We also proved that miR-29b is the direct target of lncRNA Gm44593. Overexpression of miR-29b completely restored the effect induced by lncRNA Gm44593. In addition, we provided evidences that WNK1 is the direct target of miR-29b.

Conclusion

our research uncovers the role of lncRNA Gm44593 in age-related hearing loss. We provide new insights into potential therapeutic targets for the amelioration of age-related hearing loss.

Introduction

Age-related hearing loss (AHL), characterized by the irreversible loss of cochlear hair cells[1], spiral ganglion neurons (SGNs) and stria vascularis cells (SVs), is the most common sensory disorder in old people[2], influencing the life quality older people by age 65 years or more[3]. AHL increases the risk of depression, cognitive impairment and dementia in aged adults[4, 5]. Thus, it is important to explore the mechanism of AHL and identify new targets for clinical treatment.

Long noncoding RNA (lncRNA) is a specific noncoding RNA without coding potential[6]. Numerous studies have revealed the function of lncRNA in different pathological process, such as cell apoptosis[7], cell cycle[8], necrosis, EMT[9]. Studies of lncRNA also involved numerous mechanisms, including ceRNA mechanism, chromosome remodeling, post-translational regulation[10, 11]. lncRNAs have been reported to function in numerous diseases, such as diabetes, cardiovascular diseases, tumor development[12]. Recent studies have revealed that dysregulated lncRNAs may lead to a new treatment for AHL[13]. There is no doubt that explore new lncRNA in AHL will help us to better understand the mechanism of AHL.

Mitochondria play an important role in maintaining cellular homeostasis[14]. Accumulated evidence demonstrated that mitochondrial dysfunction was related with aging diseases, including neurodegenerative disease and AHL[2, 15], although whether this is causal or simply correlative has yet to be established. Mitochondrial biogenesis is a highly regulated process to maintain normal mitochondrial function, thus regulate different physiological process[16].

In this study, we aimed to explore the function and mechanism of lncRNA Gm44593 in AHL. lncRNA Gm44593 was firstly identified as an upregulated lncRNA in the cochlea of aged C57BL/6 mice[2]. To date, there is no report about the role of lncRNA Gm44593 in AHL. We performed comprehensive functional analysis to assess the role of lncRNA Gm44593, such as cell proliferation, cell death, mitochondrial function. Bioinformatics analysis predicted numerous miRNAs may bind with lncRNA Gm44593, considering the location of lncRNA Gm44593. Rescue experiment further proved that miR-29b can reverse the effect of lncRNA Gm44593. Our research uncovers the role of lncRNA Gm44593 in age-related hearing loss. We provide new insights into potential therapeutic targets for the amelioration of age-related hearing loss.

Materials And Methods

Cell culture and transfection

The House Ear Institute-Organ of Corti 1 (HEI-OC1) cells were cultured in high glucose Dulbecco's Modified Eagle Medium (DMEM, Gibco) containing 10% fetal bovine serum (FBS, Gibco, #10437028) without antibiotics at 33 °C and 10% CO₂.

lncRNA Gm44593 overexpression plasmid was purchased from Gene Pharma (Shanghai, China). Wild type and mutant lncRNA Gm44593 vector were constructed by Gene Pharma (Shanghai, China). miR-29b mimics and inhibitor were purchased from Gene Pharma (Shanghai, China). All transfection experiments were performed via lipo 3000 reagent (Invitrogen) according to the manufacturer's protocol.

Proliferation assay

HEI-OC1 cells were transfected with lncRNA Gm44593 overexpression vector or NC vector. Cells were seeded into 96-well plates with density of 2×10^4 cells/ml and incubated overnight at 33 °C and 10% CO₂. After 48 h transfection, CCK-8 assay was performed following the manufacturer's protocol. The optical density was measured at 450 nm after 2 h incubation.

Cell death rates

HEI-OC1 cells were transfected with lncRNA Gm44593 overexpression vector or NC vector. After cell confluence reach 90%, cells were digested and pipetted thoroughly. Trypan blue staining (Beyotime, C0011, Shanghai, China) was used to measure the cell death rates.

TUNEL

Terminal deoxytransferase-mediated dUTP-biotin nick end labeling (TUNEL) assay was performed to detect the level of apoptosis. Briefly, HEI-OC1 cells were transfected with lncRNA Gm44593 overexpression vector or NC vector. After H₂O₂ exposure, Cells were washed with PBS for twice and immobilized by 4% paraformaldehyde. The apoptotic cells were visualized with the TUNEL staining following the manufacturer's instructions (Beyotime, C1090, Shanghai, China). The fluorescence density was assessed using Image J software.

JC-1 mitochondrial membrane potential

JC-1 at a concentration of 10 µg/ml (Beyotime, C2006, Shanghai, China) was added to the medium for 10 minutes. Cells were then washed twice in PBS. Fluorescence emission was filtered at 485 and 580 nm. All experiments were repeated for three times.

ATP contents

ATP assay was performed using an ATP Assay Kit (Beyotime, Shanghai, China) following the manufacturer's protocols. Chemiluminescence of samples and standards were measured with a SpectraMax M5 microplate reader (Molecular Devices, USA). The levels of ATP were calculated based on the standard curve and normalized to the protein content.

Dual-luciferase reporter assay

HEK293 cells were co-transfected with target Gm44593 or mutant Gm44593 with binding sites for miR-29b. Dual-luciferase reporter assay were conducted using Dual Luciferase Reporter Assay System (Promega, USA) following the manufacturer's protocol. Luciferase activity was measured with Multiskan Spectrum (Thermo Fisher, USA).

RIP

HEI-OC1 cells were transfected with lncRNA Gm44593 overexpression vector and miR-29b mimics. The antibody against human Ago2 was used for RNA immunoprecipitation. RIP experiments were performed using Magna RIP Kit. After Ago2 immunoprecipitation, the RNA was isolated, the expression level of lncRNA Gm44593 and miR-29b was detected by real-time PCR.

Real-time PCR

Total RNA was extracted and lysed by TRIzol reagent (Thermo Fisher, USA). RNA reverse transcription was performed using a PrimeScript™ RT reagent Kit with gDNA eraser (Takara, Japan) according to the manufacturer's instructions, and cDNA was performed by qRT-PCR using SYBR® Premix Ex Taq™ (Takara, Japan). The data were normalized using β-actin levels and further analyzed by the 2^{-ΔΔCT} method.

Statistical analysis

Data are presented as the mean ± sd, and statistical analyses were performed using ANOVA or unpaired Student's t-test with GraphPad Prism 8. P < 0.05 was considered to be significant.

Results

Biological features of lncRNA Gm44593

To study the function of lncRNA Gm44593, we analyzed the expression of lncRNA Gm44593 upon H₂O₂ and starvation treatment. As shown in Figure 1A-B, the expression of lncRNA Gm44593 was significantly increased upon H₂O₂ and starvation. Besides, we verified the location of lncRNA Gm44593 through PARIS kit. Our results showed that lncRNA Gm44593 located in cytoplasm, while GAPDH and U6 served as positive control (Figure 1C).

Overexpression of lncRNA Gm44593 maintained cell function upon H₂O₂ treatment

To study the function of lncRNA Gm44593, we constructed overexpression vector. As shown in Figure 2A, the overexpression efficiency was confirmed by real-time PCR. Next, we performed comprehensive analysis to assess the function of lncRNA Gm44593 in mitochondrial. Overexpression had no effect on cell proliferation via CCK-8 (Figure 2B). However, when we detected cell death rates, to our surprise, overexpression of lncRNA Gm44593 significantly reduced the cell death rates compared with NC group upon H₂O₂ treatment (Figure 2C). Next, we analyzed cell apoptosis rates via TUNEL assay (Figure 2D). Our results showed that overexpression of lncRNA Gm44593 remarkable reduced the cell apoptosis. Quantitative data also showed the same effect of lncRNA Gm44593 (Figure 2E). Mitochondrial function is the most important process in the aging. Thus, we assessed the cell mitochondrial function through JC-1, which assessed the membrane potential, ATP and mtDNA copy number, which assessed the cell mitochondrial function. Our results showed that lncRNA Gm44593 could maintained the cell mitochondrial function upon H₂O₂ treatment (Figure 2F-H). Thus, our results demonstrated that overexpression of lncRNA Gm44593 may exert beneficial effects during aging process.

miR-29b is the direct target of lncRNA Gm44593

To study the mechanism of lncRNA Gm44593 in the aging, and considering that lncRNA Gm44593 located in cytoplasm, we used starbase and miRNA data base to predict the potential binding miRNAs. As shown in Figure 3A, only miR-29b was decreased in the lncRNA Gm44593 overexpression group. Thus, we further analyzed the binding sequence between miR-29b and lncRNA Gm44593 (Figure 3B). To examine the target effect of miR-29b, we overexpressed miR-29b via mimics in HEI-OC1 and HEK cells. The overexpression efficiency was confirmed via real-time PCR (Figure 3C). Dual-luciferase reporter assay results showed that overexpression of miR-29b significantly reduced the luciferase activity of target gene in the wide-type group, while there is no significant difference in the mutant group (Figure 3D). Next, we performed comprehensive analysis to assess the relationship between miR-29b and lncRNA Gm44593. Overexpression of lncRNA Gm44593 significantly decreased the expression of miR-29b, while knockdown of lncRNA Gm44593 increased the expression of miR-29b (Figure 3E). We also analyzed the expression of miR-29b in the aging process, as indicated in Figure 3F-G, which is downregulated upon starvation and

H₂O₂ treatment (Figure 3F-G). Lastly, lncRNA Gm44593 negatively regulate the expression of miR-29b (Figure 3H).

WNK1 is the direct target of miR-29b

To further evaluate the mechanism of miR-29b, we used starbase to predict the potential downstream target. We predicted WNK1 is the possible target, as indicated in Figure 4A. Next, we used dual-luciferase reporter assay to assess the binding relationship in HEK cells. The relative luciferase activity was significantly downregulating in the miR-29b overexpression group, while there is no significant difference in the mutant group (Figure 4B). Next, we performed comprehensive analysis to assess the relationship between miR-29b and WNK1. Overexpression of miR-29b significantly decreased the expression of WNK1, while knockdown of miR-29b increased the expression of WNK1 (Figure 4C). We also analyzed the expression of WNK1 in the aging process, as indicated in Figure 4D-E, which is upregulated upon starvation and H₂O₂ treatment.

lncRNA Gm44593 positively regulate WNK1

To date, there is no report about the relationship between lncRNA Gm44593 and WNK1. Firstly, when we overexpressed lncRNA Gm44593, we found that the expression of WNK1 was upregulated in the lncRNA Gm44593 overexpression group, while miR-29b can completely reversed the effect induced by lncRNA Gm44593 (Figure 5A). Our results revealed that lncRNA Gm44593 positively regulate WNK1 (Figure 5B). Lastly, we used RIP experiment to further confirm the binding between miR-29b and lncRNA Gm44593. As shown in Figure 5C, the expression of miR-29b and lncRNA Gm44593 was significantly upregulated in anti-Ago2 group.

Re-introduction of miR-29b reversed the effect of lncRNA Gm44593

To date, we have analyzed the function of lncRNA Gm44593 in the aging process and verified the downstream target. To confirm our results, we performed comprehensive rescue experiment. As shown in Figure 6A, overexpression of lncRNA Gm44593 significantly reduced the cell death rates, while re-introduction of miR-29b completely reversed such effect. Besides, miR-29b also rescued the effect induced by lncRNA Gm44593 in mitochondrial function, as indicated by decreased mitochondrial membrane potential, ATP contents and mtDNA copy numbers (Figure 6B-D). Thus, our results provided evidences that lncRNA Gm44593 can protect cells against aging through regulating miR-29b and WNK1 (Figure 7).

Discussion

In this study, we aimed to explore the function and mechanism of lncRNA Gm44593 in AHL. lncRNA Gm44593 was firstly identified as an upregulated lncRNA in the cochlea of aged C57BL/6 mice. To date, there is no report about the role of lncRNA Gm44593 in AHL. We performed comprehensive functional analysis to assess the role of lncRNA Gm44593, such as cell proliferation, cell death, mitochondrial

function. Bioinformatics analysis predicted numerous miRNAs may bind with lncRNA Gm44593, considering the location of lncRNA Gm44593. Rescue experiment further proved that miR-29b can reverse the effect of lncRNA Gm44593. Our research uncovers the role of lncRNA Gm44593 in age-related hearing loss. We provide new insights into potential therapeutic targets for the amelioration of age-related hearing loss.

Several previous studies have shown that AHL is commonly associated with ROS accumulation[17], which induces mitochondrial depolarization and initiates apoptosis. ROS can induce cellular defense pathways, including autophagy, which can recycle unnecessary or dysfunctional cellular components[18]. Mitochondria have a vital role in maintaining cellular homeostasis[19]. Growing evidence suggests that mitochondrial dysfunction participates in aging diseases, such as diabetes, neurodegenerative disease[20] and AHL[21]. Mitochondrial biogenesis is a tightly regulated process to generate new mitochondria and plays an important role in maintaining normal mitochondrial function[22]. Mitochondrial biogenesis takes place under basal condition and is an adaptive response induced by cells to maintain energy demands. Mitochondria are the cellular center for energy production as well as the major source of ROS[23]. The accumulation of ROS could impair antioxidant functions and damage macromolecules, such as nuclear DNA, mitochondrial DNA (mtDNA), membranes, and proteins. Moreover, accumulation of macromolecule mutations can cause apoptosis of hair cells and thus promote the development of AHL. In our study, we revealed that lncRNA Gm44593 reduced the ROS contents and alleviate aging induced apoptosis.

Accumulating evidences showed that lncRNA play essential role in the regulation of gene expression and participate in multiple biological and pathological process, including cell proliferation, cell apoptosis, cell differentiation, EMT and m6A regulation[24]. In the inner ear, epigenetic modifications might also be related to inner ear development and have a significant role in hearing loss, hearing protection, and regeneration of functional cells[25]. A recent study revealed differential lncRNA profile between two developmental stages of the mouse inner ear sensory epithelium of the cochlea and vestibule, suggesting a possible role for lncRNAs in regulating hearing and balance[26]. In our study, we used bioinformatics method to predict the binding target of lncRNA Gm44593 and confirmed the binding relationship between lncRNA Gm44593 and miR-29b. Further rescue experiment proved that lncRNA Gm44593 alleviate aging induced apoptosis through negatively regulating miR-29b.

In summary, we analyzed the function and mechanism of lncRNA Gm44593 in AHL. Our results provided new clues about lncRNA in AHL and help understand the development of AHL. Overexpression of lncRNA Gm44593 may provide new intervention sights for AHL.

Declarations

Conflict of interest:

The authors declare no interests existed.

Funding acknowledgement

The project was funded by High-technology and Innovative project of Henan Province (2017-HI-1331)

References

- [1] Gray DT and Barnes CA. Experiments in macaque monkeys provide critical insights into age-associated changes in cognitive and sensory function. *Proceedings of the National Academy of Sciences of the United States of America* 2019;
- [2] Su Z, Xiong H, Pang J, Lin H, Lai L, Zhang H, Zhang W and Zheng Y. LncRNA AW112010 Promotes Mitochondrial Biogenesis and Hair Cell Survival: Implications for Age-Related Hearing Loss. *Oxidative medicine and cellular longevity* 2019; 2019: 6150148.
- [3] Caballero JA and Brown SP. Engagement, not personal characteristics, was associated with the seriousness of regulatory adjudication decisions about physicians: a cross-sectional study. *BMC medicine* 2019; 17: 211.
- [4] Yong M, Willink A, McMahon C, McPherson B, Nieman CL, Reed NS and Lin FR. Access to adults' hearing aids: policies and technologies used in eight countries. *Bulletin of the World Health Organization* 2019; 97: 699-710.
- [5] Suen JJ, Bhatnagar K, Emmett SD, Marrone N, Kleindienst Robler S, Swanepoel W, Wong A and Nieman CL. Hearing care across the life course provided in the community. *Bulletin of the World Health Organization* 2019; 97: 681-690.
- [6] Martone J, Mariani D, Desideri F and Ballarino M. Non-coding RNAs Shaping Muscle. *Frontiers in cell and developmental biology* 2019; 7: 394.
- [7] Xia P, Gu R, Zhang W and Sun YF. lncRNA CEBPA-AS1 Overexpression Inhibits Proliferation and Migration and Stimulates Apoptosis of OS Cells via Notch Signaling. *Molecular therapy. Nucleic acids* 2020; 19: 1470-1481.
- [8] Cao Y, Xiong JB, Zhang GY, Liu Y, Jie ZG and Li ZR. Long Noncoding RNA UCA1 Regulates PRL-3 Expression by Sponging MicroRNA-495 to Promote the Progression of Gastric Cancer. *Molecular therapy. Nucleic acids* 2020; 19: 853-864.
- [9] Hong W, Ying H, Lin F, Ding R, Wang W and Zhang M. lncRNA LINC00460 Silencing Represses EMT in Colon Cancer through Downregulation of ANXA2 via Upregulating miR-433-3p. *Molecular therapy. Nucleic acids* 2020; 19: 1209-1218.
- [10] Ou C, Sun Z, He X, Li X, Fan S, Zheng X, Peng Q, Li G, Li X and Ma J. Targeting YAP1/LINC00152/FSCN1 Signaling Axis Prevents the Progression of Colorectal Cancer. *Advanced science (Weinheim, Baden-Wurtemberg, Germany)* 2020; 7: 1901380.

- [11] Xiu B, Chi Y, Liu L, Chi W, Zhang Q, Chen J, Guo R, Si J, Li L, Xue J, Shao ZM, Wu ZH, Huang S and Wu J. LINC02273 drives breast cancer metastasis by epigenetically increasing AGR2 transcription. *Molecular cancer* 2019; 18: 187.
- [12] Li N, Yang G, Luo L, Ling L, Wang X, Shi L, Lan J, Jia X, Zhang Q, Long Z, Liu J, Hu W, He Z, Liu H, Liu W and Zheng G. THAP9-AS1 lncRNA Promotes Pancreatic Ductal Adenocarcinoma Growth and Leads to a Poor Clinical Outcome via Sponging miR-484 and Interacting with YAP. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2020; 26: 1736-1748.
- [13] T Zhao, X Liu, Z Sun, J Zhang, X Zhang, C Wang, R Geng, T Zheng, B Li and QY Zheng. RNA-seq analysis of potential lncRNAs for age-related hearing loss in a mouse model. *Aging* 2020; 12: 7491-7510.
- [14] Kraja AT, Liu C, Fetterman JL, Graff M, Have CT, Gu C, Yanek LR, Feitosa MF, Arking DE, Chasman DI, Young K, Ligthart S, Hill WD, Weiss S, Luan J, Giulianini F, Li-Gao R, Hartwig FP, Lin SJ, Wang L, Richardson TG, Yao J, Fernandez EP, Ghanbari M, Wojczynski MK, Lee WJ, Argos M, Armasu SM, Barve RA, Ryan KA, An P, Baranski TJ, Bielinski SJ, Bowden DW, Broeckel U, Christensen K, Chu AY, Corley J, Cox SR, Uitterlinden AG, Rivadeneira F, Cropp CD, Daw EW, van Heemst D, de Las Fuentes L, Gao H, Tzoulaki I, Ahluwalia TS, de Mutsert R, Emery LS, Erzurumluoglu AM, Perry JA, Fu M, Forouhi NG, Gu Z, Hai Y, Harris SE, Hemani G, Hunt SC, Irvin MR, Jonsson AE, Justice AE, Kerrison ND, Larson NB, Lin KH, Love-Gregory LD, Mathias RA, Lee JH, Nauck M, Noordam R, Ong KK, Pankow J, Patki A, Pattie A, Petersmann A, Qi Q, Ribel-Madsen R, Rohde R, Sandow K, Schnurr TM, Sofer T, Starr JM, Taylor AM, Teumer A, Timpson NJ, de Haan HG, Wang Y, Weeke PE, Williams C, Wu H, Yang W, Zeng D, Witte DR, Weir BS, Wareham NJ, Vestergaard H, Turner ST, Torp-Pedersen C, Stergiakouli E, Sheu WH, Rosendaal FR, Ikram MA, Franco OH, Ridker PM, Perls TT, Pedersen O, Nohr EA, Newman AB, Linneberg A, Langenberg C, Kilpeläinen TO, Kardina SLR, Jørgensen ME, Jørgensen T, Sørensen TIA, Homuth G, Hansen T, Goodarzi MO, Deary IJ, Christensen C, Chen YI, Chakravarti A, Brandslund I, Bonnelykke K, Taylor KD, Wilson JG, Rodriguez S, Davies G, Horta BL, Thyagarajan B, Rao DC, Grarup N, Davila-Roman VG, Hudson G, Guo X, Arnett DK, Hayward C, Vaidya D, Mook-Kanamori DO, Tiwari HK, Levy D, Loos RJF, Dehghan A, Elliott P, Malik AN, Scott RA, Becker DM, de Andrade M, Province MA, Meigs JB, Rotter JI and North KE. Associations of Mitochondrial and Nuclear Mitochondrial Variants and Genes with Seven Metabolic Traits. *American journal of human genetics* 2019; 104: 112-138.
- [15] Kline RA, Dissanayake KN, Hurtado ML, Martínez NW, Ahl A, Mole AJ, Lamont DJ, Court FA, Ribchester RR, Wishart TM and Murray LM. Altered mitochondrial bioenergetics are responsible for the delay in Wallerian degeneration observed in neonatal mice. *Neurobiology of disease* 2019; 130: 104496.
- [16] Kwon DN, Park WJ, Choi YJ, Gurunathan S and Kim JH. Oxidative stress and ROS metabolism via down-regulation of sirtuin 3 expression in Cmah-null mice affect hearing loss. *Aging* 2015; 7: 579-594.
- [17] Someya S, Yu W, Hallows WC, Xu J, Vann JM, Leeuwenburgh C, Tanokura M, Denu JM and Prolla TA. Sirt3 mediates reduction of oxidative damage and prevention of age-related hearing loss under caloric restriction. *Cell* 2010; 143: 802-812.

- [18] Someya S and Prolla TA. Mitochondrial oxidative damage and apoptosis in age-related hearing loss. *Mechanisms of ageing and development* 2010; 131: 480-486.
- [19] Kondadi AK, Anand R and Reichert AS. Functional Interplay between Cristae Biogenesis, Mitochondrial Dynamics and Mitochondrial DNA Integrity. *International journal of molecular sciences* 2019; 20:
- [20] Palmeira CM, Teodoro JS, Amorim JA, Steegborn C, Sinclair DA and Rolo AP. Mitohormesis and metabolic health: The interplay between ROS, cAMP and sirtuins. *Free radical biology & medicine* 2019; 141: 483-491.
- [21] Palikaras K and Tavernarakis N. Mitochondrial homeostasis: the interplay between mitophagy and mitochondrial biogenesis. *Experimental gerontology* 2014; 56: 182-188.
- [22] Dornas W and Schuppan D. Mitochondrial oxidative injury: A key player in nonalcoholic fatty liver disease. *American journal of physiology. Gastrointestinal and liver physiology* 2020;
- [23] Someya S, Xu J, Kondo K, Ding D, Salvi RJ, Yamasoba T, Rabinovitch PS, Weindruch R, Leeuwenburgh C, Tanokura M and Prolla TA. Age-related hearing loss in C57BL/6J mice is mediated by Bak-dependent mitochondrial apoptosis. *Proceedings of the National Academy of Sciences of the United States of America* 2009; 106: 19432-19437.
- [24] Zhu LY, Zhu YR, Dai DJ, Wang X and Jin HC. Epigenetic regulation of alternative splicing. *American journal of cancer research* 2018; 8: 2346-2358.
- [25] Yamasoba T, Someya S, Yamada C, Weindruch R, Prolla TA and Tanokura M. Role of mitochondrial dysfunction and mitochondrial DNA mutations in age-related hearing loss. *Hearing research* 2007; 226: 185-193.
- [26] Ushakov K, Koffler-Brill T, Rom A, Perl K, Ulitsky I and Avraham KB. Genome-wide identification and expression profiling of long non-coding RNAs in auditory and vestibular systems. *Scientific reports* 2017; 7: 8637.

Figures




Figure 1

Biological features of lncRNA Gm44593. A. The expression of lncRNA Gm44593 was upregulated upon H₂O₂ treatment. B. The expression of lncRNA Gm44593 was increased upon starvation treatment. C. lncRNA Gm44593 located in cytoplasm via PARIS kit.




Figure 2

Overexpression of lncRNA Gm44593 maintained cell function. A. The overexpression efficiency was confirmed via real-time PCR. B. lncRNA Gm44593 had no effect on proliferation. C. Overexpression of lncRNA Gm44593 significantly reduced the cell death rates. D. TUNEL assay showed that overexpression of lncRNA Gm44593 significantly reduced the cell apoptosis rates. E. Quantitative data of TUNEL assay. F. JC-1 mitochondrial membrane potential results showed that lncRNA Gm44593 can maintain the cell membrane potential. G. lncRNA Gm44593 maintained the ATP content compared with NC group upon H₂O₂ treatment. H. Upon H₂O₂ treatment, lncRNA Gm44593 increased the mtDNA number compared with NC group.




Figure 3

miR-29b is the direct target of lncRNA Gm44593. A. 5 miRNAs were predicted to work with lncRNA Gm44593. Only miR-29b was downregulated. B. The potential binding sequence between miR-29b and lncRNA Gm44593 were showed. C. The overexpression efficiency was confirmed via real-time PCR in HEI-OC1 and HEK cells. D. Dual-luciferase reporter assay showed that the relative activity was significantly downregulated in miR-29b group. E. Overexpression of lncRNA Gm44593 reduced the expression of miR-29b, while knockdown of lncRNA Gm44593 increased the expression of miR-29b. F. The expression of miR-29b was significantly downregulated upon starvation treatment. G. The expression of miR-29b was significantly downregulated upon H2O2 treatment. H. lncRNA Gm44593 negatively regulate the expression of miR-29b.




Figure 4

WNK1 is the direct target of miR-29b. A. Bio-informatics analysis predicted that WNK1 was the direct target of miR-29b. The potential binding sequence was showed. B. The dual-luciferase activity was significantly downregulated in miR-29b overexpression group in HEK cells. C. Overexpression of miR-29b significantly reduced the expression of WNK1, while miR-29b inhibitor treatment increased the expression of WNK1. D. The expression of WNK1 was significantly upregulated upon starvation treatment. E. The expression of WNK1 was significantly increased upon H2O2 treatment.




Figure 5

lncRNA Gm44593 positively regulate WNK1. A. Overexpression of lncRNA Gm44593 increased the expression of WNK1, and miR-29b reversed such effect. B. lncRNA Gm44593 positively regulate WNK1. C. RIP experiment confirmed the binding between miR-29b and lncRNA Gm44593.




Figure 6

Re-introduction of miR-29b reversed the effect of lncRNA Gm44593. A. Re-introduction of miR-29b completely reversed the protective role of lncRNA Gm44593. B. Overexpression of miR-29b restored the effect of lncRNA Gm44593 in mitochondrial membrane potential. C. ATP content was measured to evaluate the restore effect of miR-29b. D. mtDNA number was assessed. Re-introduction of miR-29b reversed the effect induced by lncRNA Gm44593.




Figure 7

Graphical scheme of lncRNA Gm44593/miR-29b/WNK1.