Insecticide Resistance in *Anopheles stephensi* in Somali Region, Eastern Ethiopia

Solomon Yared (✉ solyar2005@yahoo.com)

Jigjiga University <https://orcid.org/0000-0003-1510-8316>

Araya Gebressielasie

Addis Ababa University

Lambodhar Damodaran

University of Georgia

Victoria Bonnell

University of North Carolina at Charlotte

Karen Lopez

University of North Carolina at Charlotte

Daniel Janies

University of North Carolina at Charlotte

Tamar Carter

Baylor University

Research

Keywords: *Anopheles stephensi*, Insecticide resistance, Kebridehar, Somali Region

Posted Date: December 4th, 2019

DOI: <https://doi.org/10.21203/rs.2.18020/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Malaria Journal on May 12th, 2020. See the published version at <https://doi.org/10.1186/s12936-020-03252-2>.

Abstract

Background: The movement of malaria vectors into new areas is a growing concern in the efforts to control malaria. The recent report of *Anopheles stephensi* in eastern Ethiopia has raised the necessity to understand the insecticide resistance status of the vector in the region to better inform vector-based interventions. The aim of this study was to evaluate insecticide resistance in *An. stephensi* in eastern Ethiopia using two approaches: 1) World Health Organization (WHO) bioassay tests in *An. stephensi* and 2) genetic analysis of insecticide resistance genes in *An. stephensi* in eastern Ethiopia.

Methods: Mosquito larva and pupae were collected from Kebridehar. Insecticide susceptibility of *An. stephensi* was tested with malathion 5%, bendiocarb 0.1%, propoxur 0.1%, deltamethrin 0.05%, permethrin 0.75%, Pirimiphos-methyl 0.25% and DDT 4%, according to WHO standard protocols.

Results: All *An. stephensi* samples were resistant to carbamates, with mortality rates 23% and 21% for bendiocarb and propoxur, respectively. Adult *An. stephensi* was also resistant to pyrethroid insecticides with mortality rates 67% for deltamethrin and 53% for permethrin. Resistance to DDT and malathion was detected in *An. stephensi* with mortality rates of 32% as well as *An. stephensi* was resistance to pirimiphos-methyl with mortality rates 14%. Analysis of the voltage gate sodium channel gene (*vgsc*) revealed the absence of *kdr* L1014 mutations.

Conclusion: Overall, these findings support that *An. stephensi* is resistant to several classes of insecticides, most notably pyrethroids. However, the absence of the *kdr* L1014 gene may suggest non-target site resistance mechanisms. Continued insecticide resistance surveillance should be carried out in the region to confirm the findings on *An. stephensi* in Ethiopia.

Background

Malaria remains a major health problem. In 2017, an estimated 219 million cases of malaria occurred worldwide and 435,000 resulted in deaths (WHO, 2018). In Ethiopia, malaria remains a major public health concern with millions of cases and thousands of deaths reported annually (WHO, 2017). Unlike most of the African continent, malaria can be caused by infection with *Plasmodium vivax* or *P. falciparum*. Efforts to control the transmission of malaria currently target *Anopheles arabiensis*, the primary malaria vector in Ethiopia, as well the secondary vectors *An. funestus*, *An. pharoensis*, and *An. nili* (Ghebreyesus *et al.*, 2006). Successes in reducing the malaria burden are now threatened by the recent detection of the South Asian urban vector *An. stephensi* in the Horn of Africa. This mosquito was first detected in the Somali Regional State of Ethiopia (Carter, Yared *et al.*, 2018) in 2016 and has subsequently been confirmed to have a broad distribution in Northeast and east Ethiopia (Balkew *et al.* 2019). *An. stephensi* was also been reported in Djibouti in 2014 (Seyfarth *et al.* 2019) and there are now concerns that this species may spread throughout the African continent (WHO, 2019).

One major obstacle to vector control in Ethiopia and elsewhere is the ever-developing insecticide resistance as a result of indiscriminate and rampant use of the synthetic chemicals in public health and agriculture (Fettene *et al.*, 2013; Reid and McKenzie, 2016; Rivero *et al.*, 2010). Pyrethroids remain the only class of insecticides recommended for the treatment of (long-lasting insecticide-treated nets) LLINs and accounted for a large proportion of the insecticide used for indoor residual spraying (IRS) in Ethiopia and elsewhere in Africa (WHO, 2015; Kenea *et al.*, 2019). This heavy reliance on a single insecticide class has caused mosquito species to develop insecticide resistance. In mosquitoes, pyrethroid resistance is mainly attributed to two major mechanisms: target-site

insensitivity and metabolic-based resistance. Target-site resistance is due to mutations in the voltage-gated sodium channel on the mosquito's neurons that prevent the insecticide's ability to interfere with the closing of sodium channel that would usually result in paralysis (knockdown) (for review see Ranson *et al.*, 2011). The knockdown (kdr) mutation L1014 has been observed across multiple Culicidae. Metabolic resistance mediated by detoxifying enzymes also plays a significant role in insecticide resistance in malaria vectors (Hemingway and Ranson, 2000; Liu, 2015). The over-expression of detoxification enzymes such as cytochrome P450 monooxygenases (P450s), carboxylcholinesterases (CCEs) and glutathione S-transferases (GSTs) in mosquitoes are frequently associated with resistance to different classes of insecticides (David *et al.*, 2013).

Insecticide resistance in *An. stephensi* has been reported in Afghanistan, Pakistan, Dubai, and India (Safi et al. 2017). In these regions, the frequency of the kdr L1014 mutation varies with strong support for metabolic resistance as well as target site resistance playing a role in *An. stephensi*. In Ethiopia, studies on *An. arabiensis* in the western portion of the country report phenotypic resistance to pyrethroids along with L1014 variant (Alemayehu et al., 2017). Data on the insecticide resistance status of malaria vectors in the eastern portion of the country is lacking, including that of the recently identified *An. stephensi*. Knowing the status of insecticide resistance of local malaria vectors can aid with vector control planning that involves the use of insecticides. Here we report the insecticide resistance status of east Ethiopian *An. stephensi* using bioassay tests and molecular analysis.

Methods And Materials

Study area

Our samples were collected in Kebri Dehar, a small town in the Somali Regional State, as previously detailed (Carter and Yared *et al.*, 2018). This town has a tropical semiarid climate with typically bimodal rainfall patterns. The area is also known to experience recurrent droughts. The population size is over 100,000 individuals, many of whom are pastoralists.

Larval sampling and rearing procedure

Larvae samples were collected from Kebri Dehar small town in Somali region in the SRS in eastern Ethiopia as previously detailed (Carter and Yared *et al.*, 2018). This region is predominantly lowland plain, with sparse vegetation including trees and shrubby. Briefly, immature stages of *An. stephensi* were collected from water reservoirs in Kebridehar town using the dipping method according to WHO guidelines and were reared to adulthood in the field laboratory (Fig. 1). Collections took place from November to December 2016. In the field laboratory, the larvae and pupae were maintained at $28 \pm 2^{\circ}\text{C}$ and $70 \pm 10\%$ relative humidity. The pupae were sorted and transferred with pipettes from the enamel trays to beakers with small amounts of water. Each beaker was placed inside a cage and was provided with 10% sugar solution for rearing them in the cage (Fig 1). After two to three days, the pupae emerged to adults and the cages were put in safe place protected from contamination, ants, and other insects. The lab reared females of *An. stephensi* were used for different insecticide susceptibility test using WHO bioassay.

Mosquitoes Identification

Species identification of *the* mosquitoes was conducted using morphological keys (Gillies, 1987, Glick and Wrbuw, 1992) and molecular analysis of ITS2 and COI loci as reported previously for *An. stephensi* (Carter and Yared *et al.*, 2018).

Insecticide susceptibility test for *An. stephensi*

Insecticide susceptibility tests were carried out following WHO insecticide susceptibility test procedure (WHO, 2013). A total of 700 non-blood fed adult female *An. stephensi* (2–3 day-old) were exposed to insecticide impregnated papers with discriminating concentrations of DDT (4%), malathion (5%), deltamethrin (0.05%), permethrin 0.75%, bendiocarb (0.1%), and propoxur (0.1%). Batches of 25 mosquitoes in four replicates were exposed to insecticide impregnated papers for 1 h in WHO test tubes. The knockdown effects for all tested insecticides were recorded at 10, 15, 20, 30, 40, 50, and 60 minutes. A control in two replicates (50 female *An. stephensi* were used for each insecticide), each with equal number of mosquitoes, exposed to papers impregnated with oil was run concurrently. Mosquitoes were then transferred into holding tubes with untreated papers, where they were supplied with 10% sucrose solution. And, mortality was recorded 24-hour post exposure. When mosquito mortality rate in the control is between 5–20%, mortality was corrected using Abbott's formula (Abbott, 1925). All survived and dead specimens following bioassay were kept individually in Eppendorf tubes in silica gel for further molecular identification and *kdr* PCR assays.

Determination of resistance was based on WHO criteria as follows: 98–100% mortality indicates susceptibility, 90–97% mortality indicates resistance candidate (more investigation is needed) and less than 90% mortality suggests resistance (WHO, 2013).

Amplification and sequencing of insecticide resistance loci

For *kdr* mutation analysis, PCR was used to amplify the region of the *vgsc* gene that housed the L1014 alleles. Mosquito legs were used as templates for DNA. Mosquitoes were PCR amplified and sequenced individually. Analysis of the *vgsc* was completed according to Singh *et al.* (2011). PCR amplification of a portion the acetylcholinesterase gene (*ace-1*), associated with resistance to organophosphates and carbamates (Weill *et al.*, 2003), was also performed for *An. stephensi* according to the protocol detailed in (Dabire *et al.*, 2009). All PCR reactions were performed at 25ul total with 2X Promega Hot Start Master Mix (Promega, Madison, W) and the primer conditions listed in Table 1. PCR products were run on 2% agarose gel for 1 hour at 100V to confirm successful PCR amplification.

Table 1. List of primer and conditions used for PCR amplification of portions of the voltage gated sodium channel and acetylcholinesterase genes.

Assay	Primer	Sequence	Annealing Temperature (°C)	Final Concentration (μM)
Voltage gated sodium channel				
	KdrF	GGACCAYGATTTGCCAAGAT	50	1.25
	VGS_1R	CGAAATTGGACAAAAGCAAGG	50	1.25
Acetylcholinesterase				
	Ex3AGdir	GATCGTGGACACCGTGTTTCG	56	1
	Ex3AGrev	AGGATGGCCCGCTGGAACAG	56	1

Analysis of vgsc and ace-1 sequences

Sequences were submitted as queries to the National Center for Biotechnology Information's (NCBI) Basic Local Alignment Search Tool (BLAST) to confirm correct loci were amplified. Sequences were then aligned to identify kdr L1014 and ace-1R G1109 mutations. The allele and genotype frequencies of these mutations were then calculated.

Results

An. stephensi insecticide resistance

Bioassay results

A total of 750 *An. stephensi* were tested with different insecticides based on WHO protocol. The results of the susceptibility status of populations of *An. stephensi* are presented in Table 2. Overall, the percent mortality after exposure to insecticides ranged from 21% (Bendiocarb) to 67% (Deltamethrin). Using the WHO mortality threshold of above 98%, *An. stephensi* demonstrated resistance to propoxur, DDT, malathion and permethrin.

Table 2. Percentage of mortality of *An. stephensi* in different insecticide in Kebridehar town

Insecticides discriminating concentration (%)	Classification	No. of <i>An. stephensi</i> tested	Mortality rate after 24hrs	Resistance? (<98%)
Bendiocarb 0.1%	Carbamates	100	23%	Yes
Propoxure 0.1%	Carbamates	100	21%	Yes
Deltamethrin 0.05%	Pyrethroid	100	67%	Yes
Permethrin 0.75%	Pyrethroid	100	53%	Yes
Malathion 5%	Organophosphates	100	32%	Yes
DDT 4%	Organochlorine	100	32%	Yes
Pirimiphos-methyl 0.25%	Organophosphates	100	14%	No

An. stephensi insecticide resistance mutations

Portions of the vgsc and aceylcholinesterase genes containing the kdr and ace-1R mutations respectively were analyzed. Amplification produced 185 bp fragment. Of the 129 *An. stephensi* specimens from Kebridehar examined for KDR mutations, none carried the L1014 mutation. Of the 30 *An. stephensi* analyzed, no ace-1R mutations were observed.

Discussion

This is the first report of *An. stephensi* in Ethiopia exhibiting insecticide resistance. What is most concerning is that *An. stephensi* showed resistance to seven insecticides included in this study highlighting a potential challenge with insecticide based vector control in this region. There is some consistency with previous studies on *An. stephensi* resistance to insecticides. As in the present study, *An. stephensi* was shown to be resistant to DDT in Iran (Gorouhi *et al.*, 2016). Similarly, a study on *An. stephensi* in Afghanistan revealed resistance to deltamethrin, malathion, permethrin and DDT (Safi *et al.* 2019). In addition, *An. stephensi* carbamate resistance was observed in a recent study in Iran (Abbasi *et al.* 2019) as observed in the present study. However, there were some difference between our findings on *An. stephensi* and previous reports, where *An. stephensi* was found to be susceptible to the pyrethroids (deltamethrin and permethrin) and malathion in Iran (Gorouhi *et al.*, 2016, Zareet *et al.*, 2016) and Pakistan (Ali Khan *et al.* 2018). These differences may reflect differences in the type and extent of insecticide use in Ethiopia compared to other countries.

One surprising finding from our study was the absence of kdr mutation with phenotypic evidence of pyrethroid resistance. The absence of kdr mutations in pyrethroid resistant *Anopheles* is rare but not unprecedented. A study conducted on *An. stephensi* collected in Afghanistan revealed a low frequency of L1014 wild-type mutation (44%) in mosquitoes and a lack of homozygotes of the mosquitoes that were resistant to deltamethrin (Ahmad *et al.*,

2016). The phenotypic presentation of resistance in the majority of *An. stephensi* specimens with the absence or low frequency of *kdr* mutations may suggest that metabolic resistance as opposed to targeted resistance is the primary resistance mechanisms in the Ethiopian *An. stephensi*. A follow-up study on cytochrome P450s, esterases, glutathione S-transferases (GSTs) and acetylcholine esterase (AChE) activities in pyrethroid resistant mosquitoes in Afghanistan further highlight the role of metabolic resistance in *An. stephensi* (Safi et al. 2017). It is also possible the other variants in the *vgsc* gene may lead to resistance. Additional sequencing and analysis of the entire *vgsc* should be conducted for identification of other mutations that could lead insecticide resistance.

There are some similarities between these findings on *An. stephensi* resistance and what has been reported in *An. arabiensis* in Ethiopia. In *An. arabiensis*, resistance has been reported for insecticides belonging to all four chemical classes approved for IRS and LLINs. These include DDT (organochlorine), malathion (organophosphate), bendiocarb and propoxur (carbamates) and alpha-cypermethrin, cyfluthrin, deltamethrin, etofenprox, lambda-cyhalothrin and permethrin (pyrethroids) (Balkew *et al.*, 2010; Yewhalaw *et al.*, 2011; Massebo *et al.*, 2013; Asale *et al.*, 2014; Messenger *et al.*, 2017). However, the frequency of KDR mutations are much higher in the Ethiopian *An. arabiensis* (Balkew *et al.*, 2010; Messenger *et al.*, 2017) than what is reported here. These findings suggest that while, insecticides may induce the development of resistance over time, mechanisms for resistance may vary across species. In depth cross-species genetic analysis for selective signatures on the *vgsc* and unidentified loci across the genome are needed and underway to further elucidate the differing mechanisms for insecticide resistance in *Anopheles* species.

In the present study, no ACE-1 mutations were observed. There is still some ambiguity around the significance of the ACE-1 mutation has been proposed to induce resistance to organophosphates and carbamates resistance (Weill *et al.*, 2003). The mutation was absent in the *An. stephensi* tested in our study this may reflect the history of the type of insecticides used in the region.

The finding of multi-insecticide resistance in *An. stephensi* in Kebri Dehar Ethiopia emphasizes the need for additional investigation in other parts of Ethiopia and the Horn of Africa. Additionally, only a portion of the available insecticides were tested in this study. Resistance investigations should include some of the newer compound classes including lambda-cyhalothrin, temephos and chlorpyrifos. The complete absence of the mutation is unexpected but may be due to the *An. stephensi* collection being limited to one site. Future studies should evaluate the frequency of this mutation in other regions of Ethiopia.

Declarations

Authors' contributions

SY: conceptualizing the study, data analysis, and manuscript preparation, reared mosquitoes, tested bioassay; AG: reared mosquitoes, tested bioassay and manuscript preparation; TC: helped with the molecular analysis, design of the molecular analysis, writing the manuscript LD, VB, KL: helped with the molecular analysis DJ: provide resources and facilitated the molecular analysis. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests. Consent for publication Not applicable.

Funding

This study was financially supported by Jigjiga University. This project was partially funded by the University of North Carolina at Charlotte Multicultural Postdoctoral Fellowship.

Acknowledgements

Our gratitude goes to Jigjiga University for the financial support. It is our great pleasure to thank Mr. Geleta Bekele for his technical support in rearing mosquito at field laboratory.

Ethics approval and consent to participate:

Not applicable

Consent for publication: Not applicable.

References

- Abbott, W.S. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* 1925, 18, 265-267.
- Ahmad M, Buhler C, Pignatelli P, Ranson H, Nahzat S M, Naseem M, Sabawoon M F, Siddiqi A M, Vink M. Status of insecticide resistance in high-risk malaria provinces in Afghanistan. *Malar J.* 2016, 15:98.
- Alemayehu, E., Asale, A., Eba, K., Getahun, K., Tushune, K., Bryon, A., Morou, E., Vontas, J. Van Leeuwen, T., Duchateau, L., and Yewhalaw, D. Mapping insecticide resistance and characterization of resistance mechanisms in *Anopheles arabiensis* (Diptera: Culicidae) in Ethiopia. *Parasite Vector* 2017, 10:407
- Ali Khan H A, Akram W, Lee S. Resistance to Selected Pyrethroid Insecticides in the Malaria Mosquito, *Anopheles stephensi* (Diptera: Culicidae), From Punjab, Pakistan. *J Med Entomol.* 2018, 55:735-738.
- Asale A, Getachew Y, Hailesilassie W, Speybroeck N, Duchateau L, Yewhalaw D. Evaluation of the efficacy of DDT indoor residual spraying and longlasting insecticidal nets against insecticide resistant populations of *Anopheles arabiensis* Patton (Diptera: Culicidae) from Ethiopia using experimental huts. *Parasit Vector.* 2014, 7:131.
- Balkew, M., Muntaser I, Koekemoer LL, Brooke BD, Engers H, Aseffa A, et al. (2010). Insecticide resistance in *Anopheles arabiensis* (Diptera: Culicidae) from villages in central, northern and south west Ethiopia and detection of kdr mutation. *Parasit Vectors.* 3:40.
- Bloomquist J R. Ion channels as target for insecticides. *Annu Rev Entomol.* 1996, 41:163–90.
- Dabiré K R, Diabaté A, Namontougou M, Djogbenou L, Kengne P, Simard F, Bass C, Baldet T. Distribution of insensitive acetylcholinesterase (ace-1R) in *Anopheles gambiae* s.l populations from Burkina Faso (West Africa). *Trop. Med. Int. Health* 2009, 14:396-403.
- David J P, Ismail H M, Chandor-Proust A, Paine M J. Role of cytochrome P450s in insecticide resistance: impact on the control of mosquito-borne diseases and use of insecticides on Earth. *Philos T Roy Soc B.* 2013, 368(1612): 20120429
- Fettene M, Olana D, Christian R N, Koekemoer L L, Coetzee M. Insecticide resistance in *Anopheles arabiensis* from Ethiopia. *Afr Entomol.* 2013, 21:89–94.

FMoH. National malaria guidelines. Addis Ababa: Federal Democratic Republic of Ethiopia Ministry of Health; 2012.

Ghebreyesus, A., Deressa, W., Witten, K., Getachew, A., and Seboxa, T. Malaria. In: Berhane, Y., Haile Marim, D. and Kloos, H (Eds). Epidemiology and Ecology of Health and Disease in Ethiopia. Shama Books, Addis Ababa, 2006 pp 556-576.

Gillies, M.T., Coetzee, M. A Supplement to the Anophelinae of Africa South of the Sahara (Afrotropical Region). Publ S Afr Inst Med Res. 1987, 55:1-143

Glick, J.I. Illustrated Key to the Female Anopheles of Southwestern Asia and Egypt (Diptera: Culicidae). Defense Technical Information Center. Walter Reed Bio systemsatics Unit, 1992

Gorouhi MA, Vatandoost H, Oshaghi MA, Raeisi A, Enayati AA, Mirhendi H, Hanafi-Bojd AA, Abai MR, Salim-Abadi Y, Rafi F. Current Susceptibility Status of *Anopheles stephensi* (Diptera: Culicidae) to Different Imagicides in a Malarious Area, Southeastern of Iran. J Arthropod Borne Dis. 2016, 10:493-500.

Hemingway J, Ranson H. Insecticide resistance in insect vectors of human disease. Annu Rev Entomol. 2000, 45:371–91.

Kenea, O., Balkew, M., Tekie, H. et al. Impact of combining indoor residual spraying and long-lasting insecticidal nets on *Anopheles arabiensis* in Ethiopia: results from a cluster randomized controlled trial. Malar J 18, 182 (2019) doi:10.1186/s12936-019-2811-1

Liu, N. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research Directions. Ann. Rev. of Entomol. 2015, 60: 537-559

Massebo F, Balkew M, Gebre-Michael T, Lindtjorn B. Blood meal origins and susceptibility of *An. arabiensis* from Chano South-West Ethiopia. Parasite Vector. 2013, 6: 44.

Messenger, Josephat Shililu, Seth R. Irish, Gedeon Yohannes Anshebo, Alemayehu Getachew Tesfaye, Yemane Ye-Ebiyo, Sheleme Chibsa, Dereje Dengela, Gunawardena Dissanayake, Estifanos Kebede, Endalew Zemene, Abebe Asale, Mekonnen Yohannes, Hiwot Solomon Taffese, Kristen George, Christen Fornadel, Aklilu Seyoum, Robert A. Wirtz and Delenasaw Yewhalaw. Insecticide resistance in *Anopheles arabiensis* from Ethiopia (2012–2016): a nationwide study for insecticide resistance monitoring. Malar J. 2017, 16:469

Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet. 2012, 380:2197–223.

Ranson H, Jensen B, Vulule JM, Wang X, Hemingway J, Collins FH. Identification of point mutation in the voltage-gated sodium channel gene of Kenyan *Anopheles gambiae* associated with resistance to DDT and pyrethroids. Insect Mol Biol. 2000, 9:491–7.

Reid M C, McKenzie F E. The contribution of agricultural insecticide use to increasing insecticide resistance in African malaria vectors. Malar J. 2016, 15:107.

- Rivero A, Vézilier J, Weill M, Read AF, Gandon S. Insecticide control of vector-borne diseases: when is insecticide resistance a problem? *PLoS Pathog.* 2010, 6(8):e1001000.
- Safi N H Z, Ahmadi A A, Nahzat S, Warusavithana S, Safi N, Valadan R, Shemshadian A, Sharifi M, Enayati A, Hemingway J. Status of insecticide resistance and its biochemical and molecular mechanisms in *Anopheles stephensi* (Diptera: Culicidae) from Afghanistan. *Malar J.* 2019, 18(1):249. doi: 10.1186/s12936-019-2884-x.
- Safi NH, Ahmadi AA, Nahzat S, Ziapour SP, Nikookar SH, Fazeli-Dinan M, Enayati A, Hemingway J. Evidence of metabolic mechanisms playing a role in multiple insecticides resistance in *Anopheles stephensi* populations from Afghanistan. *Malar J.* 2017,16:100.
- Seyfarth M, Khaireh BA, Abdi AA, Bouh SM, Faulde MK. Five years following first detection of *Anopheles stephensi* (Diptera: Culicidae) in Djibouti, Horn of Africa: populations established-malaria emerging. *Parasitol Res.* 2019, 118:725-732.
- Taffese H S, Elizabeth Hemming-Schroeder, Cristian Koepfli, Gezahegn Tesfaye, Ming-chieh Lee, James Kazura, Gui-Yun Yan and Guo-Fa Zhou. Malaria epidemiology and interventions in Ethiopia from 2001 to 2016. *Infect Dis Poverty* 2018, 7:103
- Vatandoost H, Oshaghi MA, Abaie MR, Shahi M, Yaaghoobi F, Baghaii M, Hanafi-Bojd AA, Zamani G, Townson H. Bionomics of *Anopheles stephensi* Liston in the malarious area of Hormozgan province, southern Iran, 2002. *Acta Tropica.* 2006, 97:196-203.
- Weill M, Lutfalla G, Mogensen K, Chandre F, Berthomieu A, Berticat C, et al. Comparative genomics: insecticide resistance in mosquito vectors. *Nature* 2003, 423:136–7.
- WHO. Global plan for insecticide resistance management in malaria vectors. Geneva, Switzerland. 2012
- WHO. Indoor residual spraying: an operational manual for indoor residual spraying (IRS) for malaria transmission control and elimination – 2nd ed. 2015
- WHO. Test procedures for insecticide resistance monitoring in malaria vector mosquitoes.
- Geneva: World Health Organization; 2013 (<http://www.who.int/malaria/publications/atoz/9789241505154/en/>, accessed 12 September 2016).
- WHO. World Malaria Report, World Health Organization, Geneva, Switzerland. Geneva: WHO. 2017
- WHO. World Malaria Report, World Health Organization, Geneva,Switzerland. Geneva: WHO. 2018
- Yewhalaw D, Van Bortel W, Denis L, Coosemans M, Duchateau L,Speybroeck N. First evidence of high knockdown resistance frequency in *Anopheles arabiensis* (Diptera: Culicidae) from Ethiopia. *Am J Trop Med Hyg.*2010, 83:122–5
- Yewhalaw D, Wassie F, Steurbaut W, Spanoghe P, Van Bortel W, Denis L, etal. Multiple insecticide resistance: an impediment to insecticide-basedmalaria vector control program. *PLoS One.* 2011, 6(1):e16066.

Figures

Figure 1

Rearing larvae to adult at field laboratory