

Radiographic Analysis in Reduction Loss After Distal Radius Fracture Fixation with Variable Angle Volar Locking Plate

Pin-Chieh Fang

School of Traditional Chinese Medicine, College of Medicine, Chang Gung University

Tak-Yu-Yubie Lo

School of Traditional Chinese Medicine, College of Medicine, Chang Gung University

Chun-Ying Cheng

Bone and Joint Research Center, Department of Orthopaedic Surgery, Chang Memorial Hospital-Linkou, Chang Gung University College of Medicine

Ying-Chao Chou

Bone and Joint Research Center, Department of Orthopaedic Surgery, Chang Gung Memorial Hospital-Linkou; Chang Gung University College of Medicine

Alvin Chao-Yu Chen (✉ alvinchen@cgmh.org.tw)

Bone and Joint Research Center, Department of Orthopaedic Surgery, Chang Gung Memorial Hospital-Linkou, and Chang Gung University College of Medicine <https://orcid.org/0000-0003-0111-490X>

Research article

Keywords: distal radius fracture, variable angle volar locking plate, radiographic parameter

Posted Date: October 8th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-86534/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Reduction loss is commonly seen even in the newly designed locking plate fixation for distal radius fractures. Our study purpose is to investigate the efficacy of the variable angle volar locking plate (VAVLP) in maintenance of fracture fixation.

Methods A total of 37 patients of unilateral distal radius fractures receiving VAVLP fixation were enrolled. Forearm radiographs immediately after surgery and those at 3 months were retrospectively reviewed by orthopedist, radiologist and medical students for analysis of radiographic parameter including radial height (RH), ulnar variance (UV), radial inclination (RI), volar tilt (VT), tear drop angle (TDA), distal dorsal cortical distance (DDD) and Soong classification (SC).

Results By comparing the 3-month measurement and normal data, the difference of RH / UV / TDA was significant (p-value of 0.000) while the difference of RI / VT was insignificant. However, there was no significant difference regarding those five parameters between postoperative and 3-month measurement. Linear regression on DDD exhibited positive dependence with p-value of 0.002 between postoperative and 3-month changes; predictability after modification was 21.3%. Postoperative SC was grade 0 in 13 patients, grade 1 in 21, and grade 2, in 3. There were 7 of Gr 0 and 2 of Gr 1 became one grade up.

Conclusion VAVLP fixation in distal radius fracture can maintain radiographic alignment without significant reduction loss for at least 3 months. Realignment within normal range was in RI and VT, but not in RH / UV / TDA.

Background

Distal radial fracture is a very common injury, which has shown a bimodal distribution of age-specific incidence with an increasing prevalence in adults and aging people [1–3]. Non-surgical treatment is generally adopted for non- or minimally displaced fractures. For displaced distal radial fractures, operative management with internal fixation using volar locking plate (VLP) is preferred over percutaneous fixation in recent years [4].

VLP can be divided into two types, one is of locking screws and pegs in fixed angles; the other is of variable angle locking design. While a fixed-angle VLP fixation can improve the stability by providing a rigid construct with screws inserted and locked in a predetermined direction [5, 6], the variable-angle volar locking plate (VAVLP) allows a greater adaptability for screw angle insertion to support the dorsal subchondral bone for weak and comminuted fractures [7]. However, significantly decreased ultimate failure moment noted at 15°-inclined screw insertion indicated potential reduction loss and recommend titanium VAVLP be used with precaution [8]. Given that reduction loss remains a critical issue after treatment of distal radius fractures [9] even in using newly designed locking plates [10], radiographic parameters anecdotally used in fracture evaluation and functional correlation [11, 12] have been currently adapted for assessment of secondary collapse after fracture fixation [13]. This study aims to evaluate the

efficacy of VAVLP fixation in the maintenance of realignment through a retrospective review and comparison of postoperative radiographs in distal radius fractures.

Methods

To evaluate the stability of VAVLP fixation of radial fractures, posteroanterior (PA) and lateral views of forearm x-ray of immediately and 3 months after operation were evaluated [14, 15]. The inclusion criteria were: (1) distal radial fracture diagnosed by forearm radiographs (including 5 Smith's type, 18 Colle's type, and 19 Barton's type), and (2) patients with VAVLP fixation (Synthes, 2.4 mm Variable Angle LCP). A total of 43 cases which underwent surgical management for distal radial fractures between October 2011 and October 2019 in our hospital were being reviewed. The exclusion criteria were: (1) patient with incomplete medical records and follow-up (5 patients), and poor quality of radiographs (1 patient). Finally, 37 patients were included in this study. Postoperative x-ray images were taken at a mean of 1 day after operation; radiographs of 3 months following the operation were taken at a mean of 84 days after operation. All and the diagnostic imaging studies were assessed by a radiologist, an orthopedist and two medical students in internship training.

Radiographic parameters and analysis

To standardize our measurement, radiological parameters were measured on standard x-rays of the wrist according to the methods introduced by Kreder et al [16]. For both PA view and lateral view, the center of radius was measured at 3 and 5 cm below the mid-region of the proximal lunate articular surface in order to form the central axis of the radius. Radial height (RH), ulnar variance (UV) and radial inclination (RI) were measured on PA view taken with both shoulder abduction and elbow flexion in 90°, forearm pronation and supination in 0° [17]. Volar tilt (VT), tear drop angle (TDA) [18], distal dorsal cortical distance (DDD) [19] and Soong classification (SC) [20] were measured on lateral view x-rays taken with shoulder abduction, elbow flexion and forearm supination, all in 90°. RH represents the distance between the two parallel lines perpendicular to the long axis of the radius where one line passing through the distal articular surface and the other one intersects the distal articular surface of the ulnar head. UV refers to the distance between the levels of radial and ulnar articular surfaces, one line passing through the distal ulnar articular surface parallel to the other one passing through the medial radial articular surface perpendicular to the central axis of the radius. RI on AP projection represents the angle between a line connecting the radial styloid tip and ulnar aspect of the distal radial articular surface and a line perpendicular to the central axis of the radius. VT refers to the angle between a line perpendicular to the central axis of the radius and a line along the distal radial articular surface at the joint margin. TDA is the angle between a line passing through the central axis of the teardrop and a line perpendicular to the central axis. DDD refers to the distance between the tip of the most distal screw and the dorsal rim of distal radius. SC represents a tangential line drawn to the volar rim, which is parallel to the diaphyseal bone of the radial shaft in order to determine the plate prominence. Plates that do not extend the line were graded as Grade 0; Plates that extend volar to the line but remain proximal to the rim were graded as

Grade 1; Plates that are distal or directly on the volar rim were graded as Grade 2. The above parameters were measured on the digital images with accuracies up to 0.01 mm and 0.1° (Fig. 1).

According to the central limit theorem, when the sample number is greater than 30 (N = 37), it tends to be normal distribution. Thus, a 2-sample paired T-test was used for comparing the postoperative radiographs with those of 3 months following operation. A 1-sample T-test was used for comparing the parameters of radiographs of 3 months following operation with the normal values quoted in past studies, in order to determine if the normal anatomy was restored. Level of significance was set at less than 0.05. Linear regression was applied for comparing the DDD immediately after operation with the change in value of 3 months following operations. Maximum likelihood ratio Chi-square test was used for comparing SC immediately after operation with those of 3 months following operation. With reference to different literatures, the average value of parameters were as follows: RH 11.6 mm; UV 1.5 mm (negative ulnar variance), RI 22.5°; VT 11.2; TDA 68°; value of DDD immediately after the operation is in correlation with the variance of DDD over the 3 months after the surgery [20]. All statistical analyses were performed with SPSS v21 (IBM Corporation, Armonk, NY).

Results

Comparison with normal values

The result of comparison between radiographic parameters of 3 months following the operation with normal values, revealed significant differences in RH, UV, TDA when compared to their own normal values, with $p = 0.000$, $p = 0.000$ and $p = 0.000$ respectively. However, there is no significant difference for RI and VT compared to the normal values, with $p = 0.184$ and $p = 0.127$ respectively (Table 1).

Table 1
Comparison of 3 months following operative radiographic parameters with normal values

		Normal values	Mean	SD	Range	<i>p-value</i>
Radial height	(mm)	11.6	9.57	3.19	3.16 to 15.8	0.000
Ulnar variance	(mm)	1.50	1.67	2.21	-3.90 to 6.06	0.000
Radial inclination	(degree)	22.5	21.35	5.17	7.10 to 33.40	0.184
Volar tilt	(degree)	11.2	12.93	12.93	-4.90 to 24.20	0.127
Tear drop angle	(degree)	25.4	54.79	9.14	35.00 to 69.30	0.000

Abbreviation: SD, standard deviation.

Comparison with values of immediately after operation

The result of comparison between radiographic parameters of immediately after operation with 3 months following the operation, showed no statistical difference was noted when comparing the above values to

the values of 3 months following the surgery, with $p = 0.207$, $p = 0.267$, $p = 0.049$, $p = 0.368$ and $p = 0.276$ respectively (Table 2).

Table 2
Comparison of 3 months following operative radiographic parameters with radiographic parameters immediately after the operation

		A mean	B mean	SD	Range	<i>p-value</i>
Radial height	(mm)	9.57	9.33	3.17	3.26 to 16.58	0.207
Ulnar variance	(mm)	1.67	1.41	2.05	-4.05 to 5.10	0.267
Radial inclination	(degree)	21.35	20.37	4.92	6.00 to 31.40	0.049
Volar tilt	(degree)	12.93	13.69	5.53	0.80 to 22.70	0.368
Tear drop angle	(degree)	54.79	56.46	8.76	25.20 to 70.20	0.276
A = Radiographic parameters at 3-month follow-up						
B = Radiographic parameters immediately after the operation						
Abbreviation: SD, standard deviation						

Comparing the DDD immediately after operation with the change in value of 3 months following operations

By comparing the DDD right after the operation with the change in values of DDD of 3 months following the operation DDD by linear regression, results showed significant dependence with $p = 0.002$, and predictability after modification is 21.3% (Fig. 2).

Comparing the grading of SC immediately after the operation with SC of 3 months following the operation

Regarding SC, 13 cases were graded as Grade 0 after operation; 6 cases remained as Grade 0 (46.2%) and the other 7 cases (53.8%) were progressed to Grade 1 in radiographs of 3-month postoperatively. The remaining 21 cases were graded as Grade 1 after operation; 19 (90.5%) of them remained as Grade 1, and the other 2 (9.5%) cases were progressed to Grade 2 in radiographs of 3 months following the operation. Three cases were recorded as Grade 2 after operation, and all 3 cases remained as Grade 2 (100%) in radiographs of 3 months following the operation.

To compare the grading of SC immediately after operation and 3 months following operation, maximum likelihood ratio Chi square was used and revealed significant dependence with $p = 0.000$ (Table 3).

Table 3

Comparing the grading of SC immediately after the operation with SC of 3 months following the operation by maximum likelihood ratio Chi-square test

		SC of 3 months following operation				
		Grade 0	Grade 1	Grade 2	Total	p-value
	Grade 0	7	6	0	13	
SC immediately after operation	Grade 1	0	19	2	21	0.000
	Grade 2	0	0	3	3	
Total		7	28	5	37	
Abbreviation: SC, Soong classification						

Discussion

This study revealed that when using VAVLP for fixation in distal radius fractures, RH, UV, TDA could not be restored to its normal values, while RI and VT were successfully restored to their normal values. The above results were noted in cases using fixed angle VLP as well [14, 15]. The potential causes behind might be due to the heterogeneity in individuals, fracture patterns and the length of treatment [21]. Another study by Pienaar et al in 2013 also stated that no matter which kind of reduction and fixation was used, TDA cannot be restored to its normal value [22].

By comparing the values of parameters immediately after operation with those of 3 months following operation, results showed that RH, UV, RI, VT, TDA were able to sustain at least a 3-month period in radioulnar variance.

In analyzing the DDD change in our cases, linear regression showed a positive dependence between the values of postoperative and 3-month radiographs. Optimization of screw/plate position by measuring DDD was first proposed in 2015 and recommended a limitation of 6 mm in maximum during surgery to avoid subsequent displacement [19]. It was even more critically concerned for elderly patients with osteoporotic bone with suggestion of DDD within 4.6 mm according to another publication in 2018 [23]. In our cases, positive dependence of postoperative DDD was noted on the DDD change at 3 months and the mean of postoperative DDD was 5.23+/- 1.70. Maintenance of realignment up to 3 months was achieved in most cases with intraoperative DDD within 6 mm. The design of variable-angled locking holes in VAVLP allows screw insertion in greater adaptability to reach subchondral bone and minimize DDD for providing superior maintenance of fracture fixation.

The SC was originally proposed to evaluate volar locking implant prominence and risks in flexion tendon rupture. A recent clinical report analyzed 400 cases receiving two different kinds of locking plates and stated that the group using variable angle LCP had a greater SC grading [23]. In analyzing the 37 patients

of our study, we found there was a trend that patients with postoperative SC grade 1 and 2 showed less subsequent displacement than those with grade 0; the difference was statistically significant ($p = 0.000$).

There are several limitations in our study. This is a radiographic analysis based on retrospective case review without control group. The sample size is small with heterogeneity in fracture pattern. Only radiographs immediately and 3 months after surgery were evaluated. In addition, no clinical correlation is documented.

Conclusions

VAVLP can maintain radiographic realignment up to 3 months after surgery. No significant reduction loss is found. Radiographic parameters are restored to normal range in RI and VT, but not in RH / UV / TDA. Variable- angled locking holes provide greater adaptability to minimize DDD facilitating maintenance of fracture fixation.

List Of Abbreviations

VAVLP variable angle volar locking plate

RH radial height

UV ulnar variance

RI radial inclination

VT volar tilt

TDA tear drop angle

DDD distal dorsal cortical distance

SC Soong classification

VLP volar locking plate

PA posteroanterior

Declarations

Ethics Approval:

Institutional review board approval (IRB no. 202000939B0) was obtained to perform a review of patient records and radiographs.

Consent for publication:

Not Applicable.

Availability of supporting data:

The datasets generated during the current study are available from the corresponding author on reasonable request.

Competing interests:

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article. The authors report no competing interests.

Funding:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions:

In this study, Pin-Chieh Fang and Tak-Yu-Yubie Lo contributed to data collection and draft writing; Yi-Hsuan Lin and Chih-Hao Chiu, contributed to outcome assessment and formal analysis; Chun-Ying Cheng, assisted in manuscript review and editing; Ying-Chao Chou, contributed to data analysis and study supervision. Alvin Chao-Yu Chen is the single surgeon for all cases and responsible for study design. All authors have read and approved the final manuscript.

Acknowledgements:

Not applicable

Authors' information:

Pin-Chieh Fang^a; Tak-Yu-Yubie Lo^a; Chun-Ying Cheng^b; Ying-Chao Chou^b; Alvin Chao-Yu Chen^b

^aSchool of Traditional Chinese Medicine, College of Medicine, Chang Gung University, Taoyuan, Taiwan

^bBone and Joint Research Center, Department of Orthopaedic Surgery, Chang Gung Memorial Hospital-Linkou and Chang Gung University College of Medicine,

References

1. Jerrhag D, Englund M, Karlsson MK, Rosengren BE. Epidemiology and time trends of distal forearm fractures in adults – a study of 11.2 million person-years in Sweden. *BMC Musculoskelet Disord* 2017;18:240. doi:10.1186/s12891-017-1596-z
2. Hevonkorpi TP, Launonen AP, Huttunen TT, Kannus P, Niemi S, Mattila VM. Incidence of distal radius fracture surgery in Finns aged 50 years or more between 1998 and 2016. *BMC Musculoskelet Disord* 2016;19:70. doi:10.1186/s12891-018-1983-0
3. Azad A, Kang HP, Alluri RK, Vakhshori V, Kay HF, Ghiassi A. Epidemiological and treatment trends of distal radius fractures across multiple age groups. *J Wrist Surg* 2019;8: 305–311. doi:10.5555/s-0039-1685205
4. Mellstrand-Navarro C, Pettersson HJ, Tornqvist H, Ponzer S. The operative treatment of fractures of the distal radius is increasing: results from a nationwide Swedish study. *Bone Joint J* 2014;96:963–969. doi:10.1302/0301-620X.96B7.33149
5. Thelen S, Grassmann JP, Jungbluth P, Windolf J. Distal radius fractures: current treatment concepts and controversies. *Chirurg* 2018;89:798–812. doi:10.1007/s00104-018-0724-0
6. Quadlbauer S, Pezzeri C, Jurkowitsch J, Rosenauer R, Pichler A, Schättin S, Hausner T, Leixnering M. Early complications and radiological outcome after distal radius fractures stabilized by volar angular stable locking plate. *Arch Orthop Trauma Surg* 2018;138:1773–1782. doi:10.1007/s00402-018-3051-5
7. Park JH, Hagopian J, Ilyas AM. Variable-angle locking screw volar plating of distal radius fractures. *Hand Clin* 2010;26:373–380, vi. doi:10.1016/j.hcl.2010.04.003
8. Lenz M, Wahl D, Gueorguiev B, Jupiter JB, Perren SM. Concept of variable angle locking—evolution and mechanical evaluation of a recent technology. *J Orthop Res* 2015;33:988–992. doi:10.1002/jor.22851
9. Vosbikian MM, Ketonis C, Huang R, Costanzo JA, Ilyas AM. Reduction loss after distal radius fracture fixation with locked volar plates. *J Surg Orthop Adv* 2017;26:160–165. PMID: 29130877
10. Lee SJ, Park JW, Kang BJ, Lee JI. Clinical and radiologic factors affecting functional outcomes after volar locking plate fixation of dorsal angulated distal radius fractures. *J Orthop Sci* 2016;21:619-624. Doi:10.1016/j.jos.2016.05.007
11. Dario P, Matteo G, Carolina C, Marco G, Cristina D, Daniele F, Andrea F. Is it really necessary to restore radial anatomic parameters after distal radius fractures? *Injury* 2014;45 Suppl 6:S21–S26. doi:10.1016/j.injury.2014.10.018
12. Ng CY, McQueen MM. What are the radiological predictors of functional outcome following fractures of the distal radius? *J Bone Joint Surg Br* 2011;93:145–150. doi:10.1302/0301-620X.93B2.25631

13. Walenkamp M, Mulders M, van Hilst J, Goslings JC, Schep N Prediction of distal radius fracture redisplacement: a validation study. *J Orthop Trauma* 2018;32:e92–e96. doi:10.1097/BOT.0000000000001105
14. Osada D, Kamei S, Masuzaki K, Takai M, Kameda M, Tamai K. Prospective study of distal radius fractures treated with a volar locking plate system. *J Hand Surg Am* 2008;33:691–700. doi:10.1016/j.jhsa.2008.01.024
15. Rozental TD, Blazar PE. Functional outcome and complications after volar plating for dorsally displaced, unstable fractures of the distal radius. *J Hand Surg Am* 2006; 31:359–365. doi:10.1016/j.jhsa.2005.10.010
16. Kreder HJ, Hanel DP, McKee M, Jupiter J, McGillivray G, Swiontkowski MF. X-ray film measurements for healed distal radius fractures. *J Hand Surg Am* 1996;21:31–39. doi:10.1016/S0363-5023(96)80151-1
17. Thuysbaert G, Ringburg A, Petronilia S, Vanden Berghe A, Hollevoet N. Measurement of ulnar variance and radial inclination on X-rays of healed distal radius fractures. With the axis of the distal radius or ulna? *Acta Orthop Belg* 2015;81:308–314. PMID: 26280972
18. Fujitani R, Omokawa S, Iida A, Santo S, Tanaka Y. Reliability and clinical importance of teardrop angle measurement in intra-articular distal radius fracture. *J Hand Surg Am* 2012;37:454–459. doi: 1016/j.jhsa.2011.10.056
19. Vosbikian MM, Ketonis C, Huang R, Ilyas AM. Optimal positioning for volar plate fixation of a distal radius fracture: determining the distal dorsal cortical distance. *Orthop Clin North Am* 2016;47:235–244. doi: 1016/j.ocl.2015.08.020
20. Soong M, Earp BE, Bishop G, Leung A, Blazar P. Volar locking plate implant prominence and flexor tendon rupture. *J Bone Joint Surg Am* 2011;93:328–335. doi:2106/JBJS.J.00193
21. Schuurman AH, Maas M, Dijkstra PF, Kauer JM. Assessment of ulnar variance: a radiological investigation in a Dutch population. *Skeletal Radiol* 2001;30:633–638. doi:1007/s002560100414
22. Pienaar G, Anley C, Ikram A. Restoration of teardrop angle (TDA) in distal radius fractures treated with volar locking plates. *SA Orthop J* 2013;12:32-34.
23. Bergsma M, Brown K, Doornberg J, Sierevelt I, Jaarsma R, Jadav B. Distal radius volar plate design and volar prominence to the watershed line in clinical practice: comparison of Soong grading of 2 common plates in 400 patients. *J Hand Surg Am* 2019;44:853–859. doi:1016/j.jhsa.2019.04.012

Figures




Figure 1

Radiographs of posteroanterior (A) and lateral view (B and C) at 3 months following operation of distal radius fracture with variable-angle volar locking plate RH= Radial height, UV= Ulnar variance, RI= Radial inclination, VT= Volar tilt, TDA= Tear drop angle, DDD= Distal dorsal cortical distance, SCA= Soong classification auxiliary-line




Figure 2

Linear regression demonstration of DDD DDD= distal dorsal cortical distance