Paenibacillus Sinensis sp. nov., a Nitrogen-fixing Species Isolated from Plant Rhizospheres

Qin Li

China Agricultural University, Beijing

Yashi Li

China Agricultural University, Beijing

Xiaomeng Liu

China Agricultural University, Beijing

San-Feng Chen (✉ chensf@cau.edu.cn)

China Agricultural University, Beijing

Research Article

Keywords: Genomic characterisations , Nitrogen-fixing bacteria , *Paenibacillus sinensis* sp. nov. , Rhizosphere of plant

Posted Date: September 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-867873/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at Antonie van Leeuwenhoek on October 31st, 2021. See the published version at <https://doi.org/10.1007/s10482-021-01677-6>.

Abstract

Two strains HN-1^T and 39 were isolated from rhizospheres of different plants grown in different regions of PR China. The two strains exhibited high nitrogenase activities and possessed almost identical 16S rRNA gene sequences. The average nucleotide identity (ANI) and digital DNA–DNA hybridization (dDDH) values between the two strains were 99.9 and 93.8% respectively, suggesting that they belong to one species. Phylogenetic analysis based on the 16S rRNA gene sequence showed that strains HN-1^T and 39 are the members of the genus *Paenibacillus* and both strains exhibited 99.5% similarity to *Paenibacillus stellifer* DSM 14472^T and the both strains represented a separate lineage from all other *Paenibacillus* species. However, the ANI of type strain HN-1^T with *P. stellifer* DSM 14472^T was 90.69, which was below the recommended threshold value (< 95–96% ANI). The dDDH showed 42.1% relatedness between strain HN-1^T and *P. stellifer* DSM 14472^T, which was lower than the recommended threshold value (dDDH < 70%). The strain HN-1^T contain anteiso-C_{15:0} as major fatty acids and MK-7 as predominant isoprenoid quinone. The major polar lipids were diphosphatidylglycerol, phosphatidylglycerol, phosphatidylethanolamine, four aminophospholipids and an unidentified glycolipid. Unlike the most closely related *P. stellifer* DSM 14472^T, strain HN-1^T or 39 was positive for catalase reaction. Distinct phenotypic and genomic characterisations from previously described taxa support the classification of strains HN-1^T or 39 as representatives of a novel species of the genus *Paenibacillus*, for which the name *Paenibacillus sinensis* is proposed, with type strains HN-1^T (= CGMCC 1.18902, JCM 34620), and reference strain 39 (= CGMCC 1.18879, JCM 34616), respectively.

Introduction

The genus *Paenibacillus* was proposed by Ash et al. (1993) and its description was emended by Shida *et al.* (Shida et al. 1997). Some species of the genus *Bacillus* were transferred to the genus *Paenibacillus* (Ash et al. 1993; Heyndrickx et al. 1996; Shida et al. 1997; Lee et al. 2004; Hu et al. 2010), and further descriptions of novel members increased the number of species of the genus *Paenibacillus*. At the time of writing, the genus comprises 256 species and four subspecies with validly published names (www.bacterio.net/paenibacillus.html). Members of the genus *Paenibacillus* are rod-shaped, aerobic or facultatively anaerobic, spore-forming bacteria with anteiso-C_{15:0} as the major cellular fatty acid and menaquinone 7 (MK-7) as the major menaquinone, and their DNA G + C contents range from 45 to 54 mol% (Ash et al. 1993; Priest 2009).

Species of the genus *Paenibacillus* have been isolated from diverse environment habitats (<https://lpsn.dsmz.de/genus/paenibacillus>) and have diverse physiological characteristics. Presently, there are only a few species with nitrogen fixation ability in these bacteria, including *Paenibacillus borealis* (Elo et al. 2001), *Paenibacillus brasilensis* (von der Weid et al. 2002), *Paenibacillus graminis* (da Mota et al. 2004), *Paenibacillus massiliensis* (Ding et al. 2005), *Paenibacillus zanthoxyli* (Ma et al. 2007a), *Paenibacillus sabinae* (Ma et al. 2007b), *Paenibacillus forsythiae* (Ma and Chen 2008), *Paenibacillus donghaensis* (Choi et al. 2008), *Paenibacillus sonchi* (Hong et al. 2009), *Paenibacillus*

riograndensis (Beneduzi et al. 2010), *Paenibacillus jilunlii* (Jin et al. 2011a), *Paenibacillus sophorae* (Jin et al. 2011b), *Paenibacillus stellifer* (Jin et al. 2011c), *Paenibacillus tritici* (Wang et al. 2013a), *Paenibacillus polymyxa* (Wang et al. 2013b), *Paenibacillus beijingensis* (Gao et al. 2012), *Paenibacillus taohuashanense* (Xie et al. 2012), *Paenibacillus brassicae* (Gao et al. 2013), *Paenibacillus bryophyllum* (Liu et al. 2018), *Paenibacillus helianthi* (Ambrosini et al. 2018), *Paenibacillus maysiensis* (Wang et al. 2018), *Paenibacillus rhizophilus* (Ripa et al. 2019), *Paenibacillus durus* (Guella et al. 2019). Most of N₂-fixing *Paenibacillus* species isolated from plant roots have been shown to play an important role in promoting plant growth by nitrogen fixation, phosphate solubilization, production of plant phytohormones and various enzymes (Xie et al. 2016; Grady et al. 2016; Li et al. 2019). Comparative genomic analysis revealed the conservation of *nif* cluster comprising 9 genes (*nifB nifH nifD nifK nifE nifN nifX hesA nifV*) in nitrogen-fixing *Paenibacillus* strains (Xie et al. 2014). In addition, some N₂-fixing *Paenibacillus* strains with additional *nif* and *nif*-like genes exhibited higher nitrogenase activities (Li et al. 2014; Li et al. 2021). Such traits make the study of the regulation of the multiple *nif* genes of N₂-fixing *Paenibacillus* under different environmental conditions and their adaptation to varying ecological niches interesting.

In this study, two nitrogen-fixing strains HN-1^T and 39 isolated from the rhizosphere of plant were characterized by a polyphasic taxonomic approach, one more presumably novel species strain belonging to the genus *Paenibacillus*.

Materials And Methods

Isolation and culture conditions

Strain 39 was isolated from a soil sample collected from arbor rhizosphere in Beijing of China (39°57'N, 116°17'E). 1 g soil sample was suspended in 9-mL sterile water, stirred for 30 min and heated at 80°C for 15 min. After that, 100 µl suspension was spread on nitrogen-free medium agar plates in triplicate. After incubation at 30°C for 3 days, single colonies were isolated by streaking plating. The nitrogen-free medium consisted 20 g sucrose, 0.1 g K₂HPO₄, 0.4 g KH₂PO₄, 0.2 g MgSO₄·7H₂O, 0.1 g NaCl, 0.01 g FeCl₃, and 0.002 g Na₂MoO₄ per liter water. The strain HN-1^T was previously isolated from the rhizosphere of rice and on nitrogen-free medium agar plates (Liu et al. 2019). Strains were routinely cultured in LD medium (per liter contains 2.5 g NaCl, 5 g yeast, and 10 g tryptone) at 30°C for further identification and study. The type strain of the genus *Paenibacillus*, *P. polymyxa* DSM 36^T, *P. sabinae* DSM 17841^T, *P. stellifer* DSM 14472^T, *P. zanthoxyli* JH29^T, *P. graminis* RSA19, *P. tritici* BJ-18^T and *P. durus* ATCC 35681^T were obtained from our bacterial collection. Bacteria were freeze-dried or frozen using a sterile glycerol solution in cryogenic tubes to preserve the samples (30% v/v), and stored at – 80°C and – 20°C.

NitrogenFixing Capability

To determine nitrogenase activity, strains HN-1^T and 39 and reference *Paenibacillus* strains, including *P. polymyxa* DSM 36^T, *P. sabinae* DSM 17841^T, *P. stellifer* DSM 14472^T, *P. zanthoxyli* JH29^T, *P. graminis* RSA19, *P. tritici*solis BJ-18^T and *P. durus* ATCC 35681^T were grown in 20 mL of LB broth medium in 50 mL flasks shaken overnight at 30°C. The cultures were collected by centrifugation, precipitations were washed three times with sterilized water and then resuspended in nitrogen-limited medium (per liter contains 10.4 g Na₂HPO₄, 3.4 g KH₂PO₄, 26 mg CaCl₂·2H₂O, 30 mg MgSO₄, 0.3 mg MnSO₄, 36 mg Ferric citrate, 7.6 mg Na₂MoO₄·2H₂O, 10 µg p-aminobenzoic acid, 5 µg biotin, 0.3 g glutamate and 4 g glucose). The nitrogenase activity was determined using the acetylene reduction assay and expressed as nmol C₂H₄ mg⁻¹ protein h⁻¹ as described previously (Wang et al. 2013b).

Genomic data analysis

Total DNA was extracted from strains HN-1^T and 39, evaluated by gel electrophoresis, and estimated using a NanoDrop 2000 (Thermo Scientific, MA, USA). The draft genome sequence was produced by using Illumina paired-end sequencing technology at the mega genomics. Assembly was conducted using SOAP de-novo v. 1.04 assembler (Li et al. 2008). Gene prediction was made using Glimmer v. 3.0 (Delcher et al. 2007). Annotation of protein-coding sequence was performed by using the Basic Local Alignment Search Tool (BLAST) against the COG, Kyoto Encyclopedia of Genes and Genomes (KEGG) databases and NCBI nr protein database.

Average nucleotide identity (ANI) were calculated in EZBioCloud [<https://www.ezbiocloud.net/tools/ani>] (Yoon et al. 2017a), using the algorithm published by Lee et al. (2016). Digital DNA–DNA hybridization (dDDH) values were computed at GGDC (Genome-to-Genome Distance Calculator) using GGDC 2.0 BLAST + and recommended formula 2 (Meier-Kolthoff et al. 2013).

Phylogenetic analysis

The 16S rRNA gene sequences of 39 were acquired from a PCR product using highly specific forward primer 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and universal reverse primer 1492R (5'-GGTTACCTTGTACGACTT-3'). The DNA sequence obtained was compared to reference 16S rRNA gene sequences available in GenBank and EMBL databases obtained from the NCBI Genome Database using BLASTN software (Altschul et al. 1990) and the EzBioCloud server (<https://www.ezbiocould.net>) (Yoon et al. 2017b). The sequences of the *gyrB* gene were obtained from the genome of HN-1^T and 39 and other type strains. Multiple sequence alignments were analysed using CLUSTAL X (Thompson et al. 1997). The phylogenetic tree calculating evolutionary distance matrices was constructed by the neighbor-joining method (Saitou and Nei, 1987) using MEGA (version 7.0) (Kumar et al. 2016). Bootstrap analysis was conducted on 1000 replications (Felsenstein, 1985).

Phenotypic Characterization

Cell morphology was obtained by scanning electrical microscopy (SEM), after incubated on endospore-forming medium agar plate [yeast extract 0.07%, tryptone 0.1%, glucose 0.1%, $(\text{NH}_4)_2\text{SO}_4$, 0.02%, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.02%, K_2HPO_4 , 0.1% (w/v), pH 7.2] for 72 h. The flagellation type was determined by transmission electron microscopy (TEM) after 48 h incubation of strain HN-1 on LD medium. Cell motility was evaluated in semi-solid (0.3% agar) LD medium after incubation at 30°C for 24 h. Physiological and biochemical characteristics were determined in comparison with *P. sabinae* DSM 17841^T and *Paenibacillus stellifer* DSM 14472^T. Most physiological and biochemical tests, including activities of catalase and oxidase, nitrate reduction, hydrolysis of starch, aesculin and tween 20, production of dextrin and indole, methy red reaction, Voges-Proskauer reaction, lysozyme test and production of acid from fermentation of different substrates were performed according to Zhao et al. (2014). Temperature range for growth were determined after incubation at 4, 10, 15, 25, 28, 30, 37, 40 and 45°C on LD agar. The pH range for growth was determined in LD broth adjusted to pH 4.0–10.0 (using increments of 1.0 pH unit) by using HCl and NaOH buffers. Growth in the absence of NaCl and in the presence of 0, 0.2, 0.5, 1.0, 2.0, 3.0 and 4.0 % (w/v) NaCl was investigated by using LD broth. A spectroscopic method of monitoring turbidity at OD₆₀₀ was used to assess the growth at various temperature, pH values and NaCl concentration. The ability of strains to assimilate different substrates were tested using Biolog GEN III MicroPlate system (Biolog Microstation™, CA, USA) following the manufacturer's instructions.

Chemotaxonomy

Strains were incubated in LD medium at 30°C for 2 days. The compositions of cellular fatty acid were analyzed according to the method described by Komagata and Suzuki (1987) using Sherlock Identification System (MIDI) (Sasser et al. 2005). Cellular menaquinones and respiratory quinones were extracted, purified, and analyzed by HPLC according to the method described by Collins (1980). Polar lipid was extracted by the method of Minnikin et al. (1979), and was identified by two-dimensional TLC as described by Collins et al. (1980).

Result And Discussion

NitrogenFixing Capability

Since bacteria in the soil sample were cultured in nitrogen-free medium on the purpose of isolating nitrogen-fixing strain, strain 39 is possible to have nitrogen-fixing capability. Strains HN-1^T isolated from the rhizosphere of rice was detected by acetylene reduction to have nitrogen-fixing capacity (Liu et al. 2019). To confirm genes encoding nitrogenase are contained in their genome, *nifH* gene was amplified and sequenced. However, the sequencing result of *nifH* fragment both displayed double peaks, which indicated that there were multiple *nifH* genes in their genome. Acetylene reduction assays were performed

to verify the nitrogenase activity of HN-1^T and 39. As shown in Table 1, strains HN-1^T and 39 exhibited very high nitrogenase activity compared to other nitrogen-fixing *Paenibacillus* species, suggesting a high efficiency of the nitrogen fixation process.

Table 1
Nitrogenase activity of strains HN-1^T and 39 in comparison
with some nitrogen-fixing species of the genus
Paenibacillus

Strain	Nitrogenase activity [nmol C ₂ H ₄ (mg protein h) ⁻¹]
<i>P. polymyxa</i> DSM 36 ^T	1355.1 ± 152.4
<i>P. stellifer</i> DSM 14472 ^T	6099.5 ± 497.3
<i>P. zanthoxyli</i> JH29 ^T	6282.4 ± 307.7
<i>P. graminis</i> RSA19 ^T	4272.9 ± 207.9
<i>P. sabinae</i> DSM 17841 ^T	7749 ± 371.8
<i>P. durus</i> ATCC 35681 ^T	5511.5 ± 260.2
<i>P. triticisoli</i> BJ-18 ^T	743.6 ± 82.9
39	7160.3 ± 584.6
HN-1 ^T	6937.2 ± 625.1

Molecular Characterization

The newly generated 16S rRNA gene sequence of strain 39 were compared with sequences of bacterial type strains in the Genbank nucleotide database. Phylogenetic trees were inferred using the neighbour-joining and maximum-parsimony methods in the software MEGA7. As no significant differences were found among the phylogenetic trees obtained by the different methods used, only the trees constructed by using the neighbour-joining method are shown. Phylogenetic analysis based on 16S rRNA gene sequences revealed that strains HN-1^T and 39 clustered with species of the genus *Paenibacillus* and formed a monophyletic cluster with *P. stellifer* DSM 14472^T, as the three strains formed a separate phylogenetic branch within the genus *Paenibacillus* with a high bootstrap resampling value of 100 % (Fig. 1). The strain 39 shares 100% 16S rRNA gene sequence identity with strain HN-1^T. The strains HN-1^T or

39 showed the highest 16S rRNA gene sequence identity to *P. stellifer* DSM 14472^T (99.5%), followed by *P. durus* ATCC 35681^T (97.4%) and *P. sabinae* DSM 17841^T (97.0%).

Generally, 98.7% sequence identity on the 16S rRNA gene are considered to be within the same species (Kim et al. 2014). However, several reports have been published showing that *Paenibacillus* species with >99 % 16S rRNA gene sequence similarity may not belong to the same species (Kamfer et al. 2017; Kim and Cha 2018; Ghio et al. 2019; Guella et al. 2019; Velazquez et al. 2020). Thus, housekeeping genes are now routinely used to complement the 16S rRNA gene analysis for species level determination (da Mota et al. 2004; Holmes et al. 2004; Rodriguez et al. 2019). Due to the low level of discrimination based on 16S rRNA gene between closely related species, the *gyrB* gene (coding for the b subunit of DNA gyrase) was used as an alternative phylogenetic marker (Wang et al. 2007). The *gyrB* genes were retrieved from the HN-1^T and 39 genomes. The *gyrB* gene clearly distinguishes HN-1^T and 39 from other *Paenibacillus* species with only 93.04% gene sequence identity to *P. stellifer* DSM 14472 (Fig. S1). Based on the 95–96% *gyrB* gene sequence similarity as the interspecies gap (Lee et al. 2008; Liu et al. 2013), strains HN-1^T and 39 could be assigned to novel species.

Phenotypic Characteristics

Strains HN-1^T and 39 were found to be Gram-positive, facultatively anaerobic, motile and rod-shaped. Colonies of strain HN-1^T and 39 grown on LD medium after 72 h of incubation at 30°C were usually 0.8–1.2 mm in diameter, circular, moist, milky and convex. The transmission electron micrographs of type strain HN-1^T showed the presence of peritrichous flagella on cell surface (Fig. 2a). Strain HN-1^T produced ellipsoidal spores in swollen sporangia in the terminal region of the cell by scanning electron microscope (Fig. 2b).

In order to determine physiological and biochemical characteristics of HN-1^T and 39 in comparison with *P. stellifer* DSM 14472^T and *P. sabinae* DSM 17841^T, a series of tests were carried out following the proposed minimal standards for describing new taxa of facultatively anaerobic, endospore-forming bacteria (Logan et al. 2009). The ability of strains to assimilate different substrates were tested using GEN III microplates by Biolog system (Biolog Microstation TM, CA, USA) (Kiran et al. 2017; Ripa et al. 2019).

The strains HN-1^T and 39 grew well in up to 4% NaCl (w/v), however, strain *P. stellifer* DSM 14472^T tolerated only 3% NaCl. The pH range for growth was 5.0–9.0 and the temperature range for growth is 15–42°C. Strains HN-1^T and 39 was determined to be negative for the Voges–Proskauer reaction, and positive for the methyl red reaction. Strains HN-1^T and 39 were positive for catalase reaction and can produce acid from rhamnose and sorbitol, which differentiated HN-1^T and 39 from the most related *P. stellifer* DSM 14472^T. The physiological and biochemical characteristics of HN-1^T and 39 are shown in Table 2. In Biolog analysis, significant differences were observed in the metabolization of different

substrates on Biolog GEN III microplate by strains HN-1^T and 39 from its closest type strain *P. stellifer* DSM 14472^T (Table 3). Strain HN-1^T and *P. stellifer* DSM 14472^T differed in the metabolization of D-Fucose, D-Maltose, 3-Methyl glucose, D-Sorbitol, Stachyose, Citric acid, α-Keto-butyric acid, Mucic acid, Methyl pyruvate, Gelatin, Inosine, D-Glucose-6-PO₄, Pectin, Aztreonam, Fusidic acid, Nalidixic acid, Vancomycin, Lithium chloride, Sodium bromate, Sodium lactate 1%, Rifamycin sv and Troleandomycin as a sole carbon source. Compared with type strains of closely related *Paenibacillus* species, strain HN-1^T and 39 exhibited nearly identical phenotypic characteristics.

Table 2

Differential phenotypic characteristics between strain HN-1^T, 39 and closely related strains.

Characteristic	HN-1 ^T	39	DSM14472 ^T	DSM 17841 ^T
Growth conditions				
Temperature (°C)	15–42	15–42	15–42	15–42
pH	5–9	5–9	5–9	5–9
NaCl	0–4%	0–4%	0–3%	0–3%
0.001% lysozyme	-	-	-	-
Oxidase	-	-	-	-
Catalase	+	+	-	+
Voges-Proskauer	-	-	-	+
Methyl red reaction	+	+	+	+
Nitrate reduction	-	-	-	+
Hydrolysis of:				
Starch	+	+	+	-
Tween 20	-	-	-	-
Tyrosine	-	-	-	-
Esculin	+	+	+	+
Production of:	-	-	-	-
Indole				
Dihydroxyacetone	-	-	-	-
Dextrin	+	+	+	-
Production of acid from:				
Glucose	+	+	+	+
Fructose	+	+	+	+
Rhamnose	+	+	-	-
Glycerol	-	-	-	-
Trehalose	+	+	+	+

Strains: 1. HN-1^T; 2. 39; 3. *P. stellifer* DSM 14472^T; 4, *P. sabinae* DSM 17841^T; +. Positive reaction; -, negative reaction.

Characteristic	HN-1 ^T	39	DSM14472 ^T	DSM 17841 ^T
Arabinose	+	+	+	-
Mannitol	-	-	-	+
Xylose	+	+	+	-
Sorbitol	+	+	-	+
Maltose	+	+	+	+
Starch	+	+	+	+
Strains: 1. HN-1 ^T ; 2. 39; 3. <i>P. stellifer</i> DSM 14472 ^T ; 4, <i>P. sabinae</i> DSM 17841 ^T ; +. Positive reaction; 2. negative reaction.				

Table 3

Phenotypic fingerprinting of 1. strain HN-1^T; 2. strain 39; 3. *P. stellifer* DSM 14472^T; 4. *P. sabinae* DSM 17841^T by oxidation of sole substrates on Biolog GEN III MicroPlates (All data from present study generated under the same conditions).

substrate	1	2	3	4		1	2	3	4
Carbohydrates:					Esters:				
n-Acetyl-β-D-mannosamine	-	-	-	-	D-Lactic acid methyl ester	-	-	-	-
n-Acetyl-D-galactosamine	-	-	-	-	Methyl pyruvate	w	w	-	+
n-Acetyl-D-glucosamine	-	-	-	-	Alcohols:				
n-Acetyl neuraminic acid	-	-	-	-	Glycerol	-	-	-	-
D-Arabitol	-	-	-	-	Amides:				
α-D-Glucose	+	+	+	+	Glucuronamide	w	w	w	-
α-D-Lactose	+	+	+	-	Amino acids:				
D-Cellobiose	+	+	+	+	L-Alanine	-	-	-	-
D-Fructose	+	+	+	+	L-Arginine	-	-	-	-
D-Fucose	-	-	w	-	L-Aspartic acid	-	-	-	-
D-Galactose	+	+	+	+	D-Aspartic acid	-	-	-	-
D-Gentiobiose	+	+	+	+	Gelatin	w	w	-	-
β-Methyl-D-glucoside	+	+	+	+	Glycyl-L-proline	-	-	-	-
D-Maltose	+	+	+	+	L-Glutamic acid	-	-	-	-
D-Mannitol	-	-	-	+	L-Histidine	-	-	-	-
D-Mannose	+	+	w	+	L-Pyroglutamic acid	-	-	-	-
D-Melibiose	w	w	+	+	L-Serine	-	-	-	-
3-Methyl glucose	+	+	w	+	D-Serine	-	-	-	-
D-Raffinose	+	+	+	+	Nucleosides				
D-Salicin	+	+	+	+	Inosine	w	w	-	-
D-Sorbitol	+	+	-	+	Phosphorylated compounds:				
D-Trehalose	+	+	+	+	D-Fructose-6-PO ₄	w	w	w	-
D-Turanose	+	+	+	+	D-Glucose-6-PO ₄	w	w	-	-

+, Positive; -, negative; w, weak reaction

substrate	1	2	3	4		1	2	3	4
L-Fucose	w	w	w	-	Polymers:				
L-Rhamnose	w	w	w	-	Dextrin	+	+	+	+
Stachyose	+	+	w	+	Pectin	+	+	w	+
Sucrose	+	+	+	+	Tween 40	-	-	-	-
Carboxylic acid:					Antibiotics:				
Acetic acid	-	-	-	-	Aztreonam	+	+	w	+
Acetoacetic acid	w	w	w	+	Fusidic acid	w	w	-	+
γ -Amino-butyric acid	-	-	-	-	Lincomycin	-	-	-	+
Bromo-succinic acid	-	-	-	-	Minocycline	-	-	-	+
Citric acid	w	w	-	-	Nalidixic acid	+	+	-	+
Formic acid	-	-	-	-	Rifamycin sv	-	-	+	+
D-Galacturonic acid	w	w	w	-	Troleandomycin	-	-	-	+
D-Gluconic acid	-	-	-	+	Vancomycin	+	+	w	+
D-Glucuronic acid	w	w	w	-	pH 5	w	w	w	w
L-Galactonic acid lactone	w	w	w	-	pH 6	+	+	+	+
α -Hydroxy-butyric acid	-	-	-	-	NaCl 4%	+	+	-	w
β -Hydroxy-D, L butyric acid	-	-	-	-	NaCl 8%	-	-	-	-
<i>p</i> -Hydroxy-phenylacetic acid	-	-	-	-	Guanidine HCl	-	-	-	-
α -Keto-butyric acid	w	w	-	w	Lithium chloride	+	+	w	+
α -Keto-glutaric acid	w	w	w	-	Myo-inositol	-	-	-	-
L-Lactic acid	-	-	-	-	Niaproof 4	-	-	-	-
D-Malic acid	-	-	-	-	Potassium tellurite	+	+	+	+
L-Malic acid	-	-	-	-	Sodium bromate	+	+	-	w
Mucic acid	w	w	-	-	Sodium butyrate	+	+	+	+
Propionic acid	-	-	-	-	Sodium lactate 1%	+	+	w	+
Quinic acid	-	-	-	-	Tetrazolium blue	-	-	w	w
D-Saccharic acid	-	-	-	-	Tetrazolium violet	w	w	+	w
+, Positive; -, negative; w, weak reaction									

Chemotaxonomic Characteristics

In order to determine the composition of cellular fatty acid, HN-1^T and type strains *P. stellifer* DSM 14472^T and *P. sabinae* DSM 17841^T were incubated in LD medium at 30°C for 2 days. Whole cell fatty acid analysis revealed that anteiso-C_{15:0}, C_{14:0}, C_{16:0}, iso-C_{14:0}, iso-C_{16:0} and iso-C_{15:0} are present as major (> 5 %) fatty acids, and anteiso-C_{15:0}, anteiso-C_{17:0}, iso-C_{17:0} and C_{18:1ω9c} are present as minor (< 5 but > 1 %) fatty acids (Table S1). Anteiso-C_{15:0} is the predominant fatty acid of members of the genus *Paenibacillus* (Ash et al. 1993), consistent with strain HN-1^T (41.14%) being a member of this genus. However, in the closely related type strains *P. stellifer* DSM 14472^T, the fatty acid C_{16:0} was found to be more abundant than anteiso-C_{15:0}. The major menaquinone of HN-1^T was MK-7, in conformity to genus *Paenibacillus*. The polar lipids detected by two-dimensional TLC are diphosphatidylglycerol (DPG), phosphatidylethanolamine (PE), phosphatidylglycerol (PG), four aminophospholipids (APL) and unidentified glycolipid (Fig. S1), in agreement with the profile of genus *Paenibacillus*.

Genome sequence and similarity analysis

Genome sequencing was performed to evaluate the genomic relatedness of the strains HN-1^T and 39 to its closely related recognized species in the genus *Paenibacillus*. Genomes of strains HN-1^T and 39 were approximately 6.32 and 6.45 Mbp, respectively. The DNA G + C content of the strains HN-1^T and 39 were 53.36 and 52.99 mol%, respectively, which is within the range 39–59 mol% reported for the members of genus *Paenibacillus* (Im et al. 2017). The total number of protein coding genes in HN-1^T and 39 were 5631 and 5782, respectively. While, the related strain *P. stellifer* DSM 14472^T had a complete genome of 5.65Mb, comprising 4958 protein coding genes with a DNA G + C content of 55.6 mol% (Genbank: CP009286). More detailed genomic information of strains HN-1^T and 39 were shown in Supplementary Table S2 and high-quality draft genomes of strains HN-1^T and 39 were deposited in GenBank under accession numbers JAHCMB000000000 and JAHBAZ000000000, respectively.

In this study, digital DNA–DNA hybridization (dDDH) values were calculated using GGDC. The dDDH value of genomes for strain HN-1^T and strain 39 was 99.8%. The level of DNA–DNA relatedness was determined to be 42.1% by strain HN-1^T with *P. stellifer* DSM 14472^T, which is lower than the threshold value of 70 % for species delineation (Chun et al. 2018).

The average nucleotide identity (ANI) is widely used to define bacterial species (Konstantinidis and Tiedje 2005; Varghese et al. 2015; Chun et al. 2018; Ciufo et al. 2018). The ANI between the whole-genome sequence of HN-1^T and that of the most closely related species *P. stellifer* DSM 14472^T was calculated on ezbiocloud (Yoon et al., 2017a). The ANI value of genomes for HN-1^T and 39 was 99.93%, but the ANI values of strain HN-1^T were less than 91% with the related species *P. stellifer* DSM 14472^T, *P. sabinae* DSM 17841^{TT} and *P. durus* 35681^T (Table 4), lower than the threshold of 95–96% for differentiating

bacterial species (Richter and Rossello-Mo'ra 2009), suggesting that the new isolate HN-1^T represents a distinctive species.

Table 4
Pairwise genome comparisons between the strains HN-1^T and 39, and their close relatives

Query Genome	Reference genome	ANI (%)	dDDH (%)
HN-1 ^T	39	99.93	99.8
HN-1 ^T	<i>P. stellifer</i> DSM 14472 ^T (CP009286)	90.69	42.1
HN-1 ^T	<i>P. sabinae</i> DSM 17841 ^T (CP004078)	76.96	22.0
HN-1 ^T	<i>P. durus</i> ATCC 35681 ^T (CP011114)	76.80	22.0
39	<i>P. stellifer</i> DSM 14472 ^T (CP009286)	90.83	42.2
39	<i>P. sabinae</i> DSM 17841 ^T (CP004078)	77.01	22.2
39	<i>P. durus</i> ATCC 35681 ^T (CP011114)	76.78	22.2

The nitrogen fixation genes of strains HN-1^T and 39 were extracted by using Prokka software from the genome sequences (Seemann 2014). The genome of strains HN-1^T and 39 contain a compact *nif* cluster comprising ten genes *nifB*, *nifH*, *nifD*, *nifK*, *nifE*, *nifN*, *nifX*, *orf1*, *hesA* and *nifV* encoding Mo-nitrogenase, which is unique features of the *Paenibacillus* nitrogen fixation system. In addition to the *nif* cluster, the two strains have *anfHDK* encoding Fe-nitrogenase and linked to additional copies of *nifBENX* genes, while the closely related species *P. stellifer* DSM 14472^T contains *anfHDK* preceding additional *nifV* gene. Beyond the *nif* and *anf* cluster, there are multiple *nifHDK*-like genes located at different sites in their genomes. The organization of *nif*, *anf* and *nif*-like genes in type strain HN-1^T and the closely related species *P. stellifer* DSM 14472^T was shown in Fig. S3. Previous studies showed that 3 *nifH* genes of *P. sabinae* DSM 17841^T are functional by complementing *K. oxytoca* $\Delta nifH$ mutant (Hong et al. 2012). Thus, the high nitrogenase activity exhibited by these strains may be due to their additional *nif* genes.

Paenibacillus durus ATCC 35681^T can fix nitrogen even in the presence of nitrate due to the absence of nitrate reductase (Seldin et al. 1984). Whole genome sequence analysis strains HN-1^T and 39 revealed that nitrate reductase gene cluster *narJHG* were not detected, which suggested these two strains can also fix nitrogen in the nitrate-enriched medium. The draft genome of strains HN-1^T and 39 harbor two sets of NAD(P)H-nitrite reductases (*nirBD*) which are involved in the reduction of nitrite to ammonium in both assimilatory and dissimilatory reduction processes. Additional searches for genes associated with nitric oxide (*nirS* or *nirK*) and nitrous oxide reduction (*norBC*) were performed, but these genes were not detected in their genomes. Therefore, strains HN-1^T and 39 may possess dissimilatory nitrate reduction to ammonium pathway, but lack denitrification pathway.

The phenotypic, phylogenetic and genomic data of strains HN-1^T and 39 showed that they are different from all other closely related species of genus *Paenibacillus*. Therefore, we conclude that strain HN-1^T or 39 should be recognised as a novel species of the genus *Paenibacillus*, for which the name *Paenibacillus sinensis* sp. nov. is proposed.

Description of *Paenibacillus sinensis* sp. nov.

Paenibacillus sinensis (sin. en'sis. L.gen. n. sinensis of China, where the type strain HN-1^T was isolated).

Cells are Gram-positive, facultative anaerobic, rod-shaped (0.4–0.5 µm×2.0–3.2 µm) and motile by means of peritrichous flagella. In slightly swollen sporangia, an ellipsoidal spore is formed and located in terminal position of cells. Colonies on LD medium are circular, convex, cream white, with diameter 1.0–2.0 mm. Nitrogen fixation positive and multiple *nifH* genes are present. The growth temperature is 15–42°C, optimal at 30°C. The growth pH range is 5.0–9.0, optimal at pH 7.0. NaCl concentration of 0–4% (w/v) is tolerable for growth, optimal at 0–0.2%. Positive tests for catalase, methyl red test, starch and aesculin hydrolysis, but negative for oxidase, Voges–Proskauer reaction, nitrate reduction. The various substrates are assimilated examined using Biolog GEN III microplates: dextrin, D-maltose, D-trehalose, D-cellobiose, D-gentiobiose, sucrose, D-turanose, stachyose, D-raffinose, α-D-lactose, D-melibiose, β-Methyl-D-glucoside, D-salicin, α-D-glucose, D-mannose, D-fructose, D-galactose, 3-methyl glucose, 1% sodium lactate, D-serine, D-sorbitol and pectin were utilized. Strains are resistant to inhibitory chemicals: aztreonam, nalidixic acid, vancomycin, lithium chloride, potassium tellurite, sodium bromate, sodium butyrate, sodium lactate 1% and sensitive to troleandomycin, lincomycin, guanidine HCl, niaproof 4, tetrazolium blue. The major menaquinone is MK-7. The predominant fatty acid is anteiso-C_{15:0}. The major polar lipids are DPG, PE, and PG. The DNA G + C contents for strains HN-1^T and 39 are 53.36 and 52.99 mol%, respectively.

The type strain, HN-1^T (= CGMCC 1.18902, JCM 34620), was isolated from the rhizosphere soil of rice in Hunan P. R. China. The GenBank (EMBL) accession number for the 16S rRNA gene sequence of strain HN-1^T is MF967304 and the GenBank accession number for the draft genome sequence is JAHCMB000000000.

Declarations

Funding

This work was supported by the the National Key Research and Development Program of China (No. 2019YFA0904700) and the National Natural Science Foundation of China (No. 32000048).

Conflict of interest

The authors declare no conflict of interest.

Availability of data and material

The GenBank accession numbers for 16S rRNA gene sequences of strains HN-1^T and 39 are MF967304 and MZ153121, respectively. The draft genome sequences of strains HN-1^T and 39 have been deposited at NCBI under the accession no. JAHCMB000000000 and JAHBAZ000000000.

Ethics approval

Not applicable.

References

1. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215; 403-410
2. Ambrosini A, Sant'Anna FH, Heinzmann J, Fernandes GD, Bach E et al (2018) *Paenibacillus helianthi* sp. nov., a nitrogen fixing species isolated from the rhizosphere of *Helianthus annuus* L. *Antonie Van Leeuwenhoek* 111(12):2463-2471
3. Ash C, Priest FG, Collins MD (1993) Molecular identification of rRNA group 3 bacilli (ash, farrow, wallbanks and collins) using a PCR probe test. Proposal for the creation of a new genus *Paenibacillus*. *Antonie Van Leeuwenhoek* 64:253–260
4. Beneduzi A, Costa PB, Parma M, Melo IS, Bodanese-Zanettini MH et al (2010) *Paenibacillus riograndensis* sp nov., a nitrogen-fixing species isolated from the rhizosphere of *Triticum aestivum*. *Int J Syst Evol Microbiol* 60:128-133
5. Choi JH, Im WT, Yoo JS, Lee SM, Moon DS et al (2008) *Paenibacillus donghaensis* sp nov., a xylan-degrading and nitrogen-fixing bacterium isolated from East Sea sediment. *J Microbiol Biotechnol* 18(2):189-193
6. Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, da Costa MS, Rooney AP, Yi H, Xu XW, De Meyer S et al (2018) Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 68: 461-466
7. Ciufo S, Kannan S, Sharma S, Badretdin A, Clark K, Turner S, Brover S, Schoch CL, Kimchi A, DiCuccio M (2018) Using average nucleotide identity to improve taxonomic assignments in prokaryotic genomes at the NCBI. *Int J Syst Evol Microbiol* 68:2386–2392
8. Collins MD, Goodfellow M, Minnikin DE (1980) Fatty acid, isoprenoid quinone and polar lipid composition in the classification of *Curtobacterium* and related taxa. *J Gen Microbiol* 118: 29-37
9. da Mota FF, Gomes EA, Paiva E, Rosado AS, Seldin L (2004) Use of *rpoB* gene analysis for identification of nitrogen-fixing *Paenibacillus* species as an alternative to the 16S rRNA gene. *Lett*

10. Delcher AL, Bratke KA, Powers EC, Salzberg SL (2007) Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23: 673-679
11. Ding Y, Wang J, Liu Y, Chen S (2005) Isolation and identification of nitrogen-fixing bacilli from plant rhizospheres in Beijing region. *J Appl Microbiol* 99(5):1271-1281
12. Elo S, Suominen I, Kampfer P, Juhanoja J, Salkinoja-Salonen M et al (2001) *Paenibacillus borealis* sp. nov., a nitrogen-fixing species isolated from spruce forest humus in Finland. *Int J Syst Evol Microbiol* 51:535-545
13. Felsenstein J (1985). Confidence -limits on phylogenies-an approach using the bootstrap. *Evolution* 39: 783-791
14. Gao M, Xie LQ, Wang YX, Chen J, Xu J et al (2012) *Paenibacillus beijingensis* sp nov., a novel nitrogen-fixing species isolated from jujube garden soil. *Antonie Van Leeuwenhoek* 102(4):689-694
15. Gao M, Yang H, Zhao J, Liu J, Sun YH et al (2013) *Paenibacillus brassicae* sp nov., isolated from cabbage rhizosphere in Beijing, China. *Antonie Van Leeuwenhoek* 103(3):647-653
16. Ghio S, Sauka DH, Ferrari AE, Piccini RE, Ontanon OM, Campos D (2019) *Paenibacillus xylanivorans* sp. nov., a xylan-degrading bacterium isolated from decaying forest soil. *Int J Syst Evol Microbiol* 69: 3818-3823
17. Grady EN, MacDonald J, Liu L, Richman A, Yuan ZC (2016) Current knowledge and perspectives of *Paenibacillus*: a review. *Microb Cell Fact* 15: 203
18. Guella F, Porto RZ, Sant'Anna FH, Ambrosini A, Passaglia LMP (2019) Genomic metrics analyses indicate that *Paenibacillus azotofixans* is not a later synonym of *Paenibacillus durus*. *Int J Syst Evol Microbiol* 69(9):2870-2876
19. Heyndrickx M, Vandemeulebroecke K, Scheldeman P, Kersters K, DeVos P, Logan NA, Aziz AM, Ali N, Berkeley RCW (1996) A polyphasic reassessment of the genus *Paenibacillus*, reclassification of *Bacillus lautus* (Nakamura 1984) as *Paenibacillus lautus* comb nov and of *Bacillus peoriae* (Montefusco et al 1993) as *Paenibacillus peoriae* comb nov, and emended descriptions of *P. lautus* and of *P. peoriae*. *Int J Syst Bacteriol* 46: 988-1003
20. Holmes DE, Nevin KP, Lovley DR (2004) Comparison of 16S rRNA, *nifD*, *recA*, *gyrB*, *rpoB* and *fusA* genes within the family *Geobacteraceae* fam. nov. *Int J Syst Evol Microbiol* 54: 1591-1599
21. Hong YY, Ma YC, Zhou YG, Gao F, Liu HC et al (2009) *Paenibacillus sonchi* sp nov., a nitrogen-fixing species isolated from the rhizosphere of *Sonchus oleraceus*. *Int J Syst Evol Microbiol* 59:2656-2661
22. Hong Y., Ma YC, Wu LX, Maki M, Qin WS, Chen SF (2012) Characterization and analysis of *nifH* genes from *Paenibacillus sabinae* T27. *Microbiol Res* 16: 596-601
23. Hu XF, Li SX, Wu JG, Wang JF, Fang QL, Chen JS (2010) Transfer of *Bacillus mucilaginosus* and *Bacillus edaphicus* to the genus *Paenibacillus* as *Paenibacillus mucilaginosus* comb. nov and *Paenibacillus edaphicus* comb. nov. *Int J Syst Evol Microbiol* 60: 8-14

24. Im WT, Yi KJ, Lee SS, Moon HI, Jeon CO, Kim DW, Kim SK (2017) *Paenibacillus konkukensis* sp nov., isolated from animal feed. Int J Syst Evol Microbiol 67: 2343-2348
25. Jin HJ, Zhou YG, Liu HC, Chen SF (2011a) *Paenibacillus jilunlii* sp nov., a nitrogen-fixing species isolated from the rhizosphere of Begonia semperflorens. Int J Syst Evol Microbiol 61:1350-1355
26. Jin HJ, Lv J, Chen SF (2011b). *Paenibacillus sophorae* sp. nov., a nitrogen-fixing species isolated from the rhizosphere of Sophora japonica. Int J Syst Evol Microbiol 61:767-771
27. Jin HJ, Tu R, Xu F, Chen SF (2011c) Identification of nitrogen-fixing *Paenibacillus* from different plant rhizospheres and a novel *nifH* gene detected in the *P. stellifer*. Microbiology 80(1):117-124
28. Kamfer P, Busse HJ, McInroy JA, Hu CH, Kloepper JW, Glaeser SP (2017) *Paenibacillus rhizoplanae* sp nov., isolated from the rhizosphere of Zea mays. Int J Syst Evol Microbiol 67: 1058-1063
29. Kim M, Oh HS, Park SC, Chun J (2014) Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. Int J Syst Evol Microbiol 64: 346-351
30. Kim YS, Cha CJ (2018) *Paenibacillus translucens* sp nov., isolated from tidal flat sediment. Int J Syst Evol Microbiol 68: 936-941
31. Kiran S, Swarnkar MK, Mayilraj S, Tewari R, Gulati A (2017) *Paenibacillus ihbetae* sp nov., a cold-adapted antimicrobial producing bacterium isolated from high altitude Suraj Tal Lake in the Indian trans-Himalayas. Syst Appl Microbiol 40: 430-439
32. Konstantinidis KT, Tiedje JM (2005) Genomic insights that advance the species definition for prokaryotes. Proc Natl Acad Sci USA 102:2567–2572
33. Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. Mol Biol Evol 33: 1870-1874
34. Lee F., Tien CJ, Tai CJ, Wang LT, Liu YC, Chern LL (2008) *Paenibacillus taichungensis* sp nov., from soil in Taiwan. Int J Syst Evol Microbiol 58: 2640-2645
35. Lee I, Kim YO, Park SC, Chun J (2016) OrthoANI: An improved algorithm and software for calculating average nucleotide identity. Int J Syst Evol Microbiol 66: 1100-1103
36. Lee JS, Pyun YR, Bae KS (2004) Transfer of *Bacillus ehimensis* and *Bacillus chitinolyticus* to the genus *Paenibacillus* with emended descriptions of *Paenibacillus ehimensis* comb. nov and *Paenibacillus chitinolyticus* comb. nov. Int J Syst Evol Microbiol 54: 929-933
37. Li Q, Zhang HW, Zhang LQ, Chen SF (2021) Functional analysis of multiple *nifB* genes of *Paenibacillus* strains in synthesis of Mo-, Fe- and V-nitrogenases. Microb Cell Fact 20(1):139-139
38. Li RQ, Li YR, Kristiansen K, Wang J (2008) SOAP: short oligonucleotide alignment program. Bioinformatics 24: 713-714
39. Liu XM, Li Q, Li YB, Guan GH, Chen SF (2019) *Paenibacillus* strains with nitrogen fixation and multiple beneficial properties for promoting plant growth. PeerJ 7: e744

40. Liu LH, Yuan T, Yang F, Liu ZW, Yang MY et al (2018) *Paenibacillus bryophyllum* sp. nov., a nitrogen-fixing species isolated from *Bryophyllum pinnatum*. *Antonie Van Leeuwenhoek* 111(12):2267-2273
41. Li XX, Deng ZP, Liu Z., Yan YL, Wang TS, Xie JB, Lin M, Cheng Q, Chen SF (2014). The genome of *Paenibacillus sabinae* T27 provides insight into evolution, organization and functional elucidation of *nif* and *nif*-like genes. *BMC Genomics* 15:723
42. Li YB, Li YL, Zhang HW, Wang MY, Chen SF (2019) Diazotrophic *Paenibacillus beijingensis* BJ-18 provides nitrogen for plant and promotes plant growth, nitrogen uptake and metabolism. *Front Microbiol* 10:1119.
43. Liu Y, Lai QL, Dong CM, Sun FQ, Wang LP, Li GY, Shao ZZ (2013) Phylogenetic diversity of the *Bacillus pumilus* group and the marine ecotype revealed by multilocus sequence analysis. *PLoS One* 8: e80097
44. Logan NA, Berge O, Bishop AH, Busse HJ, De Vos P, Fritze D, Heyndrickx M, Kampfer P, Rabinovitch L, Salkinoja-Salonen MS et al (2009) Proposed minimal standards for describing new taxa of aerobic, endospore-forming bacteria. *Int J Syst Evol Microbiol* 59: 2114-2121
45. Ma YC, Zhang J, Chen SF (2007a). *Paenibacillus zanthoxyli* sp nov, a novel nitrogen-fixing species isolated from the rhizosphere of *Zanthoxylum simulans*. *Int J Syst Evol Microbiol* 57:873-877
46. Ma YC, Xia ZQ, Liu XM, Chen SF (2007b) *Paenibacillus sabinae* sp nov., a nitrogen-fixing species isolated from the rhizosphere soils of shrubs. *Int J Syst Evol Microbiol* 57: 6-11
47. Ma YC, Chen SF (2008) *Paenibacillus forsythiae* sp nov., a nitrogen-fixing species isolated from rhizosphere soil of *Forsythia mira*. *Int J Syst Evol Microbiol* 58:319-323
48. Meier-Kolthoff JP, Auch AF, Klenk HP, Goker M (2013) Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14: 60
49. Minnikin DE, Collins MD, Goodfellow M (1979) Fatty acid and polar lipid composition in the classification of *Cellulomonas*, *Oerskovia* and related taxa. *J Appl Bacteriol* 47: 87-95
50. Priest FG (2009) Genus I, *Paenibacillus*. In: De Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, Schleifer KH, Whitman WB (eds) *The firmicutes*, *Bergey's manual of systematic bacteriology*, vol 2, 2nd edn. Springer, New York, pp 269–296
51. Richter M, Rossello-Mora R (2009) Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* 106: 19126-19131
52. Ripa FA, Tong S, Cao WD, Wang ET, Wang TY, Liu HC, Gao JL, Sun JG (2019) *Paenibacillus rhizophilus* sp. nov., a nitrogen-fixing bacterium isolated from the rhizosphere of wheat (*Triticum aestivum* L.). *Int J Syst Evol Microbiol* 69: 3689-3695
53. Rodriguez M, Reina JC, Bejar V, Llamas I (2019) *Paenibacillus lutrae* sp. nov., a chitinolytic species Isolated from a river otter in castril natural park, Granada, Spain. *Microorganisms* 7:637
54. Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4(4):406–425

55. Sasser M, Kunitsky C, Jackoway G, Ezzell JW, Teska JD, Harper B, Parker S, Barden D, Blair H, Breezee J et al (2005) Identification of *Bacillus anthracis* from culture using gas chromatographic analysis of fatty acid methyl esters. J AOAC Int 88: 178-181
56. Seemann T (2014) Prokka: rapid prokaryotic genome annotation. Bioinformatics 30: 2068-2069.
57. Seldin L, Vanelsas JD, Penido EGC (1984) *Bacillus azotofixans* sp. nov. a nitrogen-fixing species from Brazilian soils and grass Roots. Int J Sys Bacteriol 34(4):451-456. 10
58. Shida O, Takagi H, Kadowaki K, Nakamura LK, Komagata K (1997) Transfer of *Bacillus alginolyticus*, *Bacillus chondroitinus*, *Bacillus curdlanolyticus*, *Bacillus glucanolyticus*, *Bacillus kobensis*, and *Bacillus thiaminolyticus* to the genus *Paenibacillus* and emended description of the genus *Paenibacillus*. Int J of Syst Bacteriol 47: 289-298
59. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Res 25, 4876-4882
60. Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K et al (2015) Microbial species delineation using whole genome sequences. Nucleic Acids Res 43:6761–6771
61. Velazquez LF, Rajbanshi S, Guan SH, Hinchee M, Welsh A (2020) *Paenibacillus ottowii* sp. nov. isolated from a fermentation system processing bovine manure. Int J Syst Evol Microbiol, 70(3):1463-1469
62. von der Weid I, Duarte GF, van Elsas JD, Seldin L (2002) *Paenibacillus brasilensis* sp nov., a novel nitrogen-fixing species isolated from the maize rhizosphere in Brazil. Int J Syst Evol Microbiol 52:2147-2153
63. Wang LT, Lee FL, Tai CJ, Kasai H (2007) Comparison of *gyrB* gene sequences, 16S rRNA gene sequences and DNA-DNA hybridization in the *Bacillus subtilis* group. Int J Syst Evol Microbiol 57: 1846-1850
64. Wang LY, Li J, Li QX, Chen SF (2013a) *Paenibacillus beijingensis* sp nov., a nitrogen-fixing species isolated from wheat rhizosphere soil. Antonie Van Leeuwenhoek 104(5):675-683
65. Wang L., Zhang LH, Liu ZZ, Zhao DH, Liu X M, Zhang B, Xie JB, Hong YY, Li PF, Chen S.F et al. (2013b). A minimal nitrogen fixation gene cluster from *Paenibacillus* sp. WLY78 enables expression of active nitrogenase in *Escherichia coli*. PLoS Genet 9: e1003865
66. Wang TS, Xie JY, Wang LY, Chen SF (2018) *Paenibacillus maysiensis* sp nov., a nitrogen-fixing species isolated from the rhizosphere soil of maize. Curr Microbiol 75(10):1267-1273
67. Xie JB, Zhang LH, Zhou YG, Liu HC, Chen SF (2012) *Paenibacillus taohuashanense* sp nov., a nitrogen-fixing species isolated from rhizosphere soil of the root of *Caragana kansuensis* Pojark. Antonie Van Leeuwenhoek 102(4):735-741
68. Xie JB, Du ZL, Bai LQ, Tian CF, Zhang YZ, Xie JY, Wang TS, Liu XM, Chen X, Cheng Q et al (2014) Comparative genomic analysis of N₂-fixing and non-N₂-fixing *Paenibacillus* spp.: organization, evolution and expression of the nitrogen fixation genes. PLoS Genet 10: e1004231

69. Xie JB, Shi HW, Du ZL, Wang TS, Liu XM et al. Comparative genomic and functional analysis reveal conservation of plant growth promoting traits in *Paenibacillus polymyxa* and its closely related species. Sci Rep 2016; 6:21329.
70. Yoon SH, Ha SM, Lim J, Kwon S, Chun J (2017a) A large-scale evaluation of algorithms to calculate average nucleotide identity. Antonie Van Leeuwenhoek 110: 1281-1286
71. Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J (2017b) Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. Int J Syst Evol Microbiol 67: 1613-1617
72. Zhao B, Lin H, He SJ (2014) Microbiology experiment, 2nd edn. Science Press, Beijing

Figures

Figure 1

Neighbor-joining phylogenetic tree based on 16Sr RNA gene sequences showing the position of strains 39 and HN-1T among species of the genus *Paenibacillus*. Bootstrap analyses were performed with 1000 cycles. Numbers (50%) at nodes are bootstrap values. Bar 0.01 substitutions per nucleotide positions.

Figure 2

Morphology of flagella and endospores, (a) Transmission electron microscopic image of flagella of strain HN-1T and (b) scanning electron microscopic image of vegetative cells and spores of strain HN-1T.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Fig.S1.tif](#)
- [Fig.S2.tif](#)
- [Fig.S3.tif](#)
- [TableS1.docx](#)
- [TableS2.docx](#)