Potential Roles for Pattern Molecule of PAMP-triggered Immunity in Improving Crop Cold Tolerance

Ye Jin

Central China Normal University

Za Khai Tuang

Central China Normal University

Yizhong Wang

Central China Normal University

Zhenjiang Wu

Central China Normal University

Wan-Nian Yang (✉ yangwn@mail.ccnu.edu.cn)

Central China Normal University <https://orcid.org/0000-0003-0056-7448>

Research Article

Keywords: freezing tolerance, flagellin 22, cross-talk, salicylic acid

Posted Date: September 16th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-874385/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Plant Cell Reports on November 24th, 2021.

See the published version at <https://doi.org/10.1007/s00299-021-02811-4>.

Abstract

Pathogen infection cross-activates cold response and increase cold tolerance of host plants. However it is not possible to use the infection to increase cold tolerance of field plants. Here flagellin 22 (flg22), the most widely-studied PAMP, was used to mimic the pathogen infection to cross-activate cold response. Flg22 treatment alleviated the injury caused by freezing in Arabidopsis, oilseed and tobacco. In Arabidopsis, flg22 activated the expression of immunity and cold-related genes. Moreover the flg22 induced alleviation of cold injury was lost in *NahG* transgenic line (SA-deficient), *sid2-2* and *npr1-1* mutant plants, and flg22-induced expression of cold tolerance-related genes, which indicating that salicylic acid signaling pathway is required for the alleviation of cold injury by flg22 treatment. In short flg22 application can be used to enhance cold tolerance in field via a salicylic acid-depended pathway.

Key Message

The application of flagellin 22 (flg22), the most widely-studied PAMP, enhance crop cold tolerance. ICE1-CBF pathway and SA signaling is involved in the alleviation of cold injury by flg22 treatment.

Introduction

Plants have worldly monitoring systems to adapt a variety of environmental disturbances including biotic and abiotic stress, such as drought, heat, cold, salinity, pathogen infection, insect feeding (Suzuki et al., 2014). Upon perceiving stress, plants trigger a series of signal transduction leading to reprogramming of gene expression to establish new balance between growth and defense response against the adverse environment to finish lifecycle and produce seeds (Zhu, 2016).

Cold stress, including chilling stress (0-15°C) and freezing stress (<0°C), restricts plant growth and development, limits plant geographical location and reduces crop production (Shi et al., 2018). C-repeat binding factors (CBFs), conserved transcription factors in many plant species, are critical for cold acclimation (Jia et al., 2016). *CBF* genes are rapidly induced by cold stress, and then activate the expression of downstream cold responsive gene (COR) to increase plant cold tolerance (Chinnusamy et al., 2007). *CBFs* are regulated by various transcription factors, which recognize the cis-elements in *CBFs*' simultaneously promoters under cold stress, including ICE1 (CBF expression1), CAMTA (calmodulin-binding transcription activator) proteins, MYB15, EIN3 (Ethylene insensitive 3) (Shi et al., 2018).

Under field conditions, cold stress affects the growth and development of plants simultaneously with other stress, such as pathogen infection. Cold stress cross activates disease resistance by activating the expression of PR1, PR2, and PR5 (Pathogenesis-Related 1, 2, and 5) through activating the function of transcription factor NAC062 (Seo et al., 2010), activating the pattern-triggered immunity in Arabidopsis plants in a HAT1 (HISTONE ACETYL TRANSFERASE 1)-dependent manner (Singh et al., 2014) and releasing CAMTA transcription factors mediated repression of SA biosynthesis (Kim et al., 2017). Our previous research showed that cold stress cross-activates basal immunity against bacterial pathogen

Pseudomonas syringae pv *tomato* DC3000 (*Pst* DC3000) through activating SA pathway and inhibiting of JA pathway (Wu et al., 2019). Interestingly this pathogen infection also cross-activates cold response and thus increase cold tolerance of host plants (Tuang et al., 2020). These investigations provide an idea to exploit immunity to increase plants abiotic stress tolerance.

Plants has a multilayered system to defense against pathogens, plant cells are able to recognize conserved epitopes of pathogen, which called pathogen-associated molecular patterns (PAMPs), such as bacterial flagellin (Wyrsh et al., 2015). Plasma membrane-localized pattern recognition receptors (PRRs) activates PAMP-triggered immunity (PTI), a mild and wide-spectrum disease resistance against various pathogens (Bigeard et al., 2015). Flagellin-22 (Flg22), the most widely-studied PAMP, has a conserved 22-amino acid domain shared by the bacterial flagellum. Flg22 is recognized by flagellin sensing 2 (FLS2) and activates a mitogen-activated protein kinase (MAPK) cascade that regulates the activation of defense-related genes (Zipfel et al., 2004), including salicylic acid (SA) pathway maker genes such as *PR1*, *PAD4*, *EDS1*, *NPR1* (Denoux et al., 2009), callose deposition (Muthamilarasan & Prasad, 2013), and production of reactive oxygen species (ROS) (Bigeard et al., 2015).

In this work, it is investigated whether flg22, a well-studied PAMP is able to cross-activate cold response and thus increase the cold tolerance of plants and our data indicated that the PAMP is a potential activator useful for improve the cold tolerance of crops.

Results

Flg22 treatment alleviated the injury caused by freezing in Arabidopsis

In previous work, it was shown that bacterial pathogen *Pst* DC3000 infection cross-activates the cold response and thus increased cellular viability under freezing (Tuang et al., 2020). As the pathogen infection leads to serious disease to plants, it is not possible to use the infection to increase freezing tolerance of field plants. Thus we investigated whether flg22, (a pathogen associated molecular patterns PAMPs), a maker peptide for PTI in freezing stress response, derived from bacterial flagellin proteins, could mimic the pathogen infection to cross-activate cold response to some extent.

Five days old Arabidopsis seedlings were treated with flg22, and 2 days later the infiltrated (2 dpi) seedlings were stressed under -4°C for 1 h, 2 h, and 2.5 h respectively. Then extracted the chlorophyll and determine the contents using UV-Spectrometer. As shown in Figure 1, both chlorophyll 'a' and 'b' of flg22 treated seedlings showed lower chlorophyll contents compared to control seedlings under control condition (22°C). However, at 1 h freezing stress, chlorophyll 'a' & 'b' in control and flg22 treated seedlings showed no difference. At 2 h freezing stressed flg22 treated seedlings showed higher contents compared to those of control seedlings (Figure 1). The results indicated that freezing stress not much effect chlorophyll contents in a test temperature and time points, but prior flg22 treatment induced chlorophyll contents upon freezing stress in Arabidopsis seedlings.

To further examine whether Flg22 treatment functions as pathogen infection under freezing stress. *Arabidopsis* plants were infiltrated by flg22, and 2 days later the infiltrated plants were stressed under freezing temperature (-4°C) for 6 h or 8 h, and the cellular viability was evaluated by trypan blue staining and ion leakage measurement. As shown in Figure 2, upon freezing, the CK (non-infiltrated), Mock (infiltrated by water), and flg22-infiltrated plants were significantly decreased in cellular viability and increased ion leakage level compared with their counterpart controls (non-freezing) (Figure 2). However, after 8 h of freezing stress, flg22-infiltrated plants showed significant higher cellular viability and lower ion leakage level than CK and mock plants (Figure 2a & c). These results indicate that prior flg22 treatment could alleviate cellular injury caused by subsequent freezing stress.

Flg22 infiltration induced chlorophyll contents of Oilseed and Tobacco seedlings upon freezing

To know whether flg22 treatment affects freezing tolerance in crops as found in *Arabidopsis*, we also treated five days old *Brassica napus* (winter Oilseed Rape) seedlings by flg22, and 2 dpi seedlings were stressed under -4°C. As shown in Figure 3 chlorophyll 'a' of flg22 treated seedlings showed higher contents than those of control seedlings in 1 h, 2 h, and 2.5 h of freezing stressed (Figure 3); and chlorophyll 'b' of flg22 treated seedlings showed higher contents than those of control seedlings in 2 h, and 2.5 h of freezing stressed (Figure 3). This result suggests that prior flg22 treatment-induced chlorophyll contents upon freezing stress.

Then we checked the chlorophyll contents of *Nicotiana tabacum* (Tobacco) seedlings upon freezing stress. Similar to Oilseed, Tobacco seedlings also showed higher chlorophyll contents in flg22 treated seedlings (Figure 4) compared to control seedlings upon freezing stress. Under normal conditions (22°C), control and flg22 treated seedlings showed no significant contents to each other. However, chlorophyll contents of flg22 prior treated seedlings showed higher contents compared to control seedlings after freezing stress (Figure 4). All chlorophyll contents were down to basal level at 2.5 h of freezing stress (Figure 3, 2.5 h). The results suggest freezing stress adversely affects the chlorophyll contents of Tobacco, and flg22 treatment indeed induced chlorophyll contents upon freezing stress in tested conditions. All these above indicates that flg22 treatment enhanced freezing tolerance in oilseed and tobacco.

Flg22 treatment activated the expression of cold-related genes

As higher cellular viability was detected in flg22-treated plants stressed by freezing (Figure 4a-4b) and our previous work showed the pathogen infection activated the expression of cold-responsive genes in host plants (Tuang et al, 2020), it was asked whether flg22 treatment also cross-induces these genes expression. We checked the expressions of some cold stress-related genes. As expected, some key genes

involved in cold tolerance were activated by flg22 at early (6 hpi) or/and late (48 hpi) stages. Interestingly, some of tested genes were only upregulated at early stage, including *ICE1*, *CBF3*, *COR15A* (Figure 5a, d and e). *CBF1*, *COR15B*, *COR27* and *COR413M2* were upregulated at both early and late stages (Figure 5b, f, g and h).

To find out the genetic mechanism on flg22 treatment increasing cold tolerance, we analyzed the expression of more genes which play positive function in cold tolerance. The expression of *CHS1*, *ATERDJ3A*, *DREB2A*, *MBF1C* (Figure 6a, c, g and i).were induced at both early and late stages. The expression of *AZI1*, *MKK2*, *DREB2C*, *EARL11*, *NAC019*, *NAC042* (Figure 6b, f, h, i, j and k).were only upregulated at early stage. The expression of flg22-activated immunity genes such as *EDS1*, *EDS5*, *EDS16* and *PR1* which were upregulated at both early and late stages (Supplement Figure 1). All these results indicate flg22 treatment, similar to pathogen injection, activates the expression of genes involved cold response pathway in host plants.

Flg22-activated cold tolerance is dependent on SA

Salicylic acid (SA) is required for the alleviation of cold injury by *pst* Dc3000, therefore we wondered whether the alleviation of cold injury by Flg22 was associated with SA accumulation. *NahG* transgenic plant, containing a salicylate hydroxylase gene *NahG*, was an SA deficient line (Tuang et al., 2020). As shown in figure 7a & c under freezing for 6 hours flg22 treated *NahG* plants showed no difference in cell viability and ion leakage with control and mock treated plants. After -4°C treatment for 8 hours flg22 infected *NahG* plants were more severely damaged by freezing than CK and mock ones (figure 7a&c), which was contrast with the results detected in wildtype. *NahG* transgenic plant, SA deficient line, lost the capacity of alleviating freezing injury by flg22 treatment, indicating SA was required in the alleviating.

To confirm the function of SA in alleviation of cold injury by flg22 treatment, the cold tolerance of flg22 treated *sid2-2* and *npr1-1* plants was checked. *SID2* plays an important role in SA accumulation, pathogen induced SA biosynthesis was significantly decreased in *sid2-2* mutant plants (Dempsey et al., 2011). The SA induced by pathogen induce the expression of downstream genes by activate NPR1 (Pajerowska-Mukhtar et al., 2013). NPR1 is the regulator of SA signaling pathway. As shown in figure 7e-l, flg22 treated *sid2-2* and *npr1-1* plants showed no difference in cell viability and ion leakage with control and mock treated plants under freezing for 6 hours. Flg22 treated *sid2-2* plants were more severely damaged by freezing for 8 hours than CK and mock ones. These above indicated the requirement of SID and NPR for the alleviation of freezing injury by flg22 treatment.

The effect of flg22 treatment on expression of cold tolerance genes was lost or compromised in *NahG*, *sid2-2*, and *npr1-1* plants

To check whether SA was required for the activation of genes which were related the cold tolerance and induced by flg22 treatment, the expression of these genes were detected in flg22 treated *NahG* transgenic, *sid2-2*, *npr1-1* plants. As shown in figure 8 the induction by flg22 infection of genes, including *CBF1*, *CBF3*, *ATERDJ3A*, *NAC042*, was lost in *NahG* transgenic plants. Only the expression of *DREB2C* and *MBF1C* was partially induced. In *sid2-2* mutant plants the induction of *CBF1*, *CBF3*, *DREB2C*, *MBF1C* was completely lost (figure 8). In *npr1-1* mutant plants the induction of *CBF1*, *CBF3*, *DREB2C*, *NAC042* and *MBF1C* was completely lost (figure 8). All these above indicated that NPR1-mediated SA signaling pathway is required for the activation of these cold tolerance related genes.

Discussion

Flagellin-22 treatment alleviates the injury caused by subsequent freezing stress

A great deal of evidences showed the crosstalk between abiotic stress and immune response, environmental factors are proved to modulate plant immunity (Saijo and Loo, 2020). For example, UV-B or excessive light stress induced the synthetize of flavonoids (Schenke et al., 2019), which weaken SA-based immunity (Serrano et al., 2012); heat stress induced the expression of HSP, which enhanced plant immunity by the accumulation and stability of pathogenesis-related (PR) proteins (Ul Haq et al., 2019); cold stress activated disease resistance through SA dependent pathway (Wu et al., 2019). Our earlier experiment showed that *pst* DC3000 infection enhanced the tolerance of freezing and heat injury in *Arabidopsis* (Tuang et al., 2020). However these findings are not possible for use in field because lack of practicability or the damage to plants.

Here, flagellin-22 treatment not only alleviated freezing injury in *Arabidopsis* (Figure 1&2) but also in oilseed (Figure 3) and tobacco (Figure 4). Flg22, a conserved peptide with 22 amino acid, has been widely used to induce PTI responses and does not lead to serious disease to plant (Wyrsh et al., 2015). All these above suggested that flg22 application may be used to enhance cold tolerance in field.

Flagellin-22 treatment activates signaling pathway of cold stress

When exposed to cold stress, plant rapidly induces the expression of many transcription factors, including the APETALLA2 (AP2)-domain proteins CBFs (*CBF1*, *CBF2* and *CBF3*). The expression of *CBF* genes was regulated by *ICE1*, a bHLH transcription factor (Chinnusamy et al., 2007). In *Arabidopsis*, *CBF2* is suggested as a negative regulator of *CBF1* and *CBF3* expression during cold response (Novillo et al., 2004). The cold-induced CBFs bind C-repeat/ dehydration-responsive elements in the promoters of downstream genes and activate the expression of downstream genes, such as numerous COR genes (Zhu 2016). Interestingly, in our work, the expression of *ICE1* was induced by flg22 treatment at early stage (6 hours after injection), the expression of *CBF1* and *CBF3* was up-regulated by flg22 and the expression

of *CBF2* was depressed by flg22 treatment at early and late stages (Figure 5). Furthermore a sum of *COR* genes were up-regulated by flg22 treatment, including *COR15A*, *COR15B*, *COR27*, *COR143M2* (Figure 5). These results suggested that the ICE1-CBF pathway is activated by flg22, thus increasing cold tolerance of host plants.

SA is required for alleviating the injury caused by cold stress

Salicylic acid (SA) has emerged as a key plant defense hormone for both pathogen-associated molecular patterns (PAMPs)-triggered immunity (PTI) and effector-triggered immunity (ETI) (Zhang and Li, 2019). SA also plays an important role in the plant responses to different abiotic stresses (Rivas-San and Plasencia, 2011). Cold was shown to induce SA biosynthesis (Kim et al., 2013 & 2017), meanwhile exogenous SA treatment was proved to provide protection against cold stress (Miura & Furumoto, 2013). Flg22 treatments caused SA accumulation through SID2-mediated SA biosynthesis (Tsuda et al., 2008). SID2, which encodes isochorismate synthase, is a SA biosynthetic enzyme (Wildermuth et al., 2001).

Our current work showed Flg22 treatment induced SA was required for the alleviation of subsequent cold injury, since both *NahG* line and *sid2-2* mutant plants lost the ability to alleviating the injury (Figure 7). Consistently, in these plants, Flg22 treatment-activated expression of tested cold stress genes was lost or compromised (Figure 8). Moreover, the alleviation of cold injury by Flg22 was lost in *npr1-1* mutant (Figure 7 & 8), which suggest SA signaling pathway also play a vital role in flg22 enhanced cold tolerance.

In summary, our work suggested that flg22 application may be used to enhance cold tolerance in field, indicated that flg22 treatment enhance of host plants to cold stress by activating cold signaling pathway ICE1-CBF. NPR1-mediated SA signaling is involved in the alleviation of cold injury by prior flg22 treatment.

Materials And Methods

Plant materials and flg22 treatment

The Arabidopsis materials including ecotype Col-0, *npr1-1* and *sid2-1*, *NahG* transgenic plants and growth conditions were described in Tuang et al (2020). The four-week-old plants with the 8th true leaf emerging were vacuum-infiltrated by 2 mg/L of Flg22 at -0.1 MPa for 1 min, followed by culturing under the normal condition for 2 days. Then, the plants were moved to -4°C for different time. The cellular viability, Chlorophyll contents and ion leakage were measured upon each time-point. Freezing treatment were performed under a dark condition.

For oilseed and tobacco materials, seeds were sown on half MS basal agar plates, and cold acclimated at 4°C for 48 h, then transferred to 22-23°C for germination. 5-day-old seedlings on growing plates were

sprayed by 2 mg/L flg22 solution, and sequentially cultured for 2 days. Then plates were moved to -4°C incubator for 1 h, 2 h, 2.5 h, and the chlorophyll contents were measured upon each time-point for evaluating the freezing damage.

Cellular viability assay and Chlorophyll contents measurement

Electrical conductance and trypan blue staining were used to evaluate the cellular viability as described in Tuang et al., (2020). Chlorophyll extraction was performed by following the method from Karan & Subudhi, (2012).

RNA isolation and Real-Time PCR

Total RNAs were extracted from Arabidopsis leaves with NucleoZol solution (Macherey-Nagel GmbH & Co. KG, Germany) following the manufacturer's instructions. Prime Script™ RT Reagent Kit with gDNA Eraser (Takara Bio Inc. China) was used to remove DNA contamination and synthesize cDNA. All qPCR reactions were performed using the UNICON™SYBR® Green master mix (Yeasen Co. Ltd, China). *TUB2* (*At5g62690*) was used as the internal control. The data were shown as the average ± SD from three biological replicates. Primers were described in our previous work (Tuang et al., 2020).

Statistics

The statistics analysis was performed using the software GraphPad Prism (Version 7.00). Student's t was used to test the significance of a single pair of sample differences. One-way ANOVA analysis of variance was used to test the significance of the differences between multiple pairs of samples, with Tukey's multiple comparisons test for a post-hoc test for pairwise significant comparison. A difference at $P < 0.05$ was considered significant.

Declarations

Author Contribution

YJ and WY conceived and designed research. ZT, YJ, YW and ZW conducted experiments. YJ wrote the manuscript. All authors read and approved the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Funding

This work was financially supported by the National Natural Science Foundation of China (grant numbers 31470365 and 31700216)

Data availability

All data generated or analysed during this study are included in this published article and its supplementary information files.

References

1. Bigeard J, Colcombet J, Hirt H. (2015). Signaling mechanisms in pattern-triggered immunity (PTI). *Molecular plant*. 8: 521-39.
2. Chinnusamy V, Zhu JH, Zhu JK. (2007) Cold stress regulation of gene expression in plants, *Trends Plant Sci*. 12: 444-51.
3. Denoux C, Galletti R, Mammarella N, Gopalan S, Werck D, De Lorenzo G, Dewdney J. (2009). Erratum: Activation of Defense Response Pathways by OGs and Flg22 Elicitors in Arabidopsis Seedlings. *Molecular plant*. 2: 838.
4. Dempsey DA, Vlot MCW, Daniel FK. (2011). Salicylic acid biosynthesis and metabolism. The Arabidopsis Book/American Society of Plant Biologists, 2011(9), e0156.
5. Karan, R., & Subudhi, P. K. (2012). A stress inducible SUMO conjugating enzyme gene (SaSce9) from a grass halophyte *Spartina alterniflora* enhances salinity and drought stress tolerance in Arabidopsis. *BMC plant biology*. 12: 187.
6. Kim YS, An C, Park S, Gilmour SJ, Wang L, Renna L, Brandizzi F, Grumet R, Thomashow MF. (2017). CAMTA-mediated regulation of salicylic acid immunity pathway genes in Arabidopsis exposed to low temperature and pathogen infection. *Plant Cell*. 29: 2465-77.
7. Kim Y, Park S, Gilmour SJ, Thomashow MF. (2013). Roles of CAMTA transcription factors and salicylic acid in configuring the low-temperature transcriptome and freezing tolerance of Arabidopsis. *Plant J*. 75: 364-76.
8. Jia Y, Ding Y, Shi Y, Zhang X, Gong Z, Yang S. (2016) The *cbfs* triple mutants reveal the essential functions of CBFs in cold acclimation and allow the definition of CBF regulons in Arabidopsis. *New Phytol*. 212: 345-53.
9. Miura K & Furumoto T. (2013). Cold signaling and cold response in plants. *Int J Mol Sci*. 14: 5312-37.
10. Muthamilarasan M & Prasad M. (2013). Plant innate immunity: an updated insight into defense mechanism. *J Biosci*. 38: 433-49.
11. Novillo F, Alonso J M, Ecker J R, Salinas J. (2004). CBF2/DREB1C is a negative regulator of CBF1/DREB1B and CBF3/DREB1A expression and plays a central role in stress tolerance in

- Arabidopsis. *Proc. Natl. Acad. Sci.* 101: 3985-90.
12. Pajerowska-Mukhtar KM, Emerine DK, Mukhtar MS. (2013). Tell me more: Roles of NPRs in plant immunity. *Trends in Plant Science*, 18: 402-11.
 13. Rivas-San VM & Plasencia J. (2011). Salicylic acid beyond defence: its role in plant growth and development. *J Exp Bot.* 62:3321-38
 14. Saijo Y & Loo EP. (2020). Plant immunity in signal integration between biotic and abiotic stress responses. *New Phytol.* 225: 87-104.
 15. Schenke D, Utami HP, Zhou Z, Gallegos M-T, Cai D. (2019). Suppression of UV-B stress induced flavonoids by biotic stress: Is there reciprocal crosstalk? *Plant Physiol. Biochem.* 134: 53-63.
 16. Seo PJ, Kim MJ, Park JY, Kim SY, Jeon J, Lee YH, Kim J, Park CM. (2010) Cold activation of a plasma membrane-tethered NAC transcription factor induces a pathogen resistance response in Arabidopsis. *Plant J.* 61: 661-71.
 17. Serrano M, Kanehara K, Torres M, Yamada K, Tintor N, Kombrink E, Schulze-Lefert P, Saijo Y. (2012). Repression of sucrose/ultraviolet B light-induced flavonoid accumulation in microbe-associated molecular pattern-triggered immunity in Arabidopsis. *Plant Physiol.* 58: 408-22.
 18. Shi Y, Ding Y, Yang S. (2018). Molecular Regulation of CBF Signaling in Cold Acclimation. *Trends Plant Sci.* 23: 623-37.
 19. Singh P, Yekondi S, Chen PW, Tsai CH, Yu CW, Wu K, Zimmerli L. (2014). Environmental history modulates Arabidopsis pattern triggered immunity in a HISTONE ACETYLTRANSFERASE1-dependent manner. *Plant Cell.* 26: 2676-88.
 20. Suzuki N, Rivero RM, Shulaev V, Blumwald E, Mittler R. (2014). Abiotic and biotic stress combinations. *New Phytol.* 203:32-43.
 21. Tsuda K, Sato M, Glazebrook J, Cohen JD, Katagiri F. (2008). Interplay between MAMP-triggered and SA-mediated defense responses. *Plant J.* 53: 763-75.
 22. Tuang ZK, Wu Z, Jin Y, Wang Y, Oo PPZ, Zuo G, Shi H, Yang W. (2020). Pst DC3000 infection alleviates subsequent freezing and heat injury to host plants via a salicylic acid-dependent pathway in Arabidopsis. *Plant Cell Environ.* 43:801-17.
 23. Ul Haq S, Khan A, Ali M, Khattak AM, Gai WX, Zhang HX, Wei AM, Gong ZH. (2019). Heat Shock Proteins: Dynamic Biomolecules to Counter Plant Biotic and Abiotic Stresses. *Int J Mol Sci.* 20: 5321.
 24. Wildermuth MC, Dewdney J, Wu G, Ausubel FM. (2001). Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature.* 414: 562-5.
 25. Wyrsh I, Domínguez-Ferreras A, Geldner N, Boller T. (2015) Tissue-specific FLAGELLIN-SENSING 2 (FLS2) expression in roots restores immune responses in Arabidopsis fls2 mutants. *New Phytol.* 206:774-84.
 26. Zhang Y, Li X. (2019). Salicylic acid: biosynthesis, perception, and contributions to plant immunity. *Curr Opin Plant Biol.* 50: 29-36.

27. Zhu JK. (2016). Abiotic Stress Signaling and Responses in Plants. *Cell*. 167: 313-24.

28. Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JD, Felix G, Boller T. (2004) Bacterial disease resistance in Arabidopsis through flagellin perception. *Nature*. 428: 764-7.

Figures

Figure 1

The effects of flg22 treatment on chlorophyll contents of Arabidopsis seedlings upon freezing stress. 2 dpi Arabidopsis seedlings were treated by freezing (-4°C) for the indicated time (0 h (Control), 1 h, 2 h, 2.5

h). The quantified values are the average $\pm$ SD from three biological repeats, and different lowercase letters, a-g, showed a significant difference ($p < 0.05$).

Figure 2

Cellular viability assay on 2 dpi of flg22 treated Arabidopsis plants (four weeks old) by trypan blue staining and ion leakage upon freezing stress. (a) Cellular viability assay by trypan blue staining upon freezing (-4°C) stress at the indicated time, 22°C (Control), -4°C(6h), -4°C(8h); (b) Cellular viability assay by ion leakage assay upon freezing (-4°C) stress at the indicated time, 22°C (Control), -4°C(6h), -4°C(8h). (c) and (d) Ratio of cold treatment value to control in (a) and (b). The quantified values in (a) and (b) are the average $\pm$ SD from three biological repeats, and different lowercase letters, a-d, showed a significant difference ($p < 0.05$).

Oilseed

Figure 3

The effects of flg22 treatment on chlorophyll contents of Oilseed seedlings upon freezing stress. 2 dpi Oilseed seedlings were treated by freezing (-4°C) for the indicated time (0 h (Control), 1 h, 2 h, 2.5 h). The quantified values are the average $\pm$ SD from three biological repeats, and different lowercase letters, a-g, showed a significant difference ($p < 0.05$).

Tobacco

Figure 4

The effects of flg22 treatment on chlorophyll contents of Tobacco seedlings upon freezing stress. 2 dpi seedlings Tobacco were treated by freezing (-4°C) for the indicated time (0 h (Control), 1 h, 2 h, 2.5 h). The quantified values are the average $\pm$ SD from three biological repeats, and different lowercase letters, a-g, showed a significant difference ($p < 0.05$).

Figure 5

RT-qPCR to check the expression of several marker genes in cold tolerance upon flg22 treatment at 6 hpi, and 48 hpi. TUB2 (AT5G62690) was used as an internal reference. The data were shown as the average $\pm$ SD of three biological repeats. The different lower letters, a, b, c, etc. in graphs indicate the significant difference between different samples ($p < 0.05$, $n = 3$).

Figure 6

RT-qPCR to check the expression of several cold-responsive genes upon flg22 treatment at 6 hpi, and 48 hpi. TUB2 (AT5G62690) was used as an internal reference. The data were shown as the average $\pm$ SD of three biological repeats. The different lower letters, a, b, c, etc. in graphs indicate the significant difference between different samples ($p < 0.05$, $n = 3$).

Figure 7

cell viability analysis on 2-dpi flg22 treated *NahG* transgenic (a–d), *sid2-2* (e–h), and *npr1-1* mutant (i–l) plants under freezing (–4°C). 2-dpi mutant plants were treated by freezing (–4°C) for the indicated time. (a), (e), and (i) Cell viability assay by trypan blue staining. (b), (f), and (j) Ion leakage assay. (c), (d), (g), (h), (k), and (l) Ratio of freezing treatment value to control in (b), (c), (g), (h), (l), and (m) respectively The

quantified values in (a), (b), (e), (f), (i), and (j) are the average $\pm$ SD from three biological repeats, and different lowercase letters, a, b, and c, showed the significant difference ($p < .05$)

Figure 8

RT-qPCR to check the expression of some pathogen-inducible abiotic stress genes in flg22 treatment NahG transgenic, sid2-2, and npr1-1 mutant plants (2 dpi). TUB2 (AT5G62690) was used as the internal reference. The data were shown as the average $\pm$ SD of three biological repeats. The different lower letters, a, b, and c, in graphs indicate the significant difference between different samples ($p < 0.05$)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementFigure1.docx](#)