

# Designing Chickpea For a Hotter Drier World

**Oluwaseun Johnson Akinlade**

University of Queensland - Saint Lucia Campus: The University of Queensland

**Kai Voss-Fels**

University of Queensland - Saint Lucia Campus: The University of Queensland

**Roy Costilla**

AgResearch Ruakura Research Centre Hamilton

**Jana Kholova**

International Crops Research Institute for the Semi-Arid Tropics

**Sunita Choudhary**

International Crops Research Institute for the Semi-Arid Tropics

**Rajeev K Varshney**

International Crops Research Institute for the Semi-Arid Tropics

**Lee Hickey** (✉ [l.hickey@uq.edu.au](mailto:l.hickey@uq.edu.au))

University of Queensland <https://orcid.org/0000-0001-6909-7101>

**Millicent R Smith**

University of Queensland - Saint Lucia Campus: The University of Queensland

---

## Research Article

**Keywords:** Cicer arietinum, abiotic stress, drought, heat, genomic prediction, high-throughput phenotyping

**Posted Date:** September 22nd, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-875442/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

Chickpea (*Cicer arietinum* L.) is one of the most important grain legumes in the world, but its current and future production is threatened due to the increased incidence of drought and heat stress. To address this challenge, an integrated crop improvement strategy encompassing breeding, genomics, physiology and agronomy is required. Here, we review the physiological traits known to confer drought and heat adaptation in chickpea and identify areas of drought and heat adaptation research that should be prioritised in the future. Furthermore, we underscore approaches to efficiently phenotype chickpea adaptation traits and highlight the significant challenges and importance of understanding the nexus between canopy and root development. Finally, we present the opportunity to adopt multi-trait genomic prediction approaches to efficiently utilise key physiological traits, that can be assayed using high-throughput phenotyping platforms, to accelerate genetic gain in drought and heat prone environments.

## Key Message

***A body of research on heat and drought stress in chickpea has built a foundation of trait knowledge that can be harnessed using new breeding technologies to accelerate genetic gain for target environments.***

## Introduction

Chickpea (*Cicer arietinum* L.) plays a crucial role in human and animal nutrition and agricultural systems by improving soil fertility and providing a disease break (Siddique et al. 2011). Chickpea is primarily grown in rainfed production systems that typically rely on the residual soil moisture from the preceding season (Ramamoorthy et al. 2017; Varshney et al. 2013b). As such, production is often affected by drought and heat, particularly when abiotic stress coincides with the reproductive phase (Gaur et al. 2018).

Yield losses due to drought are variable and depend on the timing and degree of water stress. For instance, yield reductions can range from 58–95% depending on severity (Fang et al. 2010; Leport et al. 2006). Heat stress also significantly impacts yield due to its impact on pod set and seed set number (Devasirvatham et al. 2013; Devasirvatham et al. 2012). For every 1°C increase in seasonal temperature, yield is predicted to decrease by 53–301 kg<sup>-1</sup> ha<sup>-1</sup> in chickpea (Kalra et al. 2008). Under climate change, yield losses to heat and drought are expected to increase. Chickpea is particularly vulnerable to climatic change given the limited genetic variation in elite germplasm (Muehlbauer and Sarker 2017).

Best management practices may not guarantee maximum performance because of in-season variability, particularly in regions without reliable access to irrigation. Therefore, germplasm that is well adapted to the target environment is required. Research and breeding efforts have been employed to improve yield in some regions over the past 30 years. However, current yield trends suggest that production is increasing due to expansion in total land area cultivated, rather than genetic improvement per se (Foyer et al. 2016).

Continued investment in research and crop improvement programs is critical to develop more productive cultivars suited to the target environments in which they are grown.

Within any crop improvement program, understanding the target environment is key to ensure genetic progress leads to productivity gains in growers fields (Cooper et al. 2020). This includes stresses that limit crop productivity such as the intensity and frequency of water deficit. To achieve this goal, envirotyping approaches can be undertaken and information used to strategically deploy a combination of traits into the right environments (Tardieu et al. 2018; Kholova et al. 2021). This process has commenced for chickpea in India (Hajjarpoor et al. 2018).

A number of studies have highlighted key physiological traits that underpin adaptation to heat and drought environments (Chen et al. 2017; Singh et al. 2008; Ramamoorthy et al. 2017). The publication of accessible genomic resources, including the reference genomes for *Cicer arietinum* and *Cicer reticulatum* (Varshney 2016; Varshney et al. 2013c), provided a major step forward for chickpea improvement. These resources and tools, combined with new breeding technologies, such as high-throughput phenotyping and genomic selection, offer new opportunities to rapidly improve germplasm by targeting physiological traits that confer yield advantages in drought and heat-prone environments (Roorkiwal et al. 2020).

This review highlights recent developments in physiological and molecular research of drought and heat adaptation in chickpea. It underscores approaches and technologies that could accelerate the development of high yielding drought and heat adapted varieties.

## Literature Analysis Reveals Opportunities For Future Research

To understand the current status of drought and heat research in chickpea and identify key research areas that may be explored into the future, we searched for articles in Scopus using the following keywords: drought; water deficit; heat; high temperature; chickpea; *Cicer arietinum*; *Cicer reticulatum*. Our search produced 365 articles published between 1970 and 29 June 2020. A total of 58% of the articles focused on drought, 38% on heat, and 4% on heat and drought combined.

In addition, we analysed all text in the abstract, title, and keywords of each article using Leximancer version 4.5. The analysis revealed that studies containing words; “gene or genetics”, “flowering” and “canopy” occurred at a lower frequency (Fig. 1). This suggests that many of the studies to date have focused on the agronomic performance of chickpea to drought and heat. This is not surprising, considering that trait dissection and gene discovery studies require long-term and sustained funding to support resources including, though not limited to, high-cost infrastructure, laboratory facilities, and specialised personnel.

Given the excellent genetic resources available for chickpea and the availability of new technologies we predict a rapid expansion of genomics research in the next decade (Thudi et al. 2021). Additionally, lessons learned from other crops will likely accelerate progress for chickpea as adaptation mechanisms are often conserved and may be translated across crops.

# Traits Conferring Drought And Heat Adaptation

Over thousands of years, terrestrial plants have successfully adapted to a dry environment characterized by temperature and light extremes. In agricultural systems, the impact of abiotic stress is influenced by genotype, environment and management interactions (Messina et al. 2009). Plants use different mechanisms to respond to challenges in the environment which allow for continued survival. Plant responses to the environment are complex as multiple abiotic or biotic stressors often affect species simultaneously (Canci and Toker 2009; Rani et al. 2020). As a result, drought and heat tolerance mechanisms operate at various spatial and temporal scales (Tardieu et al. 2018). The extent to which drought and heat impact on plants is primarily dependent on the timing and duration of the stress (Carrão et al. 2016). These factors cumulatively make it challenging to study and improve plants for drought and heat adaptation.

Although the timing and severity of stress are dependent on the environment, the majority of research suggests that drought mostly impacts the reproductive phase in several crops, including chickpea (Lamaoui et al. 2018; Ramamoorthy et al. 2017; Daryanto et al. 2015; Rani et al. 2020). As the canopy develops and water demand increases, residual soil moisture is often depleted during the critical reproductive stage. Under drought stress, pod and flower abscission increases in response to reduced assimilate supply, pod set reduces and consequently, seed yield is impacted (Leport et al. 2006; Fang et al. 2010). High temperature, has a similar impact during the reproductive phase, affecting pod set and pod filling stages leading to a yield penalty (Devasirvatham et al. 2013). Due to the nature of abiotic stress, heat and drought stress often occur simultaneously. Unsurprisingly, many traits associated with adaptation to heat stress also confer an advantage under drought. For instance, early flowering can avoid both ends of season drought and heat.

Traits vary in their importance for drought and heat adaptation and the impact that each has on agronomic performance is context dependent. QTL have been identified for traits involved in drought and heat adaptation in chickpea (Table 1). Table 1 primarily includes traits associated with phenology, yield components and root system architecture. This highlights an opportunity to understand the genetic controls of canopy development and other key traits associated with water use efficiency.

Table 1  
Summary of QTL identified for drought and heat adaptation traits

Trait	QTL	Location	Reference
Root length density (cm cm <sup>-3</sup> )	<i>QR3rld01</i>	CaLG04, 62.56–73.06 cM bin_4_13239546 bin_4_13378761	(Varshney et al. 2014)
	<i>QR3rld01</i>	CaLG04 ( 3.23–5.37 cM)	(Jaganathan et al. 2015)
Total dry root weight/total plant dry weight ratio (RTR, %)	<i>Ca_04586</i> ,	CaLG04 (2.7 Mb)	(Singh et al. 2016)
	<i>QR3rtr01</i> (16.67 %)	CaLG04 (5 cM)	(Varshney et al. 2014)
	<i>QR3rtr01</i> ( 10.85–13.56 %)	CaLG04 (1.81–5.37 cM)	(Jaganathan et al. 2015)
Primary root branches (PBS)	<i>QR3pbs02</i> (12.92 %)	CaLG08, bin_8_6034209 bin_8_5984553	(Jaganathan et al. 2015)
100-Seed weight (g)	<i>QR4100sdw02</i> ,	CAM2093-ICCM0349	(Varshney et al. 2014)
	<i>Ca_04364</i> (28.61%), <i>Ca_04607</i> (19.25%)	CaLG04 (1.13 cM), CaLG04 (1.36 cM)	(Singh et al. 2016)
	<i>QR3100sdw03</i> (60.41%),	CaLG04, flanked CaM2093–ICCM0249	(Varshney et al. 2014)
	<i>QR3sdw01</i> ( 10.78–26.91 %)	CaLG04 ( 0.73–5.91cM)	(Jaganathan et al. 2015)
	<i>qSW-1</i> (11.54%), <i>qSW-2</i> (19.24%)	LG3 (CEST129- CESSR221)	(Gupta et al. 2015)
Total number of seeds per plot	Qts02_5 (12.0%)	Ca5_44667768, Ca5_46955940, Ca6_7846335, Ca6_14353624	(Paul et al. 2018)
Percentage Pod Setting	q%podset06_5 (11.51%), q%podset08_6 (8.44%)	Ca5_44667768, Ca5_46955940, Ca6_7846335, Ca6_14353624	(Paul et al. 2018)
Grain Yield per Plot	qgy02_5 (16.04%), qgy03_6 (4.43%)	Ca5_44667768, Ca5_46955940, Ca6_7846335, Ca6_14353624	(Paul et al. 2018)

\*% is the phenotypic variation explained (PVE).

Trait	QTL	Location	Reference
Pod/plant	<i>QR3pod01</i> , <i>QR4pod02</i>	CaLG04, bin_4_13239546 bin_4_13378761, CaM0772- TS45	(Varshney et al. 2014)
Filled pods per plot	qfpod02_5 (11.57%), qfpod03_6 (6.56%)	Ca5_44667768, Ca5_46955940, Ca6_7846335, Ca6_14353624	(Paul et al. 2018)
Plant height (cm)	<i>QR3pht01</i> ,	CaLG04	(Varshney et al. 2014)
	<i>QR3pht06</i> , <i>QR3pht08</i>	CaLG04	(Varshney et al. 2014)
	<i>qPH-1</i> (13.98%), <i>qPH-2</i> (12.17%),	LG6 (NCPGR199- NCPGR202), LG7 (CEST47- NCPGR34)	(Gupta et al. 2015)
	<i>QR3pht03</i> ( 14.36- 76.54 %)	CaLG04	(Varshney et al. 2014)
Days to 50% flowering (DF)	<i>QR3df04</i> (67.71%),	CaLG08 (1.81 cM), bin_8_6034209 bin_8_5984553,,	(Jaganathan et al. 2015)
	<i>QR3df01</i> (26.87%), <i>QR4df06</i>	NCPGR164 Ca8_3050452, TA103II-TA122	(Varshney et al. 2014)
Early flowering	<i>efl-1</i> , <i>Efl-1</i>	ICCV 2 X JG 62 (RIL)	(Kumar and van Rheenen 2000)
Days to maturity	<i>QR3dm01</i> (47.43%)	CaLG06 7.33 cM	(Jaganathan et al. 2015)
Visual Score on podding behaviour	qvs05_6 (11.07%)	Ca6_7846335, Ca6_14353624	(Paul et al. 2018)
*% is the phenotypic variation explained (PVE).			

There are two main ways to adapt a crop to drought; 1) optimise the timing of water use across development and 2) improve access to water.

A determinant of water demand in chickpea is the rate and extent of canopy development (Sivasakthi et al. 2017). This is highly dependent on the timing of water use, which among other traits, is associated with phenology (Zaman-Allah et al. 2011). Canopy architecture traits such as leaf area development, canopy size and canopy conductance influence transpiration rate and in some environments, may improve drought adaptation (Thudi et al. 2014).

To improve access to water, root architecture traits have been a major target in chickpea crop improvement over many decades (Siddique et al. 2011; Saxena et al. 1993). Significant genetic variation

in root traits have been reported (Kashiwagi et al. 2006; Chen et al. 2017; Purushothaman et al. 2017; Serraj et al. 2004). Moreover, many studies report that improvements in root traits, specifically root length density and rooting depth, have an overall positive impact on adaptation to drought (Ludlow and Muchow 1990; Krishnamurthy et al. 2003; Kashiwagi et al. 2005; Kashiwagi et al. 2015; Gaur et al. 2008; Ramamoorthy et al. 2017; Saxena et al. 1993). Rooting depth can exceed 100 cm (Kashiwagi et al. 2006), and water uptake from soil layers of 90–120 cm is a feature of many drought adapted genotypes (Purushothaman et al. 2017). Serraj et al. (2004) noted that increases in root depth and root length density led to greater water use translating to higher yields. Similarly, a prolific root system was determined to positively affect seed yield under drought (Kashiwagi et al. 2006; Varshney et al. 2013a).

However, changes in root architecture may have a tradeoff in some environments. Increases in rooting depth and biomass may not necessarily lead to increases in grain yields due to the metabolic cost of increased biomass partitioning and energy loss through respiration (Vadez et al. 2008; Kashiwagi et al. 2015; Ramamoorthy et al. 2017). Additionally, it is now also recognised in chickpea that temporal changes in root growth influence the effective use of available water during the crop cycle (Vadez et al. 2008; Zaman-Allah et al. 2011; Vadez et al. 2007). There is a strong association between aboveground biomass and profligate water uptake, leading to higher water use and a yield penalty in some contexts as water availability during pod initiation is critical for yield development (Fig. 2). During the vegetative period, conservative water use can improve water availability during pod initiation, which is a critical period for determining yield. Conservative water use is a function of the root system and aboveground traits such as canopy development.

The importance of considering root traits in terms of their spatial and temporal characteristics is supported by studies that describe genotypes suitable for water-limited environments as those with root growth vigour and deeper soil root proliferation at the beginning of the reproductive stage (Kell 2011; Singh et al. 1995). Furthermore, an optimal root system for efficient uptake of soil water is unlikely to be optimal for nutrient uptake. To avoid this tradeoff, there may be genetic variation in the root system architecture that could be exploited.

## Understanding The Root-shoot Nexus

The relationship between canopy development and the root system is crucial and can be investigated to avoid yield tradeoffs in drought and heat adapted varieties. Similar to the principle of supply and demand, the root system supplies the shoot with water and soil nutrients according to demand as they operate within the same hydraulic continuum (Vadez 2014; Vadez et al. 2014). Sivasakthi et al. (2017) reported that root hydraulic characteristics influence chickpea transpiration rate under different vapour pressure deficit conditions and in turn affects its vigour and water use efficiency. This highlights the strong connection between root and canopy traits. This makes sense from an evolutionary perspective as it is how plants regulate the balance of resources relative to growth and reproductive success. However, this nexus could get in the way when selecting for traits associated with improved water use and uptake. For example, it is difficult to match water uptake and water use efficiency with time because root and

canopy development are closely related (Ratnakumar et al. 2009; Maurel et al. 2010; Zaman-Allah et al. 2011; Bouteille et al. 2012). Identifying genetic variants that deviate from this relationship may provide novel pathways to design chickpea varieties suited to drought and heat for different environments.

A strong connection between root and canopy development has been reported in several species. For example, in wheat and barley a key gene in the flowering pathway, *VERNALIZATION1* (*VRN1*) (Deng et al. 2015) also modulates root architecture (Voss-Fels et al. 2018). Similar findings have been reported in maize, where 62 of 133 QTL for nodal root number were also associated with flowering time (Zhang et al. 2018). To date, little is known about the relationship between above- and below-ground traits in chickpea. Nevertheless, there is a need to assemble unique combinations of root and canopy architectural traits. If these cannot be found in existing germplasm collections, they may be engineered through gene editing approaches by targeting the key pathways involved (Massel et al. 2021).

## Approaches To Efficiently Measure Traits

Targeted, novel phenotyping methods are required to understand phenotype-genotype relationships (Cobb et al. 2013) and support crop improvement (Tester and Langridge 2010; Rahaman et al. 2015). Ideally, phenotyping approaches will be hypothesis-driven, accurate, rapid, and cost-effective (Jang et al. 2020; de Castro et al. 2019; Kholova et al. 2021). This will make them suitable for evaluating large mapping populations which are required to dissect the genetic architecture of adaptive traits and more likely to be integrated into breeding programs (Rutkoski et al. 2016).

The success of molecular and traditional breeding may be amplified when integrated with high-throughput phenotyping tools (Xiao et al. 2017; Ren et al. 2020). Phenotyping complex traits underpinning adaptation to drought and heat can be time-consuming, labour intensive and often expensive. For instance, measuring mature root system architecture may give precise information for physiological insight, but only a small number of genotypes can be screened and the technique is usually destructive. This is not complementary with modern plant breeding approaches that seek to evaluate large numbers of lines in a non-destructive fashion.

Emerging phenotyping technologies enable rapid, non-invasive trait capture of large populations directly in the field (Smith et al. 2021). These technologies can utilise mechanistic physiological understanding of abiotic stress response for applied breeding applications. For instance, unmanned aerial vehicle (UAV) imaging platforms are evolving as a powerful phenotyping tool and may offer an efficient, field-based, non-invasive approach to evaluate drought and heat traits such as integrative end-of-season traits and canopy temperature (Reynolds et al. 2006; Lopes and Reynolds 2010). While for cereals, plot-level indices derived from UAV platforms have been validated (Pinto and Reynolds 2015; Deery et al. 2014; Deery et al. 2016) this must be first demonstrated prior to implementation in chickpea breeding programs. Leaf, canopy and plot scale traits are likely to be very different for chickpea compared to cereals, such as wheat due to differences in leaf size, shape, canopy architecture and development.

Dedicated phenotyping facilities allow for the precise evaluation of traits in a high-throughput manner under controlled conditions (Honsdorf et al. 2014; Asif et al. 2021) or conditions similar to the field (Vadez et al. 2015). Phenotyping platforms reach their full potential when they are used to evaluate well-designed populations important to the breeding program and phenotypes are analysed with other datasets in a context specific manner. Such technologies have been used in chickpea and combined with genomics have led to the discovery of QTL associated with adaptation in water limited environments (Sivasakthi et al. 2018). Phenotyping technologies such as the LeasyScan (Vadez et al. 2015) are a unique and powerful tool to discover novel trait variations that may be subsequently used in breeding programs.

## Mining And Exploiting Genetic Variation

As with most modern crops, elite chickpea germplasm has a narrow genetic base as a result of several genetic bottlenecks, including domestication and intensive selection in breeding programs (Chandora et al. 2020). Globally, there are approximately 87,341 chickpea accessions in genebanks, with the International Crop Research Institute for the Semi-arid Tropics (ICRISAT) genebank housing 20,764 accessions (308 wild and 20,456 cultivated) and the International Centre for Agricultural Research in the Dry Areas (ICARDA) housing 15,734 accessions (540 wild and 15,194 cultivated) (Abbo et al. 2003). These accessions include wild relatives, landraces and historical breeding material.

While genebank accessions may carry useful traits for heat and drought adaptation, there are many challenges associated with using these materials in a breeding context. This germplasm is typically not adapted to the target environment and therefore requires a high degree of pre-breeding to effectively transfer the novel traits into adapted germplasm. This can be a time consuming process, involving many cycles of backcrossing. Furthermore, without markers linked to useful traits, to identify suitable donors, genebank accessions must be screened through phenotyping platforms which is a major bottleneck. These challenges make the genebank-to-farm process a lengthy, laborious and costly one. A further consideration is a need for a benefit-sharing agreement and Nagoya compliance for some historical collection of certain biological materials (Sherman and Henry 2020; Smith et al. 2018).

Despite the wealth of genetic resources, this genetic variation is underutilised in most breeding programs. To date, QTL mapping studies (Table 1) have largely focused on bi-parental populations. Such populations offer high statistical power but lack mapping resolution and only examine two alleles at any one locus (Ongom and Ejeta 2018; Jannink 2007). To identify useful germplasm, Focused Identification of Germplasm Strategies (FIGS) concept was developed to create a subset of genetic material for easy evaluation. For example, FIGS was successfully applied to discover materials needed for drought tolerance in maize (Bari et al. 2012) and aphid resistance in wheat (El Bouhssini et al. 2012). Similarly, this method may be applied to scout for parental lines with novel alleles for drought and heat adaptation in chickpea. Incorporating available supporting software for decision making in specific traits would enhance the use of FIGS in chickpea (Bari et al. 2016). The concept of core reference sets can also be used to finely characterise and identify important germplasm from a pool of materials available across genebanks (Glaszmann et al. 2010). These core reference sets are used to represent available diversity,

evaluate for phenotypic variation and dissect traits for genes underlying their functions (Glaszmann et al. 2010; Emma Huang et al. 2013). In general, core collections help to improve the use of genetic resources for crop improvements (Upadhyaya and Ortiz 2001).

Once promising accessions have been identified, populations can be created to understand the genetic control of relevant traits. Creating multi-parent populations allows for the inclusion of multiple donor parents potentially carrying multiple alleles at any one locus (Thudi et al. 2014; Bandillo et al. 2013) increasing the likelihood of discovering novel, allelic variation that could be harnessed in breeding programs (Ladejobi et al. 2016). For instance, Multiparent Advanced Generation Intercross (MAGIC) and Nested Association Mapping (NAM) populations have been generated to study a range of economically important traits in a variety of crops such as; sorghum (Ongom and Ejeta 2018), maize (Huang et al. 2015; Holland 2015; Ren et al. 2020), barley (Mathew et al. 2018; Hemshrot et al. 2019; Afsharyan et al. 2020), and cowpea (Huynh et al. 2018). Recently a chickpea MAGIC population was created at ICRISAT using eight parents (Attri 2018; Gaur et al. 2019). This represents a useful resource to study the genetic control of heat and drought adaptation. To complement this effort a NAM population could be created to harness diversity from genebanks and continually updated over time (Bari et al. 2016).

Identifying alleles underpinning complex traits and transferring them into elite material is challenging and time-consuming. Therefore, we propose an integrated approach for chickpea (Fig. 3). High-throughput phenotyping could be used to rapidly screen germplasm with desirable heat and drought traits for pre-breeding activities that could exploit multi-parent populations for trait dissection. Identified alleles may be incorporated into elite germplasm for the target environment. This integrated approach requires a multidisciplinary team and cutting edge pre-breeding and breeding approaches.

## **Latest Breeding Approaches: Tools For Faster Crop Improvement**

Understanding the genetic architecture of traits in diverse germplasm gives insight into key QTL and novel alleles that can be targeted in breeding programs. Selection is an essential step in plant breeding programs for improvement of target traits of interest. A concept of genomics-assisted breeding has been suggested for integrating advanced genome discoveries in crop improvement (Varshney et al. 2005; Varshney et al. 2021). For instance, marker assisted selection (MAS) involves applying selection for a molecular marker associated with a trait of interest. MAS has been used in the past to hasten the process of selection and can reduce the years of pre-breeding and breeding activities for simply inherited traits underpinned by single or few genes. When combined with backcrossing, i.e. marker assisted backcrossing (MABC) the technique can facilitate the transfer of a trait into an elite genetic background, for example the introgression of a QTL region into elite chickpea cultivars to improve drought tolerance (Varshney et al. 2013a; Bharadwaj et al. 2021). However, MAS or MABC is not the preferred approach for complex traits governed by many genes (Charmet et al. 2020; Wang et al. 2018). Alternatively, genomic prediction (GP) is well-suited to improve complex traits as it considers the effect of all loci on the trait of interest (Cossa et al. 2017).

Genomic prediction (GP), the use of genome-wide DNA polymorphisms to estimate breeding values, was proposed to better select individuals when dealing with complex traits. GP does not rely on QTL identification and detection (Meuwissen et al. 2001; Meuwissen et al. 2016). Application in a breeding program involves building a prediction model based on a reference population with genome-wide markers and phenotype information (Wang et al. 2018; Heffner et al. 2009; Kushwah et al. 2020). GP can increase genetic gain per year and reduces the length of time required for variety development (Wang et al. 2018). By performing GP on selection candidates, a plant breeder can make selections in early generations without the time-consuming step of phenotyping across multiple years and seasons. This saves time in the breeding cycle and can help improve selection intensity and accuracy when combined with performance data. GP has many applications including a role in exploiting novel allelic variation. For instance, Crossa et al. (2017) reported that application of GP during pre-breeding can accelerate the identification and flow of genes from genebank accessions into elite lines.

In chickpea, GP holds potential to increase the rate of genetic gain in breeding programs. This has been demonstrated in chickpea for yield related traits, however, the prediction accuracies must be improved for selection implementation in breeding programmes (Roorkiwal et al. 2016; Roorkiwal et al. 2018). GP has been effectively applied in other crops including sorghum (Velazco et al. 2019), wheat (Montesinos-Lopez et al. 2019) and barley (Bhatta et al. 2020). To achieve this for chickpea, the accuracy of the GP model is the most important consideration for successful implementation. Accuracy of GP depends on the magnitude of the trait heritability and the size of the reference population. Using a multi-genetic background population and multivariate GP models can also increase prediction accuracy (Crossa et al. 2017).

In drought and heat prone environments, the heritability of yield traits is typically lower compared to environments where resources are not limiting. This source of variation and lower heritability, impacts on prediction accuracy. To counter this, traits correlated with yield (secondary traits) may be included in the prediction model to increase prediction accuracy (Rutkoski et al. 2016; Pszczola et al. 2013). For instance, key physiological traits underpinning performance may be incorporated into the prediction framework (Watson et al. 2019; Rutkoski et al. 2016). This strategy may be employed to enhance GP for chickpea performance in drought and heat environments (Santantonio et al. 2020). Traits such as those outlined in this review, particularly those that can be phenotyped in high-throughput platforms, could be targeted for inclusion in GP frameworks. To implement this successfully, an understanding of trait value in the target environment is critical to prioritise trait selection.

## Conclusion

Drought and heat stress are the major abiotic stresses impacting chickpea yield globally. It is clear, that climate variability and erratic rainfall distribution will worsen, and the effects of such scenarios will challenge crop productivity. Fortunately, previous physiological and genetic studies in chickpea have provided a strong foundation for future crop improvement. While good progress has been made to identify traits important for yield in drought and heat environments the underlying mechanisms are yet to

be fully elucidated. To create novel varieties that are able to balance resource capture and use, a deeper understanding of the relationship between canopy development and root architecture is required. The availability of genomic resources and high-throughput phenotyping platforms presents opportunities to accelerate the design of new chickpea varieties better suited to target environments. A multidisciplinary team is required to implement an integrated approach incorporating a strong understanding of physiological mechanisms driving adaptation in target environments, the latest phenotyping platforms, utilization of genebank resources, trait dissection and breeding technologies.

## Declarations

## Conflicts of interest

Authors declare no conflicts of interests

## Authors' Contributions

OJA, LTH and MRS conceived and designed the manuscript. RC, KVF, JK, SC, and RKV contributed additional concepts and academic expertise. All authors reviewed the article and approved the submitted version.

## Acknowledgments

OJA was supported by a UQ Research Training Scholarship

## Data availability

Data used to perform literature analysis is available in the Scopus database ([www.scopus.com](http://www.scopus.com)).

## References

1. Abbo S, Shtienberg D, Lichtenzveig J, Lev-Yadun S, Gopher A (2003) The chickpea, summer cropping, and a new model for pulse domestication in the ancient near east. *The Quarterly review of biology* 78 (4):435-448. doi:10.1086/378927
2. Afsharyan NP, Sannemann W, Leon J, Ballvora A (2020) Effect of epistasis and environment on flowering time in barley reveals a novel flowering-delaying QTL allele. *J Exp Bot* 71 (3):893-906. doi:10.1093/jxb/erz477
3. Asif MA, Garcia M, Tilbrook J, Brien C, Dowling K, Berger B, Schilling RK, Short L, Trittermann C, Gilliham M, Fleury D, Roy SJ, Pearson AS (2021) Identification of salt tolerance QTL in a wheat RIL

- mapping population using destructive and non-destructive phenotyping %J *Functional Plant Biology*. 48 (2):131-140. doi:<https://doi.org/10.1071/FP20167>
4. Attri H (2018) Evaluation of F4 derived F5 MaGIC population for yield and its contributing traits in chickpea.
  5. Bandillo N, Raghavan C, Muyco PA, Sevilla MA, Lobina IT, Dilla-Ermita CJ, Tung CW, McCouch S, Thomson M, Mauleon R, Singh RK, Gregorio G, Redona E, Leung H (2013) Multi-parent advanced generation inter-cross (MAGIC) populations in rice: progress and potential for genetics research and breeding. *Rice (N Y)* 6 (1):11. doi:10.1186/1939-8433-6-11
  6. Bari A, Damania AB, Mackay M, Dayanandan S (2016) Applied Mathematics and Omics to Assess Crop Genetic Resources for Climate Change Adaptive Traits.
  7. Bari A, Street K, Mackay M, Endresen DTF, De Pauw E, Amri A (2012) Focused identification of germplasm strategy (FIGS) detects wheat stem rust resistance linked to environmental variables. *Genetic Resources and Crop Evolution* 59 (7):1465-1481. doi:10.1007/s10722-011-9775-5
  8. Bharadwaj C, Tripathi S, Soren KR, Thudi M, Singh RK, Sheoran S, Roorkiwal M, Patil BS, Chitikineni A, Palakurthi R, Vemula A, Rathore A, Kumar Y, Chaturvedi SK, Mondal B, Shanmugavadivel PS, Srivastava AK, Dixit GP, Singh NP, Varshney RK (2021) Introgression of “QTL-hotspot” region enhances drought tolerance and grain yield in three elite chickpea cultivars. 14 (1):e20076. doi:<https://doi.org/10.1002/tpg2.20076>
  9. Bhatta M, Gutierrez L, Cammarota L, Cardozo F, German S, Gomez-Guerrero B, Pardo MF, Lanaro V, Sayas M, Castro AJ (2020) Multi-trait Genomic Prediction Model Increased the Predictive Ability for Agronomic and Malting Quality Traits in Barley (*Hordeum vulgare* L.). *G3 (Bethesda)* 10 (3):1113-1124. doi:10.1534/g3.119.400968
  10. Bouteille M, Rolland G, Balsera C, Loudet O, Muller B (2012) Disentangling the intertwined genetic bases of root and shoot growth in *Arabidopsis*. *PLoS One* 7 (2):e32319. doi:10.1371/journal.pone.0032319
  11. Canci H, Toker C (2009) Evaluation of Yield Criteria for Drought and Heat Resistance in Chickpea (*Cicer arietinum* L.). *Journal of Agronomy and Crop Science* 195 (1):47-54. doi:10.1111/j.1439-037X.2008.00345.x
  12. Carrão H, Naumann G, Barbosa P (2016) Mapping global patterns of drought risk: An empirical framework based on sub-national estimates of hazard, exposure and vulnerability. *Global Environmental Change* 39:108-124. doi:10.1016/j.gloenvcha.2016.04.012
  13. Chandora R, Gayacharan, Shekhawat N, Malhotra N (2020) Chapter 3 - Chickpea genetic resources: collection, conservation, characterization, and maintenance. In: Singh M (ed) *Chickpea: Crop Wild Relatives for Enhancing Genetic Gains*. Academic Press, pp 37-61. doi:<https://doi.org/10.1016/B978-0-12-818299-4.00003-8>
  14. Charmet G, Tran LG, Auzanneau J, Rincant R, Bouchet S (2020) BWGS: A R package for genomic selection and its application to a wheat breeding programme. *PLoS One* 15 (4):e0222733. doi:10.1371/journal.pone.0222733

15. Chen Y, Ghanem ME, Siddique KH (2017) Characterising root trait variability in chickpea (*Cicer arietinum* L.) germplasm. *J Exp Bot* 68 (8):1987-1999. doi:10.1093/jxb/erw368
16. Cobb JN, Declerck G, Greenberg A, Clark R, McCouch S (2013) Next-generation phenotyping: requirements and strategies for enhancing our understanding of genotype-phenotype relationships and its relevance to crop improvement. *Theor Appl Genet* 126 (4):867-887. doi:10.1007/s00122-013-2066-0
17. Cooper M, Tang T, Gho C, Hart T, Hammer G, Messina C (2020) Integrating genetic gain and gap analysis to predict improvements in crop productivity. *Crop Science* 60 (2):582-604. doi:10.1002/csc2.20109
18. Crossa J, Perez-Rodriguez P, Cuevas J, Montesinos-Lopez O, Jarquin D, de Los Campos G, Burgueno J, Gonzalez-Camacho JM, Perez-Elizalde S, Beyene Y, Dreisigacker S, Singh R, Zhang X, Gowda M, Roorkiwal M, Rutkoski J, Varshney RK (2017) Genomic Selection in Plant Breeding: Methods, Models, and Perspectives. *Trends Plant Sci* 22 (11):961-975. doi:10.1016/j.tplants.2017.08.011
19. Daryanto S, Wang L, Jacinthe PA (2015) Global Synthesis of Drought Effects on Food Legume Production. *PLoS One* 10 (6):e0127401. doi:10.1371/journal.pone.0127401
20. de Castro AI, Rallo P, Suarez MP, Torres-Sanchez J, Casanova L, Jimenez-Brenes FM, Morales-Sillero A, Jimenez MR, Lopez-Granados F (2019) High-Throughput System for the Early Quantification of Major Architectural Traits in Olive Breeding Trials Using UAV Images and OBIA Techniques. *Front Plant Sci* 10:1472. doi:10.3389/fpls.2019.01472
21. Deery D, Jimenez-Berni J, Jones H, Sirault X, Furbank R (2014) Proximal Remote Sensing Buggies and Potential Applications for Field-Based Phenotyping. *4* (3):349-379
22. Deery DM, Rebetzke GJ, Jimenez-Berni JA, James RA, Condon AG, Bovill WD, Hutchinson P, Scarrow J, Davy R, Furbank RT (2016) Methodology for High-Throughput Field Phenotyping of Canopy Temperature Using Airborne Thermography. *Frontiers in plant science* 7:1808. doi:10.3389/fpls.2016.01808
23. Deng W, Casao MC, Wang P, Sato K, Hayes PM, Finnegan EJ, Trevaskis B (2015) Direct links between the vernalization response and other key traits of cereal crops. *Nat Commun* 6:5882. doi:10.1038/ncomms6882
24. Devasirvatham V, Gaur PM, Mallikarjuna N, Raju TN, Trethowan RM, Tan DKY (2013) Reproductive biology of chickpea response to heat stress in the field is associated with the performance in controlled environments. *Field Crops Research* 142:9-19. doi:10.1016/j.fcr.2012.11.011
25. Devasirvatham V, Tan DKY, Gaur PM, Raju TN, Trethowan RM (2012) High temperature tolerance in chickpea and its implications for plant improvement. *Crop and Pasture Science* 63 (5). doi:10.1071/cp11218
26. El Bouhssini M, Ogbonnaya FC, Chen M, Lhaloui S, Rihawi F, Dabbous A (2012) Sources of resistance in primary synthetic hexaploid wheat (*Triticum aestivum* L.) to insect pests: Hessian fly, Russian wheat aphid and Sunn pest in the fertile crescent. *Genetic Resources and Crop Evolution* 60 (2):621-627. doi:10.1007/s10722-012-9861-3

27. Emma Huang B, Clifford D, Cavanagh C (2013) Selecting subsets of genotyped experimental populations for phenotyping to maximize genetic diversity. *Theor Appl Genet* 126 (2):379-388. doi:10.1007/s00122-012-1986-4
28. Fang X, Turner NC, Yan G, Li F, Siddique KH (2010) Flower numbers, pod production, pollen viability, and pistil function are reduced and flower and pod abortion increased in chickpea (*Cicer arietinum* L.) under terminal drought. *J Exp Bot* 61 (2):335-345. doi:10.1093/jxb/erp307
29. Foyer CH, Lam H-M, Nguyen HT, Siddique KHM, Varshney RK, Colmer TD, Cowling W, Bramley H, Mori TA, Hodgson JM, Cooper JW, Miller AJ, Kunert K, Vorster J, Cullis C, Ozga JA, Wahlqvist ML, Liang Y, Shou H, Shi K, Yu J, Fodor N, Kaiser BN, Wong F-L, Valliyodan B, Conside MJ (2016) Neglecting legumes has compromised human health and sustainable food production. *Nature Plants* 2 (8):16112. doi:10.1038/nplants.2016.112
30. Gaur PM, Krishnamurthy L, Kashiwagi J (2008) Improving drought-avoidance root traits in chickpea (*Cicer arietinum* L.) -current status of research at ICRISAT. *Plant Production Science* 11 (1):3-11. doi:10.1626/pps.11.3
31. Gaur PM, Samineni S, Thudi M, Tripathi S, Sajja SB, Jayalakshmi V, Mannur DM, Vijayakumar AG, Ganga Rao NVPR, Ojiewo C, Fikre A, Kimurto P, Kileo RO, Girma N, Chaturvedi SK, Varshney Rajeev K, Dixit GP, Link W (2018) Integrated breeding approaches for improving drought and heat adaptation in chickpea (*Cicer arietinum* L.). *Plant Breeding* 138 (4):389-400. doi:10.1111/pbr.12641
32. Gaur PM, Samineni S, Thudi M, Tripathi S, Sajja SB, Jayalakshmi V, Mannur DM, Vijayakumar AG, Rao N, Ojiewo C, Fikre A, Kimurto P, Kileo RO, Girma N, Chaturvedi SK, Varshney RK, Dixit GP (2019) Integrated breeding approaches for improving drought and heat adaptation in chickpea (*Cicer arietinum* L.). *Plant Breeding* 138 (4):389-400. doi:10.1111/pbr.12641
33. Glaszmann JC, Kilian B, Upadhyaya HD, Varshney RK (2010) Accessing genetic diversity for crop improvement. *Curr Opin Plant Biol* 13 (2):167-173. doi:10.1016/j.pbi.2010.01.004
34. Gupta S, Kumar T, Verma S, Bharadwaj C, Bhatia S (2015) Development of gene-based markers for use in construction of the chickpea (*Cicer arietinum* L.) genetic linkage map and identification of QTLs associated with seed weight and plant height. *Mol Biol Rep* 42 (11):1571-1580. doi:10.1007/s11033-015-3925-3
35. Hajjarpoor A, Vadez V, Soltani A, Gaur P, Whitbread A, Suresh Babu D, Gumma MK, Diancoumba M, Kholová J (2018) Characterization of the main chickpea cropping systems in India using a yield gap analysis approach. *Field Crops Research* 223:93-104. doi:<https://doi.org/10.1016/j.fcr.2018.03.023>
36. Heffner EL, Sorrells ME, Jannink J-L (2009) Genomic Selection for Crop Improvement. *Crop Science* 49 (1):1-12. doi:10.2135/cropsci2008.08.0512
37. Hemshrot A, Poets AM, Tyagi P, Lei L, Carter CK, Hirsch CN, Li L, Brown-Guedira G, Morrell PL, Muehlbauer GJ, Smith KP (2019) Development of a Multiparent Population for Genetic Mapping and Allele Discovery in Six-Row Barley. *Genetics* 213 (2):595-613. doi:10.1534/genetics.119.302046
38. Holland JB (2015) MAGIC maize: a new resource for plant genetics. *Genome Biol* 16:163. doi:10.1186/s13059-015-0713-2

39. Honsdorf N, March TJ, Berger B, Tester M, Pillen K (2014) High-Throughput Phenotyping to Detect Drought Tolerance QTL in Wild Barley Introgression Lines. *PLOS ONE* 9 (5):e97047. doi:10.1371/journal.pone.0097047
40. Huang BE, Verbyla KL, Verbyla AP, Raghavan C, Singh VK, Gaur P, Leung H, Varshney RK, Cavanagh CR (2015) MAGIC populations in crops: current status and future prospects. *Theor Appl Genet* 128 (6):999-1017. doi:10.1007/s00122-015-2506-0
41. Huynh BL, Ehlers JD, Huang BE, Munoz-Amatriain M, Lonardi S, Santos JRP, Ndeve A, Batiemo BJ, Boukar O, Cisse N, Drabo I, Fatokun C, Kusi F, Agyare RY, Guo YN, Herniter I, Lo S, Wanamaker SI, Xu S, Close TJ, Roberts PA (2018) A multi-parent advanced generation inter-cross (MAGIC) population for genetic analysis and improvement of cowpea (*Vigna unguiculata* L. Walp.). *Plant J* 93 (6):1129-1142. doi:10.1111/tpj.13827
42. Jaganathan D, Thudi M, Kale S, Azam S, Roorkiwal M, Gaur PM, Kishor PB, Nguyen H, Sutton T, Varshney RK (2015) Genotyping-by-sequencing based intra-specific genetic map refines a "QTL-hotspot" region for drought tolerance in chickpea. *Mol Genet Genomics* 290 (2):559-571. doi:10.1007/s00438-014-0932-3
43. Jang G, Kim J, Yu J-K, Kim H-J, Kim Y, Kim D-W, Kim K-H, Lee CW, Chung YS (2020) Review: Cost-Effective Unmanned Aerial Vehicle (UAV) Platform for Field Plant Breeding Application. *Remote Sensing* 12 (6). doi:10.3390/rs12060998
44. Jannink J-LJG (2007) Identifying quantitative trait locus by genetic background interactions in association studies. *176* (1):553-561
45. Kalra N, Chakraborty D, Sharma A, Rai HK, Jolly M, Chander S, Kumar PR, Bhadraray S, Barman D, Mittal RB, Lal M, Sehgal M (2008) Effect of increasing temperature on yield of some winter crops in northwest India. *Current Science* 94 (1):82-88
46. Kashiwagi J, Krishnamurthy L, Purushothaman R, Upadhyaya HD, Gaur PM, Gowda CLL, Ito O, Varshney RK (2015) Scope for improvement of yield under drought through the root traits in chickpea (*Cicer arietinum* L.). *Field Crops Research* 170:47-54. doi:10.1016/j.fcr.2014.10.003
47. Kashiwagi J, Krishnamurthy L, Upadhyaya HD, Krishna H, Chandra S, Vadez V, Serraj R (2006) Genetic variability of drought-avoidance root traits in the mini-core germplasm collection of chickpea (*Cicer arietinum* L.). *Euphytica* 146 (3):213-222. doi:10.1007/s10681-005-9007-1
48. Kashiwagi J, Krishnamurthy L, Upadhyaya HD, Krishna H, Chandra S, Vadez V, Serraj RJE (2005) Genetic variability of drought-avoidance root traits in the mini-core germplasm collection of chickpea (*Cicer arietinum* L.). *146* (3):213-222
49. Kell DBJAoB (2011) Breeding crop plants with deep roots: their role in sustainable carbon, nutrient and water sequestration. *108* (3):407-418
50. Kholova J, Urban MO, Cock J, Arcos J, Arnaud E, Aytekin D, Azevedo V, Barnes AP, Ceccarelli S, Chavarriaga P, Cobb JN, Connor D, Cooper M, Craufurd P, Debouck D, Fungo R, Grando S, Hammer GL, Jara CE, Messina C, Mosquera G, Nchanji E, Ng EH, Prager S, Sankaran S, Selvaraj M, Tardieu F,

- Thornton P, Valdes-Gutierrez SP, van Etten J, Wenzl P, Xu Y (2021) In pursuit of a better world: crop improvement and the CGIAR. *J Exp Bot* 72 (14):5158-5179. doi:10.1093/jxb/erab226
51. Krishnamurthy L, Kashiwagi J, Upadhyaya H, Serraj RJIC, Newsletter P (2003) Genetic diversity of drought-avoidance root traits in the mini-core germplasm collection of chickpea. (10):21-24
52. Kumar J, van Rheenen H (2000) Brief communication. A major gene for time of flowering in chickpea. *Journal of Heredity* 91 (1):67-68. doi:10.1093/jhered/91.1.67 %J Journal of Heredity
53. Kushwah A, Gupta S, Bindra S, Johal N, Singh I, Bharadwaj C, Dixit GP, Gaur PM, Nayyar H, Singh S (2020) Chapter 6 - Gene pyramiding and multiple character breeding. In: Singh M (ed) *Chickpea: Crop Wild Relatives for Enhancing Genetic Gains*. Academic Press, pp 131-165.  
doi:<https://doi.org/10.1016/B978-0-12-818299-4.00006-3>
54. Ladejobi O, Elderfield J, Gardner KA, Gaynor RC, Hickey J, Hibberd JM, Mackay IJ, Bentley AR (2016) Maximizing the potential of multi-parental crop populations. *Appl Transl Genom* 11:9-17.  
doi:10.1016/j.atg.2016.10.002
55. Lamaoui M, Jemo M, Datla R, Bekkaoui F (2018) Heat and Drought Stresses in Crops and Approaches for Their Mitigation. *Front Chem* 6:26. doi:10.3389/fchem.2018.00026
56. Leport L, Turner NC, Davies SL, Siddique KHM (2006) Variation in pod production and abortion among chickpea cultivars under terminal drought. *European Journal of Agronomy* 24 (3):236-246.  
doi:10.1016/j.eja.2005.08.005
57. Lopes MS, Reynolds MPJFPB (2010) Partitioning of assimilates to deeper roots is associated with cooler canopies and increased yield under drought in wheat. 37 (2):147-156
58. Ludlow M, Muchow RJAia (1990) A critical evaluation of traits for improving crop yields in water-limited environments. 43:107-153
59. Massel K, Lam Y, Wong ACS, Hickey LT, Borrell AK, Godwin ID (2021) Hotter, drier, CRISPR: the latest edit on climate change. *Theoretical and Applied Genetics* 134 (6):1691-1709. doi:10.1007/s00122-020-03764-0
60. Mathew B, Leon J, Sannemann W, Sillanpaa MJ (2018) Detection of Epistasis for Flowering Time Using Bayesian Multilocus Estimation in a Barley MAGIC Population. *Genetics* 208 (2):525-536.  
doi:10.1534/genetics.117.300546
61. Maurel C, Simonneau T, Sutka M (2010) The significance of roots as hydraulic rheostats. *J Exp Bot* 61 (12):3191-3198. doi:10.1093/jxb/erq150
62. Messina C, Hammer G, Dong Z, Podlich D, Cooper M (2009) Chapter 10 - Modelling Crop Improvement in a G×E×M Framework via Gene–Trait–Phenotype Relationships. In: Sadras V, Calderini D (eds) *Crop Physiology*. Academic Press, San Diego, pp 235-581.  
doi:<https://doi.org/10.1016/B978-0-12-374431-9.00010-4>
63. Meuwissen T, Hayes B, Goddard M (2016) Genomic selection: A paradigm shift in animal breeding. *Animal Frontiers* 6 (1):6-14. doi:10.2527/af.2016-0002
64. Meuwissen TH, Hayes BJ, Goddard ME (2001) Prediction of total genetic value using genome-wide dense marker maps. *Genetics* 157 (4):1819-1829

65. Montesinos-Lopez OA, Montesinos-Lopez A, Tuberosa R, Maccaferri M, Sciara G, Ammar K, Crossa J (2019) Multi-Trait, Multi-Environment Genomic Prediction of Durum Wheat With Genomic Best Linear Unbiased Predictor and Deep Learning Methods. *Front Plant Sci* 10:1311. doi:10.3389/fpls.2019.01311
66. Muehlbauer FJ, Sarker A (2017) Economic Importance of Chickpea: Production, Value, and World Trade. In: Varshney RK, Thudi M, Muehlbauer F (eds) *The Chickpea Genome*. Springer International Publishing, Cham, pp 5-12. doi:10.1007/978-3-319-66117-9\_2
67. Ongom PO, Ejeta G (2018) Mating Design and Genetic Structure of a Multi-Parent Advanced Generation Intercross (MAGIC) Population of Sorghum (*Sorghum bicolor* (L.) Moench). *G3 (Bethesda)* 8 (1):331-341. doi:10.1534/g3.117.300248
68. Paul PJ, Samineni S, Thudi M, Sajja SB, Rathore A, Das RR, Khan AW, Chaturvedi SK, Lavanya GR, Varshney RK, Gaur PM (2018) Molecular Mapping of QTLs for Heat Tolerance in Chickpea. *Int J Mol Sci* 19 (8). doi:10.3390/ijms19082166
69. Pinto RS, Reynolds MP (2015) Common genetic basis for canopy temperature depression under heat and drought stress associated with optimized root distribution in bread wheat. *Theor Appl Genet* 128 (4):575-585. doi:10.1007/s00122-015-2453-9
70. Pszczola M, Veerkamp R, De Haas Y, Wall E, Strabel T, Calus MJA (2013) Effect of predictor traits on accuracy of genomic breeding values for feed intake based on a limited cow reference population. *J Anim Sci* 117 (11):1759-1768
71. Purushothaman R, Krishnamurthy L, Upadhyaya HD, Vadez V, Varshney RK (2017) Genotypic variation in soil water use and root distribution and their implications for drought tolerance in chickpea. *Functional Plant Biology* 44 (2). doi:10.1071/fp16154
72. Rahaman MM, Chen D, Gillani Z, Klukas C, Chen M (2015) Advanced phenotyping and phenotype data analysis for the study of plant growth and development. *Front Plant Sci* 6:619. doi:10.3389/fpls.2015.00619
73. Ramamoorthy P, Lakshmanan K, Upadhyaya HD, Vadez V, Varshney RK (2017) Root traits confer grain yield advantages under terminal drought in chickpea (*Cicer arietinum* L.). *Field Crops Res* 201:146-161. doi:10.1016/j.fcr.2016.11.004
74. Rani A, Devi P, Jha UC, Sharma KD, Siddique KHM, Nayyar H (2020) Developing Climate-Resilient Chickpea Involving Physiological and Molecular Approaches With a Focus on Temperature and Drought Stresses. *Frontiers in Plant Science* 10. doi:10.3389/fpls.2019.01759
75. Ratnakumar P, Vadez V, Nigam SN, Krishnamurthy L (2009) Assessment of transpiration efficiency in peanut (*Arachis hypogaea* L.) under drought using a lysimetric system. *Plant Biol (Stuttg)* 11 Suppl 1:124-130. doi:10.1111/j.1438-8677.2009.00260.x
76. Ren W, Gong X, Li K, Zhang H, Chen F, Pan Q (2020) Recombination Pattern Characterization via Simulation Using Different Maize Populations. *Int J Mol Sci* 21 (6). doi:10.3390/ijms21062222
77. Reynolds M, Dreccer F, Trethowan R (2006) Drought-adaptive traits derived from wheat wild relatives and landraces. *Journal of Experimental Botany* 58 (2):177-186. doi:10.1093/jxb/erl250


78. Roorkiwal M, Bharadwaj C, Barmukh R, Dixit GP, Thudi M, Gaur PM, Chaturvedi SK, Fikre A, Hamwiah A, Kumar S, Sachdeva S, Ojiewo CO, Tar'an B, Wordofa NG, Singh NP, Siddique KHM, Varshney RK (2020) Integrating genomics for chickpea improvement: achievements and opportunities. *Theor Appl Genet* 133 (5):1703-1720. doi:10.1007/s00122-020-03584-2
79. Roorkiwal M, Jarquin D, Singh MK, Gaur PM, Bharadwaj C, Rathore A, Howard R, Srinivasan S, Jain A, Garg V, Kale S, Chitikineni A, Tripathi S, Jones E, Robbins KR, Crossa J, Varshney RK (2018) Genomic-enabled prediction models using multi-environment trials to estimate the effect of genotype x environment interaction on prediction accuracy in chickpea. *Sci Rep* 8 (1):11701. doi:10.1038/s41598-018-30027-2
80. Roorkiwal M, Rathore A, Das RR, Singh MK, Jain A, Srinivasan S, Gaur PM, Chellapilla B, Tripathi S, Li Y, Hickey JM, Lorenz A, Sutton T, Crossa J, Jannink J-L, Varshney RK (2016) Genome-Enabled Prediction Models for Yield Related Traits in Chickpea. *7* (1666). doi:10.3389/fpls.2016.01666
81. Rutkoski J, Poland J, Mondal S, Autrique E, Perez LG, Crossa J, Reynolds M, Singh R (2016) Canopy Temperature and Vegetation Indices from High-Throughput Phenotyping Improve Accuracy of Pedigree and Genomic Selection for Grain Yield in Wheat. *G3 (Bethesda)* 6 (9):2799-2808. doi:10.1534/g3.116.032888
82. Santantonio N, Atanda SA, Beyene Y, Varshney RK, Olsen M, Jones E, Roorkiwal M, Gowda M, Bharadwaj C, Gaur PM, Zhang X, Dreher K, Ayala-Hernández C, Crossa J, Pérez-Rodríguez P, Rathore A, Gao SY, McCouch S, Robbins KR (2020) Strategies for Effective Use of Genomic Information in Crop Breeding Programs Serving Africa and South Asia. *11* (353). doi:10.3389/fpls.2020.00353
83. Saxena NP, Krishnamurthy L, Johansen CJCS (1993) Registration of a drought-resistant chickpea germplasm. *33* (6):1424-1424
84. Serraj R, Krishnamurthy L, Kashiwagi J, Kumar J, Chandra S, Crouch JH (2004) Variation in root traits of chickpea (*Cicer arietinum* L.) grown under terminal drought. *Field Crops Research* 88 (2):115-127. doi:<https://doi.org/10.1016/j.fcr.2003.12.001>
85. Sherman B, Henry RJ (2020) The Nagoya Protocol and historical collections of plants. *Nat Plants* 6 (5):430-432. doi:10.1038/s41477-020-0657-8
86. Siddique KHM, Johansen C, Turner NC, Jeuffroy M-H, Hashem A, Sakar D, Gan Y, Alghamdi SS (2011) Innovations in agronomy for food legumes. A review. *Agronomy for Sustainable Development* 32 (1):45-64. doi:10.1007/s13593-011-0021-5
87. Singh K, Bejiga G, Saxena M, Singh MJJoA, Science C (1995) Transferability of chickpea selection indices from normal to drought-prone growing conditions in a Mediterranean environment. *175* (1):57-63
88. Singh R, Sharma P, Varshney RK, Sharma SK, Singh NK (2008) Chickpea improvement: role of wild species and genetic markers. *Biotechnol Genet Eng Rev* 25:267-313. doi:10.5661/bger-25-267
89. Singh VK, Khan AW, Jaganathan D, Thudi M, Roorkiwal M, Takagi H, Garg V, Kumar V, Chitikineni A, Gaur PM, Sutton T, Terauchi R, Varshney RK (2016) QTL-seq for rapid identification of candidate

- genes for 100-seed weight and root/total plant dry weight ratio under rainfed conditions in chickpea. *Plant Biotechnol J* 14 (11):2110-2119. doi:10.1111/pbi.12567
90. Sivasakthi K, Tharanya M, Kholova J, Wangari Muriuki R, Thirunalasundari T, Vadez V (2017) Chickpea Genotypes Contrasting for Vigor and Canopy Conductance Also Differ in Their Dependence on Different Water Transport Pathways. *Front Plant Sci* 8:1663. doi:10.3389/fpls.2017.01663
91. Sivasakthi K, Thudi M, Tharanya M, Kale SM, Kholova J, Halime MH, Jaganathan D, Baddam R, Thirunalasundari T, Gaur PM, Varshney RK, Vadez V (2018) Plant vigour QTLs co-map with an earlier reported QTL hotspot for drought tolerance while water saving QTLs map in other regions of the chickpea genome. *BMC Plant Biol* 18 (1):29. doi:10.1186/s12870-018-1245-1
92. Smith D, Hinz H, Mulema J, Weyl P, Ryan MJ (2018) Biological control and the Nagoya Protocol on access and benefit sharing – a case of effective due diligence. *Biocontrol Science and Technology* 28 (10):914-926. doi:10.1080/09583157.2018.1460317
93. Smith DT, Potgieter AB, Chapman SC (2021) Scaling up high-throughput phenotyping for abiotic stress selection in the field. *Theoretical and Applied Genetics* 134 (6):1845-1866. doi:10.1007/s00122-021-03864-5
94. Tardieu F, Simonneau T, Muller B (2018) The Physiological Basis of Drought Tolerance in Crop Plants: A Scenario-Dependent Probabilistic Approach. *Annu Rev Plant Biol* 69:733-759. doi:10.1146/annurev-arplant-042817-040218
95. Tester M, Langridge P (2010) Breeding technologies to increase crop production in a changing world. *Science (New York, NY)* 327 (5967):818-822. doi:10.1126/science.1183700
96. Thudi M, Palakurthi R, Schnable JC, Chitikineni A, Dreisigacker S, Mace E, Srivastava RK, Satyavathi CT, Odeny D, Tiwari VK, Lam HM, Hong YB, Singh VK, Li G, Xu Y, Chen X, Kaila S, Nguyen H, Sivasankar S, Jackson SA, Close TJ, Shubo W, Varshney RK (2021) Genomic resources in plant breeding for sustainable agriculture. *J Plant Physiol* 257:153351. doi:10.1016/j.jplph.2020.153351
97. Thudi M, Upadhyaya HD, Rathore A, Gaur PM, Krishnamurthy L, Roorkiwal M, Nayak SN, Chaturvedi SK, Basu PS, Gangarao NV, Fikre A, Kimurto P, Sharma PC, Sheshashayee MS, Tobita S, Kashiwagi J, Ito O, Killian A, Varshney RK (2014) Genetic dissection of drought and heat tolerance in chickpea through genome-wide and candidate gene-based association mapping approaches. *PLoS One* 9 (5):e96758. doi:10.1371/journal.pone.0096758
98. Upadhyaya HD, Ortiz R (2001) A mini core subset for capturing diversity and promoting utilization of chickpea genetic resources in crop improvement. *Theoretical and Applied Genetics* 102 (8):1292-1298. doi:10.1007/s00122-001-0556-y
99. Vadez V (2014) Root hydraulics: The forgotten side of roots in drought adaptation. *Field Crops Research* 165:15-24. doi:<https://doi.org/10.1016/j.fcr.2014.03.017>
100. Vadez V, Kholova J, Hummel G, Zhokhavets U, Gupta SK, Hash CT (2015) LeasyScan: a novel concept combining 3D imaging and lysimetry for high-throughput phenotyping of traits controlling plant water budget. *J Exp Bot* 66 (18):5581-5593. doi:10.1093/jxb/erv251

101. Vadez V, Kholova J, Medina S, Kakker A, Anderberg H (2014) Transpiration efficiency: new insights into an old story. *J Exp Bot* 65 (21):6141-6153. doi:10.1093/jxb/eru040
102. Vadez V, Krishnamurthy L, Serraj R, Gaur PM, Upadhyaya HD, Hoisington DA, Varshney RK, Turner NC, Siddique KHM (2007) Large variation in salinity tolerance in chickpea is explained by differences in sensitivity at the reproductive stage. *Field Crops Research* 104 (1):123-129. doi:<https://doi.org/10.1016/j.fcr.2007.05.014>
103. Vadez V, Rao S, Kholova J, Krishnamurthy L, Kashiwagi J, Ratnakumar P, Sharma K, Bhatnagar-Mathur P, Basu PJJofL (2008) Root research for drought tolerance in legumes: Quo vadis? *J Exp Bot* 21 (2):77-85
104. Varshney RK (2016) Exciting journey of 10 years from genomes to fields and markets: Some success stories of genomics-assisted breeding in chickpea, pigeonpea and groundnut. *Plant Sci* 242:98-107. doi:10.1016/j.plantsci.2015.09.009
105. Varshney RK, Bohra A, Yu J, Graner A, Zhang Q, Sorrells ME (2021) Designing Future Crops: Genomics-Assisted Breeding Comes of Age. *Trends in Plant Science* 26 (6):631-649. doi:10.1016/j.tplants.2021.03.010
106. Varshney RK, Gaur PM, Chamarthi SK, Krishnamurthy L, Tripathi S, Kashiwagi J, Samineni S, Singh VK, Thudi M, Jaganathan D (2013a) Fast-Track Introgression of “ for Root Traits and Other Drought Tolerance Traits in JG 11, an Elite and Leading Variety of Chickpea. *The Plant Genome* 6 (3). doi:10.3835/plantgenome2013.07.0022
107. Varshney RK, Graner A, Sorrells ME (2005) Genomics-assisted breeding for crop improvement. *Trends Plant Sci* 10 (12):621-630. doi:10.1016/j.tplants.2005.10.004
108. Varshney RK, Mohan SM, Gaur PM, Gangarao NV, Pandey MK, Bohra A, Sawargaonkar SL, Chitkineni A, Kimurto PK, Janila P, Saxena KB, Fikre A, Sharma M, Rathore A, Pratap A, Tripathi S, Datta S, Chaturvedi SK, Mallikarjuna N, Anuradha G, Babbar A, Choudhary AK, Mhase MB, Bharadwaj C, Mannur DM, Harer PN, Guo B, Liang X, Nadarajan N, Gowda CL (2013b) Achievements and prospects of genomics-assisted breeding in three legume crops of the semi-arid tropics. *Biotechnology advances* 31 (8):1120-1134. doi:10.1016/j.biotechadv.2013.01.001
109. Varshney RK, Song C, Saxena RK, Azam S, Yu S, Sharpe AG, Cannon S, Baek J, Rosen BD, Tar'an B, Millan T, Zhang X, Ramsay LD, Iwata A, Wang Y, Nelson W, Farmer AD, Gaur PM, Soderlund C, Penmetsa RV, Xu C, Bharti AK, He W, Winter P, Zhao S, Hane JK, Carrasquilla-Garcia N, Condie JA, Upadhyaya HD, Luo M-C, Thudi M, Gowda CLL, Singh NP, Lichtenzveig J, Gali KK, Rubio J, Nadarajan N, Dolezel J, Bansal KC, Xu X, Edwards D, Zhang G, Kahl G, Gil J, Singh KB, Datta SK, Jackson SA, Wang J, Cook DR (2013c) Draft genome sequence of chickpea (*Cicer arietinum*) provides a resource for trait improvement. *Nature Biotechnology* 31 (3):240-246. doi:10.1038/nbt.2491
110. Varshney RK, Thudi M, Nayak SN, Gaur PM, Kashiwagi J, Krishnamurthy L, Jaganathan D, Koppolu J, Bohra A, Tripathi S, Rathore A, Jukanti AK, Jayalakshmi V, Vemula A, Singh SJ, Yasin M, Sheshshayee MS, Viswanatha KP (2014) Genetic dissection of drought tolerance in chickpea (*Cicer arietinum* L.). *Theor Appl Genet* 127 (2):445-462. doi:10.1007/s00122-013-2230-6


111. Velazco JG, Jordan DR, Mace ES, Hunt CH, Malosetti M, van Eeuwijk FA (2019) Genomic Prediction of Grain Yield and Drought-Adaptation Capacity in Sorghum Is Enhanced by Multi-Trait Analysis. *Front Plant Sci* 10:997. doi:10.3389/fpls.2019.00997
112. Voss-Fels KP, Robinson H, Mudge SR, Richard C, Newman S, Wittkop B, Stahl A, Friedt W, Frisch M, Gabur I, Miller-Cooper A, Campbell BC, Kelly A, Fox G, Christopher J, Christopher M, Chenu K, Franckowiak J, Mace ES, Borrell AK, Eagles H, Jordan DR, Botella JR, Hammer G, Godwin ID, Trevaskis B, Snowdon RJ, Hickey LT (2018) VERNALIZATION1 Modulates Root System Architecture in Wheat and Barley. *Mol Plant* 11 (1):226-229. doi:10.1016/j.molp.2017.10.005
113. Wang X, Xu Y, Hu Z, Xu C (2018) Genomic selection methods for crop improvement: Current status and prospects. *The Crop Journal* 6 (4):330-340. doi:10.1016/j.cj.2018.03.001
114. Watson A, Hickey LT, Christopher J, Rutkoski J, Poland J, Hayes BJ (2019) Multivariate Genomic Selection and Potential of Rapid Indirect Selection with Speed Breeding in Spring Wheat. *Crop Science* 59 (5). doi:10.2135/cropsci2018.12.0757
115. Xiao Y, Liu H, Wu L, Warburton M, Yan J (2017) Genome-wide Association Studies in Maize: Praise and Stargaze. *Mol Plant* 10 (3):359-374. doi:10.1016/j.molp.2016.12.008
116. Zaman-Allah M, Jenkinson DM, Vadez V (2011) A conservative pattern of water use, rather than deep or profuse rooting, is critical for the terminal drought tolerance of chickpea. *J Exp Bot* 62 (12):4239-4252. doi:10.1093/jxb/err139
117. Zhang Z, Zhang X, Lin Z, Wang J, Xu M, Lai J, Yu J, Lin Z (2018) The genetic architecture of nodal root number in maize. *Plant J* 93 (6):1032-1044. doi:10.1111/tpj.13828

## Figures




**Figure 1**

Analysis of 365 independent articles research on heat and drought adaptation between 1970 and June 2020 sourced from Scopus. (A) Pie chart of the 365 articles showing the percentage of research articles focusing on drought alone, heat alone and combination of drought and heat. The blue colour represents the drought articles, red represents the heat articles, while the yellow represents the articles focusing on the combined effects of drought and heat (B) Text analysis result from Leximancer version 4.5 showing the number of times a specific word is used and ranked them according to the number of times they appeared in the titles, abstracts, and keywords.



**Figure 2**

Aboveground biomass associated with water use A) Profligate water use leading to yield penalty B) Conservative water use leading to yield benefit.



**Figure 3**

Schematic diagram of how genetic variation in gene bank material could be exploited, using pre-breeding techniques such as FIGS, multi-parental population with high throughput phenotyping platforms and techniques for variety development.