Development of a Lifestyle Intervention Program for Overweight or Obese Preconception, Pregnant and Postnatal Women: a Qualitative Study

Chee Wai Ku (✉ kucheewai@gmail.com)

Duke-NUS Medical School

Shu Hui Leow

University of Cambridge

Lay See Ong

Duke-NUS Medical School

Christina Erwin

Duke-NUS Medical School

Isabella Ong

National University of Singapore

Xiang Wen Ng

KK Women's and Children's Hospital

Jacinth JX Tan

Singapore Management University

Fabian Yap

KK Women's and Children's Hospital

Jerry Kok Yen Chan

KK Women's and Children's Hospital

See Ling Loy

KK Women's and Children's Hospital

Research Article

Keywords: lifestyle intervention program, overweight or obese preconception, pregnant and postnatal women, qualitative study, lifestyle habits

Posted Date: September 9th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-877807/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The time period before, during and after pregnancy represents a unique opportunity for interventions to cultivate sustained healthy lifestyle behaviors to improve the metabolic health of mothers and their offspring. However, the success of a lifestyle intervention is dependent on uptake and continued compliance. To identify enablers and barriers towards engagement with a lifestyle intervention, thematic analysis of 15 in-depth interviews with overweight or obese women in the preconception, pregnancy or postpartum periods was undertaken, using the integrated–Promoting Action on Research Implementation in Health Services (i-PARIHS) framework as a guide to systematically chart factors influencing adoption of a novel lifestyle intervention. Barrier factors include time constraints, poor baseline knowledge, family culture, food accessibility, and lack of relevant data sources. Enabling factors were motivation to be healthy for themselves and their offspring, family and social support, a holistic delivery platform providing desired information delivered at appropriate times, regular feedback, goal setting, and nudges. From the findings of this study, we propose components of an idealized lifestyle intervention including (i) taking a holistic life-course approach to education, (ii) using mobile health platforms to reduce barriers, provide personalized feedback and promote goal-setting, and (iii) health nudges to cultivate sustained lifestyle habits.

Introduction

Overweight or obese women in the reproductive age group have an increased lifetime risk of adverse metabolic sequelae, along with subfertility, adverse obstetric outcomes in the mother and long-term adverse health outcomes for the child [1–9]. In line with the Developmental Origins of Health and Disease (DOHaD) paradigm, where early-life developmental influences are associated with later-life health and disease, optimizing the health of the mother can achieve the greatest benefit for mother and child [10]. Healthy diet and physical activity are key factors in addressing the obesity pandemic and related health complications [11].

Various lifestyle interventions have been implemented in the preconception or pregnancy phases, which represent unique opportunities for cultivating sustained healthy lifestyle behaviors [12–16]. However, large randomized controlled studies of lifestyle interventions, such as dietary education, physical activity tracking, counselling workshops, and various combinations of these, have failed to achieve sustainable results [17, 18]. In particular, lifestyle interventions initiated during pregnancy have been unsuccessful in preventing gestational diabetes or large for gestational age babies [19]. Few weight loss interventions have been conducted for women before conception [20]. While some have been shown to improve fertility, they did not result in significant improvements in metabolic outcomes [21–23]. Although there is considerable research and clear consensus regarding the goals of effective lifestyle interventions [20], difficulty translating the evidence into clinical practice is widely documented. Key challenges to achieving outcomes in lifestyle interventions lie in successful engagement of participants and continued compliance with the intervention [20, 24]. Enablers and barriers to behavioral change may arise from family and community characteristics, the local setting, and by social, environmental and policy contexts

[13–16, 25, 26]. However, few studies have engaged with implementation science theories and integrated research findings to develop interventions that enable high engagement and sustained behavioral change. Hence, the optimal components of a lifestyle intervention for pregnancy that is acceptable, sustainable, and tailored to the concerns and needs of the target population needs to be determined.

The integrated – Promoting Action on Research Implementation in Health Services (i-PARIHS) framework supports successful implementation of evidence-based outcomes into real-world practice [27]. Specifically, it was designed to guide complex interventions in healthcare, providing a framework to systematically chart factors affecting implementation. By identifying barriers and enablers as perceived by users, it has previously been identified as a useful tool for iterative planning of the implementation of patient-reported outcome measures (PROM) into routine oncology care [28]. As the eventual goal of this study is to create a lifestyle intervention program targeted at women with high body mass index (BMI), utilizing this framework will help to identify both enabler and barrier factors that should be resolved when designing a successful behavioral change intervention. Hence, the i-PARIHS framework was applied as a guide to explore the perceived enablers and barriers influencing the adoption of a lifestyle intervention program for the overweight or obese among preconception, pregnant and postnatal women.

Methods

As the goal is to develop an intervention program that serves to accompany women throughout their preconception, pregnancy and postnatal journey, qualitative research was conducted as it provides medical researchers with deep insight into their patients' life-worlds [29]. Participants were recruited through a quota sampling method. In total, 15 in-depth interviews were conducted at KK Women's and Children's Hospital (KKH), Singapore, between September 2020 to December 2020. Participants were females between 21–40 years of age with BMI ≥ 25 kg/m². They were either in the preconception (n = 5), pregnancy (n = 5), or postnatal (n = 5) phases of their journey. Information needs and the concept of data saturation were used to guide the sample size determination [30].

Interview protocol

Data was gathered via two methods in each participant: by questionnaire and an in-depth interview session. Before the start of the interview session, participants filled out a short questionnaire that captured demographic data including age, ethnicity, BMI, smoking exposure, alcohol use, supplement use, educational attainment and employment status.

A semi-structured interview guide was used during the in-depth interview session to provide systematic coverage of topics, allowing flexibility to explore the responses of participants when needed. This guide was developed based on a preliminary literature review into the knowledge, attitudes and beliefs regarding nutrition and lifestyle behavior among overweight and obese women before, during and after pregnancy [13, 13–16, 25, 31–33]. The interview guide (Additional file 1) consisted of a standardized set of open-ended questions focusing on the knowledge, attitudes and perceptions of women towards a

lifestyle intervention program, followed by further probing questions. These aimed to find out the perceived importance of their own and their offspring's health; their baseline understanding of healthy behaviors, weight and health goals; motivators and barriers to achieving their goals; and acceptable dietary or other lifestyle interventions that they would be willing to undertake.

The in-depth interview sessions were conducted by three researchers who developed the interview guide. The interview sessions lasted around 60 minutes, were conducted in person using the English language, audio-recorded and transcribed verbatim. A team of research assistants transcribed the audio-recordings, which were mutually cross-checked for accuracy. Two independent researchers subsequently verified the transcripts for accuracy.

Analysis

The in-depth interview transcripts were analyzed inductively using thematic analysis [34]. All transcripts were analyzed using NVivo (release 1.4.1, QSR International), a qualitative data analysis software [35]. Two study team members coded the transcripts independently. Subsequently, these codes were then discussed between the two study team members, resulting in identified themes. The development of codes and themes was an iterative and reflexive process where the team members shared their perspectives. Any discordances were resolved through consensus discussion.

The identified themes were subsequently mapped onto the i-PARIHS framework, which has four key constructs at its core: *Recipient*, *Context*, *Innovation* and *Facilitation*. These constructs were identified to be important considerations for a successful implementation of research findings [36]. The *Recipient* construct refers to a person's characteristics that may affect implementation of the intervention [27]. The *Context* construct refers to non-people factors that can act to enable or constrain implementation, such as culture, leadership, the wider health system or organization [27]. Within this, we delved into how manipulating the environment surrounding a potential user may influence engagement with an intervention. The *Innovation* construct refers to the circumstantial interpretation of how the evidence should be applied to practice, which often involves tailoring to *Recipient* and *Context* factors [27]. The *Facilitation* construct refers to implementation processes that comprises of a facilitator (role) applying a process (a set of strategies and actions) [27]. As the facilitators apply the innovation on the recipients in the given context, each of the other constructs interact with the *Facilitation* construct.

Initial mapping was done independently by two study team members with discordance resolved through consensus discussions with a third study team member. Across the four i-PARIHS constructs, several barrier and enabler factors were identified. The finalized thematic map and its codes can be found in Additional file 2.

Results

1. Characteristics of the participants

Table 1 shows the participants' baseline characteristics. Of the 15 participants, majority were Chinese, had completed tertiary education, were employed full-time, did not smoke, rarely or were non-alcohol drinkers, and consumed health supplements. They had a mean age of 33.8 ± 5.1 years and BMI of 33.7 ± 5.7 kg/m².

Table 1
Characteristics of study participants

Demographic	Total (n = 15)
Age	
<30 years	3 (20%)
30–34 years	6 (40%)
≥35 years	6 (40%)
Ethnicity	
Chinese	9 (60%)
Malay	2 (13%)
Indian	4 (27%)
Highest Level of Education	
Secondary Education	5 (33%)
Tertiary Education	10 (67%)
Employment	
Not Employed	4 (27%)
Part-Time Employed	1 (7%)
Full-Time Employed	10 (67%)
Smoking status	
Never	11 (73%)
Have Stopped Smoking	3 (20%)
Currently Smoking	1 (7%)
Alcohol intake	
Never or less than 1 per month	12 (80%)
1–3 times per month	3 (20%)
Supplement intake	
No	3 (20%)
Yes	12 (80%)

2. Qualitative findings

Ten basic themes representing participants' perceived barriers and enablers to program participation and engagement were identified. They were organized into four main themes according to the constructs in the i-PARIHS framework, namely *Recipient*, *Context*, *Innovation* and *Facilitation*.

2.1. Perceived barriers

Recipient: **Time constraints**

Participants indicated that time pressure was a major challenge preventing them from engaging in healthy behaviors. They explained that the multiple roles and duties in their lives restrict them from having time to adopt healthy lifestyle habits such as exercising. *"As a mother, as a housewife, as a daughter-in-law, and as a person, I need to find time to do it all, to do everything. That is going to be very challenging"* (Interviewee 02). Another participant said: *"Taking care of the kids and working, we really (sic) didn't have time to exercise"* (Interviewee 06).

Recipient: **Lack of knowledge about impact of obesity on health**

Not all participants were aware of the health implications of being overweight and obese. *"Other than my weight, I don't think there are any other health concerns"* (Interviewee 14). Similar responses were given when participants were probed further on the potential health risks of obesity. *"Author: Do you have any thoughts about the impact of having a high BMI? Interviewee 02: Err no. Author: But specifically, do you know whether there are any negative effects? Interviewee 02: I don't think so."* Participants were also unaware about the implications of obesity on their offspring's health. *"I feel that despite being overweight and obese, the risk to baby is minimal to none"* (Interviewee 01).

Context: **Lack of relevant data sources**

Participants are not able to access data sources relevant to their context. They pointed out that currently available information on guiding maternal healthy living, which are not relatable to the local culture, hinder them from making lifestyle changes. *"There are apps out there that track and tell you what to expect in pregnancy, what foods to eat, and what exercises to do, but they are only representative of western countries. I cannot find anything that is applicable to the local context so that's why I'm not motivated to learn or follow the advice"* (Interviewee 07). Some participants commented that existing sources are mostly contextualized in western cultural settings, which are not easily adopted in the local context nor appealing to Asian women. *"I think it's because (these lifestyle apps are) are tailored to the angmoh^ context. Those apps are designed in western countries, (like) pumpkin soup (which) is not very Singaporean"* (Interviewee 08). (^Angmoh is a colloquial term referring to Caucasians, and may be used as an adjective that references western culture)

Context: **Family culture**

Participants mentioned that the traditional mindset from elder family members plays a major part in determining their daily lifestyle practices. *"I think the old people are heavily influenced by their parents'*

mindset. They tell me not to eat certain foods... Or to do certain things" (Interviewee 04). They expressed difficulty in communicating with their elderly in the family. *"My mother-in-law helped during my confinement period. We were totally lost. She assumed that what she did was the norm. But we have new products in the market now to help with childcare. She doesn't know about them and she doesn't believe in them. So it's a bit tough with a bit of a clash, (and) my husband had to be the middleman"* (Interviewee 07). Eating together is a social practice in Asia [37]. During social gatherings and activities, individuals may adopt different eating behaviors that are not aligned with healthy diets. *"...when we are not meeting with our families, it's probably easier to stick to the diet but when you meet during weekends, you're happy, and there's so much food, good food, to order in so we caved in many times"* (Interviewee 06). In addition, family members living in the same household often share meals and lifestyle habits which are difficult to modify individually, separate from other household members. *"To be honest, because I don't prepare my own food, I'm at the mercy of what my mom feels like cooking that day"* (Interviewee 09).

Context: **Food accessibility**

Finally, there is a lack of an enabling environment to support positive lifestyle and dietary changes. Accessible and convenient unhealthy food options is a factor that perpetuates unhealthy diets. *"because I'm late from work and I'm so hungry, because of the clinic schedule, we look at which (sic) shop doesn't have a queue. Old Chang Kee* for example is the fastest, so its tempting to just buy (from there)"* (Interviewee 13). (*Old Chang Kee is an established food retail chain in Singapore that sells deep fried snacks). This participant also added: *"...because every day when I come to work I'll buy a lot of breakfast snacks, like from McDonalds probably (sic). So I just get one meal and sometimes I'll order extra hash browns as well"* (Interviewee 13).

2.2. Perceived enablers

Recipient: **Motivation to change**

Participants expressed motivation to make a healthy change, with the desire to be in better physical health for their offspring(s) when they grow up. *"Having a baby motivates (me) a lot, because I want to be there to see them grow up and I want to lose weight for my four children"* (Interviewee 02). Another participant said: *"Yes, I can live for a long time (sic) to see my child get married (and) to have grandchildren"* (Interviewee 05). Participants were also motivated to change for their children. *"It's my child who changed my die. As my toddler joins me for meals, I (must) make sure the food is not fried, is not (sic) high in fat or seasoning, so I'm motivated by my child's need(s) to eat healthily."* (Interviewee 08).

Context: **Family and social support**

Within the immediate environment, participants reported that family and social support would be a strong enabler. In particular, partner engagement with lifestyle changes alongside the prospective mother enabled mutual motivation. *"We were at our fittest together and we both encourage each other. So we both motivate each other"* (Interviewee 06). One participant highlighted the importance of engaging their partner in the program. *"(The program) should not just reach out to the mother, but also reach out to the*

father. (It) will be helpful if the partner, the father, also knows (the content of the program)” (Interviewee 07). Support of the immediate family in the care of a young baby was also pivotal for new mothers to engage in healthy behaviors. *“My mother helps me to take care of the baby for about two hours, so that I can hit the gym or have a walk”* (Interviewee 04). Participants in our study also valued making social connections with peers to form a community of people going through similar experiences and provide mutual support. *“Mummies groups (sic) are good. Every pregnancy is different. Each baby is different. Everything is different. So with sharing by (sic) different mummies, it will give (me) confidence to hear that someone’s pregnancy (experience) is similar to mine. Then, I’m not the odd one out”* (Interviewee 07). Another participant stated: *“Because there are things you cannot tell your parents, you cannot tell your other half. So telling people in the mummies group who understand you, then you will (sic) feel like, oh I’m not alone. And you do make good friends. Then (sic) you go for outings with the baby and learn from each other’s parenting. So I think it’s quite a good one”* (Interviewee 05).

Innovation: **Desired and timely information**

Participants indicated that concrete information about specific topics, such as healthy eating, exercise guide, mental health and breastfeeding support, should be included in the intervention program. *“I want to learn more about diet intake (sic), the type of exercises I can do, how to breastfeed, and also sleep training for the baby. I also want more assurance on the type of exercises that I can do, and for how long”* (Interviewee 08). Another participant added: *“More on (the) breastfeeding issue ... I still see (many) mummies struggling with breastfeeding”* (Interviewee 05). Participants commented it was important that the information delivered was compatible with the specific needs and challenges faced at different phases of the journey from preconception to postpartum, necessitating different information, and intervention intensity. *“Before delivery you can provide (information) so they can be more well-equipped and at the same time, during the pregnancy, they can still go back to it to read up again”* (Interviewee 09). Participants were also keen to receive information about the postpartum period as early as preconception. *“Ideally, before I get pregnant, I want to learn about breastfeeding and sleep training. I only got in touch with breastfeeding and sleep training during pregnancy. I find that is too late and as a new mom, I don’t know how to analyze and filter all the information, so if there is some information on this during preconception, I can be mentally prepared for motherhood”* (Interviewee 08). The desired frequency and intensity of intervention was highlighted by a participant: *“(During) preconception, I would prefer to receive information every month, or every 2 months. As for health tips or nudges, I don’t mind receiving such information every day. During pregnancy, I would prefer receiving information every week. (In the) Postpartum period, I am happy to receive weekly newsletters”* (Interviewee 08).

Innovation: **Holistic delivery platform**

Participants wanted a single platform to deliver preconception, pregnancy and postnatal care which allows users to continue using the tool throughout the journey. *“There are apps online, but none are three-in-one - before pregnancy, pregnancy and postnatal, the whole journey. If you have an app that puts everything together, this app would be very widely used”* (Interviewee 05). Most of the participants

preferred a mobile application to deliver the intervention, with one of them saying: *“Mobile application, like an all-in-one application that I can refer to as and when I want to”* (Interviewee 11). Another participant expressed a similar sentiment. *“Yes I will still prefer the app. Because you might be working. You might not have time to attend. I think the most efficient method is to have weekly classes. Because you need to know what to expect each week, what needs to be done. But no one will have time to attend a physical class every week. So I think an app is still better”* (Interviewee 08). However, there were others who were hesitant to embrace new technology. *“I prefer physical classes that are face to face because I cannot concentrate when classes are online”* (Interviewee 15).

Facilitation: **Frequent engagement**

Some participants suggested the use of a monitoring logbook to monitor their health and feedback reports to guide healthy behaviors. *“Monitor our diet intake and exercise, that may keep us more engaged”* (Interviewee 11). This would help to provide ongoing motivation and enable them to troubleshoot their lifestyle. *“To improve engagement, you can have a monitoring logbook, get a gift or something to motivate (us)”* (Interviewee 02). The desire for an interactive process was echoed by another participant: *“Yeah. Especially when there’s a section where you can actually (sic) ask a question, and get an answer, maybe not instantaneously but you know you may get one within the day, or maybe within a week, but there will be someone who answers the question. Or it could be become like a thumb’s up thing, where most voted questions get answered faster”* (Interviewee 09). Participants also suggested nudges to support regular re-engagement, which would provide reminders and ongoing motivation to promote sustained behavior change. *“I feel like it would be good if there are notification pushes periodically maybe once or twice a week, but also if I were interested and I were to search it up within the app itself, its listed there as well. So, if I want to read forward, beyond those week by week pregnancy information pushes. For example what size (the baby bump) will be the following week, I can find that information (on the app) if I’m interested”* (Interviewee 09). Another participant added: *“Each of these nudges serves as a reminder when it’s time to run. Perhaps a nudge for the parents to think about whether they have any questions to ask the doctor”* (Interviewee 06).

Discussion

Ten enabler and barrier factors emerged from the interview process. These were framed in the lens of the i-PARIHS framework to understand how perceived barriers and enablers towards adoption of lifestyle changes shape participation and engagement with intervention programs. Among women of childbearing age who were overweight or obese, at different stages from preconception to postnatal, factors surrounding individual women, the sociocultural context, and key characteristics determining the uptake and continued efficacy of a lifestyle intervention were identified. Not tailoring the intervention sufficiently to these factors could explain the variable efficacy of existing interventions that aim to effect behavioral changes, such as diet modifications and increased physical activity [18, 21, 23, 38–40]. Hence, we propose an intervention that mitigates these barriers and capitalizes on enabling factors to ensure successful implementation and sustainable impact.

1. Education across the life-course to address knowledge gaps

Despite strong motivation among mothers and prospective mothers to adopt healthier lifestyles for their babies, knowledge gaps and misconceptions are barriers that prevent the adoption of such behaviors [15, 25, 33]. Specifically, women are unaware that obesity before and during pregnancy has lifelong impact on maternal and child health. This may stem from insufficient guidance on healthy lifestyle behaviors applicable to the local context [13, 16, 25]. Addressing these through education will enable participants to perceive value in optimizing health for themselves and their children in the long-run, while providing practical suggestions that are culturally convenient, so women can act [41]. These form the considerations in designing an educational package as part of a lifestyle intervention program, aligned with our findings in this study.

First, the intervention should address the spectrum of needs and concerns beyond obesity and physical health, including physical activity and exercise, breastfeeding, and infant feeding practices. Participants of other studies have brought up similar concerns that are beyond the scope of the clinician's agenda of targeting obesity or a specific health problem. Some topics of interest include support for miscarriage, sleep, mental health, postpartum care and baby care in the first year of life [42–44]. Notably, this indicates that participants desired a holistic intervention, which extends beyond their medical needs or a specific health concern to address their personal concerns as a woman and mother-to-be.

Second, the educational package should be relevant to users throughout the journey from preconception to postnatal phases. Information should be tailored to each phase as motivation and concerns for women differ at each of these phases. Before conceiving, the primary concern for prospective mothers is getting pregnant and optimizing fertility, hence information targeted at these topics are most likely to catch their interest. During pregnancy, women are concerned about how their lifestyle habits or symptoms affect the wellbeing of their fetus. Pregnancy-safe exercise plans at different trimesters are an example of information that can be provided. After delivery, women may worry about their recovery and care for their newborn, so topics such as breastfeeding and sleep training may be more relevant. In addition, women experience different time constraints during each phase. They have the least time postpartum as they are caring for their newborns, and the most time preconception where there are few medical visits and no fatigue from pregnancy. Thus, the intensity should also be tailored to these time constraints, with the greatest intensity before pregnancy, easing up during and after pregnancy.

There are many applications already available on the market, however these largely target a specific phase or aspect of the journey. Pregnant women currently use a potpourri of these apps [42, 45], but desire a comprehensive application that would serve all of their needs throughout the journey from preconception to postnatal. Taking a life-course approach achieves this by recognizing that an individual's needs and concerns evolve over time, and tailors the intervention to suit their needs at different phases of the journey. In addition, initiating the intervention before conception gives women enough time to acquire nutrition knowledge and develop exercise habits [46], enabling long-lasting lifestyle changes that are sustained before, during, and beyond pregnancy [20, 46]. Continuing to support

healthy behaviors in the postnatal period [44] improves metabolic health when entering a subsequent pregnancy, and later in life.

Third, the identified barriers of contextual factors coupled with a lack of reliable locally relevant data sources show that difficulty integrating recommendations for lifestyle changes into local practices and habits prevent engagement with an intervention. Thus, the educational package must be designed with local practices, culture, and diet in mind. In particular, the influence of family culture on a person's diet should be mitigated by engaging family members, especially partners, in the intervention program, which can be done by providing tips for exercising together or meal plans for the whole family. Not only does this promote healthy eating and lifestyle habits, but it also helps to establish spousal support throughout the journey of preconception, pregnancy and postpartum care. As it is a common practice for locals to purchase foods from food courts and food retailers (43), there is a need to incorporate education in identifying healthier food choices while eating out. This can integrate and augment current government policies and public health messaging such as the Healthier Dining Programme and Healthier Choice Symbol [47, 48].

2. Promoting sustainable engagement with healthy lifestyle changes

The ultimate goal of lifestyle interventions is to cultivate sustained, long-term changes in behavior [20]. Strategies to enable these are a topic of interest and research [20, 24]. Social support has been identified to be vital in promoting healthy behavior both in this study and others [42, 45]. Meanwhile, time is a well-documented barrier to healthy lifestyle behaviors (e.g. exercising), particularly during the time surrounding pregnancy and motherhood [16, 26, 43, 45]. Okesene-Gafa and colleagues (2016) found that perceived time constraint was cited as a likely reason for predicted non-participation if a nutritional intervention was created [16]. A longitudinal study conducted on Australian women showed that women's perception of time pressure increased with number of children, as well as working hours they experienced [49]. Notably, majority of our participants are employed (73%) and thus our data provided a reflection of working women's experiences and time constraints. For a lifestyle intervention to successfully enable behavioral change, time as a barrier needs to be sufficiently addressed.

Nudge interventions are increasingly used to influence health behavior to regularly re-engage users, and as promising adjuncts to educational interventions to counteract the adverse effects of decision biases that inhibit people from acting rationally [50, 51]. Nudges can be delivered in the form of push notifications or short text messages, e.g. via social media that provide information or guidance, motivational messages, activity reminders, or promote interaction with the intervention [52]. This facilitates short and regular interactions with the lifestyle intervention, continuously re-engaging the user and providing consistent reminders for habit formation to drive long-term behavioral change [44]. Importantly, the intervention needs to be responsive to users' needs. Incorporating built-in self-evaluation and feedback processes would enable users to support change independently, thereby enhancing self-efficacy and self-control [53]. It also allows personalization of healthy messages and nudges through

goal-setting and feedback, which have previously been found to be more effective than non-customized messages in improving self-efficacy and dietary behaviors [54].

Taken together, a sustainable intervention needs to incorporate a range of complex features providing guidance and education throughout the pregnancy life-course, while enabling a high degree of personalization to individual needs with feedback, goal setting and regular nudges. This is best achieved through a mobile health (mHealth) platform. mHealth is broadly defined as medical and public health practice supported by mobile devices such as smartphones [55]. The main advantage of using a mHealth platform compatible with smartphones is that it can be accessed anywhere at any time [45]. The intervention can be delivered through small but frequent interactions with the mobile app that can be completed in pockets of free time, which fits in well with the modern lifestyle of city-dwelling women, who have also expressed that they do not want to be given too much information at once [42]. Further, each interaction with the intervention can be tailored to the amount of available time, for instance completing short exercise sessions or consuming bite-sized information [42], allowing changes to be flexibly incorporated into daily life.

Strengths and Limitations

A major strength of this study is the inclusion of the important phases in a woman's life journey, from preconception to pregnancy and postpartum, and with recruitment from community and hospital settings. The codes and themes that arose from women in the preconception, pregnancy or postpartum phases were similar and is likely to represent women's attitudes towards lifestyle interventions throughout these phases in general. However, this study only interrogates the perspectives of end-users, and does not take into consideration the perspective of other stakeholders. A lifestyle intervention is a multidisciplinary and multistakeholder effort, with constraints and barriers from other perspectives that would influence the success of the intervention aside from that of end-users [14, 24, 27]. However, there is a paucity of research that integrates the interplay of perspectives of multiple stakeholders in a lifestyle intervention, which should be a topic of further research.

Conclusion

Obesity caused by poor lifestyle behaviors, including dietary choices and lack of physical activity, has long-term adverse health effects in women and their children. The preconception, pregnancy and postpartum time period represents a unique opportunity for intervening to cultivate sustained healthy lifestyle behaviors. Although most pregnancies are planned and mothers are intrinsically driven to provide the best for their offspring, they may not be equipped with knowledge about preconception, pregnancy and postnatal care to put this into effect. The success of a lifestyle intervention is heavily dependent on uptake and requires continued compliance, which has been a challenge in previous studies. Using the i-PARIHS framework to explore the perceived enablers and barriers of adopting a lifestyle intervention program among preconception, pregnant and postpartum women who are overweight or obese, we propose components of an idealized lifestyle intervention to improve metabolic health. This includes a

holistic education package taking a life-course approach to improve women's baseline knowledge of health; using a mobile health platform to reduce barriers to adopting lifestyle changes, provide personalized feedback and goal-setting; and regular health nudges to cultivate sustained lifestyle habits.

Declarations

Data availability

Please contact the corresponding author for more information.

Acknowledgements

The authors would like to thank KK Women's and Children's Hospital for the institutional support received during this study. Special thanks go out to Ms Rachael Loo who provided invaluable administrative support and coordination. We also thank the team who worked tirelessly to transcribe interview recordings into text, Ms Aqilah Dariah Mohd Zulkarnain, Ms Kelly Lim Zhi Qi, Mr Ng Ding Yi, Ms Nur Nadirah Lim, Ms Tan Wei Mun Amanda, Ms Tan Yi Wan and Mr Ku Chee Onn. The authors would also like to express their gratitude towards all study participants for their time and assistance.

Authors' contributions

CWK, FY, JKYC and SLL conceived and designed the study. IO reviewed literature for the semi-structured interview guides. CWK, SLL and XWN designed and conducted the interviews. CSE and IO were part of the team responsible for transcribing the interviews. CWK and SLL reviewed the interview transcripts and performed thematic analysis. CWK, SLL and SHL mapped themes onto the i-PARIHS framework. CSE analyzed data from the questionnaire. SHL, CWK and LSO prepared the manuscript. JJXT, FY, JKYC and SLL edited the manuscript. All authors reviewed and approved the final manuscript. All authors have agreed to be personally accountable for their contributions and ensure that questions related to the accuracy or integrity of any part of the work, even ones they were not personally involved, are appropriately investigated, resolved and the resolution documented in the literature.

Competing interests

The authors declare that they have no competing interests.

Ethics approval and consent to participate

The current study was reviewed and approved by the SingHealth Centralized Institutional Review Board (CIRB Ref: 2020/2530). All participants signed an informed consent form before participating in the study. Identifying information was removed from the data prior to the analysis and all methods were performed in accordance with the relevant guidelines and regulations.

Consent for publication

The authors declare that they provide consent for publication.

Funding

This research was supported by the National Medical Research Council Singapore (Grant Number NMRC/CG/M003/2017). The funding body did not influence either the data collection and analysis or the writing and the decision to submit the manuscript. KCW and JCKY received salary support from Singapore's Ministry of Health's National Medical Research Council (NMRC/MOH-000596-00 and NMRC/CSA-SI-008-2016 respectively).

References

1. Law, D. C. G., Macle hose, R. F. & Longnecker, M. P. Obesity and time to pregnancy. *Hum. Reprod. Oxf. Engl.* **22**, 414–420 (2007).
2. Boots, C. E., Bernardi, L. A. & Stephenson, M. D. Frequency of euploid miscarriage is increased in obese women with recurrent early pregnancy loss. *Fertil. Steril.* **102**, 455–459 (2014).
3. Lashen, H., Fear, K. & Sturdee, D. W. Obesity is associated with increased risk of first trimester and recurrent miscarriage: matched case-control study. *Hum. Reprod. Oxf. Engl.* **19**, 1644–1646 (2004).
4. Aune, D., Saugstad, O. D., Henriksen, T. & Tonstad, S. Maternal body mass index and the risk of fetal death, stillbirth, and infant death: a systematic review and meta-analysis. *JAMA* **311**, 1536–1546 (2014).
5. Sohlberg, S., Stephansson, O., Cnattingius, S. & Wikström, A.-K. Maternal body mass index, height, and risks of preeclampsia. *Am. J. Hypertens.* **25**, 120–125 (2012).
6. Kim, S. S. *et al.* Obstetric and neonatal risks among obese women without chronic disease. *Obstet. Gynecol.* **128**, 104–112 (2016).
7. Price, S. A., Sumithran, P., Nankervis, A., Permezel, M. & Proietto, J. Preconception management of women with obesity: A systematic review. *Obes. Rev.* obr.12804 (2018) doi:10.1111/obr.12804.
8. Cnattingius, S. *et al.* Maternal obesity and risk of preterm delivery. *JAMA* **309**, 2362–2370 (2013).
9. Nuthalapaty, F. S., Rouse, D. J. & Owen, J. The association of maternal weight with cesarean risk, labor duration, and cervical dilation rate during labor induction. *Obstet. Gynecol.* **103**, 452–456 (2004).
10. Fleming, T. P. *et al.* Origins of lifetime health around the time of conception: causes and consequences. *The Lancet* **391**, 1842–1852 (2018).
11. OECD. *The Heavy Burden of Obesity: The Economics of Prevention*. (OECD, 2019). doi:10.1787/67450d67-en.
12. Beluska-Turkan, K. *et al.* Nutritional gaps and supplementation in the first 1000 days. *Nutrients* **11**, 2891 (2019).
13. Blondin, J. H. & LoGiudice, J. A. Pregnant women's knowledge and awareness of nutrition. *Appl. Nurs. Res.* **39**, 167–174 (2018).

14. Lee, A., Newton, M., Radcliffe, J. & Belski, R. Pregnancy nutrition knowledge and experiences of pregnant women and antenatal care clinicians: a mixed methods approach. *Women Birth* **31**, 269–277 (2018).
15. Funnell, G., Naicker, K., Chang, J., Hill, N. & Kayyali, R. A cross-sectional survey investigating women's information sources, behaviour, expectations, knowledge and level of satisfaction on advice received about diet and supplements before and during pregnancy. *BMC Pregnancy Childbirth* **18**, 182 (2018).
16. Okesene-Gafa, K., Chelimo, C., Chua, S., Henning, M. & McCowan, L. Knowledge and beliefs about nutrition and physical activity during pregnancy in women from South Auckland region, New Zealand. *Aust. N. Z. J. Obstet. Gynaecol.* **56**, 471–483 (2016).
17. Poston, L. *et al.* Preconceptional and maternal obesity: epidemiology and health consequences. *Lancet Diabetes Endocrinol.* **4**, 1025–1036 (2016).
18. Simmons, D. *et al.* Effect of physical activity and/or healthy eating on GDM risk: the DALI lifestyle study. *J. Clin. Endocrinol. Metab.* **102**, 903–913 (2017).
19. Oteng-Ntim, E., Varma, R., Croker, H., Poston, L. & Doyle, P. Lifestyle interventions for overweight and obese pregnant women to improve pregnancy outcome: systematic review and meta-analysis. *BMC Med.* **10**, 47 (2012).
20. Hanson, M. *et al.* Interventions to prevent maternal obesity before conception, during pregnancy, and post partum. *Lancet Diabetes Endocrinol.* **5**, 65–76 (2017).
21. Mutsaerts, M. A. Q. *et al.* The LIFESTYLE study: costs and effects of a structured lifestyle program in overweight and obese subfertile women to reduce the need for fertility treatment and improve reproductive outcome. A randomised controlled trial. *BMC Womens Health* **10**, 22 (2010).
22. Wekker, V. *et al.* Long-term effects of a preconception lifestyle intervention on cardiometabolic health of overweight and obese women. *Eur. J. Public Health* **29**, 308–314 (2019).
23. van Elten, T. *et al.* Preconception lifestyle and cardiovascular health in the offspring of overweight and obese women. *Nutrients* **11**, 2446 (2019).
24. Dadich, A., Piper, A. & Coates, D. Implementation science in maternity care: a scoping review. *Implement. Sci.* **16**, 16 (2021).
25. Forbes, L., Graham, J., Berglund, C. & Bell, R. Dietary change during pregnancy and women's reasons for change. *Nutrients* **10**, 1032 (2018).
26. Lim, S. L. *et al.* Attitudes & beliefs that influence healthy eating behaviours among mothers of young children in Singapore: a cross-sectional study. *Appetite* **148**, 104555 (2020).
27. Harvey, G. & Kitson, A. PARIHS revisited: from heuristic to integrated framework for the successful implementation of knowledge into practice. *Implement. Sci.* **11**, 33 (2015).
28. Roberts, N. A. *et al.* The utility of the implementation science framework “Integrated Promoting Action on Research Implementation in Health Services” (i-PARIHS) and the facilitator role for introducing patient-reported outcome measures (PROMs) in a medical oncology outpatient department. *Qual. Life Res.* (2020) doi:10.1007/s11136-020-02669-1.

29. Braun, V. & Clarke, V. Novel insights into patients' life-worlds: the value of qualitative research. *Lancet Psychiatry* **6**, 720–721 (2019).
30. Polit, D. F. & Beck, C. T. *Essentials of Nursing Research: Appraising Evidence for Nursing Practice*. (Wolters Kluwer Health/Lippincott Williams & Wilkins, 2010).
31. Vander Wyst, K. B. *et al.* A social media intervention to improve nutrition knowledge and behaviors of low income, pregnant adolescents and adult women. *PLOS ONE* **14**, e0223120 (2019).
32. Arrish, J., Yeatman, H. & Williamson, M. Australian midwives and provision of nutrition education during pregnancy: a cross sectional survey of nutrition knowledge, attitudes, and confidence. *Women Birth* **29**, 455–464 (2016).
33. Aktaç, S., Sabuncular, G., Kargin, D. & Gunes, F. E. Evaluation of nutrition knowledge of pregnant women before and after nutrition education according to sociodemographic characteristics. *Ecol. Food Nutr.* **57**, 441–455 (2018).
34. Braun, V. & Clarke, V. Using thematic analysis in psychology. *Qual. Res. Psychol.* **3**, 77–101 (2006).
35. NVivo. (QSR International Pty Ltd, 2020).
36. Kitson, A. L. *et al.* Evaluating the successful implementation of evidence into practice using the PARiHS framework: theoretical and practical challenges. *Implement. Sci.* **3**, 1 (2008).
37. Reddy, G. & van Dam, R. M. Food, culture, and identity in multicultural societies: Insights from Singapore. *Appetite* **149**, 104633 (2020).
38. Farpour-Lambert, N. J., Ells, L. J., Martinez de Tejada, B. & Scott, C. Obesity and weight gain in pregnancy and postpartum: an evidence review of lifestyle interventions to inform maternal and child health policies. *Front. Endocrinol.* **9**, 546 (2018).
39. Birdsall, K. M., Vyas, S., Khazaezadeh, N. & Oteng-Ntim, E. Maternal obesity: a review of interventions. *Int. J. Clin. Pract.* **63**, 494–507 (2009).
40. Lan, L. *et al.* Systematic review and meta-analysis of the impact of preconception lifestyle interventions on fertility, obstetric, fetal, anthropometric and metabolic outcomes in men and women. *Hum. Reprod.* **32**, 1925–1940 (2017).
41. van Dijk, M. R. *et al.* A mobile app lifestyle intervention to improve healthy nutrition in women before and during early pregnancy: single-center randomized controlled trial. *J. Med. Internet Res.* **22**, e15773 (2020).
42. Goetz, M. *et al.* Perceptions of patient engagement applications during pregnancy: a qualitative assessment of the patient's perspective. *JMIR MHealth UHealth* **5**, e73 (2017).
43. Halili, L. *et al.* Development and pilot evaluation of a pregnancy-specific mobile health tool: a qualitative investigation of SmartMoms Canada. *BMC Med. Inform. Decis. Mak.* **18**, 95 (2018).
44. Soltani, H. *et al.* Maternal obesity management using mobile technology: a feasibility study to evaluate a text messaging based complex intervention during pregnancy. *J. Obes.* **2015**, 1–10 (2015).

45. Lau, Y. *et al.* Development of a healthy lifestyle mobile app for overweight pregnant women: qualitative study. *JMIR MHealth UHealth* **6**, e91 (2018).
46. Blau, L. E. & Hormes, J. M. Preventing excess gestational weight gain and obesity in pregnancy: the potential of targeting psychological mechanisms. *Curr. Obes. Rep.* **9**, 522–529 (2020).
47. Healthy Hawker Food. *Health Hub* <https://www.healthhub.sg/live-healthy/24/healthyhawkerfood> (2020).
48. Health Promotion Board. Healthier choice symbol. <https://www.hpb.gov.sg/food-beverage/healthier-choice-symbol> (2021).
49. Otterbach, S. *et al.* The effect of motherhood and work on women's time pressure: a cohort analysis using the Australian longitudinal study on women's health. *Scand. J. Work. Environ. Health* **42**, 500–509 (2016).
50. Thaler, R. H. & Sunstein, C. R. *Nudge: improving decisions about health, wealth, and happiness*. (Penguin Books, 2009).
51. Möllenkamp, M., Zeppernick, M. & Schreyögg, J. The effectiveness of nudges in improving the self-management of patients with chronic diseases: a systematic literature review. *Health Policy* **123**, 1199–1209 (2019).
52. Webb, T. L., Joseph, J., Yardley, L. & Michie, S. Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *J. Med. Internet Res.* **12**, e4 (2010).
53. Tang, J., Abraham, C., Stamp, E. & Greaves, C. How can weight-loss app designers' best engage and support users? A qualitative investigation. *Br. J. Health Psychol.* **20**, 151–171 (2015).
54. McCarroll, M. L. *et al.* Feasibility of a lifestyle intervention for overweight/obese endometrial and breast cancer survivors using an interactive mobile application. *Gynecol. Oncol.* **137**, 508–515 (2015).
55. WHO Global Observatory for eHealth. *MHealth: New Horizons for Health Through Mobile Technologies*. (World Health Organization, 2011).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)
- [Additionalfile2.docx](#)