

The Effect of Co-exposure to Glyphosate, Cadmium, and Arsenic on Chronic Kidney Disease

Junne-Ming Sung

National Cheng Kung University Medical Center: National Cheng Kung University Hospital

Wei-Hsiang Chang

National Cheng Kung University

Kuan-Hung Liu

National Cheng Kung University College of Medicine

Chung Yu Chen

Chang Jung Christian University

Trias Mahmudiono

Airlangga University: Universitas Airlangga

Wan-Ru Wang

National Cheng Kung University

Ho-Chi Hsu

National Cheng Kung University

Zhen-Yi Li

National Cheng Kung University

Hsiu-Ling Chen (✉ hsiulinchen@mail.ncku.edu.tw)

National Cheng Kung University <https://orcid.org/0000-0002-1209-6789>

Research Article

Keywords: glyphosate, AMPA, metal, As, Cd, CKD

Posted Date: October 7th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-882942/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Exposure and Health on January 15th, 2022. See the published version at <https://doi.org/10.1007/s12403-021-00451-3>.

Abstract

The usage of glyphosate is increasing worldwide. Glyphosate and its major metabolite, aminomethylphosphonic acid (AMPA), are of potential toxicological concern in unknown chronic kidney disease (CKDu). As with Cd and other elements, glyphosate exposure has been reported as risk factor for CKDu in farmers. This study aimed to evaluate the influence of co-exposure to glyphosate and heavy metals in chronic kidney disease. In this study, the urine samples from 55 patients with CKD and 100 participants without CKD were analyzed for glyphosate, As, Cd, and Pb concentrations, and eGFR. Negative associations between glyphosate, AMPA, As, and Cd concentrations in the urine and eGFR were found for study subjects ($p < 0.05$). With regard to the effect of co-exposure, the odds ratios (OR) for subjects with an eGFR of $< 60 \text{ mL/min/1.73 m}^2$ was significant because of the high Cd concentration ($> 1 \text{ } \mu\text{g/g}$ creatinine; OR = 7.57, 95% CI = 1.91–29.95). With regard to the effect of co-exposure, the OR for subjects with an eGFR $< 45 \text{ mL/min/1.73 m}^2$ was significant at high glyphosate concentration ($> 1 \text{ } \mu\text{g/g}$ creatinine; OR = 1.57, 95% CI = 1.13–2.16) and As concentration ($> 1 \text{ } \mu\text{g/g}$ creatinine; OR = 1.01, 95% CI = 1.00–1.02). These results showed that glyphosate, AMPA, As, and Cd have an effect on CKD; notably, Cd, As, and glyphosate exposure can be important risk factors after stage 3a of CKD, and that there was a co-exposure effect of As and glyphosate in CKD after stage 3b. The potential health impacts of glyphosate should be considered, especial for patients with CKD and eGFR below $45 \text{ mL/min/1.73 m}^2$.

Introduction

Since the early 1990s, an unknown chronic kidney disease (CKDu) with compelling tubulointerstitial presentations, called chronic interstitial nephritis, has been reported in agricultural areas of various tropical countries (Jayasumana et al. 2015b; Jayasumana et al. 2015c; Ruwanpathirana et al. 2019), particularly in developing countries without certain chronic etiologies, such as diabetes, hypertension, and glomerulonephritis (Jha et al. 2013). CKDu has spread across rural communities in South Asia, China, and Central America (Correa-Rotter et al. 2014; Jayasumana et al. 2014; Smpokou et al. 2019; Wang, D. et al. 2019). Notably, the overall prevalence of CKDu in Sri Lanka has reached 10% , and is as high as 22.9% in several communities; it is the cause of more than 20,000 deaths annually (Jayasumana et al. 2015a).

Glyphosate, an organophosphorus pesticide, is used in relatively high amounts worldwide owing to the increased planting of genetically modified seeds (Duke et al. 2008; Myers et al. 2016); further, the contamination of the crops, leaf, and seed by glyphosate can be detected in the air, water, and rain (Chang, F. C. et al. 2011; Krüger et al. 2014). Owing to its low vapor pressure (25°C , $9.8 \times 10^{-8} \text{ mm-Hg}$), glyphosate is hardly vaporized into air, which can be adsorbed by the leaves of plants through the soil, and then passed down the phloem to enter the root (Helander et al. 2012). In the eco-environmental system, glyphosate and its major metabolite, aminomethylphosphonic acid (AMPA) (Bai et al. 2016), are of potential toxicological concern, mainly as a result of the accumulation of residues in topsoil (Silva et al. 2018; Yang et al. 2015) and the food chain (Bai et al. 2016). Therefore, the major exposure pathway in humans is through the ingestion of the residuals of glyphosate and AMPA in food (Bai et al. 2016; FSA 2018); the other routes of exposure are inhalation and dermal contact in occupational workers and farmers (Abdul et al. 2021; Cai et al. 2017; Jayasumana et al. 2015c).

A previous study showed that glyphosate has tumor-promoting potential in mouse skin (George et al. 2010), and the International Agency for Research on Cancer defined glyphosate as Group 2A (probably carcinogenic to humans) in 2015 (IARC 2015). However, the European Food Safety Authority has suggested that glyphosate does not appear to be genotoxic and would not be a carcinogen (EFSA 2015). Many recent studies have declaimed that glyphosate has

adverse effects on human health (Chang, Ellen T. et al. 2016; EFSA 2015; Samsel et al. 2013), including impacts on antioxidants, reproductive hormones, and gut microbiome (Ruuskanen et al. 2020), and it may cause acute kidney effects or chronic disease following short-term or long-term exposure (Schaeffer et al. 2020; Tsai et al. 2018; Wimalawansa, Shehani et al. 2014).

In sugarcane farmers with kidney dysfunction in CKDu-emerging regions of Sri Lanka, urinary beta 2-microglobulin and serum cystatin C levels were significantly correlated with urinary glyphosate levels, which is potentially relevant to the subsequent decline in kidney function, as indicated by estimated glomerular filtration rate (eGFR), and albumin creatinine ratio, and neutrophil gelatinase-associated lipocalin (Abdul et al. 2021). Herrera-Valdes (2019) suggested that glyphosate specifically affected the kidneys in which farmers/farmworkers were highly exposed. Moderate to severe coagulative necrosis of hepatocytes and glomerular and renal tubular necrosis were observed when 4.4–750 mg/kg of oral glyphosate was administered daily for 36 weeks in rats (Tizhe et al. 2020). In an *in vitro* study, glyphosate was found to reduce cell viability and induce apoptosis and oxidative stress in a dose-dependent manner in a human renal proximal tubule cell line; this was attributed to the similarity of the chemical structures of glyphosate and AMPA to glycine and glutamate, which are agonists of the N-methyl-D-aspartate receptor, and further to result to an imbalance of oxidant and antioxidative products involved in glyphosate-induced renal proximal tubule epithelium apoptosis (Gao et al. 2019). In addition, glyphosate consumption and acute kidney injury were also observed in toxic epidermal necrolysis (Indirakshi et al. 2017). Therefore, the adverse effects of glyphosate exposure on renal dysfunction should be considered.

Metal exposure has been proven to be associated with kidney dysfunction (Sabath et al. 2012; Tsai et al. 2018; Wimalawansa, S. J. 2016). In Taiwan, the importance of zinc (Zn) and chemical oxygen demand in rivers was demonstrated in a regression model of CKD; a high CKD prevalence was related to arsenic (As) contamination in groundwater in Taiwan (Chang, Kuan et al. 2018); and a high As level of $\geq 50 \mu\text{g/L}$ in drinking water was a risk factor for end-stage renal disease (Cheng et al. 2018). The higher heavy metal concentrations in farms close to patients' residences were associated with a higher risk of progression to end-stage kidney disease (Tsai et al. 2018).

Jayasumana et al. (2015b) first reported a new form of CKD among paddy farmers, termed Sri Lankan Agricultural Nephropathy in 1994. They found that multiple heavy metals and glyphosate may have a synergistic nephrotoxicity. Meanwhile, they also found that phosphate fertilizers were a major source of inorganic As, which is relevant to CKDu in the endemic areas of Sri Lanka (Jayasumana et al. 2015a). Moreover, Babich et al. (2020) showed that metals in drinking water, even at safe levels, can impede kidney development from an early age, which potentiates increased susceptibility to other agrochemicals, such as glyphosate. The effects of drinking water contaminants on mitochondria can contribute to the progression of kidney dysfunction. However, no evidence was found for the loss of kidney function in participants at risk of mesoamerican nephropathy (MeN) (Smpokou et al. 2019). However, other studies pointed out that there may have been a synergistic effect of glyphosate and hard water on renal injury through mitogen-activated protein kinases /cytosolic phospholipase A2/arachidonic acid and their downstream factors (Wang, R. et al. 2021; Zhang et al. 2021).

Above all, the concern of the risk of kidney disease caused by glyphosate is very important when glyphosate exposure is increasing annually. The aim of this study was to evaluate the influence of co-exposure to glyphosate and heavy metals on CKD. The impact of glyphosate residues on health is warranted in subgroups sensitive to kidney dysfunction, especially in those exposed to metals, which are also related to kidney illness.

Materials And Methods

Subject enrollment

For this cross-sectional study, 55 patients with chronic kidney disease were recruited from the Division of Nephrology of National Cheng Kung University Hospital (NCKUS) and 100 participants with healthy kidney function were recruited from Taiwan Biobank (TWB). The study was approved by the Ethics Committee of NCKUS (Tainan, Taiwan, encoded: A-ER-108-189) and TWB (Taipei, Taiwan, encoded: TWBR10811-05). All participants signed a consent form before sampling started.

Interviewer-administered questionnaire

Demographic information was obtained using a face-to-face questionnaire-based interview in NCKUS, and the information on the 100 healthy participants was provided from TWB. Personal characteristics (including sex, age, height, weight, occupational history, neighborhood geography, and socioeconomics), lifestyle factors (alcohol consumption, smoking habits, drinking other liquids, etc.), and dietary patterns (consumption frequency and quantities) were included in the questionnaires. Meanwhile, histories of familial disease were also recorded in the questionnaires.

Blood serum and urine sampling

In 55 patients with CKD, 1 mL samples of plasma and 7–8 mL of urine were obtained in hospital, stored at 4°C, and then kept at –80°C until analysis. For the 100 study participants from TWB, 0.8 mL plasma and 2 mL urine were stored in glass tubes in the dark and kept at –80°C before metal analysis.

Analysis of urine glyphosate concentrations

Liquid chromatography-tandem mass spectrometry (LC-MS/MS) was used to analyze the concentration of glyphosate and AMPA. The concentrations of glyphosate and its metabolite, AMPA, in urine samples were determined after processing through a liquid–liquid extraction in acidic conditions and analysis using a validated LC-MS/MS method. In addition, creatinine content was measured to correct for diuresis. Then, 1 mL of urine sample and 100 µL of the internal standard (IS) solution (containing 1 ng/mL of each IS) were directly extracted with 1 mL deionized water containing 2% formic acid. The mixture was then vortexed for 3 min, and 1 mL of the mixture was filtered through a 0.2-µm PTFE membrane filter before LC–MS/MS analysis. The LC-MS/MS analysis was performed using an Agilent 1200 Infinity HPLC system coupled with an Agilent 6410 triple quadrupole mass spectrometer (Agilent Technologies, Inc., Palo Alto, CA, USA). The LC separation was conducted using an IC-Pak Anion HR 6 µm, 4.6 mm × 75 mm LC column (Waters, Milford, MA, USA), at a flow rate of 200 µL/min. The mobile phases were water (A) and ACN (B), both were in 2% HCOOH. The injection volume was 20 µL. The initial gradient was 10% B; increased to 41% B at 3 min; to 70% B at 4 min; maintained at 70% B for 4 min; and then returned to 10% B at 10 min. Multiple reaction monitoring data were acquired and processed in negative and positive ESI modes. The following transitions (quantification transitions are underlined) were found to be optimal for the detection: glyphosate, $170 > \underline{88}/170 > 60$; ^{13}C , ^{15}N glyphosate, $172 > \underline{90.1}/172 > 62.1$; AMPA, $110 > \underline{63.1}/110 > 79.1$; and ^{13}C , ^{15}N AMPA, $114 > \underline{63.0}/114 > 79$. The MS-MS nebulizer was set at 40 psi, the dry gas (N_2 99.9% pure) flow was set at 10 L/min, and the source temperature and capillary voltage were kept at 350°C and 4 kV, respectively. Glyphosate and AMPA was quantified using a developed linear calibration, which spanned over two orders of magnitude, with a working range from the

lowest reportable value (0.5 ng/mL) to the highest standard (50 ng/mL); $R^2 > 0.998$. Each analysis batch (10 samples) contained a laboratory blank, a pooled sample, a spiked pooled sample, and a repeatedly spiked sample. The concentration of glyphosate and AMPA in analytical blanks had to be lower than half of the method detection limit (MDL), at 0.003 and 0.012 $\mu\text{g/L}$, respectively. A QC check standard was analyzed for each batch to verify that the instrument remained properly calibrated and the recovery rates were 90%–111%. The recoveries of glyphosate and AMPA in the spiked samples ranged from 89%–114% and 87%–104%, respectively. The coefficients of variation of glyphosate and AMPA for the repeatedly spiked samples were both $<10\%$.

Analysis of metal concentrations in the serum and urine samples

Blood sampling and analysis details for metal concentrations are available elsewhere (Batista et al. 2009; Palmer et al. 2006). Metal contents were analyzed using inductively coupled plasma-mass spectroscopy (ICP-MS; ICP-MS-ELAN DRC II, PerkinElmer). The recovery efficiencies for Pb, Cd, and As were measured by the addition of a standard solution to samples, and the recovery rates in blood were 97% (Pb), 101% (Cd), and 102% (As) in urine, as well as MDLs were 0.047 mg/L (Pb), 0.014 mg/L (Cd), and 0.02 mg/L (As).

Statistical analysis

Metal, glyphosate, and AMPA concentrations were reported in units of $\mu\text{g/g}$ creatinine in urine and $\mu\text{g/L}$ in blood. SPSS 26 (IBM SPSS Statistics) were used for data management and statistical analysis. Chi-squared tests were used to examine the frequency distributions of dichotomous variables—gender, smoking status, and drinking status—in four groups of patients with different stages of CKD. The Kruskal-Wallis and Jonckheere-Terpstra test were used to compare the differences and trends in age, metal concentrations in the blood, in the urine, and glyphosate and AMPA concentrations in the four CKD groups. In addition, to assess the effects of co-exposure to glyphosate and metals, we compared, using logistic regression, the odds ratios (ORs) of different CKDs for participants with different levels of exposure. Statistical significance was set at $p < 0.05$.

Results

Demographics and the description of biomarker levels of 155 subjects

The average age of the 105 men and 50 women in the study was 53.1 years of age; 43 subjects were smokers and 15 subjects were drinkers (Table 1). The median glyphosate concentration in urine samples was 0.33 $\mu\text{g/g}$ creatinine (0–12.13 $\mu\text{g/g}$ creatinine), and the median AMPA concentration was 0.17 $\mu\text{g/g}$ creatinine (0–14.64 $\mu\text{g/g}$ creatinine). For the exposure biomarkers, the median concentration of As was 34.4 $\mu\text{g/g}$ creatinine (1.12–1020.73 $\mu\text{g/g}$ creatinine), the Cd concentration was 0.41 $\mu\text{g/g}$ creatinine (0.07–4.66 $\mu\text{g/g}$ creatinine), and the Pb concentration was 4.59 $\mu\text{g/g}$ creatinine (0.18–62.24 $\mu\text{g/L}$). As a biomarker of kidney function, the average eGFR was 85.12 mL/min/1.73 m² (standard deviation: 40.57).

The relationship between exposure biomarkers and eGFR

We further analyzed the correlation between glyphosate, AMPA, metal concentrations, and biomarkers of kidney function in Table 2. Negative correlations were shown with glyphosate ($\beta=-0.521$), AMPA ($\beta=-0.541$), As ($\beta=-0.388$), and Cd ($\beta=-0.580$) concentrations and eGFR in the urine samples from the study subjects ($p < 0.05$), separately (Table 2). In addition, significant negative associations were also found between the decrease in eGFR and glyphosate, AMPA, Cd, and As concentrations, even after adjustment for age, sex, and BMI (Table 3).

The relationship between exposure biomarkers and eGFR

Therefore, in the second stage, we categorized all participants into two groups, comprising: subjects without CKD (eGFR >90 mL/min/1.73 m²), non-CKD; patients with CKD stage 1–3a (60 \leq eGFR <90 mL/min/1.73 m²); stage 3b (45 \leq eGFR <60 mL/min/1.73 m²); and stage 4–5 (eGFR <45 mL/min/1.73 m²). Significant differences in age and smoking status were found among the four groups, but without a consistent increase or decrease in trends (Table 4).

For exposure biomarkers (Table 5), the average glyphosate concentrations in urine samples in the non-CKD, and stage 1–3a, 3b, and 4–5 groups were 0.38, 2.67, 2.29, and 3.62 μ g/g creatinine, respectively; the AMPA concentrations were 0.25, 2.35, 2.02, and 2.09 μ g/g creatinine, the average concentrations of As were 35.11, 95.96, 161.04, and 118.49 μ g/g creatinine, and the average concentrations of Cd were 0.41, 1.11, 1.35, and 2.04 μ g/g creatinine, respectively. There were significant differences between all four groups ($p < 0.05$). In the test for trends in biomarker exposure among these four groups, trends were found for glyphosate and Cd among these four CKD stages (Figure 1).

Further, dichotomized groups were categorized for eGFR <60 , or <45 mL/min/1.73 m², and ORs of subjects with eGFR below or above the two groups were calculated from logistic regression analysis. Higher glyphosate values were significantly related to an increased risk of a decrease in eGFR compared with the glyphosate lower group after adjustment for age, sex, BMI, and the interaction terms of glyphosate, Cd, and As concentrations. In Model 1, the OR for eGFR <60 was significant because of the high Cd concentration (> 1 mg/g creatinine; OR = 7.57, 95% CI = 1.91–29.95) and the OR was greater than 1 for high glyphosate concentrations (> 1 mg/g creatinine; OR = 1.39, 95% CI = 0.90–2.15), but the association was not significant. In Model 2, the OR for eGFR <45 was significant because of the high glyphosate (> 1 mg/g creatinine; OR = 1.57, 95% CI = 1.13–2.16) and As (> 1 mg/g creatinine; OR = 1.01, 95% CI = 1.00–1.02) concentrations, but the high Cd concentration (> 1 mg/g creatinine; OR = 1.85, 95% CI = 0.83–4.11) was not significant. The OR for the decrease in eGFR varied because of the different CKD stages.

Discussion

Biomarker of glyphosate exposure

In the metabolism analysis, the maximum concentrations of glyphosate and AMPA were observed at 2.42–5.16 h after the intravenous injection of glyphosate (Anadon et al. 2009). A study was designed in which 12 participants consumed a test meal with a known concentration of glyphosate residue and a lower concentration of AMPA, and the results showed that the estimated elimination half-life for glyphosate was 9 h (Zoller et al. 2020). Therefore, the measurements of glyphosate and AMPA in urine samples can be a short-term marker for external exposure.

Mills et al. (2017) have reported that the average glyphosate and AMPA concentrations in the general population of USA were 0.024 μ g/L and 0.314 μ g/L, increasing from 0.008 μ g/L to 0.401 μ g/L in 1993–1996 and 2014–2016 in the aging healthy population in the United States, whereas a decreasing trend in glyphosate residues in urine in Germany youngsters was found (Conrad et al. 2017). This study and the other one all found that chronically ill patients

had significantly higher urinary concentrations of glyphosate residues than healthy individuals (Krüger et al. 2014). Moreover, the glyphosate and AMPA concentrations in urine samples from adults and elders in this study were higher than those found in other countries (Krüger et al. 2014; Mills et al. 2017).

Although a review paper that collected data from seven studies revealed no health concerns because the glyphosate exposure estimation for general population was far below than “acceptable daily intake” or “acceptable operator exposure level”, exposure was predominantly resulted from the occupational and dietary exposure routes in Europe and North America (Niemann et al. 2015).

Many researchers have reported that glyphosate and AMPA residues were present in soy-based infant formula, maize-derived food, beer, wine, fruit juice (González-Ortega et al. 2017; Jansons et al. 2018; Rodrigues et al. 2018), and that glyphosate was detectable in nearly all honey samples in Switzerland (Zoller et al. 2018) and in American mothers’ breastmilk (Honeycutt et al. 2014). Therefore, the evaluation of glyphosate exposure via food consumption in patients with CKD is important.

With regard to As, higher exposures were found in this study compared with biological monitoring data from urine samples of other countries (Aguilera et al. 2008; Feng et al. 2015; Morton et al. 2014). For Cd exposure, the urine Cd concentrations in this study were clearly higher than those of other studies in Western countries (Aguilera et al. 2008; Baeyens et al. 2014; Heitland et al. 2006; Morton et al. 2014; Tellez-Plaza et al. 2008) and in Thailand (Nishijo et al. 2014), but were equal to those reported by Liao et al. in Taiwan (Liao et al. 2019). Overall, the concentrations of bio-exposure markers, such as glyphosate, AMPA, Cd, and As, were higher than those of other countries.

Metals, glyphosate exposure and renal function

In discussing the relationship between exposure biomarkers and eGFR, the variation in climate, temperature, air quality, water quality, and drought, and exposure to fertilizers, soil conditioners, herbicides, fungicides, and pesticides have been considered as contributing factors for the development of CKD in South Asia (Wilke et al. 2019). For example, in Thailand, the serum creatinine concentrations were associated with glyphosate use and pesticide exposure index in the occupational group (Mueangkhiao et al. 2020), which indicated that glyphosate exposure might be related to renal dysfunction. Meanwhile, an acute kidney injury developed after the ingestion of glyphosate-based herbicide, indicating epithelial injury in proximal tubules, and glyphosate mitochondrial toxicity was also found (Kimura et al. 2020). Other environmental contaminants, such as heavy metals (e.g., Cd, As, and Pb) and organic pesticides (e.g., glyphosate) in the drinking water, even at safe levels, can impair kidney development at an early age; and may play a role in the childhood onset and progression of kidney dysfunction (Babich et al. 2020). However, in three cohorts across different phases of child development, the authors confirmed detectable glyphosate in children’s urine at various ages and stages of development, but found no evidence for renal injury in children exposed to low concentrations of glyphosate (Trasande et al. 2020). Meanwhile, Gunatilake (2019) suggested that glyphosate’s synergistic health effects when combined with paraquat, and the continuous high temperatures of lowland tropical regions could result in renal damage. Glyphosate exposure, such as enhancing the growth of Clostridia species and ruminal metabolism *in vitro* (Riede et al. 2016), promotes As toxicity in renal dysfunction (Jayasumana et al. 2015a), and the low-dose exposure of glyphosate-based herbicides disrupted the urine metabolome and its interaction with gut microbiota has been dysregulated in related diseases through the commensal microbiome (Hu et al. 2021). Overall, several pathologies associated with MeN, a type of CKDu, may be linked to glyphosate exposure, such as altered gut microbiota (Rueda-Ruzafa et al. 2019), increased As toxicity, suppressed synthesis of adrenocorticotrophic hormone, disruption of fructose metabolism, and promotion of dehydration and high serum urate (Seneff et al. 2018).

For metal exposure, the higher concentrations of Zn and nickel (Ni) in farms close to the residence of patients with CKD were associated with a higher risk of progression to end-stage kidney disease in Taiwan (Tsai et al. 2018). An epidemiology study has suggested that high plasma selenium (Se) and low red blood cell Pb levels or Cd levels can interact to increase the eGFR (20.70, 15.56–26.01 mL/min/1.73 m²) in CKD (Wu et al. 2019). However, a report showed that the non-association between glyphosate, aluminum (Al), and As exposure and decreased kidney function in 350 young adults living in area of Central America with an epidemic of MeNs (Smpokou et al. 2019). Meanwhile, the exposure and risk assessment showed that there was no treatment risk with glyphosate (Honeycutt et al. 2014; Krüger et al. 2014; Niemann et al. 2015). All of the above studies were retrospective, and used biological biomarker data for external exposure, and APMA was not included in the above studies. However, in our study, the co-exposure of As and glyphosate was found in patients after CKD stage 3b. The alternative importance of glyphosate and Cd or As exposure in the progression of CKD have been seen in patients with CKD stage 3 or above. These results can respond to the conclusion from Seneff et al. (2018), who reported that the most likely way to prevent end-stage renal failure in sugarcane workers was to stop the use of glyphosate in Brazil, and that the progression of patients with CKD into end-stage renal failure may be prolonged by a reduction in glyphosate exposure.

Conclusions

In this study, we observed that glyphosate, AMPA, As, and Cd affected CKD; notably, Cd exposure may be an important risk factor after CKD stage 3a, and As and glyphosate have a synergistic effect following co-exposure in patents with CKD beyond stage 3b. Therefore, the potential health risks of glyphosate must be considered, especial for patients with CKD and eGFR values below 45 mL/min/1.73 m².

However, a comprehensive analysis of chemical contaminants in the drinking water and the effects of these compounds and their mixtures on kidney development and function is missing. Therefore, this study is a starting point for the discussion of the etiology and the progression of kidney disease; and analytical models for alternative agro-chemicals can provide other insights in future. Furthermore, the relationships should be followed up in a large population of patients with CKD to increase the power of the statistical analysis and allow consideration of more factors that co-influence CKD progression.

Declarations

Acknowledgements

We are in great debt to Division of Nephrology, Department of Internal Medicine, National Cheng Kung University Hospital for sampling assistance and our colleagues at the Research Center of Environmental Trace Toxic Substances, National Cheng Kung University Medical College for analytical assistance.

Authors' contributions

Junne-Ming Sung contributed to conceptualization, methodology, validation, subjects' recruitment; Wei-Hsiang Chang, Chung Yu Chen contributed to formal analysis, writing; Kuan-Hung Liu contributed to subjects' recruitment, methodology; Trias Mahmudiono, Ho-Chi Hsu contributed to review, sampling and formal analysis, Wan-Ru Wang, Zhen-Yi Li contributed to data curation, visualization; Hsiu-Ling Chen contributed to conceptualization, methodology, validation, writing – original draft, editing and funding acquisition.

Funding

No.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical approval The study protocol was approved by the Ethics Committee of NCKUS (Tainan, Taiwan, encoded: A-ER-108-189) and TWB (Taipei, Taiwan, encoded: TWBR10811-05) in accordance with the Ethical Principles for Medical Research Involving Human Subjects.

Consent to participate

All study participants provided written informed consent.

Consent for publication

No need.

References

1. Abdul KSM, De Silva P, Ekanayake E, Thakshila W, Gunarathna SD, Gunasekara T et al (2021) Occupational Paraquat and Glyphosate Exposure May Decline Renal Functions among Rural Farming Communities in Sri Lanka. *Int J Environ Res Public Health* 18 (6): 3278. <https://10.3390/ijerph18063278>
2. Aguilera I, Daponte A, Gil F, Hernández AF, Godoy P, Pla A et al (2008) Biomonitoring of urinary metals in a population living in the vicinity of industrial sources: a comparison with the general population of Andalusia, Spain. *Sci Total Environ* 407 (1): 669-678.
3. Anadon A, Martinez-Larranaga M, Martinez M, Castellano V, Martínez M, Martin M et al (2009) Toxicokinetics of glyphosate and its metabolite aminomethyl phosphonic acid in rats. *Toxicology Lett* 190 (1): 91-95.
4. Babich R, Ulrich JC, Ekanayake E, Massarsky A, De Silva P, Manage PM et al (2020) Kidney developmental effects of metal-herbicide mixtures: Implications for chronic kidney disease of unknown etiology. *Environ Int* 144: 106019. <https://10.1016/j.envint.2020.106019>
5. Baeyens W, Vrijens J, Gao Y, Croes K, Schoeters G, Den Hond E et al (2014) Trace metals in blood and urine of newborn/mother pairs, adolescents and adults of the Flemish population (2007–2011). *Int J Hyg Environ Health* 217 (8): 878-890.

6. Bai SH, Ogbourne SM (2016) Glyphosate: environmental contamination, toxicity and potential risks to human health via food contamination. *Environ Sci Pollut Res Int* 23 (19): 18988-19001. <https://doi.org/10.1007/s11356-016-7425-3>
7. Batista BL, Rodrigues JL, Nunes JA, de Oliveira Souza VC, Barbosa F (2009) Exploiting dynamic reaction cell inductively coupled plasma mass spectrometry (DRC-ICP-MS) for sequential determination of trace elements in blood using a dilute-and-shoot procedure. *Anal Chim Acta* 639 (1): 13-18. <https://doi.org/10.1016/j.aca.2009.03.016>
8. Cai R, Zheng YF, Bu JG, Zhang YY, Fu SL, Wang XG et al (2017) Effects of blood lead and cadmium levels on homocysteine level in plasma. *Eur Rev Med Pharmacol Sci* 21 (1): 162-166.
9. Chang ET, Delzell E (2016) Systematic review and meta-analysis of glyphosate exposure and risk of lymphohematopoietic cancers. *J Environ Sci Health - B Pestic Food Contam Agric Wastes* 51 (6): 402-434. <https://doi.org/10.1080/03601234.2016.1142748>
10. Chang FC, Simcik MF, Capel PD (2011) Occurrence and fate of the herbicide glyphosate and its degradate aminomethylphosphonic acid in the atmosphere. *Environ Toxicol Chem* 30 (3): 548-555. <https://doi.org/10.1002/etc.431>
11. Chang K, Wu IW, Huang BR, Juang JG, Wu JC, Chang SW et al (2018) Associations between Water Quality Measures and Chronic Kidney Disease Prevalence in Taiwan. *Int J Environ Res Public Health* 15 (12): 2726.
12. Cheng YY, Chang YT, Cheng HL, Shen KH, Sung JM, Guo HR (2018) Associations between arsenic in drinking water and occurrence of end-stage renal disease with modifications by comorbidities: A nationwide population-based study in Taiwan. *Sci Total Environ* 626 581-591. <https://doi.org/10.1016/j.scitotenv.2018.01.043>
13. Conrad A, Schröter-Kermani C, Hoppe HW, Rütther M, Pieper S, Kolossa-Gehring M (2017) Glyphosate in German adults—Time trend (2001 to 2015) of human exposure to a widely used herbicide. *Int J Hyg Environ Health* 220 (1): 8-16.
14. Correa-Rotter R, Wesseling C, Johnson RJ (2014) CKD of unknown origin in Central America: the case for a Mesoamerican nephropathy. *Am J Kidney Dis* 63 (3): 506-520.
15. Duke SO, Powles SB (2008) Glyphosate: a once-in-a-century herbicide. *Pest Manag Sci* 64 (4): 319-325. <https://doi.org/10.1002/ps.1518>
16. EFSA (2015). "Glyphosate: EFSA updates toxicological profile." from <https://www.efsa.europa.eu/en/press/news/151112>
17. Feng W, Cui X, Liu B, Liu C, Xiao Y, Lu W et al (2015) Association of urinary metal profiles with altered glucose levels and diabetes risk: a population-based study in China. *PloS one* 10 (4): e0123742.
18. FSA (2018). "Pesticides in food." from <https://www.food.gov.uk/business-guidance/pesticides-in-food>.
19. Gao H, Chen J, Ding F, Chou X, Zhang X, Wan Y et al (2019) Activation of the N-methyl-D-aspartate receptor is involved in glyphosate-induced renal proximal tubule cell apoptosis. *J Appl Toxicol* 39 (8): 1096-1107. <https://doi.org/10.1002/jat.3795>
20. George J, Prasad S, Mahmood Z, Shukla Y (2010) Studies on glyphosate-induced carcinogenicity in mouse skin: A proteomic approach. *J Proteom* 73 (5): 951-964. <https://doi.org/10.1016/j.jprot.2009.12.008>
21. González-Ortega E, Piñeyro-Nelson A, Gómez-Hernández E, Monterrubio-Vázquez E, Arleo M, Dávila-Velderrain J et al (2017) Pervasive presence of transgenes and glyphosate in maize-derived food in Mexico. *J Agroecol Sust Food Syst* 41 (9-10): 1146-1161.
22. Gunatilake S, Seneff S, Orlando L (2019) Glyphosate's Synergistic Toxicity in Combination with Other Factors as a Cause of Chronic Kidney Disease of Unknown Origin. *Int J Environ Res Public Health* 16 (15): <https://doi.org/10.3390/ijerph16152734>

23. Heitland P, Köster HD (2006) Biomonitoring of 30 trace elements in urine of children and adults by ICP-MS. *Clin Chim Acta* 365 (1-2): 310-318.
24. Helander M, Saloniemi I, Saikkonen K (2012) Glyphosate in northern ecosystems. *Trends Plant Sci* 17 (10): 569-574. <https://10.1016/j.tplants.2012.05.008>
25. Herrera-Valdes R, Almaguer-Lopez MA, Orantes-Navarro CM, Lopez-Marin L, Brizuela-Diaz EG, Bayarre-Vea H et al (2019) Epidemic of Chronic Kidney Disease of Nontraditional Etiology in El Salvador: Integrated Health Sector Action and South-South Cooperation. *MEDICC Rev* 21 (4): 46-52.
26. Honeycutt Z, Rowlands H (2014) Glyphosate testing report: findings in American mothers' breast milk, urine and water. *Moms Across America and Sustainable Pulse*.
27. Hu J, Lesseur C, Miao Y, Manservigi F, Panzacchi S, Mandrioli D et al (2021) Low-dose exposure of glyphosate-based herbicides disrupt the urine metabolome and its interaction with gut microbiota. *Sci Rep* 11 (1): 3265. <https://10.1038/s41598-021-82552-2>
28. Indirakshi J, Sunnesh A, Aruna M, Reddy MH, Kumar AC, Chandra VS et al (2017) Toxic Epidermal Necrolysis and Acute Kidney Injury due to Glyphosate Ingestion. *Indian J Crit Care Med* 21 (3): 167-169. https://10.4103/ijccm.IJCCM_423_16
29. Jansons M, Pugajeva I, Bartkevics V (2018) Occurrence of glyphosate in beer from the Latvian market. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 35 (9): 1767-1775. <https://10.1080/19440049.2018.1469051>
30. Jayasumana C, Fonseka S, Fernando A, Jayalath K, Amarasinghe M, Siribaddana S et al (2015a) Phosphate fertilizer is a main source of arsenic in areas affected with chronic kidney disease of unknown etiology in Sri Lanka. *Springerplus* 24 (4): 90. <https://10.1186/s40064-015-0868-z>
31. Jayasumana C, Gunatilake S, Senanayake P (2014) Glyphosate, hard water and nephrotoxic metals: are they the culprits behind the epidemic of chronic kidney disease of unknown etiology in Sri Lanka? *Int J Environ Res Public Health* 11 (2): 2125-2147. <https://10.3390/ijerph110202125>
32. Jayasumana C, Gunatilake S, Siribaddana S (2015b) Simultaneous exposure to multiple heavy metals and glyphosate may contribute to Sri Lankan agricultural nephropathy. *BMC Nephrol* 16 103. <https://10.1186/s12882-015-0109-2>
33. Jayasumana C, Paranagama P, Agampodi S, Wijewardane C, Gunatilake S, Siribaddana S (2015c) Drinking well water and occupational exposure to Herbicides is associated with chronic kidney disease, in Padavi-Sripura, Sri Lanka. *Environ Health* 14 6. <https://10.1186/1476-069X-14-6>
34. Jha V, Garcia-Garcia G, Iseki K, Li Z, Naicker S, Plattner B et al (2013) Chronic kidney disease: global dimension and perspectives. *The Lancet* 382 (9888): 260-272. [https://doi.org/10.1016/S0140-6736\(13\)60687-X](https://doi.org/10.1016/S0140-6736(13)60687-X)
35. Kimura T, Yokoyama T, Tanemoto M (2020) Renal tubular injury by glyphosate-based herbicide. *Clin Exp Nephrol* 24 (12): 1186. <https://10.1007/s10157-020-01962-0>
36. Krüger M, Schledorn P, Schrödl W, Hoppe H-W, Lutz W, Shehata AA (2014) Detection of glyphosate residues in animals and humans. *J Environ Anal Toxicol* 4 (2): 10. <https://DOI: 10.4172/2161-0525.1000210>
37. Liao KW, Pan WH, Liou SH, Sun CW, Huang PC, Wang SL (2019) Levels and temporal variations of urinary lead, cadmium, cobalt, and copper exposure in the general population of Taiwan. *Environ Sci Pollut Res* 26 (6): 6048-6064. <https://doi: 10.1007/s11356-018-3911-0>
38. Mills PJ, Kania-Korwel I, Fagan J, McEvoy LK, Laughlin GA, Barrett-Connor E (2017) Excretion of the Herbicide Glyphosate in Older Adults Between 1993 and 2016. *JAMA* 318 (16): 1610-1611. <https://10.1001/jama.2017.11726>

39. Morton J, Tan E, Leese E, Cocker J (2014) Determination of 61 elements in urine samples collected from a non-occupationally exposed UK adult population. *Toxicol Lett* 231 (2): 179-193.
40. Mueangkhiao P, Siviroj P, Sapbamrer R, Khacha-Ananda S, Lungkaphin A, Seesen M et al (2020) Biological variation in kidney injury and kidney function biomarkers among farmers in Lamphun province, Thailand. *Environ Sci Pollut Res Int* 27 (11): 12386-12394. <https://doi.org/10.1007/s11356-020-07661-3>
41. Myers JP, Antoniou MN, Blumberg B, Carroll L, Colborn T, Everett LG et al (2016) Concerns over use of glyphosate-based herbicides and risks associated with exposures: a consensus statement. *J Environ Health* 15 (1): 19.
42. Niemann L, Sieke C, Pfeil R, Solecki R (2015) A critical review of glyphosate findings in human urine samples and comparison with the exposure of operators and consumers. *Journal für Verbraucherschutz und Lebensmittelsicherheit* 10 (1): 3-12. <https://doi.org/10.1007/s00003-014-0927-3>
43. Nishijo M, Suwazono Y, Ruangyuttikarn W, Nambunmee K, Swaddiwudhipong W, Nogawa K et al (2014) Risk assessment for Thai population: benchmark dose of urinary and blood cadmium levels for renal effects by hybrid approach of inhabitants living in polluted and non-polluted areas in Thailand. *BMC Public Health* 14 (1): 702.
44. Palmer CD, Lewis ME, Geraghty CM, Barbosa F, Parsons PJ (2006) Determination of lead, cadmium and mercury in blood for assessment of environmental exposure: A comparison between inductively coupled plasma–mass spectrometry and atomic absorption spectrometry. *Spectrochim Acta Part B At Spectrosc* 61 (8): 980-990. <https://doi.org/10.1016/j.sab.2006.09.001>
45. Riede S, Toboldt A, Breves G, Metzner M, Köhler B, Bräunig J et al (2016) Investigations on the possible impact of a glyphosate-containing herbicide on ruminal metabolism and bacteria in vitro by means of the “Rumen Simulation Technique”. *J Appl Microbiol* 121 <https://doi.org/10.1111/jam.13190>
46. Rodrigues NR, de Souza APF (2018) Occurrence of glyphosate and AMPA residues in soy-based infant formula sold in Brazil. *Food Addit Contam A* 35 (4): 723-730.
47. Rueda-Ruzafa L, Cruz F, Roman P, Cardona D (2019) Gut microbiota and neurological effects of glyphosate. *Neurotoxicology* 75 1-8. <https://doi.org/10.1016/j.neuro.2019.08.006>
48. Ruuskanen S, Rainio MJ, Gómez-Gallego C, Selenius O, Salminen S, Collado MC et al (2020) Glyphosate-based herbicides influence antioxidants, reproductive hormones and gut microbiome but not reproduction: A long-term experiment in an avian model. *Environ Pollut* 266: 115108. <https://doi.org/10.1016/j.envpol.2020.115108>
49. Ruwanpathirana T, Senanayake S, Gunawardana N, Munasinghe A, Ginige S, Gamage D et al (2019) Prevalence and risk factors for impaired kidney function in the district of Anuradhapura, Sri Lanka: a cross-sectional population-representative survey in those at risk of chronic kidney disease of unknown aetiology. *BMC Public Health* 19 (1): 763. <https://doi.org/10.1186/s12889-019-7117-2>
50. Sabath E, Robles-Osorio ML (2012) Renal health and the environment: heavy metal nephrotoxicity. *Nefrologia* 32 (3): 279-286. <https://doi.org/10.3265/Nefrologia.pre2012.Jan.10928>
51. Samsel A, Seneff S (2013) Glyphosate, pathways to modern diseases II: Celiac sprue and gluten intolerance. *Interdiscip Toxicol* 6 (4): 159-184. <https://doi.org/10.2478/intox-2013-0026>
52. Schaeffer JW, Adgate JL, Reynolds SJ, Butler-Dawson J, Krisher L, Dally M et al (2020) A Pilot Study to Assess Inhalation Exposures among Sugarcane Workers in Guatemala: Implications for Chronic Kidney Disease of Unknown Origin. *Int J Environ Res Public Health* 17 (16): 5708. <https://doi.org/10.3390/ijerph17165708>
53. Seneff S, Orlando L (2018) Is glyphosate a key factor in Mesoamerican nephropathy. *J Environ Anal Toxicol* 8 (1): 1-10. <https://doi.org/10.4172/2161-0525.1000542>

54. Silva V, Montanarella L, Jones A, Fernández-Ugalde O, Mol HGJ, Ritsema CJ et al (2018) Distribution of glyphosate and aminomethylphosphonic acid (AMPA) in agricultural topsoils of the European Union. *Sci Total Environ* 15 (621): 1352-1359. <https://doi.org/10.1016/j.scitotenv.2017.10.093>
55. Smpokou ET, Gonzalez-Quiroz M, Martins C, Alvito P, Le Blond J, Glaser J et al (2019) Environmental exposures in young adults with declining kidney function in a population at risk of Mesoamerican nephropathy. *Occup Environ Med* 76 (12): 920-926. <https://doi.org/10.1136/oemed-2019-105772>
56. Tellez-Plaza M, Navas-Acien A, Crainiceanu CM, Guallar E (2008) Cadmium exposure and hypertension in the 1999–2004 National Health and Nutrition Examination Survey (NHANES). *Environ Health Perspect* 116 (1): 51-56. <https://doi.org/10.1289/ehp.10764>.
57. Tizhe EV, Ibrahim ND, Fatihu MY, Ambali SF, Igbokwe IO, Tizhe UD (2020) Effect of zinc supplementation on chronic hepatorenal toxicity following oral exposure to glyphosate-based herbicide (Bushfire(R)) in rats. *J Int Med Res* 48 (5): 1-15. <https://doi.org/10.1177/0300060520925343>
58. Trasande L, Aldana SI, Trachtman H, Kannan K, Morrison D, Christakis DA et al (2020) Glyphosate exposures and kidney injury biomarkers in infants and young children. *Environ Pollut* 256: 113334. <https://doi.org/10.1016/j.envpol.2019.113334>
59. Tsai CC, Wu CL, Kor CT, Lian IB, Chang CH, Chang TH et al (2018) Prospective associations between environmental heavy metal exposure and renal outcomes in adults with chronic kidney disease. *Nephrol* 23 (9): 830-836. <https://doi.org/10.1111/nep.13089>
60. Wang D, Zhang G, Zhang W, Luo J, Zhu L, Hu J (2019) Successful extracorporeal membrane oxygenation support for severe acute diquat and glyphosate poisoning: A case report. *Medicine (Baltimore)* 98 (6): e14414. <https://doi.org/10.1097/MD.00000000000014414>
61. Wang R, Chen J, Ding F, Zhang L, Wu X, Wan Y et al (2021) Renal tubular injury induced by glyphosate combined with hard water: the role of cytosolic phospholipase A2. *Ann Transl Med* 9 (2): 130. <https://doi.org/10.21037/atm-20-7739>
62. Wilke RA, Qamar M, Lupu RA, Gu S, Zhao J (2019) Chronic Kidney Disease in Agricultural Communities. *Am J Med* 132 (10): e727-e732. <https://doi.org/10.1016/j.amjmed.2019.03.036>
63. Wimalawansa S, Wimalawansa S (2014) Impact of changing agricultural practices on human health: Chronic kidney disease of multi-factorial origin in Sri Lanka. *Wudpecker J Agric Res* 3(5): 110-124.
64. Wimalawansa SJ (2016) The role of ions, heavy metals, fluoride, and agrochemicals: critical evaluation of potential aetiological factors of chronic kidney disease of multifactorial origin (CKDmfo/CKDu) and recommendations for its eradication. *Environ Geochem Health* 38 (3): 639-678. <https://doi.org/10.1007/s10653-015-9768-y>
65. Wu CY, Wong CS, Chung CJ, Wu MY, Huang YL, Ao PL et al (2019) The association between plasma selenium and chronic kidney disease related to lead, cadmium and arsenic exposure in a Taiwanese population. *J Hazard Mater* 375: 224-232. <https://doi.org/10.1016/j.jhazmat.2019.04.082>
66. Yang X, Wang F, Bento CPM, Xue S, Gai L, van Dam R et al (2015) Short-term transport of glyphosate with erosion in Chinese loess soil – A flume experiment. *Sci Total Environ* 512-513: 406-414. <https://doi.org/10.1016/j.scitotenv.2015.01.071>
67. Zhang L, Ding F, Wang R, Wu X, Wan Y, Hu J et al (2021) Involvement of mitochondrial fission in renal tubular pyroptosis in mice exposed to high and environmental levels of glyphosate combined with hard water. *Environ Pollut* 283: 117082. <https://doi.org/10.1016/j.envpol.2021.117082>

68. Zoller O, Rhyn P, Rupp H, Zarn JA, Geiser C (2018) Glyphosate residues in Swiss market foods: monitoring and risk evaluation. *Food Addit Contam Part B Surveil* 11 (2): 83-91. <https://10.1080/19393210.2017.1419509>
69. Zoller O, Rhyn P, Zarn JA, Dudler V (2020) Urine glyphosate level as a quantitative biomarker of oral exposure. *Int J Hyg Environ Health* 228: 113526. [https://https://doi.org/10.1016/j.ijheh.2020.113526](https://doi.org/10.1016/j.ijheh.2020.113526)

Tables

Table 1

Demographic results of all participants

	Population (N=155)
Demographics	
Sex	
Men	105 (67.7) ^a
Women	50 (32.3) ^a
Age	53.19±16.09 ^b
BMI	25.21±4.46 ^b
Body Weight	69.48±16.14 ^b
Smoking	43 (27.7) ^a
Drinking	15 (9.7) ^a
CKD stage	
Non CKD	99 (63.9) ^a
1	3 (5.45) ^a
2	10 (6.45) ^a
3a	11 (7.10) ^a
3b	15 (9.68) ^a
4	10 (6.45) ^a
5	7 (1.02) ^a
Exposure indicator	
Glyphosate(μg/L)	1.10 (0, 11.28) ^c
Glyphosate (μg/g creatinine)	1.16 (0, 12.13) ^c
AMPA(μg/L)	0.79 (0.01, 6.73) ^c
AMPA(μg/g creatinine)	0.86 (0, 14.64) ^c
As (μg/g creatinine)	62.34 (1.12, 508.82) ^c
Cd (μg/g creatinine)	0.78 (0.07, 4.66) ^c
Pb (μg/L)	6.11 (0.08, 62.24) ^c
Biomarker of kidney function	
eGFR (mL/min/1.73m ²)	85.06±40.70 ^b
Creatinine_urine (g/L)	1.68 (0.33, 3.81) ^c

Creatinine_serum (mg/dL)	1.38 (0.40, 9.15) ^c
^a Number and parentheses with percentage	
^b Mean±SD	
^c Median and parentheses with Minimum and Maximum	

Table 2

The correlation among glyphosate, metals and biomarkers of kidney function

N=155	Glyphosate	AMPA	As	Cd	Pb	Creatinine	eGFR
r [@]	(µg/g creatinine)	(µg/g creatinine)	(µg/g creatinine)	(µg/g creatinine)	(µg/L)	(mg/dL)	(mL/min/1.73m ²)
Glyphosate		0.569**	0.477**	0.723**	0.033	0.538**	-0.526**
AMPA			0.471**	0.653**	0.115	0.495**	-0.540**
As				0.580**	0.064	0.356**	-0.384**
Cd					0.009	0.550**	-0.585**
Pb						-0.058	0.067

[@] Spearman correlation coefficient * p<0.05, ** p<0.001

Table 3

The association between different exposure biomarkers and eGFR

Independent variable	β	p-value [@]
Glyphosate (µg/g creatinine) [¶]	-5.216	0.001*
AMPA (µg/g creatinine) [¶]	-2.315	0.070
As (µg/g creatinine) [¶]	-0.107	0.002*
Cd (µg/g creatinine) [¶]	-18.348	0.001*

[¶] Adjustment for age, sex, BMI
[@] Linear regression model, * p<0.05

Table 4

Demographic results of participants in different CKD stage

CKD stages	Non-CKD	1-3a	3b	4-5	P-value
N=	99	25	14	17	
Sex					0.183 ^c
Men	64 (64.6) ^a	15 (60.0) ^a	12 (85.7) ^a	14 (82.4) ^a	
Women	35 (35.4) ^a	10 (40.0) ^a	2 (14.3) ^a	3 (17.6) ^a	
Age	45.98±12.51 ^b	65.04±11.02 ^b	71.14±15.84 ^b	63.00±14.90 ^b	<0.001 ^{*d}
BMI	24.78±4.53 ^b	25.66±4.80 ^b	25.58±4.16 ^b	26.72±3.59 ^b	0.362 ^d
Body Weight	69.83±17.52 ^b	67.98±13.68 ^b	67.67±11.27 ^b	71.17±15.23 ^b	0.893 ^d
Smoking	33 (33.3) ^a	2 (8.0) ^a	1 (7.1) ^a	7 (41.2) ^a	0.012 ^{*c}
Drinking	9 (9.1) ^a	2 (8.0) ^a	1 (7.1) ^a	3 (17.6) ^a	0.692 ^c
^a Number and parentheses with percentage ^b Mean±SD ^c Chi-square test ^d One-way ANOVA * p<0.05					

Table 5

Difference of biomarker levels in different CKD stage

CKD stages	Non-CKD	1-3a	3b	4-5	P-value@
N=	99	25	14	17	
Glyphosate (µg/g creatinine)	0.31±0.22	2.28±2.57	2.33±2.37	3.62±3.64	<0.001**
AMPA (µg/g creatinine)	0.22±0.57	1.95±2.96	1.99±2.34	2.09±3.62	<0.001**
As (µg/g creatinine)	34.12±34.63	109.66±120.55	112.63±61.83	118.49±93.06	<0.001**
Cd (µg/g creatinine)	0.36±0.21	1.46±0.96	1.01±0.70	2.04±1.25	<0.001**
Pb (µg/L)	5.55±3.65	6.77±8.12	6.29±5.65	8.3±14.84	0.751
eGFR (mL/min/1.73m ²)	109.15±24.86	64.58±20.74	36.43±4.64	15.88±8.02	<0.001**
Creatinine (mg/dL)	0.78±0.21	1.12±0.33	1.81±0.29	4.80±2.69	<0.001**
Data showed as Mean ± SD, ** means statistical significant different					
@ Kruskal Wallis-test					

Table 6

The odds ratio (OR) of eGFR decrease by glyphosate, AMPA, Cd and As levels

N=155	OR (95% CI)	p-value@
Model 1:eGFR \geq 60 vs. <60 ¶		
Glyphosate (µg/g creatinine)	1.388 (0.896-2.150)	0.142
AMPA (µg/g creatinine)	0.959 (0.646-1.424)	0.836
Cd (µg/g creatinine)	7.567 (1.912-29.949)	0.004*
As (µg/g creatinine)	1.006 (0.997-1.015)	0.208
Model 2:eGFR \geq 45 vs. <45 ¶		
Glyphosate (µg/g creatinine)	1.566 (1.134-2.162)	0.006*
AMPA (µg/g creatinine)	0.998 (0.762-1.307)	0.987
Cd (µg/g creatinine)	1.851 (0.833-4.111)	0.131
As (µg/g creatinine)	1.008 (1.000-1.015)	0.038*
¶Adjustment for age, sex, BMI and interaction terms of glyphosate, Cd and As levels		
@Logistic regression model		

Figures




Figure 1-1




Figure 1-2




Figure 1-3




Figure 1-4




Figure 1-5

Figure 1

Trend of biomarkers of all participants in different CKD stage Note: The values with the same superscript letters are significantly different by Jonckheere Trend-test ($p < 0.05$).