Denonvilliers' Fascia Acts as the Fulcrum and Hammock for Continence After Radical Prostatectomy

Xuwei Lu

Minhang Hospital, FUDAN University, Shanghai

Chang He

Minhang Hospital, FUDAN University, Shanghai

Sihong Zhang

Zhongshan Hospital, FUDAN University, Shanghai 200032

Fan Yang

Minhang Hospital, FUDAN University, Shanghai

Zhuifeng Guo

Minhang Hospital, FUDAN University, Shanghai

Jiaqi Huang

Minhang Hospital, FUDAN University, Shanghai

Minke He

Minhang Hospital, FUDAN University, Shanghai

Jiawen Wu

Minhang Hospital, FUDAN University, Shanghai

Xia Sheng

Minhang Hospital, FUDAN University, Shanghai

Wenyao Lin

Xuhui Hospital, FUDAN University, Shanghai 200031

Jie Cheng

Xuhui Hospital, FUDAN University, Shanghai 200031

Jianming Guo

Zhongshan Hospital, FUDAN University, Shanghai 200032

Hang Wang (✉ wang.hang@zs-hospital.sh.cn)

Zhongshan Hospital, FUDAN University, Shanghai 200032

Research Article

Keywords: prostate cancer, prostatectomy, Denonvilliers' fascia, urinary continence, positive surgical margin

Posted Date: September 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-883609/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Urology on December 1st, 2021. See the published version at <https://doi.org/10.1186/s12894-021-00943-z>.

Abstract

Background: Radical prostatectomy (RP) is the primary treatment of localized prostate cancer. Immediate urinary incontinence post-RP was still common and depressing without specific reason.

Methods: A multicenter cohort of 154 consecutive patients from 2018 to 2020, who was diagnosed with localized prostate cancer underwent either modified mini-incision retropubic radical prostatectomy (Mmi-RRP) or laparoscopic radical prostatectomy (LRP) or robotic-assisted radical prostatectomy (RARP). Seventy-two patients with DF spared were included in DFS group. Whereas eighty-two patients with DF completely or partially dissected were set as Group Control. The primary outcome was Immediate continence (ImC). Continuous data and categorical data were analyzed with *t*-test and *Chi*-square test, respectively. Odds ratios (*ORs*) were calculated with logistic regression.

Results: Urinary continence of Group DFS was significantly better than that of Group Control at each time point within one year after operation. Incidence rate of continence in Group DFS and Group Control were 83.3% vs 13.4% ($P<0.001$) for ImC, 90.3% vs 30.5% ($P<0.001$) at 3months, 91.7% vs 64.6% ($P<0.001$) at 6months, and 93.1% vs 80.5% ($P=0.023$) at 1year after operation, respectively. Positive surgical margin (PSM) showed no significant difference (20.8% vs 20.7%, $P=0.988$). In multivariate analysis, DFS showed importance for ImC post RP ($OR=26.418$, $P<0.001$).

Conclusions: Denonvilliers' fascia acted as the fulcrum and hammock for continence post RP. Preservation of DF contributed to better continence after RP without increase of PSM.

Trail registration: Our research was conducted retrospectively and approved by the ethical committees of Minhang Hospital, but not registered.

1. Introduction:

Prostate cancer (PCa) is the top-ranking malignant tumor for men worldwide^[1]. Radical prostatectomy (RP) is the primary therapeutic choice for localized PCa. Dramatic improvement of surgery technique and more application of newer devices have reduced the incidence of severe complications for RP. Unfortunately, incontinence after RP, which haunted patients extremely hard, is still a huge challenge. The morbidity of incontinence one-year after RP remains up to 21%, regardless whether open radical prostatectomy (ORP) or laparoscopic radical prostatectomy (LRP) or robotic-assisted radical prostatectomy (RARP) has been taken^[2]. Although many researchers have proposed many factors that might matter such as sphincter, length of membrane urethra, bladder neck preservation, neurovascular bundle (NVB) sparing, and Denonvilliers' fascia (DF) as well, that is still uncertainty about the mechanism of post RP continence^[3].

Furthermore, most literatures reported continent rate of short-term, mid-term and long-term post RP. As to immediate continence (ImC), few reports could be found. We have applied a new technique as DF-sparing

(DFS) since 2018 in RP with excellent urinary continence, especially the ImC. Because of the encouraging outcome of urinary control, there must be something worth thinking about.

2. Methods:

From January 2018 to October 2020, 154 consecutive patients with localized prostate cancer underwent RP in Zhongshan Hospital, Minhang Hospital and Xuhui Hospital, FUDAN University. Inclusion criteria were that patients must: (1) be diagnosed as localized prostate cancer (cT1 or cT2) and agree to choose surgery as their treatment; (2) have an estimated life expectancy of 10 years or more; (3) have no history of incontinence. Exclusion criteria were: (1) evidence of non-localized prostate cancer (cT3 or cT4); (2) previous pelvic radiotherapy or major pelvic surgery; (3) previous surgery for urothelial, prostate or bladder neck; (4) previous history of head injury, dementia, or psychiatric illness. Seventy-two cases that underwent either modified mini-incision retropubic radical prostatectomy (Mmi-RRP) or LRP or RARP with DFS were included in Group DFS. Group Control included 82 cases received Mmi-RRP/LRP/RARP with DF completely or partially dissected.

All the procedures were completed by three experienced surgeons. The demographic and tumor characteristics of patients are shown in Table-1. Urinary catheter was removed 5 ~ 7days after operation. All the patients were followed up for 2 ~ 26 (14.51 ± 6.85) months.

The primary outcome was ImC. Continent status reported by patients was collected at 1-week, 3-month, 6-month, and 1-year after urethral catheter was discarded. We defined continence as usage of ≤ 1 pad per 24 hours. Considering that urine leakage caused by multiple factors was prevalent shortly after catheter removal, most of which was transient and could recover soon without treatment, we defined ImC as regaining continence within 1 week after urinary catheter removal^[4].

2.1 Surgery and post-surgery protocol:

All the procedures were operated out of peritoneal under general anesthesia. Mmi-RRP was operated with an approximate 7cm longitudinal suprapubic incision, whereas LRP or RARP was with 3 ~ 4 poles. DF was thought to be composed of multiple layers and ensheathed both seminal vesicles and vas deferens into the anterior layer. After isolation of seminal vesicles, anterior layer of DF was isolated at the level of seminal vesicle triangle, and the dissection was located between prostate capsule (PC) and DF towards prostate apex in order to keep DF intact (Figure-1, Figure-2 f-h). For the sake of cavernous nerves protection, blunt dissection was preferred. In consideration of potential nerve fiber damage, energy devices were seldom used during dissection near prostate.

Unfortunately, it was difficult to keep DF intact sometimes. In locally advanced PCa cases, the tight junction between DF and prostate added extra difficulty to LRP procedures. In these situations, DF would be resected partially or completely together with prostate (Figure-2, a-d). As to NVB sparing, careful dissection along PC in post-lateral direction should be enforced in some suitable cases.

No bladder neck reconstruction was preferred in all cases. It was necessary to dissect close to prostate and keep bladder neck intact. But some situations, for example, locally advanced PCa, big prostate volume, and irregularly shaped prostate could make it hard to keep bladder neck intact. Under these circumstances, reconstruction of bladder neck was made to restore anastomosis, always by suturing bladder neck at 6 o'clock position.

2.2 Statistical analysis:

The pathology images were processed with K-viewer version 1.5.3.1 (KFBIO, Zhejiang, CHINA). The continuous data and categorical data were analyzed with *t*-test and *Chi*-square test, respectively. Kaplan-Meier method was used to compute urinary continence curve. Binary logistic regression analysis was applied to screen predictors for ImC. All data analyses were performed using SPSS 22.0 statistical software (IBM SPSS, Chicago, IL, USA). In all analysis, $P < 0.05$ was considered statistically significant.

3. Results:

All the 154 consecutive patients completed RRP successfully and were followed-up for (14.51 ± 6.85) months, ranging from 2 to 26 months. Details of procedures and pathology post-surgery were shown in Table 2. There was shorter duration of surgery ($P = 0.001$), more NVB sparing ($P = 0.001$), and less seminal vesicle invasion ($P = 0.04$) in Group DFS vs Group Control. Especially, the post-surgery continence condition was better in Group DFS than in Group Control at each time point from ImC to 3 months, 6 months ($P < 0.001$) and 12 months ($P = 0.023$). In Group DFS-pilot, even better ImC (87%) and lower PSM (8.7%) were achieved (Table-2, Figure-3).

Table 1
Demographics and tumor characteristics of patients

	Group DFS (n = 72)	Group Control (n = 82)	<i>P</i>
Mean(Range)			
Age, years	67.83 ± 5.02 (58–79)	69.27 ± 6.51 (54–80)	0.125
BMI	24.49 ± 2.55 (17.36–30.85)	24.53 ± 2.74 (16.79–30.32)	0.919
Diagnosis PSA, ng/ml	16.02 ± 12.47 (0.38–60.96)	38.89 ± 114.67 (4.06–946.60)	0.077
Prostate Volume, ml	49.20 ± 15.75 (13.5-100.7)	51.49 ± 18.49 (16.7-122.5)	0.416
Clinical stage, n (%)			0.820
T1	30 (41.7)	33 (40.2)	
T2	38 (52.8)	46 (56.1)	
T3	4 (5.6)	3 (3.7)	
Diagnosis Gleason, n (%)			0.210
≤ 6	31 (43.1)	26 (31.7)	
7	31 (43.1)	37 (45.1)	
≥ 8	10 (13.9)	19 (23.2)	
Diagnosis method, n (%)			0.516
Biopsy	66 (91.7)	78 (95.1)	
TURP	6 (8.3)	4 (4.9)	
IIEF-5 Q ₂ ≥ 3 pre-surgery, n (%)	49 (68.1)	47 (57.3)	0.170
Comorbid condition, n (%)			
CAD	9 (12.5)	10 (12.2)	0.954
DM	12 (16.7)	17 (20.7)	0.520

BMI: body mass index; CAD: coronary artery disease; COPD: chronic obstruction pulmonary diseases; CVD: cerebrovascular diseases; DM: diabetes mellitus; HBP: high blood pressure; TURP: transurethral resection of prostate.

IIEF-5 Q2: International Index of Erectile Function-5, question NO.2. "When you had erections with sexual stimulation, how often was your erection hard enough for penetration during the last 3 months?" The following responses were available: "No sexual activity" (0); "Almost never or never" (1); "A few times (much less than half the time)" (2); "Sometimes (about half the time)" (3); "Most times (much more than half the time)" (4); and "Almost always or always" (5).

	Group DFS (n = 72)	Group Control (n = 82)	<i>P</i>
HBP	38 (52.8)	39 (47.6)	0.518
COPD	8 (11.1)	4 (4.9)	0.150
CVD	4 (5.6)	6 (7.3)	0.751
BMI: body mass index; CAD: coronary artery disease; COPD: chronic obstruction pulmonary diseases; CVD: cerebrovascular diseases; DM: diabetes mellitus; HBP: high blood pressure; TURP: transurethral resection of prostate.			
IIEF-5 Q2: International Index of Erectile Function-5, question NO.2. "When you had erections with sexual stimulation, how often was your erection hard enough for penetration during the last 3 months?" The following responses were available: "No sexual activity" (0); "Almost never or never" (1); "A few times (much less than half the time)" (2); "Sometimes (about half the time)" (3); "Most times (much more than half the time)" (4); and "Almost always or always" (5).			

Table 2
Outcomes of surgery

Group	DFS, n (%)	Control, n (%)	P
Mean(Range)			
Surgery duration, min	153.19 ± 66.41 (100–400)	188.29 ± 65.06 (90–480)	0.001
Blood loss, ml	143.06 ± 107.24 (20–800)	132.44 ± 136.88 (20-1000)	0.597
NVB sparing			0.001
Bilateral	27 (37.5)	24 (29.3)	
Unilateral	13 (18.1)	2 (2.4)	
None	32 (44.4)	56 (68.3)	
LND	63 (87.5)	74 (90.2)	0.588
pT			0.472
T0	1 (1.4)	0 (0)	
T2	53 (73.6)	57 (69.5)	
T3	18 (25.0)	25 (30.5)	
Gleason scores			0.097
0	1 (1.4)	0 (0)	
≤ 6	17 (23.6)	9 (11)	
7	38 (52.8)	50 (61)	
≥ 8	16 (22.2)	23 (28)	
positive LN	0 (0)	3 (3.7)	0.248
SV invasion	7 (9.7)	18 (22)	0.040
PSM	15 (20.8)	17 (20.7)	0.988
Sites of PSM			0.344
One-site (<i>location; %</i>)	13 (<i>5 basal, 7 apical, 1 post-lateral; 18%</i>)	11 (<i>2 basal, 8 apical, 1 post-lateral; 13.4%</i>)	
Multiple-sites (<i>locations; %</i>)	2 (<i>2 apical and post-lateral; 2.8%</i>)	6 (<i>5 apical and basal, 1 apical, basal and post-lateral; 7.3%</i>)	
Post-lateral PSM	3 (4.2%)	2 (2.4%)	0.665

LND: lymph node dissection; NVB: nerve vascular bundle; PSM: positive surgical margin; SV: seminal vesicles.

Group	DFS, n (%)	Control, n (%)	<i>P</i>
IIEF-5 Q ₂ ≥ 3 post-surgery	25 (34.7)	14 (17.1)	0.012
Continence			< 0.001
Immediate-	60 (83.3)	11 (13.4)	< 0.001
3months-	65 (90.3)	25 (30.5)	< 0.001
6months-	66 (91.7)	53 (64.6)	< 0.001
12months-	67 (93.1)	66 (80.5)	0.023
LND: lymph node dissection; NVB: nerve vascular bundle; PSM: positive surgical margin; SV: seminal vesicles.			

Table 3
Logistic regression analysis for predictors of immediate continence

ImC	Univariate	Multi-variate		
	OR (95%CI)	P	OR (95%CI)	P
Surgery type		< 0.001		0.551
Mmi-RRP vs LRP	12.995 (5.685–29.703)	< 0.001	1.851 (0.394–8.687)	0.435
RARP vs LRP	1.226 (0.379–3.967)	0.734	0.762 (0.162–3.586)	0.731
LND, Yes vs No	0.735 (0.268–2.018)	0.550		
Gleason score	0.659 (0.458–0.947)	0.024	0.924 (0.533–1.603)	0.779
SV invasion, Yes vs No	0.239 (0.084–0.674)	0.007	0.257 (0.051–1.310)	0.102
PSM, Yes vs No	0.887 (0.405–1.942)	0.764		
TURP before, Yes vs No	1.182 (0.328–4.260)	0.798		
Diagnosis PSA	0.998 (0.992–1.003)	0.439		
IIEF-5 Q ₂ pre-surgery	1.392 (1.039–1.865)	0.027	1.576 (0.924–2.688)	0.095
Prostate volume	0.996 (0.977–1.014)	0.645		
Age	0.961 (0.910–1.016)	0.159	1.033 (0.929–1.148)	0.552
BMI	0.943 (0.836–1.065)	0.343		
Surgery duration	0.986 (0.979–0.992)	< 0.001	0.991 (0.981–1.002)	0.108
Blood loss	0.999 (0.996–1.001)	0.327		
NVB sparing		0.012		0.924
Unilateral vs No	4.812 (1.415–16.367)	0.012	0.763 (0.135–4.316)	0.759
Bilateral vs No	2.130 (1.056–4.299)	0.035	1.089 (0.310–3.830)	0.894
DF sparing, Yes vs No	32.273 (13.288–78.384)	< 0.001	26.418 (8.382–83.263)	< 0.001
BMI: body mass index; CI: confidence interval; ImC: immediate continence; LND: lymph node dissection; NVB: nerve vascular bundle; OR: odds ratio; PSM: positive surgical margin; SV: seminal vesicles.				

In univariate analysis, surgery type (OR = 1.226 ~ 12.995, $P < 0.001$), surgery duration (OR = 0.986, $P < 0.001$), IIEF-5 Q₂ pre-surgery (OR = 1.392, $P = 0.027$), Gleason score (OR = 0.659, $P = 0.024$), SV invasion (OR = 0.239, $P = 0.007$), NVB sparing (OR = 2.130 ~ 4.812, $P = 0.012$) and DF sparing (OR = 32.273, $P < 0.001$) were significantly related with ImC. While in multivariate analysis, only DF sparing was proved to be statistically significant for ImC, OR = 26.418, $P < 0.001$ (Table-3).

As to definition of sexual dysfunction, we used cutoff value between response 2 and 3 for question NO.2 in IIEF-5^[2]. There were more potent patients in Group DFS than in Group Control at 12months post-surgery, (34.7% vs 17.1%, $P = 0.012$). No statistical difference of PSM was found between two groups (20.8% vs 20.7%, $P = 0.988$, Table-1).

4. Discussion:

Tumor control, urinary continence and sexual potency make up the Trifecta of RP. Post-surgery incontinence is one of the most functionally devastating complications, which could make patients too depressed to maintain social relationship and can even lead to suicide. Albkri A *et al*^[5] figured out that 28.1% patients regretted for RP decision because of complications. Unfortunately, there is so far no universal definition of continence up to now, which makes the morbidity of continence post-RP vary^[6]. As literatures reported, ImC varied from 18%^[7] to 63.5%^[8], while continence at 3 months from 45%^[9] to 82%^[10], and continence at 12months from 79.8%^[2] to 91%^[11]. In consideration of urinary consequences on record, morbidity of incontinence after RP in real world could probably be higher than reported^[12]. Grivas N *et al*^[13] pointed out that continence at 12months post-surgery was only 49.2% with strict definition.

The debate between cancer free and continence still lacks conclusion. Routinely, the more tissue near the tumor incised, the better tumor control. But it also means more potential functional structures to be cut, and incontinence is more likely to happen. Therefore, the discovery of landmark structure for urinary continence is needed. There are many hypotheses, such as membrane urethra length, urethral sphincter complex, detrusor apron of bladder neck, pelvic floor musculature, DVC, and NVB^[3], *etc*. Asimakopoulos AD *et al*^[14] preferred complete periprostatic anatomy preservation to protect maximal functional structure and got 100% continence (≤ 1 pad per day) just at the time of catheter removal. However, this procedure may not be suitable for patients with higher tumor stage. Here we suppose DF to be the answer, particularly for the early-term and mid-term urinary control after RP.

DF has been demonstrated to be a multiple-layer structure both in cadaveric studies^[15, 16] and histologic studies^[17, 18] on specimen of RP. At the central-post direction of prostate and proximal of urethra, DF's fascicles tend to fuse and adhere with PC to form a dorsal raphe (DR)^[16, 18]. The tendinous DR, which extends distally to prostate apex and ends at perineal tendon, may act as a fulcrum to support prostate and proximal urethra^[19]. At the lateral-post direction of prostate, DF disperses to connect with LAF (Figure-2). Dalpiaz O *et al*^[19] described DF as part of musculofascial suspension system, stabilizing prostate apex and proximal urethra. Interestingly, there was similar view on stress urinary incontinence (SUI) of females. DeLancey^[20] delineated endopelvic fascia and anterior vaginal wall as the hammock-like supportive layer of urethra, which could stabilize and help to close urethra during cough. Sling, placing a supportive material behind urethra, had been a staple procedure for SUI and withstood the test of time^[21]. Back to continence after RP, we hypothesized "hammock theory" still work on and preferred DF to be the critical structure (Figure-1, 2 and Figure-4), which could uplift vesicourethral anastomosis. Meanwhile, DR just acted as the fulcrum beneath urethra and contributed to closure of it when abdomen

pressure increased. Hence, together with DR (also part of DF), DF was supposed to be the critical landmark for continence post RP. Our research also proved that DFS could significantly improve ImC ($OR = 26.418$, $P < 0.001$). However, because of individual difference, DF or DR could not be kept intact in every operation. So, we proposed a grading system for DFS procedure and our study showed that different grades can indicate different continence prognosis after RP (Figure-4).

Considering that the scattering neurovascular plexus, which innervates corpora cavernosa and sphincter complex, is embedded in the multiple-layers of DF^[3, 22], there was a minority of nerve fibers located in front of DF which would be dissected inevitably^[23], while most of them are located in post-lateral and post regions of prostate^[3] (Figure-1). Tewari *et al*^[24] proposed a nerve-sparing (NS) grading system based on “landmark vein (LV)” lateral to prostate. An intra-fascial dissection was defined as grade 1, while an extra-fascial dissection was defined as grade 4. Srivastava *et al*^[25] demonstrated that grade 1 achieved more early continence than grade 4 (71.8% vs 43.5%), which may account for incomplete continence situation with intact DF.

DF-sparing technique doesn't equal to complete periprosthetic anatomy preservation or intra-fascial dissection. DFS procedure emphasizes dissecting before anterior layer of DF to protect the whole DF fascicles and the majority of neurovascular fibers, which is a procedure with a specific target to be spared, making the incision area selective but thorough. Whereas complete peri-prostate sparing technique aims to preserve functional structure as much as possible^[14]. However, good functional recovery may sacrifice safety of tumor control. DF-targeted sparing procedure provides a compromised solution to the dilemma of urinary- or tumor-control. In our research, Group DFS achieved 83.3% ImC, 90.3% continence in 3 months, 91.7% in 6 months, 93.1% in 12 months, better than Group Control at every time point with no difference in PSM (20.8% vs 20.7%, $P = 0.988$), especially no difference of PSM in post-lateral direction of prostate (4.2% vs 2.4%, $P = 0.665$).

Another key point to protect neurovascular plexus is the limited usage of energy devices, such as ultrasonic or electronic scalpels, bipolar coagulator, especially during prostate apex anatomy^[26], in which nerve fibers penetrate DF or periprosthetic tissue to innervate urethral sphincter. The electrical and thermal conduction may cause extra damage to neurovascular fibers nearby. Thereby, “blunt”, or “cold” dissection for prostate apex with finger, forceps, or scissors is preferred. The NS grading system proposed by Patel *et al*^[27] proved the idea. In Patel's system, grade 5 NS was performed between “landmark artery (LA)” lateral to prostate and PC with neither sharp dissection nor energy devices, which resulted in complete nerve sparing (> 95%). Grade 4 NS (75%) was also performed between LA and PC with sharp dissection but without energy devices.

Last but not least, we preferred to pay more attention to ImC post RP. Temml C *et al*^[28] testified that QoL wasn't related to duration of incontinence but frequency and degree of incontinence, need for pads, *etc*. Coyne KS *et al*^[29] pointed out remarkable prevalence of 19.1% anxiety and 6.6% depression in patients

with urinary incontinence (UI). Anxiety may trigger and aggravate UI in return. So better ImC may guarantee not only “extra” better continence in the future but also positive mental health.

Indeed, there are limitations in present research including small sample size, retrospective, and lack of randomization. Therefore, a prospective randomized controlled study with larger sample size should be organized for further verification.

5. Conclusion:

Denonvilliers’ fascia was supposed to act as the fulcrum and hammock for urinary control after RP. We recommend to preserve DF, minimize utility of energy devices during dissecting, which could probably contribute to best continence after RP without increase of PSM.

Abbreviations

BMI: body mass index

CAD: coronary artery disease

COPD: chronic obstruction pulmonary diseases

CVD: cerebrovascular diseases

DM: diabetes mellitus

DF: Denonvillier’s fascia

DFS: DF-sparing

DR: dorsal raphe

DVC: dorsal vascular complex

HBP: high blood pressure

HE: hematoxylin and eosin

IFT: inter fascial tissue

IIEF-5: International Index of Erectile Function-5

ImC: immediate continence

LAF: levator ani fascia

LND: lymph node dissection

LRP: laparoscopic radical prostatectomy

Mmi-RRP: mini-incision retropubic radical prostatectomy

NVB: neurovascular bundle

ORP: open radical prostatectomy

PC: prostate capsule

PSM: Positive surgical margin

RARP: robotic-assisted radical prostatectomy

RP: Radical prostatectomy

TURP: transurethral resection of prostate

SUI: stress urinary incontinence

SV: seminal vesicles

VD: vas deferens

Declarations

Ethics approval and consent to participate: The study was performed in accordance with the Declaration of Helsinki. The ethical committees of Minhang Hospital approved the retrospective multicentric study. Due to the retrospective nature of the study, informed consent was waived.

Consent for publication: The authors confirm that the work described has not been published before.

Availability of data and material: All data generated or analyzed during this study are included in this published article.

Competing interests: The authors declare that they have no competing interests.

Funding: Not applicable.

Authors' Contribution: Project developed by HW, JMG. Manuscript written by XWL, edited by HW, CH and SHZ. Data collected by FY, ZFG, WYL and JC, analyzed by JQH, MKH and JWW. Pathological section managed and analyzed by XS.

Acknowledgements: We sincerely thank Ms. Miao Gong for production of pathological sections, and Dr. Yuan Ji for support in pathological image scanning and enlightening advices.

Authors' information:

1. Xuwei Lu, MD, Department of Urology, Minhang Hospital, FUDAN University, Shanghai 201199.
2. Chang He, MD, Department of Urology, Minhang Hospital, FUDAN University, Shanghai 201199.
3. Sihong Zhang, MD, Department of Urology, Zhongshan Hospital, FUDAN University, Shanghai 200032.
4. Fan Yang, MD, Department of Urology, Minhang Hospital, FUDAN University, Shanghai 201199.
5. Zhuifeng Guo, MD, Department of Urology, Minhang Hospital, FUDAN University, Shanghai 201199.
6. Jiaqi Huang, MD, Department of Urology, Minhang Hospital, FUDAN University, Shanghai 201199.
7. Minke He, MD, Department of Urology, Minhang Hospital, FUDAN University, Shanghai 201199.
8. Jiawen Wu, MD, Department of Urology, Minhang Hospital, FUDAN University, Shanghai 201199.
9. Xia Sheng, PhD, Department of Pathology, Minhang Hospital, FUDAN University, Shanghai 201199.
10. Wenyao Lin, MD, Department of Urology, Xuhui Hospital, FUDAN University, Shanghai 200031.
11. Jie Cheng, MD, Department of Urology, Xuhui Hospital, FUDAN University, Shanghai 200031.
12. Jianming Guo, PhD, Department of Urology, Zhongshan Hospital, FUDAN University, Shanghai 200032, E-mail: guo.jianming@zs-hospital.sh.cn (J. GUO).
13. Hang Wang, PhD, Department of Urology, Zhongshan Hospital, FUDAN University, Shanghai 200032, E-mail: wang.hang@zs-hospital.sh.cn (H. Wang).

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin*. 2019. 69(1): 7–34. DOI: 10.3322/caac.21551.
2. Haglind E, Carlsson S, Stranne J, Wallerstedt A, Wilderäng U, Thorsteinsdottir T, et al. Urinary Incontinence and Erectile Dysfunction After Robotic Versus Open Radical Prostatectomy: A Prospective, Controlled, Nonrandomised Trial. *Eur Urol*. 2015. 68(2): 216–225. DOI: 10.1016/j.eururo.2015.02.029.
3. Walz J, Epstein JI, Ganzer R, Graefen M, Guazzoni G, Kaouk J, et al. A Critical Analysis of the Current Knowledge of Surgical Anatomy of the Prostate Related to Optimisation of Cancer Control and Preservation of Continence and Erection in Candidates for Radical Prostatectomy: An Update. *Eur Urol*. 2016. 70(2): 301–311. DOI: 10.1016/j.eururo.2016.01.026.
4. Grasso AA, Mistretta FA, Sandri M, Cozzi G, De Lorenzis E, Rosso M, et al. Posterior musculofascial reconstruction after radical prostatectomy: an updated systematic review and a meta-analysis. *BJU Int*. 2016. 118(1): 20–34. DOI: 10.1111/bju.13480.
5. Albkri A, Girier D, Mestre A, Costa P, Droupy S, Chevrot A. Urinary Incontinence, Patient Satisfaction, and Decisional Regret after Prostate Cancer Treatment: A French National Study. *Urol Int*. 2018. 100(1): 50–56. DOI: 10.1159/000484616.

6. Liss MA, Osann K, Canvasser N, Chu W, Chang A, Gan J, et al. Continence definition after radical prostatectomy using urinary quality of life: evaluation of patient reported validated questionnaires. *J Urol*. 2010. 183(4): 1464–1468. DOI: 10.1016/j.juro.2009.12.009.
7. John H, Hauri D. Seminal vesicle-sparing radical prostatectomy: a novel concept to restore early urinary continence. *Urology*. 2000. 55(6): 820–824. DOI: 10.1016/s0090-4295(00)00547-1.
8. Moinzadeh A, Shunaigat AN, Libertino JA. Urinary incontinence after radical retropubic prostatectomy: the outcome of a surgical technique. *BJU Int*. 2003. 92(4): 355–359. DOI: 10.1046/j.1464-410x.2003.04348.x.
9. Hollabaugh RS, Dmochowski RR, Kneib TG, Steiner MS. Preservation of putative continence nerves during radical retropubic prostatectomy leads to more rapid return of urinary continence. *Urology*. 1998. 51(6): 960–967. DOI: 10.1016/s0090-4295(98)00128-9.
10. Walsh PC, Marschke PL. Intussusception of the reconstructed bladder neck leads to earlier continence after radical prostatectomy. *Urology*. 2002. 59(6): 934–938. DOI: 10.1016/s0090-4295(02)01596-0.
11. Coughlin GD, Yaxley JW, Chambers SK, Occhipinti S, Samaratunga H, Zajdlewicz L, et al. Robot-assisted laparoscopic prostatectomy versus open radical retropubic prostatectomy: 24-month outcomes from a randomised controlled study. *Lancet Oncol*. 2018. 19(8): 1051–1060. DOI: 10.1016/S1470-2045(18)30357-7.
12. Nelson JB. The Ongoing Challenge of Urinary Incontinence after Radical Prostatectomy. *J Urol*. 2017. 198(6): 1223–1225. DOI: 10.1016/j.juro.2017.09.053.
13. Grivas N, van der Roest R, Schouten D, Cavicchioli F, Tillier C, Bex A, et al. Quantitative assessment of fascia preservation improves the prediction of membranous urethral length and inner levator distance on continence outcome after robot-assisted radical prostatectomy. *Neurourol Urodyn*. 2018. 37(1): 417–425. DOI: 10.1002/nau.23318.
14. Asimakopoulos AD, Annino F, D'Orazio A, Pereira CF, Mugnier C, Hoepffner JL, et al. Complete periprostatic anatomy preservation during robot-assisted laparoscopic radical prostatectomy (RALP): the new pubovesical complex-sparing technique. *Eur Urol*. 2010. 58(3): 407–417. DOI: 10.1016/j.eururo.2010.04.032.
15. Ghareeb WM, Wang X, Chi P, Wang W. The 'multilayer' theory of Denonvilliers' fascia: anatomical dissection of cadavers with the aim to improve neurovascular bundle preservation during rectal mobilization. *Colorectal Dis*. 2020. 22(2): 195–202. DOI: 10.1111/codi.14850.
16. Xu Z, Chapuis PH, Bokey L, Zhang M. Denonvilliers' fascia in men: a sheet plastination and confocal microscopy study of the prerectal space and the presence of an optimal anterior plane when mobilizing the rectum for cancer. *Colorectal Dis*. 2017. DOI: 10.1111/codi.13906.
17. Secin FP, Karanikolas N, Gopalan A, Bianco FJ, Shayegan B, Touijer K, et al. The anterior layer of Denonvilliers' fascia: a common misconception in the laparoscopic prostatectomy literature. *J Urol*. 2007. 177(2): 521–525. DOI: 10.1016/j.juro.2006.09.028.

18. Kiyoshima K, Yokomizo A, Yoshida T, Tomita K, Yonemasu H, Nakamura M, et al. Anatomical features of periprostatic tissue and its surroundings: a histological analysis of 79 radical retropubic prostatectomy specimens. *Jpn J Clin Oncol*. 2004. 34(8): 463–468. DOI: 10.1093/jjco/hyh078.
19. Dalpiaz O, Anderhuber F. The fascial suspension of the prostate: A cadaveric study. *Neurourol Urodyn*. 2017. 36(4): 1131–1135. DOI: 10.1002/nau.23073.
20. DeLancey JO. Structural support of the urethra as it relates to stress urinary incontinence: the hammock hypothesis. *Am J Obstet Gynecol*. 1994. 170(6): 1713-20; discussion 1720–1723. DOI: 10.1016/s0002-9378(94)70346-9.
21. Schreiner G, Beltran R, Lockwood G, Takacs EB. A timeline of female stress urinary incontinence: how technology defined theory and advanced treatment. *Neurourol Urodyn*. 2020. 39(6): 1862–1867. DOI: 10.1002/nau.24407.
22. Muraoka K, Hinata N, Morizane S, Honda M, Sejima T, Murakami G, et al. Site-dependent and interindividual variations in Denonvilliers' fascia: a histological study using donated elderly male cadavers. *BMC Urol*. 2015. 15: 42. DOI: 10.1186/s12894-015-0034-5.
23. Kinugasa Y, Murakami G, Uchimoto K, Takenaka A, Yajima T, Sugihara K. Operating behind Denonvilliers' fascia for reliable preservation of urogenital autonomic nerves in total mesorectal excision: a histologic study using cadaveric specimens, including a surgical experiment using fresh cadaveric models. *Dis Colon Rectum*. 2006. 49(7): 1024–1032. DOI: 10.1007/s10350-006-0557-7.
24. Tewari AK, Srivastava A, Huang MW, Robinson BD, Shevchuk MM, Durand M, et al. Anatomical grades of nerve sparing: a risk-stratified approach to neural-hammock sparing during robot-assisted radical prostatectomy (RARP). *BJU Int*. 2011. 108(6 Pt 2): 984–992. DOI: 10.1111/j.1464-410X.2011.10565.x.
25. Srivastava A, Chopra S, Pham A, Sooriakumaran P, Durand M, Chughtai B, et al. Effect of a risk-stratified grade of nerve-sparing technique on early return of continence after robot-assisted laparoscopic radical prostatectomy. *Eur Urol*. 2013. 63(3): 438–444. DOI: 10.1016/j.eururo.2012.07.009.
26. Tewari A, Takenaka A, Mtui E, Horninger W, Peschel R, Bartsch G, et al. The proximal neurovascular plate and the tri-zonal neural architecture around the prostate gland: importance in the athermal robotic technique of nerve-sparing prostatectomy. *BJU Int*. 2006. 98(2): 314–323. DOI: 10.1111/j.1464-410X.2006.06266.x.
27. Schatloff O, Chauhan S, Sivaraman A, Kameh D, Palmer KJ, Patel VR. Anatomic grading of nerve sparing during robot-assisted radical prostatectomy. *Eur Urol*. 2012. 61(4): 796–802. DOI: 10.1016/j.eururo.2011.12.048.
28. Temml C, Haidinger G, Schmidbauer J, Schatzl G, Madersbacher S. Urinary incontinence in both sexes: prevalence rates and impact on quality of life and sexual life. *Neurourol Urodyn*. 2000. 19(3): 259–271. DOI: 10.1002/(sici)1520-6777(2000)19:3<259::aid-nau7>3.0.co;2-u.
29. Coyne KS, Kvasz M, Ireland AM, Milsom I, Kopp ZS, Chapple CR. Urinary incontinence and its relationship to mental health and health-related quality of life in men and women in Sweden, the

Figures

Figure 1

Scenes of Mmi-RRP procedure with DFS technique and pathological section through paramedian of prostate, stained by HE. (a): DF (white dot line area) kept intact after prostate removal, also shown on schematic drawing (e). (b): Anterior layer of DF cut open at seminal vesicles triangle level. Dissection was

kept between DF and PC towards apex of prostate. Black arrows exhibited the cut-line of DF's anterior layer. (c): No DF structure in the backward direction of prostate. (d): The cut-line of DF on the pathological section (red dot line circle), fused with seminal vesicle fascia propria (black dot line circle). (e-f): Schematic drawing OF dissecting bladder and prostate junction (green dot line). B: bladder; HE: hematoxylin and eosin; IFT: inter fascial tissue; N: nerve; P: prostate; PB: pubic bone; PC: prostate capsule; R: rectum; SV: seminal vesicles; VD: vas deferens. Black arrow: DF; Black arrow head: PC; White arrow: seminal vesicle fascia propria.

Figure 2

Transverse sections at mid-prostate in Group Control (a-d) and Group DFS (f-h), both stained by HE. (a): Overall vision of pathological section from Group Control. (b): In 6-o'clock direction of prostate, two layers of DF were dissected with prostate. (c): In 4-o'clock direction, anterior layer of DF extended anterolaterally alongside with prostate capsule, sometime fused together (show in black dot line circle). Middle layer of DF and NVB were removed with prostate. (d): Anterior layer of DF extended anterolaterally and connected with levator ani fascia (LAF) at 3-o'clock direction. (e): Schematic drawing simulated peri-prostate fascia structure including DF and LAF. (f): Overall vision of pathological section from Group DFS. (g): In 6-o'clock direction of prostate, few DR fragments without DF layers. (h): In 8-o'clock direction, DF was spared successfully with no DF structure visible in the specimen. DR: dorsal raphe; DVC: dorsal vascular complex; IFT: inter fascial tissue; LAF: levator ani fascia; N: nerve; P: prostate; PB: pubic bone; R: rectum; V: vessel. Black arrow: DF; Black arrow head: PC; Black rectangle: anterior layer of DF adhered with LAF; Red dot line: cut line of LAF.

Figure 3

Kaplan-Meier analysis for time to urinary continence post-surgery

Figure 4

Different grades of DFS in RP, stained by hematoxylin and eosin. (a): In Grade-1 DF sparing procedure, neither DF nor DR was cut with prostate. (b): In Grade-2 DFS procedure, DF was preserved while DR fragments removed with prostate. (c-d): In Grade-2 DFS procedure, DF spared successfully while DR was totally cut away. (e-f): Grade-3 DFS was defined as neither DF nor DR spared. Black arrow head: prostate capsule; Ca: prostate cancer; DR: dorsal raphe (show in blue dot line circle); P: prostate; U: urethra. Black arrow: DF; Black arrow head: PC.