Non-linear Relationships Between Density and Demographic Traits of Aedes

Logan A. Sauers (✉ Lsauers@ilstu.edu)

Illinois State University

Kelsey E. Hawes

Illinois State University

Steven A. Juliano

Illinois State University

Research Article

Keywords: Density-dependence, population dynamics, per capita growth, *Aedes triseriatus*, *Aedes aegypti*, *Aedes albopictus*

Posted Date: September 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-883857/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Understanding the relationship of population dynamics to density is central to many ecological investigations. Despite the importance of density-dependence in determining population growth, the empirical relationship between density and *per capita* growth remains understudied in most systems and is often assumed to be linear. In experimental studies of interspecific competition, investigators often evaluate the predicted outcomes by assuming such linear relationships, fitting linear functions, and estimating parameters of competition models. In this paper, we tested experimentally the shape of the relationship between estimated population rate of change and initial density using laboratory-reared populations of three mosquito species. We estimated *per capita* growth rate for these experimental populations over a thirty-fold range of larval densities at a standard resource abundance. We then compared fits of linear models and several different nonlinear models for the relationship of estimated rate of change and density. We find that the relationship between density and *per capita* growth is strongly non-linear in all three mosquitoes. Components of population growth (survivorship, development time, adult size) are also nonlinearly related to initial density. The causes and consequences of this nonlinearity are likely to be important issues for population and community ecology.

Introduction

A population's rate of change is central to theory and empirical studies of ecology [1–5], evolution [6–8], and disease transmission [9, 10]. Many populations will have population growth that is affected by population density [11], and density dependent effects on population rate of change are thus central to understanding population dynamics. Density dependence may result from resource competition [12, 13], availability of mates [14], social interactions [15], or dispersal [16]. Resource limitation is likely the most postulated process producing density dependent effects via increasing death rates, decreasing birth rates, or decreasing individual growth rates [17, 18]. Ultimately, when these density dependent effects act on individuals the population's *per capita* growth rate is influenced. Such density dependent effects, acting on individuals, ultimately influence *per capita* growth rate [19, 20].

Understanding density dependent effects on population growth can be especially important for investigations of disease vectors [18, 21–26]. Vectors are often targets of control efforts that may change population density, and thus may alter survival, growth, development, or fecundity. For vector mosquitoes, both field and laboratory studies indicate that density dependent effects on larvae can be strong, and can impact survival, individual growth rate (and resulting size and fecundity), and development rate (and resulting age at first reproduction) [e.g., 21, 23, 24, 27–33].

Ecological models of population *per capita* rate of change, such as Verhulst's logistic or Lotka-Volterra models, postulate a linear relationship of *per capita* growth to intra- or inter-specific density [1, 34, 35]; however, for most organisms the relationship between density and *per capita* growth rate is not well known [36–38], and a linear relationship is only one possibility. Many of the relationships between density and the demographic components of population dynamics (e.g., survivorship, fecundity,

reproductive age) are non-linear [34]. Multiple studies point out the apparent flaw in the assumption of linearity, and among the few studies explicitly testing the form of the relationship between density and *per capita* growth rates, nonlinearity has been documented repeatedly [e.g., 11, 35, 39–41]. Additionally, individual based models of resource competition suggest that multiple aspects of species biology can produce nonlinear relationships of *per capita* growth to population density [4]. Whether relationships between *per capita* growth and population density are linear is an important question, as nonlinearity would indicate that the effects of population density and resource use on population growth rate are not uniform across density (as they are modeled in simple logistic growth). Thus, the effects of increasing density may depress growth most strongly at low or high densities, and therefore may relate to rates of population recovery from low density [11, 42].

Determining the relationship between population density and *per capita* growth is likely to improve predictions and models of vector populations. Models assuming linear effects of intraspecific density on *per capita* rate of increase have been used to estimate logistic growth models of vector mosquitoes [34, 43], or to test theoretical predictions about responses to mortality [5, 45]. Models assuming linear effects of both intra- and interspecific densities have been used often to evaluate potential for coexistence of competing vector species [24, 45–51]. Among these investigations, only [51] tested explicitly alternative models for the relationship of estimated growth rate to densities; they found the best model to be one with linear relationships of estimated finite *per capita* rate of increase to density for *A. aegypti*.

To investigate the relationship between density and *per capita* growth we use three important vector species: *Aedes aegypti*, *Aedes albopictus*, and *Aedes triseriatus*. These species are an excellent model system for studies of population dynamics and density dependence for several reasons. First, these container-dwelling mosquitoes appear to be often impacted by density dependent effects in nature [e.g., 21, 28, 31–33, 52]. As container-dwelling mosquitoes they are easily reared in the laboratory in conditions that simulate natural environments in a realistic way. All three have often been the subject of population studies using Livdahl and Sugihara's composite index of performance [34], which uses a life table approach to estimate *per capita* rate of change for large experimental cohorts [34]. Use of this index facilitates rearing of high-density cohorts with substantial replication because it removes the necessity of following adult females for their entire lives to obtain *per capita* rate of increase from a full life table [34]. Two separate laboratory studies have shown that this index is highly correlated with *per capita* rate of increase estimated from a full life table following reproduction of 100% of a cohort of females [41, 53]. Finally, all three species are important vectors of human diseases and have been targets of control efforts that may alter densities of larvae, thus altering density-dependent effects on population growth, production of adults, and other traits relevant to vectorial capacities [26].

Here, we investigate the relationship between larval density and the estimated *per capita* rate of increase as calculated by the index of performance, and the component variables that are used to calculate the index: survivorship to adulthood; adult female development time; and adult female size as a predictor of female fecundity. The primary objective was to determine whether the relationship of estimated *per capita* rate of increase to larval density is best fit by a linear logistic growth model, or alternative

nonlinear models: θ logistic; Gompertz; or polynomial. We hypothesized that the relationship between density and *per capita* growth is in fact non-linear. Additionally, we tested for nonlinearities in the relationships of component variables (survivorship, development time, adult size) to larval density and evaluated how those components may contribute to any nonlinear relationship observed for estimated *per capita* rate of increase.

Methods

Laboratory Populations and Rearing

All colonies originated from field collected larvae (origins in Table S1) and had been maintained in the laboratory for several generations. Colony larvae were raised in 30 x 15 cm pans and fed bovine liver powder weekly. Colonies of *A. triseriatus* and *A. aegypti* were raised in screened cages with a volume of 0.6 m³. These containers were held in an insectary at ~ 24°C with a 17:7 h L:D cycle and an 0.75 h dawn or dusk phase at the beginning and end of the light phase. *Aedes albopictus* were raised in 0.3 m³ plastic cages with at a constant temperature of 24°C, with a 14:10 h L:D cycle. Each cage contained an oviposition cup with egg paper and multiple cotton stoppered vials containing 20% sucrose solution. Colonies were blood fed weekly from mice or guinea pigs anesthetized with a 9:1 ketamine:xylazine mixture (Illinois State University IACUC protocol #842043). Egg papers were replaced weekly and stored at high humidity. See [44] for further details on rearing.

Overall experimental design

The experiment was run as a blocked design with one complete replicate for a species constituting a temporal block. One day before hatching larvae, 1 L plastic containers received 1.0 g of dried white oak (*Quercus alba*) leaves and 0.10 g of dried, crushed crickets (*Gryllodes sigillatus*) as detritus sources that support the microbial assemblages that are food for larvae, and 900 mL of Nanopure® water. Containers were then incubated in an environmental chamber at 24°C 14:10 h L:D cycle. These containers resemble typical aquatic systems colonized by these species, both in size and in detritus composition [54]. Each container was assigned a density treatment (10, 30, 50, 80, 110, 140, 170, 200, 250, or 300 first instar larvae) and a species treatment. Mosquito eggs from lab-reared colonies of *A. triseriatus*, *A. aegypti*, and *A. albopictus* were hatched in deionized water containing 0.25 g/L of nutrient broth. Twenty-four hours after hatching, larvae were counted into the 10 density treatments and added to the containers. Containers were then returned to the environmental chamber and held under the same conditions.

Containers were checked daily for pupae by pouring into a shallow 30 x 15 cm pan. Pupae were placed into 0.92 mL glass vials with cotton stoppers and allowed to emerge as adults. Pans were then rinsed back into the 1 L containers and any evaporated or lost water replaced with Nanopure® water. After adults eclosed, residual water was removed from the vial, sexes determined, and date of eclosion recorded before being placed in a drying oven at 50°C. Wing length of females was measured by placing removed wings on a microscope slide with coverslip to flatten for imaging. Images of each wing were

taken using Epiphan Capture Tool (Epiphan Systems Inc.) with a Wild M32 microscope and Sony DXC-970MD series microscope camera. Wing lengths were measured from the images using ImageJ software. Wing lengths were used to predict female fecundity using regressions (see below).

Index of Performance

The composite index of performance r' [34] was used to estimate *per capita* rate of change for each replicate cohort for each of the species.

$$r' = \frac{\ln [1/N_0) S_x A_x f(w_x)]}{D + [S_x \times A_x f(w_x) / S_x A_x f(w_x)]} \quad (\text{Eq. 1})$$

Where N_0 is the initial number of females in the cohort (assumed to be $\frac{1}{2}$ the initial number of larvae), x is number of days since hatching, A_x is the number of females from the cohort eclosing as adults on day x , w_x is mean wing length of females from the cohort eclosing on day x , $f(w_x)$ is predicted number of female eggs produced by females with mean wing length w_x based on regressions (Supplemental Table S2), and D is the estimated number of days from eclosion to oviposition (Supplemental Table S2). This index combines these measurements in a manner that is similar to life table estimates of R_0 (net reproductive rate) and T_c (cohort generation time), allowing for estimation of population growth, estimated as $r = \ln(R_0) / T_c$ [34], for a cohort of larvae when following reproduction of multiple replicated cohorts is infeasible. One problem with r' is that when no females survive to adulthood, either because of negative effects of high larval density (e.g., 300 larvae) or because of stochastic variation in number of females in small cohorts (e.g., 10 larvae), r' is not estimable. These cohorts cannot be ignored, particularly when larval density is high, as lack of survivors is then likely a result of high density. This problem can be solved by pooling replicate cohorts of the same density and species with and without surviving females to yield a combined cohort with larger N_0 , but having been reared at the nominal larval densities and *per capita* resources [24]. Thus, when no females survive in our data set this is the approach that we take. When data from two or more replicate cohorts at the same initial density were combined, we designate the combined data as coming from a new replicate block, identified with the numbers from the original blocks that were combined (e.g., combining cohorts from replicate blocks 1 and 3 yields a new replicate block 13).

Statistical analysis

Relationships between r' and initial density (N) were fit using the SAS statistical software (version 9.3) of the SAS system for windows (copyright © 2011 SAS Institute Inc). The fits included a θ logistic relationship, a Gompertz relationship, and linear, quadratic, and cubic relationships (Supplemental Table S3), all implemented using a generalised nonlinear mixed model (PROC NL MIXED), with a normal distribution of error and an identity link function, with replicate block as a random effect. The replicate block effect was modeled as affecting each of the parameters of the models (Supplemental Table S3).

AICc values were calculated and used to determine which relationship provided the best description of the data.

Relationships of survival to adulthood to initial density were analysed by fitting a generalised nonlinear mixed model (PROC NLMIXED), with a binomial distribution of error and logit link function, with replicate block as a random effect, of surviving adults (S) vs. initial density (N):

$$S = aN / [1 + (aN/K)^d] \text{ (Eq. 2)}$$

[55, 56]. The parameters a , K , and d determine the shape of the relationship of S to N . When the parameter $d > 1.0$, S peaks at a low N and declines at high N (specifically $N \geq K$). When $d = 1.0$, S approaches an asymptote equal to aK as N increases. When $d < 1.0$, S is a monotonic increasing function of N . When $d = 0$, S increases linearly with N . The parameter a is interpreted as the proportion surviving as N approaches 0 (i.e., as $N \rightarrow 0$, $S/N \rightarrow a$). When $N = K$, $S = aN/2$. Different models included and tested for random variation among replicate blocks in each of the parameters a , K , and d in Eq. 2. Because the number of replicate containers at each N was small, only models postulating a random effect of replicate block on one parameter (a , K , or d) at a time, or no parameters were tested. Attempts to fit models with random variation in multiple parameters at the same time resulted in either nonconvergence of the iterative solution, or inestimable parameters. Best models for each species were determined by AICc.

Adult female development time (= median days to eclosion from a replicate container) and female size (= mean wing length from a replicate container) were analysed by generalised linear model ANOVA (SAS PROC GLIMMIX) using initial density N as a class variable with block as a random effect. For female mean wing length and female median development time the models used normal and negative binomial distributions of error, respectively, and identity and log link functions, respectively. Significant ANOVA results were followed by Tukey pairwise multiple comparisons among least-squares means

Results

r' vs. initial density. Gompertz models with no random effect for block for either parameter (r_0 or b) proved to be the best descriptor of the relationship of r' to initial density N for each of the three *Aedes* species, based on AICc (Table 1). The best 3 functions in each case were always alternative versions of the Gompertz or Quadratic function (Table 1) and all linear functions had DAICc > 20 and model weights < 10^{-4} , usually much less (Table 1). The results clearly and strongly refute the hypothesis of a linear relationship and are consistent with the hypothesis of a nonlinear relationship that is concave upward (Fig. 1). Best fit parameters of the Gompertz equation for each of the three species (Fig. 1) were similar and did not differ significantly in combined analysis.

Table 1

AICc values comparing the top three models of r' vs. initial density N for *Aedes aegypti*, *Aedes albopictus*, and *Aedes triseriatus*. Models that failed to converge excluded, and those beyond the top three are included in the supplemental material (Table S2). For each species the best model is highlighted in **bold face type**. Functions given in Supplemental Table S4

Model	Total Parameters	AICc	Δ AIC	$e^{(-.5*\Delta)}$	Model Weight	Evidence ratio
<i>Aedes aegypti</i> $n = 29$						
Gompertz, no random var.	4	-127.5	0	1	0.707	1
Gompertz, random b	5	-125.0	2.5	0.287	0.202	3.49
Quadratic, no random var.	5	-123.4	4.1	0.129	0.091	7.77
<i>Aedes albopictus</i> $n = 27$						
Gompertz, no random var.	4	-127.5	0	1	0.989	1
Quadratic, no random var.	5	-117.6	9.9	0.007	0.007	141.17
Gompertz, random b	5	-116.3	11.2	0.004	0.004	270.43
<i>Aedes triseriatus</i> $n = 43$						
Gompertz, no random var.	4	-202.9	0	1	0.725	1
Gompertz, random r_0	5	-200.5	2.4	0.301	0.218	3.32
Quadratic, no random var.	5	-197.6	5.3	0.071	0.051	14.15

Surviving adults vs. initial density. The nonlinear relationship from Eq. 2 generally fit the data on surviving adults vs. initial density (Fig. 2), but the three species differed in how the random effect of replicate block affected the relationship, with each species' best model having a different parameter impacted by that random variation (Table 2). The best nonlinear description for each species yielded d significantly greater than 1.0 (Table 3), indicating a peak in adult production at low density and lesser adult production at greater densities (Fig. 2). Estimates of d for the three species were similar, falling between 2.30 and 2.37 (Table 3) indicating a fairly sharp decline in adult production as density increased (Fig. 2). The species differed more in a and K , with a lowest for *A. triseriatus* and greatest for *A. albopictus*, and K lowest for *A. aegypti* and similar for *A. albopictus* and *A. triseriatus* (Table 3).

Table 2

AICc values comparing models of surviving adults S vs. initial density N for *Aedes aegypti*, *Aedes albopictus*, and *Aedes triseriatus*. Models that failed to converge excluded. For each species the best model is highlighted in **bold face type**.

Model: Parameter affected by random replicate block	Total Parameters	AICc	ΔAIC	$e^{(-.5*\Delta)}$	Model Weight	Evidence ratio
<i>Aedes aegypti</i>						
d random	5	225.0	0	1	0.802	1
<i>K</i> random	5	227.8	2.8	0.247	0.198	4.06
<i>a</i> random	5	260.1	35.1	2.39×10^{-8}	1.91×10^{-8}	4.19×10^7
no random effects	4	292.7	67.7	1.99×10^{-15}	1.60×10^{-15}	5.02×10^{14}
<i>Aedes albopictus</i>						
K random	5	184.8	0	1	0.996	1
<i>a</i> random	5	195.7	10.9	0.004	0.004	232.76
<i>d</i> random	5	211.7	26.9	1.44×10^{-6}	1.44×10^{-6}	6.94×10^5
no random effects	4	213.1	28.3	7.16×10^{-7}	7.13×10^{-7}	1.40×10^4
<i>Aedes triseriatus</i>						
a random	5	386.6	0	1	0.997	1
<i>K</i> random	5	398.2	11.6	0.003	0.003	330.30
no random effects	4	408.8	22.2	1.51×10^{-5}	1.51×10^{-5}	66171.16
<i>d</i> random	5	411.1	24.5	4.79×10^{-6}	4.77×10^{-6}	2.09×10^5

Table 3

Best fit parameters for the relationships of surviving adults S vs. initial density N (Eq. 2) for *Aedes aegypti*, *Aedes albopictus*, and *Aedes triseriatus* that are shown in Fig. 2. See Table 2 for model selection via AICc. For each species the parameter that varied among replicate blocks is indicated with an *.

Species (n replicate blocks)	Parameter	Estimate $\pm$ SE	95% confidence interval
<i>Aedes aegypti</i> (4)	a	0.83 ± 0.07	0.59–1.06
	K	29.2 ± 3.0	19.5–38.8
	d^*	2.30 ± 0.29	1.37–3.23
<i>Aedes albopictus</i> (3)	a	0.92 ± 0.04	0.73–1.12
	K^*	43.0 ± 5.8	18.2–67.8
	d	2.37 ± 0.13	1.82–2.91
<i>Aedes triseriatus</i> (5)	a^*	0.69 ± 0.09	0.46–0.93
	K	43.1 ± 3.3	31.1–52.2
	d	2.32 ± 0.12	1.99–2.64

Development time and wing length vs. initial density. Female size was significantly affected by initial density for *A. aegypti* ($F_{9,13}=7.43$, $P = 0.0007$), *A. albopictus* ($F_{9,14}=21.98$, $P < 0.0001$), and *A. triseriatus* ($F_{9,25}=5.72$, $P = 0.0003$). All three species showed a pattern of greatest mean female wing length at the lowest density and significantly lower, and statistically indistinguishable mean wing lengths at densities ≥ 50 larvae (Fig. 3).

Female development time was significantly affected by initial density for *A. aegypti* ($F_{9,14}=7.40$, $P = 0.0005$), *A. albopictus* ($F_{9,14}=7.48$, $P = 0.0005$), and *A. triseriatus* ($F_{9,26}=9.55$, $P < 0.0001$). All three species showed a pattern of lowest and statistically indistinguishable mean female days to adulthood at the two lowest densities, steadily increasing mean days to adulthood at densities of 50 and 80 larvae, and significantly greater mean days to adulthood that were largely statistically indistinguishable at densities ≥ 110 larvae (Fig. 4).

Discussion

We found that the relationship between density and the estimated *per capita* rate of increase is clearly non-linear, as the most plausible models for our data were in all three cases the Gompertz function, with no random variation among replicate blocks. In the cases of *A. aegypti* and *A. triseriatus* the second-most plausible models, by AICc, were in both cases Gompertz functions with random variation in the parameter b , but these models only attained model weights of ~ 0.21 indicating they were rather unlikely to be the correct model for these data. All other models yielded very small model weights (< 0.10 ; Table 1). All linear

models tested had model weights $< 10^{-6}$. Thus, these results are inconsistent with the common assumption that rate of increase declines linearly with density. This represents one of just a few experimental tests of the assumption of linearity of rate of increase vs. density in animal model systems [11, 35, 39–41, 51].

Our estimates of *per capita* growth rate are derived from components of: survivorship to adulthood; female development time to adulthood (indicating age at reproduction); and adult female size (indicating fecundity), all combined in an approach similar to standard life table calculations [34]. All of these components were strongly nonlinearly related to initial population density. The relationships of *per capita* growth to density were all concave upward, indicating a rapid decline in growth rate across a range of low densities, and a slowing of the decline as densities become greater (Fig. 1). This pattern of density dependent growth thus seems to be associated with relatively little change in number of adults produced, female development time, and female size over a range of high densities, contrasting with the rapid change in all components as densities rise from the lowest levels in the experiment (Figs. 2, 3, 4).

Analysis of individual based models of a single species competing for resources gave insight into how life history and biological traits of organisms produce nonlinearity in *per capita* growth [4]. That research showed that both scramble and contest competition for resources could result in nonlinear relationships of *per capita* growth to density, and that whether that relationship was concave upward, or concave downward depended on maximum individual reproductive output, and the range of resource consumption over which reaching reproductive maturity is possible (i.e., plasticity in response to resource consumption). Greater maximum fecundity, and greater plasticity in response to resource consumption (specifically ability to reproduce at low resource intake) produced a relationship of *per capita* growth vs. population density that is concave upward, whereas lower maximum fecundity and lesser plasticity in response to resource consumption (specifically a high minimum resource intake requirement) produced a relationship of *per capita* growth vs. population density that is concave downward [4]. For the *Aedes* species we investigated, individual maximum fecundity can be moderately high (i.e., 50–100 eggs per reproductive cycle; See Fig. 3 for mean sizes, and Supplemental Table S2, for size fecundity relationships), but these *Aedes* are well known for their high plasticity in development time to and size at adulthood in response to food availability and competition [57–60] and ability to survive and to complete development with very low food availability [61]. These same traits likely contribute to the concave relationships we observed, and in particular the shallow decline in *per capita* growth over the high experimental densities.

The non-linearity of the relationship of adults produced vs. density indicates that overcompensatory morality is likely [5, 44, 62, 63]. Overcompensation occurs when extrinsic mortality imposed on a population decreases density and negative effects of density on survival, resulting in increased adult production with mortality. The estimated values of the parameter d in Eq. 2 are all significantly > 1.0 , which is indicative of overcompensation in response to mortality [56, 55]. This situation, if it occurs for these species in nature, could be important for mosquito control in the interests of public health [26]. Historically, pest management has targeted populations of pests with mortality with limited attention to

how that mortality interacts with density dependence to affect population dynamics [64]. The results from our experiments show that reductions of larval populations could lead to greater production of adults, in addition to increased *per capita* growth rates.

Here we have demonstrated that the relationship between density and estimated *per capita* growth for these mosquitoes is non-linear. As models postulating a linear relationship have often been used with these mosquitoes to quantify expected effects of intra- and inter-specific competition [24, 34, 43, 45–51] our results raise the question: what conclusions from using linear models may be misleading? If data are from experiments at relatively low densities, relative to resources (i.e., densities ≤ 80 larvae in our experiments; Fig. 1), it is likely that a linear relationship would be a fairly accurate approximation of the effects of intraspecific competition (i.e., estimating carrying capacity) or interspecific competition (i.e., estimating interspecific competition coefficients). The concave upward shape of the curves in Fig. 1 indicate that carrying capacity will be underestimated by such a linear approximation. In contrast, if data are from relatively high densities relative to resources (i.e., densities ≥ 140 larvae in our experiment; Fig. 1), the shallow slope in this range is likely to yield underestimates of the maximum *per capita* rate of increase for a population at low density. This suggests that populations of these *Aedes* species have a great potential for rapid increases from rarity caused by reductions via mosquito control or associated with being a newly introduced species. *Aedes* in general, and *A. albopictus* and *A. aegypti* in particular, have been among the most successful invasive species worldwide [65, 66], and this suggestion of very high potential population growth when rare seems likely to be a contributing factor to this invasion success. When experimental data are from a wide range of densities relative to resources, using linear models is most likely to yield inaccurate estimates of population growth parameters. Further, the concave upward shape of this relationship suggests that populations may have approximately 0 growth over a relatively wide range of high densities.

Declarations

Data Availability

The data generated from this experiment will be available at Figshare.

Acknowledgements

We thank Shawn Mohammed for aiding in identification and removal of pupae, Kristina McIntire, Zoey Neale, and Geoff Ower for aiding with maintenance of the laboratory colonies, and Molly Schumacher, Katie Westby, Kris McIntire, Kim Medley, and Phil Lounibos for assistance with collecting for our laboratory colonies, and Cynthia Lord for comments on the manuscript. This research was funded by US NIAID award number 1R15AI124005-1 to SAJ.

Author Contributions

SAJ designed the experiment, LAS and KEH ran the experiment and collected the data. SAJ analyzed the data. LAS wrote the first draft of the manuscript and all authors aided in revising the final manuscript.

Competing Interests

The authors declare no competing interests.

Declaration on the Ethical Use of Vertebrate Animals

Use of live vertebrate animals, female mice [ICR (CD1) strain]] and male guinea pigs [Hartley strain], for blood feeding mosquito populations is approved by Illinois State University's Institutional Animal Care and Use Committee (IACUC protocol #842043). Individuals working with live vertebrate animals were trained on the procedures outlined in this protocol and underwent additional webinar training. All experiments were performed in accordance with the guidelines and regulations for IACUC protocol #842043, and the relevant methods relating to the use of these live animals are in accordance with ARRIVE guidelines.

References

1. Hutchinson G.E. *An Introduction to Population Ecology*. New Haven, CN, USA: Yale University Press; (1978).
2. Fussman, G.F., and Heber, G. Food web complexity and chaotic population dynamics. *Ecol. Lett.* **5**, 394-401 (1978).
3. Maron, J.L., and Crone, E. Herbivory: effects on plant abundance, distribution, and population growth. *Proc. Roy. Soc. B.* **272**, 2575-2584 (1978).
4. Johst, K., Berryman, A., and Lima, M. From individual interactions to population dynamics: individual resource partitioning simulation exposes the causes of nonlinear intra-specific competition. *Pop. Ecol.* **50**, 79–90 (2008).
5. McIntire, K.M., and Juliano, S.A. How can mortality increase population size? A test of two hypotheses. *Ecology*. **99**, 1660-1670 (2018).
6. Mylius, S.D., and Deikmann, O. On evolutionary stable life histories, optimization and the need to be specific about density dependence. *Oikos*. **74**, 218-224 (1995).
7. Courchamp, F., Clutton-Brock, T., and Grenfell, B. Inverse density dependence and the Allee effect. *Trends Ecol. Evol.* **14**, 405-410 (1999).
8. MacLean, R.C., and Gudelj, I. Resource competition and social conflict in experimental populations of yeast. *Nature*. **44**, 498-501 (2006).
9. Khatchikian, C.E., *et al.* Recent and rapid population growth and range expansion of the Lyme disease tick vector, *Ixodes scapularis*, in North America. *Evolution*. **69**, 1678-1689, (2015).
10. Lafferty, K.D., and Holt, R.D. How should environmental stress affect the population dynamics of disease? *Ecol. Lett.* **6**, 654-664 (2003).

11. Sibley, R.M., Barker, D., Denham, M.C., Hone, J., and Pagel, M. On the regulation of populations of mammals, birds, fish, and insects. *Science*. **309**, 607–610, (2005).
12. Bjorndal, K., Bolten, A.B., and Chaloupka, M.Y. Green turtle somatic growth model: evidence for density-dependence. *Ecol. App.* **10**, 269-282 (2000).
13. Lamb, J.S., Satgé, Y.G., and Jodice, P.G.R. Influence of density-dependent competition on foraging and migratory behavior of a subtropical colonial seabird. *Ecol. Evol.* **7**, 6469-6481 (2017).
14. Kobayashi, K. Sexual selection sustains biodiversity via producing negative density-dependent population growth. *Jour. Ecol.* **107**, 1433-1438 (2018).
15. López-Sepulcre, A., and Kokko, H. Territorial Defense, Territory Size, and Population Regulation. *The Amer. Nat.* **166**, 317-325 (2005).
16. Maag, N., Cozzi, G., Clutton-Brock, T., and Ozgul, A. Density-dependent dispersal strategies in a cooperative breeder. *Ecology*. **99**, 1932-1941 (2018).
17. Bonenfant, C., *et al.* Empirical evidence of density- dependence in populations of large herbivores. *Adv. Ecol. Res.* **41**, 313-357 (2009).
18. Legros, M., Lloyd, A.L., Huang, Y., and Gould, F. Density-dependent intraspecific competition in the larval stage of *Aedes aegypti* (Diptera: Culicidae): revisiting the current paradigm. *J. Med. Entomol.* **46**, 409-419 (2009).
19. Hixon, M.A., and Jones, G.P. Competition, predation, and density-dependent mortality in demersal marine fishes. *Ecology*. **86**, 2847-2859 (2006).
20. Vonesh, J.R., and De La Cruz, O. Complex life cycles and density dependence: assessing the contribution of egg mortality to amphibian declines. *Oecologia*. **133**, 325-333 (2002).
21. Southwood, T.R., Murdie, G., Yasuno, M., Tonn, R.J., and Reader, P.M. Studies on the life budget of *Ae. aegypti* in Wat Samphaya, Bangkok, Thailand. *Bull. World Health Organ.* **46**, 211–226 (1972).
22. Dye, C. Intraspecific competition amongst larval *Aedes aegypti*: food exploitation or chemical interference. *Ecol. Entomol.* **7**, 39– 46 (1982).
23. Dye, C. Models for the population dynamics of the yellow fever mosquito, *Aedes aegypti*. *J. Anim. Ecol.* **53**, 247–268 (1984).
24. Livdahl, T.P., and Willey, M.S. Prospects for an invasion: competition between *Aedes albopictus* and native *Aedes triseriatus*. *Science*. **253**, 189–191 (1991).
25. Alto, B.W., Lounibos, L.P., Higgs, S., and Juliano, S.A. Larval competition differentially affects arbovirus infection in *Aedes* mosquito. *Ecology*. **86**, 3279-3288 (2005).
26. Juliano, S.A. Population dynamics. *J. Am. Mosq. Control Assoc.* **23**, 265-275 (2007).
27. Gilpin, M.E., and McClelland, G.A.H. Systems analysis of the yellow fever mosquito *Aedes aegypti*. *Fortschritte der Zoologie*. **25**, 355– 388 (1979).
28. Juliano, S.A. Species introduction and replacement among mosquitoes: Interspecific resource competition or apparent competition? *Ecology*. **79**, 255-268 (1998).

29. Lord, C.C. Density Dependence in Larval *Aedes albopictus* (Diptera: Culicidae). *J. Med. Entomol.* **35**, 825-829 (1998).
30. Agnew, P., Hide, M., Sidobre, C., and Michalakis, Y. A minimalist approach to the effects of density-dependent competition on insect life-history traits. *Ecol. Entomol.* **27**, 396-402 (2002).
31. Walsh, R.K., Facchinelli, L., Ramsey, J.M., Bond, J.G., and Gould, F. Assessing the impact of density dependence in field populations of *Aedes aegypti*. *J. Vect. Ecol.* **36**, 300-307 (2011).
32. Walsh, R.K., Bradley, C., Apperson, C.S., and Gould, F. An experimental field study of delayed density dependence in natural populations of *Aedes albopictus*. *PLoS ONE*. **7**, e35959 (2012).
33. Walsh, R.K., et al. Regulation of *Aedes aegypti* Population Dynamics in Field Systems: Quantifying Direct and Delayed Density Dependence. *Am. J. Trop. Med. Hyg.* **89**, 68 – 77 (2013).
34. Livdahl, T. P., and Sugihara, G. Non-linear interactions of populations and the importance of estimating per capita rates of change. *J. Anim. Ecol.* **53**, 573-580 (1984).
35. Getz, W.M. A hypothesis regarding the abruptness of density dependence and the growth rate of populations. *Ecology*. **77**, 2014-2026 (1996).
36. Tenan, S., Tavecchia, G., Oro, D., and Pradel, R. Assessing the effect of density on population growth when modeling individual encounter data. *Ecology*. **100**, e02595 (2019).
37. Arditi, R., Bersier, L., Rohr, R.P. The perfect mixing paradox and the logistic equation: Verhulst vs. Lotka. *Ecosphere*. **7**, e01599 (2016).
38. Cortés, E. Perspectives on the intrinsic rate of population growth. *Meth. Ecol. Evol.* **7**, 1136-1145 (2016).
39. Smith, F.E. Population dynamics in *Daphnia magna* and a new model for population growth. *Ecology*. **4**, 651-663 (1963).
40. Ayala, F.J., Gilpin, M.E., and Ehrenfeld, J.G. Competition between species: Theoretical models and experimental tests. *Theor. Pop. Biol.* **4**, 331-356 (1973).
41. Borlestean, A., Frost, P.C., and Murray, D.L. A mechanistic analysis of density dependence in algal population dynamics. *Front. Ecol. Evol.* **3**, 37 (2015).
42. Clark, F., Brook, B.W., Delean, S., Akçakaya, H.R., and Bradshaw, C.J.A. The theta-logistic is unreliable for modelling most census data. *Meth. Ecol. Evol.* **1**, 253–262 (2010).
43. Chmielewski, M.W., Khatchikian, C., and Livdahl, T. Estimating the per capita rate of population change: How well do life-history surrogates perform? *Ann. Entomol. Soc. Am.* **103**, 734–741 (2010).
44. Neale, J.T., and Juliano, S.A. Finding the sweet spot: What levels of larval mortality lead to compensation or overcompensation in adult production? *Ecosphere*. **10**, e02855 (2019).
45. Armistead, J.S., Arias, J.R., Nishimura, N., and Lounibos, L.P. Interspecific larval competition between *Aedes albopictus* and *Aedes japonicus* (Diptera: Culicidae) in northern Virginia. *J. Med. Entomol.* **45**, 629–37 (2008).
46. Kaplan, L., Kendell, D., Robertson, D., Livdahl, T., and Khatchikian, C. *Aedes aegypti* and *Aedes albopictus* in Bermuda: Extinction, invasion, invasion and extinction. *Bio. Invasions*. **12**, 3277–3288

(2010).

47. Juliano, S.A. Coexistence, exclusion, or neutrality? A meta-analysis of competition between *Aedes albopictus* and resident mosquitoes. *Isr. J. Ecol. Evol.* **56**, 325–351 (2010).
48. Murrell, E.G., and Juliano, S.A. Competitive abilities in experimental microcosms are accurately predicted by a demographic index for R^* . *PLoS One*. **7**, e43458 (2012).
49. Leisnham, P.T., and Juliano, S.A. Interpopulation differences in competitive effect and response of the mosquito *Aedes aegypti* and resistance to invasion of a superior competitor. *Oecologia*. **164**, 221–230 (2010).
50. Leisnham, P.T., Lounibos, L.P., O'Meara, G.F., and Juliano, S.A. Interpopulation divergence in competitive interactions of the mosquito *Aedes albopictus*. *Ecology*. **90**, 2405–2413 (2009).
51. Evans, M.V., Drake, J.M., Jones, L., and Murdock, C.C. Assessing temperature-dependent competition between two invasive mosquito species. *Ecol. Appl.* **31**, e02334 (2021).
52. Léonard, P.M., and Juliano, S.A. Effects of leaf litter and density on fitness and population performance of the hole mosquito *Aedes triseriatus*. *Ecol. Entomol.* **20**, 125-136 (1995).
53. Chandrasegaran, K., and Juliano, S.A. How Do Trait-Mediated Non-Lethal Effects of Predation Affect Population-Level Performance of Mosquitoes? *Front. Ecol. Evol.* **7**, 25 (2019).
54. Yee, D.A., Kaufman, M.G., and Juliano, S.A. The significance of ratios of detritus types and microorganism productivity to competitive interactions between aquatic insect detritivores. *J. Anim. Ecol.* **76**, 1105-1115 (2007).
55. Osenberg, C.W., *et al.* Rethinking ecological inference: density dependence in reef fishes. *Ecol. Lett.* **5**, 715–721 (2002).
56. Schmitt, R.J., Holbrook, S.J., Osenberg, C.W. Quantifying the effects of multiple processes on local abundance: a cohort approach for open populations. *Ecol. Lett.* **2**, 294-303 (1999).
57. Fish, D. An analysis of adult size variation within natural mosquito population. In: *Ecology of Mosquitoes: Proceedings of a Workshop*. (ed. Lounibos, L.P., Rey, J.R., Frank, J.H.) 419-429 (Vero Beach, FL: Medical Entomology Laboratory, 1985)
58. Schneider, J.R., Chadee, D.D., Mori, A., Romero-Severson, J., and Severson, D.W. Heritability and adaptive phenotypic plasticity of adult body size in the mosquito *Aedes aegypti* with implications for dengue vector competence. *Infect. Genet. Evol.* **11**, 11–16 (2011).
59. Wormington, J.D., and Juliano, S.A. Sexually dimorphic body size and development time plasticity in *Aedes* mosquitoes (Diptera: Culicidae). *Evol. Ecol. Res.* **16**, 1–12 (2014).
60. Steinwascher, K. Competition and growth among *Aedes aegypti* larvae: Effects of distributing food inputs over time. *PLoS One*. **15**, e0234676 (2020).
61. Barrera, R. Competition and resistance to starvation in larvae of container-inhabiting *Aedes* mosquitoes. *Ecol. Entomol.* **21**, 117–127 (1996).
62. Servanty, S., *et al.* Assessing whether mortality is additive using marked animals: a Bayesian state-space modeling approach. *Ecology*. **91**, 1916-1923 (2010).

63. Wolfe, M.L., *et al.* Is anthropogenic cougar mortality compensated by changes in natural mortality in Utah? Insights from long-term studies. *Biol. Conserv.* **182**, 187-196 (2015).
64. Kogan, M. Integrated pest management: historical perspectives and contemporary developments. *Ann. Rev. Entomol.* **43**, 243-270 (1998).
65. Lounibos, L.P. Invasions by insect vectors of human diseases. *Ann. Rev. Entomol.* **47**, 233–66 (2002).
66. Juliano, S.A., and Lounibos, L.P. Ecology of invasive mosquitoes: effects on resident species and on human health. *Ecol. Lett.* **8**, 558–574 (2005).

Figures

Figure 1

Relationship between estimated per capita rate of change r' and initial density N for *Aedes aegypti*, *Aedes albopictus*, and *Aedes triseriatus*, as described by the best models (Table 1), which were Gompertz models without random variation due to blocks. The solid black line is the predictions from the model averages, and the grey area represents 95% confidence interval on the prediction. Best fit parameters (+SE) for *A. aegypti*, *A. albopictus*, and *A. triseriatus* were: $r_0 = 0.167+0.017$, $0.223+0.019$, and $0.0173+0.014$, respectively, and $b = -0.0363+0.0038$, $-0.0472+0.0041$, and $-0.0358+0.0032$, respectively. The shape and color of data points corresponds to the different replicate blocks for each species; cases of combined cohorts from different blocks are indicated with multiple numbers (e.g., replicate blocks 13, 123, etc.)

Figure 2

Survivorship to adulthood S versus initial density N for *Aedes aegypti*, *Aedes albopictus*, and *Aedes triseriatus*. The solid black line is the prediction for the regression of number of surviving adults vs. density (eq. 2), and the dotted lines indicate 95% confidence interval on the prediction. Best fit parameters for *A. aegypti*, *A. albopictus*, and *A. triseriatus* are given in Table 3. The color of data points corresponds to the different replicate blocks for each species.

Figure 3

Least squares mean+SE of cohort mean female wing length for *Aedes aegypti*, *Aedes albopictus*, and *Aedes aegypti*, averaged over replicate runs for each species. Means for a species associated with the same letters are not significantly different by Tukey's multiple comparisons.

Figure 4

Least squares mean+SE of cohort median female time to adulthood for *Aedes aegypti*, *Aedes albopictus*, and *Aedes aegypti*, averaged over replicate runs for each species. Least squares means for a species associated with the same letters are not significantly different by Tukey’s multiple comparisons.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SauersHawesJulianoSupplements.docx](#)