Evaluating the relevance of sequence conservation in the prediction of pathogenic missense variants

Emidio Capriotti (✉ emidio.capriotti@unibo.it)

University of Bologna: Università di Bologna <https://orcid.org/0000-0002-2323-0963>

Piero Fariselli

University of Turin: Università degli Studi di Torino

Research Article

Keywords: Variant interpretation, Pathogenic missense variants, evolutionary information, conservation score

Posted Date: September 9th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-884099/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Human Genetics on January 31st, 2022. See the published version at <https://doi.org/10.1007/s00439-021-02419-4>.

Abstract

Evolutionary information is the primary tool for detecting functional conservation in nucleic acid and protein. This information has been extensively used to predict structure, interactions and functions in macromolecules. Pathogenicity prediction models rely on multiple sequence alignment information at different levels. However, most accurate genome-wide variant deleteriousness ranking algorithms consider different features to assess the impact of variants. Here, we analyze three different ways of extracting evolutionary information from sequence alignments in the context of pathogenicity predictions at DNA and protein levels. We showed that protein sequence-based information is slightly more informative in the annotation of Clinvar missense variants than those obtained at the DNA level. Furthermore, to achieve the performance of state-of-the-art methods, such as CADD, the conservation of reference and variant, encoded as frequencies of reference/alternate alleles or wild-type/mutant residues, should be included. Our results on a large set of missense variants show that a basic method based on three input features derived from the protein sequence profile performs similarly to the CADD algorithm which uses hundreds of genomic features. This observation indicates that for missense variants, evolutionary information, when properly encoded, plays the primary role in ranking pathogenicity.

Introduction

High-throughput sequencing technologies have changed our daily research by rapidly accumulating genomic data and helping to profile patient genomes (MacArthur *et al.*, 2014; Claussnitzer *et al.*, 2020). These studies make variant interpretation a fundamental challenge in precision medicine (Fernald *et al.*, 2011; Capriotti *et al.*, 2012; McInnes *et al.*, 2021). Missense variants by changing a single amino acid in a protein sequence can be neutral or induce loss of function.

In the last two decades several methods have been developed to prioritize functional missense variants relying on protein sequence/structure information (Ancien *et al.*, 2018; Tennessen *et al.*, 2012; Niroula and Vihinen, 2016; Petrosino *et al.*, 2021) and the protein interaction networks (Rost *et al.*, 2016; Capriotti *et al.*, 2019; Ozturk and Carter, 2021).

It is widely accepted that evolutionary information encoded in multiple sequence alignments of DNAs and proteins is a major resource for scoring variant pathogenicity. This paper evaluates the relevance of this information for missense variant predictions by comparing simple scores and simple predictors with the widely used and well-performing Combined Annotation-Dependent Depletion (CADD) algorithm (Rentzsch *et al.*, 2019).

We computed the conservation scores on DNA (*PhastCons100way* and *PhyloP100way*), the frequencies of the reference and alternative alleles in the genome, and frequencies of the wild-type and mutant residues in protein multiple alignments. Our analysis showed that a machine learning method trained on a few sequence conservation features at DNA or protein levels, achieves similar performance of a state-of-the-art algorithm. In this work we compared the performance of CADD with those reached by three different basic gradient boosting algorithms on a set of missense variants from the *Clinvar* database. Our result indicates

that the evolutionary information provides the main features for scoring the pathogenicity of missense variants.

Material And Methods

Datasets

To evaluate the performance of different machine learning methods for predicting the pathogenicity of missense variants we collected two datasets from the *Clinvar* database (Landrum *et al.*, 2020). For building the two datasets we considered two versions of *Clinvar* released in June 2020 and August 2021, respectively. The first dataset (*CommonClinvar*) consists of the missense variants annotated as *Pathogenic* and *Benign* in both versions of the database while the second dataset (*NewClinvar*) collects the new missense variants reported in the last version of *Clinvar* since June 2020 (Fig. S1). The variants reported in the older version of *Clinvar* not confirmed in the last version were discarded. Thus, the *CommonClinvar* consists of 36,751 missense variants from 7,582 proteins 53.5% of which are annotated as *Benign* and the remaining ones (46.5%) as *Pathogenic*. *NewClinvar*, which includes only the newly annotated variants, is composed of 5,172 from 1,855 proteins 43.4% of which are reported as *Benign* and 56.6% as *Pathogenic*. The composition of the two datasets is summarized in Table S1. Both *CommonClinvar* and *NewClinvar* datasets are available as supplementary files.

Conservation features

In this work we analyzed the performance in the prediction of pathogenic variants using three basic methods based on sequence conservation features. Each method considers only three input features, which are described below.

The first two methods are based on features obtained from a genome level multiple sequence alignment made available through the UCSC genome browser (Kent *et al.*, 2002), which evaluates conservation along the DNA multiple alignments. The conservation scores considered for the first method are calculated by *PhastCons* (Siepel *et al.*, 2005) and *PhyloP* (Pollard *et al.*, 2010) algorithms, while the features of the second method are extracted from the *multiz100way* multiple alignments. The *PhastCons100* and *PhyloP100way* scores as well as the *multiz100way* alignments for the hg38 human reference genome are available at <https://hgdownload.cse.ucsc.edu/goldenpath/hg38/>.

The three features of the last method are generated from a protein sequence profile calculated on the results of a *BLAST* (Altschul *et al.*, 1997) search on the *UniRef90* database (Suzek *et al.*, 2007) released in June 2020. For the *BLAST* search we used an e-value cutoff of 10^{-9} as suggested in previous works (Capriotti *et al.*, 2006; Calabrese *et al.*, 2009; Capriotti *et al.*, 2013).

In detail, we consider the *PhastCons100way* and *PhyloP100way* scores for each mutated loci, the frequencies of the reference and alternative alleles from the *multiz100way* multiple sequence alignment, and the total number of aligned genomic sequences of the mutated loci. To map the missense variant at

the protein level, we calculated the wild-type and mutant residue frequencies from the output of a *BLAST* search and the number of aligned proteins in the mutated site.

Machine learning algorithms

Using the eight features described above, we develop three binary classifiers (*PPScores*, *DNAProf*, *ProtProf*) using the following groups of three features:

- *PPScore*: *PhastCons100way* (*PC*) and *PhyloP100way* (*PP*) scores, and number aligned genomic sequences (N_g) in *multiz100way*
- *DNAProf*: Frequencies of the reference (f_{ref}) and alternative (f_{alt}) alleles, and number aligned genomic sequences in *multiz100way* (N_g).
- *ProtProf*: Frequencies of the wild-type (f_{wt}) and mutant (f_{mut}) residues, and number aligned protein sequences (N_p) from a *BLAST* search on *UniRef90*.

For each group of features defined above we developed a binary classifier based on the gradient boosting algorithm as implemented in the *scikit-learn* package (Pedregosa, F. *et al.*, 2011). The proposed groups of features are summarized in Table 1.

Table 1
Three groups of features used

Group	Features		
PPScores	PC	PP	N_g
DNAProf	f_{ref}	f_{alt}	N_g
ProtProf	f_{wt}	f_{mut}	N_p
for the development of the binary classifiers.			

Training and testing procedure

We first evaluated the performance of each method on *CommonClinvar* using a 10-fold cross-validation procedure for a fair evaluation of the proposed method performance. To reduce at the minimum the possible overfitting we mapped each missense variant on the relative protein sequence and we cluestter all the sequences using the *blastclust* algorithm (<https://ftp.ncbi.nih.gov/blast/documents/blastclust.html>) with a sequence identity threshold of 25% and a coverage of 50%. Using the clustering based on sequence similarity we perform a 10-fold cross-validation procedure keeping all the variants belonging to the same cluster in the same subset. A second test is performed considering the *NewClinvar* dataset. In this case the impact of the variants of a given protein are predicted excluding from the training set (*CommonClinvar*) all the variants belonging to proteins of the same cluster. We extracted a balanced set of *Pathogenic* and *Benign* variants from *CommonClinvar* and *NewClinvar* dataset for each test, randomly downscaling the

most abundant class. The reported scoring measures for all the methods are averaged over ten randomly selected sets.

Benchmarking and performance measures

To characterize the prediction power of the main conservation features described in this work, for each of them we developed a single feature binary classifier based on a single threshold. For each feature the classification threshold is optimized on the *CommonClinvar* dataset maximizing both the true positive and true negative rates. The optimized threshold is tested in the classification of the *NewClinvar* variant dataset. The same procedure is used to optimize the raw score output threshold of the CADD algorithm (Rentzsch *et al.*, 2019) as binary classifier.

Finally, the performances of all the binary classifiers described above are compared with those achieved by the CADD algorithm. All the measures considered for scoring the performance of the methods are defined in Supplementary Materials.

Results

Feature analysis and single feature classification

In the first part of our work, we analyzed the distributions of the main features used for the classification task. We focused on the six conservation features ($PC, PP, f_{ref}, f_{alt}, f_{wt}, f_{mut}$) comparing their distributions for the subsets of *Pathogenic* and *Benign* variants. The average, median and standard deviation of such distributions are reported in Table S2. As observed in previous works (Kircher *et al.*, 2014; Capriotti and Fariselli, 2017) the distribution of the *PhyloP100way* score (PP) in mutated loci associated with *Pathogenic* and *Benign* variants are significantly different (Fig. 1). Indeed, the two distributions show median values of 7.5 and 1.5, respectively, with a Kolmogorov-Smirnov distance (KSD) of 0.57 (Fig. 1B and Table S2). This distance is greater than the KSD observed for the *PhastCons100way* score (PC).

A higher difference between the distributions of the conservation scores for the subset of *Pathogenic* and *Benign* variants is observed when the frequencies in sequence profile from genomic and proteins are considered. The most remarkable differences are generally detected when comparing the distributions of the frequency of the alternative allele (f_{alt}) and the mutant residue (f_{mut}) for which the KSD is ~ 0.60 . Analyzing the frequencies of the reference allele (f_{ref}) and wild-type residue (f_{wt}) their KSD is 0.58 and 0.55, respectively (Table S2). The distributions of the four types of frequencies (f_{ref}, f_{alt}, f_{wt} and f_{mut}) for the subsets of *Pathogenic* and *Benign* variants are plotted in Fig. 2

This observation agrees with the results obtained in the prediction of *Pathogenic* variants using a classification threshold on a single feature. The classification threshold is optimized on the *CommonClinvar* dataset maximizing both the true positive and negative rates (Table S3). Applying the optimized thresholds on the prediction of the variants in the *NewClinvar* dataset, we found that a simple classifier based on the frequency of the mutant residue extracted from a protein sequence profile achieve

81% overall accuracy (Q2), 0.63 Matthews correlation coefficient (MC) and an Area Under the Receiver Operating Characteristic Curve (AUC) of 0.86 (Table 2).

Table 2

Performance of basic predictors based on a single feature on the *NewClinvar* dataset Prediction threshold are optimized on the *CommonClinvar* dataset. Q2: Overall Accuracy, TNR: True negative rate, NPV: Negative predicted value, TPR: True Positive Rate, PPV: Positive Predicted Value, MC: Matthews Correlation Coefficient, F1: harmonic mean of precision and sensitivity, AUC: Area Under the Receiver Operator Characteristic Curve, AUP Area under the Precision Recall Curve. All the performance measures are defined in Supplementary Materials.

Feature	Threshold	Q2	TNR	NPV	TPR	PPV	MC	F1	AUC	AUP
<i>PC</i>	1.000	0.737	0.611	0.816	0.862	0.689	0.489	0.766	0.755	0.815
<i>PP</i>	4.704	0.769	0.796	0.756	0.743	0.784	0.539	0.763	0.841	0.828
f_{ref}	0.977	0.779	0.815	0.760	0.742	0.801	0.559	0.770	0.836	0.843
f_{alt}	0.000	0.794	0.750	0.821	0.837	0.770	0.589	0.802	0.828	0.863
f_{wt}	0.702	0.769	0.806	0.750	0.731	0.791	0.539	0.759	0.844	0.836
f_{mut}	0.005	0.815	0.819	0.812	0.810	0.817	0.629	0.814	0.857	0.856

According to the previous observation, the *PhastCons100way* score (*PC*) is the least discriminating feature. When using the optimized threshold on the classification of the *NewClinvar* variants, the method based on a single *PC* threshold achieves 74% overall accuracy, 0.49 MC and 0.75 AUC (Table 2). Slightly lower performances are obtained when the frequencies of the reference allele and the wild-type residue in the sequence profile are considered. In this case the method based on a single f_{ref} threshold results in 78% Q2, 0.56 MC and 0.84 AUC. These results can also be observed plotting the Receiving Operating Characteristic (ROC) and Precision-Recall (PR) curves reported in Fig. S2.

Assessment of the machine learning methods

Starting from the previous observations, we developed three machine learning approaches based on the different groups of conservation features. The *PPScore* method is based on the *PhastCons100way*, *PhyloP100way* scores representing unique conservation measures not describing the type of nucleotides observed in the mutated loci. The other two methods consider the frequencies of the nucleotides or residues in the original and new sequences that correspond to f_{ref} , f_{alt} and f_{wt} , f_{mut} for *DNAProf* and *ProtProf*, respectively. To these groups of measures we added a third feature representing the total number of sequences aligned in the mutated loci (N_g, N_p). We implemented three machine learning methods for predicting *Pathogenic* variants based on the gradient boosting algorithm with these groups of features. First, the performance of these methods is tested with a 10-fold cross-validation procedure on the *CommonClinvar* dataset. To avoid possible overfitting we clustered all the proteins based on the sequence identity and grouped all their variants in a unique subset. The average performance of *PPScore*, *DNAProf* and *ProtProf* on a balanced set of *Pathogenic* and *Benign* variants are reported in Table S4. The results

show that among the three methods *ProtProf*, which is based on protein sequence profile, achieved the highest performance reaching 83% overall accuracy (Q2), 0.67 Matthews correlation coefficient and 0.91 Area Under the Receiver Operating Characteristic Curve (AUC). *PPScore* which is based on *PhastCons100way*, *PhyloP100way* show the lowest performance resulting in ~ 4% lower AUC and ~ 9% lower MC. An intermediate level of performance is achieved by *DNAProf* which results in ~ 2% lower AUC and ~ 3% lower MC with respect to *ProtProf*. Similar results are obtained when assessing the performance of the three methods on the *NewClinvar* dataset. Also in this case we predicted the impact of each variant removing from the training set all the variants in the *CommonClinvar* training set belonging to the same cluster of proteins. The performance of *PPScore*, *DNAProf* and *ProtProf* on a balanced set of variants from the *NewClinvar* dataset are summarized in Table 3.

Table 3

Prediction in cross-validation of the *NewClinvar* variant dataset. Q2: Overall Accuracy, TNR: True negative rate, NPV: Negative predicted value, TPR: True Positive Rate, PPV: Positive Predicted Value, MC: Matthews Correlation Coefficient, F1: harmonic mean of precision and sensitivity, AUC: Area Under the Receiver Operator Characteristic Curve, AUP Area Under the Precision Recall Curve. All the performance measures are defined in Supplementary Materials. For CADD a raw score classification threshold of 3.1 was considered.

Method	Q2	TNR	NPV	TPR	PPV	MC	F1	AUC	AUP
CADD	0.844	0.821	0.860	0.867	0.829	0.688	0.847	0.911	0.905
<i>ProtProf</i>	0.831	0.865	0.809	0.796	0.855	0.662	0.824	0.910	0.905
<i>DNAProf</i>	0.812	0.780	0.834	0.845	0.794	0.626	0.818	0.881	0.873
<i>PPScore</i>	0.771	0.776	0.769	0.767	0.774	0.543	0.770	0.855	0.846

Comparison with CADD algorithm

In the final part of our analysis we compared the performance of our simple gradient boosting-based algorithms with those obtained with CADD (Rentzsch *et al.*, 2019). CADD is one of the most accurate methods for predicting *Pathogenic* variants in coding and non-coding regions (Benevenuta *et al.*, 2021). This method, which is based on more than hundreds of genomic features, was trained on more than 30 million variants. To use CADD as a binary classifier we considered the raw output of the program and we selected the threshold that maximizes the true positive and negative rates on the *CommonClinvar* dataset. The performance of CADD at the optimal raw score classification threshold of 3.1 is reported in Table S4. This threshold was used for the classification of the variants in the *NewClinvar* dataset. The performance of CADD on the *NewClinvar* dataset is summarized in Table 3. This analysis shows that CADD and *ProtProf* algorithms result in a similar performance in the classification of *Pathogenic* missense variants in terms of Area Under the Receiver Operating Characteristic (AUC) and Precision-Recall (AUP) curves on both *CommonClinvar* and *NewClinvar* datasets. We can also observe that *DNAProf* which is based on the sequence profile extracted from the *multiz100way* sequence alignments results only in ~ 3% lower AUC and AUP. The Receiver Operating Characteristic and Precision-Recall curves for CADD and the three methods presented in this manuscript are plotted in Fig. 3.

Conclusion And Discussion

Here we analyzed different evolutionary information encodings for missense variant pathogenicity predictions. We compared the encoding at DNA and protein levels, where different multiple alignments techniques apply. Multiple alignments of protein sequence include more sequence and more remote homologs for many genes than pre-calculated genome alignments from the UCSC genome browser. This condition can be the reason why the performance of a method trained using the protein-based information is slightly better. With these simple inputs based on evolutionary information, a machine learning method can perform comparably to CADD, which uses more sophisticated inputs. This result shows that, at least for the missense variants, an input based on evolutionary information of the wild-type and mutated residue plays the most relevant role in scoring pathogenic variants.

Recently, it has been suggested that protein positions have a significant role and can act as *Neutral*, *Toggle* or *Rheostat* (Miller *et al.*, 2019). Here we indicate an alternative view of protein positions that can be seen as a non-linear combination of the frequencies of wild-type/mutant residues at protein level or reference/alternative allele at DNA level. The results of our analysis suggest that the performances of new and more sophisticated machine learning algorithms should always be compared with those achieved by simple conservation-based methods. As recently proposed (Walsh *et al.*, 2021), the design of such benchmark tests should consider the adoption of specific guidelines for avoiding bias in the training and testing sets. This procedure is important to exclude overfitting on the context-dependent features (Grimm *et al.*, 2015) and identify new important features for improving the performance of variant scoring algorithms.

Declarations

Funding

This work was supported by the PRIN project, “Integrative tools for defining the molecular basis of the diseases: Computational and Experimental methods for Protein Variant Interpretation” of the Ministero Istruzione, Università e Ricerca (PRIN201744NR8S).

Conflicts of interest/Competing interests

None

Availability of data and material

Additional information is available in Supplementary Materials. The *CommonClinvar* and *NewClinvar* datasets are provided as supplementary files.

Authors' contributions

EC developed and collected and analyzed the data. Both the authors designed the research and contributed to the writing of the manuscript.

Acknowledgments

PF thanks the Italian Ministry for Education, University and Research for the programme “Dipartimenti di Eccellenza 20182022D15D18000410001”.

References

1. Altschul,S.F. *et al.* (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*, **25**, 3389–402.
2. Ancien,F. *et al.* (2018) Prediction and interpretation of deleterious coding variants in terms of protein structural stability. *Sci. Rep.*, **8**, 4480.
3. Benevenuta,S. *et al.* (2021) Calibrating variant-scoring methods for clinical decision making. *Bioinformatics*, (In press).
4. Calabrese,R. *et al.* (2009) Functional annotations improve the predictive score of human disease-related mutations in proteins. *Hum Mutat*, **30**, 1237–44.
5. Capriotti,E. *et al.* (2012) Bioinformatics for personal genome interpretation. *Brief Bioinform*, **13**, 495–512.
6. Capriotti,E. *et al.* (2019) Integrating molecular networks with genetic variant interpretation for precision medicine. *Wiley Interdiscip Rev Syst Biol Med*, **11**, e1443.
7. Capriotti,E. *et al.* (2006) Predicting the insurgence of human genetic diseases associated to single point protein mutations with support vector machines and evolutionary information. *Bioinformatics*, **22**, 2729–34.
8. Capriotti,E. *et al.* (2013) WS-SNPs&GO: a web server for predicting the deleterious effect of human protein variants using functional annotation. *BMC Genomics*, **14 Suppl 3**, S6.
9. Capriotti,E. and Fariselli,P. (2017) PhD-SNPg: a webserver and lightweight tool for scoring single nucleotide variants. *Nucleic Acids Res*, **45**, W247–W252.
10. Claussnitzer,M. *et al.* (2020) A brief history of human disease genetics. *Nature*, **577**, 179–189.
11. Fernald,G.H. *et al.* (2011) Bioinformatics challenges for personalized medicine. *Bioinformatics*, **27**, 1741–8.
12. Grimm,D.G. *et al.* (2015) The Evaluation of Tools Used to Predict the Impact of Missense Variants Is Hindered by Two Types of Circularity. *Hum. Mutat.*, **36**, 513–523.
13. Kent,W.J. *et al.* (2002) The human genome browser at UCSC. *Genome Res.*, **12**, 996–1006.
14. Kircher,M. *et al.* (2014) A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet*, **46**, 310–5.
15. Landrum,M.J. *et al.* (2020) ClinVar: improvements to accessing data. *Nucleic Acids Res.*, **48**, D835–D844.
16. MacArthur,D.G. *et al.* (2014) Guidelines for investigating causality of sequence variants in human disease. *Nature*, **508**, 469–76.

17. McInnes,G. *et al.* (2021) Opportunities and challenges for the computational interpretation of rare variation in clinically important genes. *Am. J. Hum. Genet.*, **108**, 535–548.
18. Miller,M. *et al.* (2019) funtrp: identifying protein positions for variation driven functional tuning. *Nucleic Acids Res.*, **47**, e142.
19. Niroula,A. and Vihinen,M. (2016) Variation Interpretation Predictors: Principles, Types, Performance, and Choice. *Hum. Mutat.*, **37**, 579–597.
20. Ozturk,K. and Carter,H. (2021) Predicting functional consequences of mutations using molecular interaction network features. *Hum. Genet.*
21. Pedregosa, F. *et al.* (2011) Scikit-learn: Machine Learning in Python. *JMLR*, **12**, 2825–2830.
22. Petrosino,M. *et al.* (2021) Analysis and Interpretation of the Impact of Missense Variants in Cancer. *Int. J. Mol. Sci.*, **22**, 5416.
23. Pollard,K.S. *et al.* (2010) Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res*, **20**, 110–21.
24. Rentzsch,P. *et al.* (2019) CADD: predicting the deleteriousness of variants throughout the human genome. *Nucleic Acids Res.*, **47**, D886–D894.
25. Rost,B. *et al.* (2016) Protein function in precision medicine: deep understanding with machine learning. *FEBS Lett.*, **590**, 2327–2341.
26. Siepel,A. *et al.* (2005) Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res*, **15**, 1034–50.
27. Suzek,B.E. *et al.* (2007) UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics*, **23**, 1282–8.
28. Tennessen,J.A. *et al.* (2012) Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science*, **337**, 64–69.
29. Walsh,I. *et al.* (2021) DOME: recommendations for supervised machine learning validation in biology. *Nat. Methods*.

Figures

Figure 1

(A) Boxplot and (B) cumulative distributions of the PhyloP100way score in the variation sites for the subsets of Pathogenic and Benign variants in the CommonClinvar dataset. The maximum distance between the two distributions is at 4.7 that corresponds to a Kolmogorov Smirnov distance of 0.57.

Figure 2

Distributions of the frequencies of references (A) / alternative (C) alleles in the variation sites and the frequencies of wild-type (B) / mutant (D) residues in protein mutation sites for the subset of Pathogenic and Benign variants in the CommonClinvar dataset.

Figure 3

Receiver Operating Characteristic (A) and Precision Recall (B) curves for the different gradient boosting algorithms (PPScore, DNAProf, ProtProf) and CADD on the NewClinvar dataset.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [20210906conservationsupmat.docx](#)
- [commonclinvardataset.tsv](#)
- [newclinvardataset.tsv](#)