Heavy Metal -Induced Oxidative Stress And Alteration In Secretory Proteins In Yeast Isolates

Gurulingaiah Bhavya

University of Mysore Department of Biotechnology

Nagaraja Geetha

University of Mysore Department of Biotechnology

Kavitha Hiremath

Karnatak University Dharwad

SUDISHA JOGAIAH (✉ jsudish@kud.ac.in)

Karnatak University Dharwad <https://orcid.org/0000-0003-2831-651X>

Research Article

Keywords: Candida sp., exoproteome (Secretory proteins), glutathione, superoxide dismutase, low molecular weight proteins

Posted Date: September 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-884162/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at Archives of Microbiology on February 14th, 2022. See the published version at <https://doi.org/10.1007/s00203-022-02756-6>.

Abstract

In the recent years, yeasts have evolved as potent bioremediative candidates for the detoxification of xenobiotic compounds found in the natural environment. *Candida* sp. are well studied apart from *Saccharomyces* in heavy metal detoxification mechanisms. In the current study, *Candida parapsilosis* strain ODBG2, *Candida* sp. strain BANG3 and *Candida viswanathii* strain ODBG4 were isolated from industrial effluents and contaminated ground water were studied for their metal tolerance. Among these three isolates, the metal tolerance was found to be more towards Lead (Pb 2mM), followed by Cadmium (Cd 1.5mM) and Chromium (Cr(VI), 1mM). On further exploring the involvement of primary defensive enzymes in these isolates exhibited towards metal tolerance, the anti-oxidative enzyme Superoxide dismutase (SOD) was found to be prominently high as 25% during first 24h of metal-isolate interaction. In the Catalase (CAT) enzyme assay, it was observed that, the increased enzyme activity at 48h also triggered the activity of peroxidases (PO), which lead to the increase in reduced glutathione (GSH) in the organism by 0.87-1.9 folds, as a metal chelator and also as a second line of defensive molecule. The exoproteome profile showed the early involvement (exponential growth phase) of secreted proteins (low molecular weight) of about ~40-45kDa under Cd and Pb stress (0.5mM). The exoproteome profiling under heavy metal stress in *Candida parapsilosis* strain ODBG2 and *Candida viswanathii* strain ODBG4 is the first report.

Introduction

Metals play an important role in homeostasis in the biological systems. Among the metals few are considered as hazardous to living biota and characterized as heavy metals (HM)s and metalloids. The HMs has become a part of the environment by geochemical cycles and as well as anthropogenic activities such as usage of pesticides in excess, industrial effluents, chemical manufacturing unit sites etc., (Satapute et al. 2019). The presence of hazardous metals in the nature exerts various kinds of deformities in the biological systems in microbes to human at the level of biochemical to molecular sites (Lin et al. 2019; Amaakwah et al. 2020). In the global scenario, the scarcity of portable water is on an higher end due to contaminants in fresh water system and as well as ground water due to geogenic origin as well as percolating water from industries, that too especially with arsenic (As). The HMs like mercury (Hg), lead (Pb), cadmium (Cd), fluoride (F), accounts as major contaminants (Mohankumar et al. 2016; Shaji et al. 2021). Certain microbes like bacteria and fungi form habitat naturally in these contaminated sites by developing strong biochemical defense mechanisms to thrive under harsh conditions (Bhavya et al. 2015; Geetha et al. 2020). Among the microbes, filamentous fungi and yeasts have evolved strongly and portraying has potent bioremediative candidates for xenobiotic compounds (Dey et al. 2020). The recent studies, have reported on *Candida* spp. especially *Candida tropicalis*, *Candida albicans*, *Candida parapsilosis*, *Candida etchellsii*, *Candida lipolytica*, as heavy metal resistant yeasts (Kachiprath et al. 2019). The fungi/yeast possess a versatile group of mechanisms such as biosorption, bioaccumulation, sequestration which aids in the detoxification of heavy metals. Apart from these mechanisms, primary way to combat the metal stress is through the action of antioxidant enzymes such as catalase (CAT),

peroxidases (PO), glutathione peroxidases (GPx), superoxide dismutase (SOD), glutathione S transferase (GSTs), etc.,. The primary targets of these enzymes are the reactive oxygen species (ROS) generated through biotic and abiotic stresses (Bandyopadhyay et al. 1999) thus reducing the toxicity as well as activation of series of enzymatic pathway and leading to production of low molecular compound like GSH, Metallothiones, oxalic acid, secondary metabolites, etc. These molecules interact with the heavy metals through -SH groups, rendering the inactivating heavy metals and rendering the homeostasis maintenance of the cell. The detoxification process is a combinatorial strategy during which, involvements of various extracellular enzymes like lipase, DNAase, laccase are identified or observed (Santos et al. 2017). During stress and infectious state, yeast secretes various kinds of metabolites to its surrounding environment to regulate its growth and maintenance. The secretory molecules are not only encompassed of primary and secondary metabolites, exopolysaccharides, but also possess various sets of proteins known as exoproteome/extracellular proteins. These metabolites and exoproteins help the fungi to adapt for the harsh changes in its living habitat. The profiling of exoproteins in fungi under heavy metal stress helps in understanding the stress response systems (Selvam et al. 2015).

Based on this knowledge, the current work was targeted to study the role of SOD and CAT, a antioxidant enzymes and reduced GSH levels at different time intervals and metal concentrations. Further, as there is a lacuna in the field of exoproteome stress biology; preliminary work was carried out to know the secretory proteome profiles on SDS-PAGE. This part of work has been reported for the first time in *Candida parapsilosis* and *Candida viswanathii* strains isolated from heavy metal contaminated water source from the prevailing study.

Materials And Methods

2.1 Isolation of yeasts from industrial effluents.

The textile effluent samples and heavy metal contaminated ground water samples were collected from Bengaluru and Odisha industrial areas respectively and brought to the laboratory, stored at ambient temperature till further utilization. Isolation of yeasts was carried out by spread plate technique using 100µl of each sample plated separately on yeast extract dextrose agar media (YPDA: yeast extract 1%, peptone and dextrose 2% each, agar 1.5%; pH 6.2) and incubated at 35°C for 48hr. The colonies were selected and pure cultures were used for further studies.

2.2 Preparation of metal stock solutions

The metal salts as a source of heavy metal (HM) ions used for the experiments were cadmium chloride (CdCl₂), lead nitrate [Pb(NO₃)₂], and potassium dichromate (K₂Cr₂O₇). Each HM stock solution (50mM) was prepared in deionized water and stored at room temperature.

2.3 Determination of MIC (Minimum inhibitory concentration) /Metal Tolerance Test.

The MIC towards the heavy metals was determined at different metal ion concentrations for the yeast isolates at a constant temperature of 35⁰C in YPD broth media. The actively growing cells (optical density 1.5 at 600nm) were transferred into YPD broth amended with heavy metal salts and the growth was recorded at 24hr interval till 120hr. The isolate growing in non-heavy metal-amended media was considered as a control. The concentration at which the cultures exhibited minimal growth was considered as MIC.

2.4 Molecular Identification of heavy meal tolerant yeast isolates/strains.

Colony PCR technique was chosen for the amplification of 5.8s rRNA gene by utilizing the universal primers, the forward primer ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and the reverse primer ITS4 (5' TCCTCCGCTTATTGATATGC-3') (White et al. 1990). In brief, the log phase growing colonies were selected and dispensed in 50µl of deionized water. From this, 2 µl was used for amplification. The PCR reaction mixture consisted of ready to use master mix 20 µl (TaKaRa Taq HS perfect Mix), forward primer (2 µl - ITS 1), reverse primer (2 µl - ITS 4), 2 µl cells as template and 14 µl of sterile water. PCR was carried out at 95⁰C for 5 min 30sec for initial denaturation followed by 30 cycles of denaturation at 94⁰C for 45sec, annealing at 56⁰C for 55sec, and extension at 72⁰C for 60sec followed by final extension for 10min at 72⁰C. PCR products (amplicons) were analyzed on 1.2% agarose gel using 1X Tris borate EDTA buffer (1X TBE). The PCR amplicons were sent for Sanger sequencing to Juniper Life science Pvt Ltd, Bengaluru for sequencing using ITS 1 (forward primer). The obtained sequences were identified by using nBLAST from NCBI (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). The sequences were deposited in Gene Bank and accession numbers were obtained. The sequence alignment and phylogenetic tree was carried out among the identified stains using MEGA X (Kumar et al. 2018).

2. 5. Oxidative stress-induced enzyme studies (CAT, SOD).

2. 5.1. Enzyme extraction:

The protein extraction was carried out according to Ilyas et al. (2016) with slight modification. In brief, cells were harvested by centrifuging at 12000rpm, washed thrice in phosphate buffer of pH 7. Cells were suspended in 1.5ml of extraction buffer containing 50mM Tris-HCl buffer of pH 7.4, 5mM EDTA, 0.5M sucrose, 100mM KCl and 1mM PMSF. Cells were sonicated for 20s with 50s interval at 25Hz for 3 cycles, and centrifuged at 10000rpm for 12min (4⁰C). Protein concentration in the sample was determined by Bradford method and BSA as a standard (Bradford, 1976). The extracts were aliquot and stored at 4⁰C until further use.

2.5.2. Enzyme activity assay

2.5.2.1. Superoxide dismutase (SOD) assay

SOD activity was determined based on indirect assay i.e., inhibition of auto oxidation of quercetin (Kengaiah et al. 2020). The reaction contained 16mM sodium phosphate buffer of pH 7, 8mM TEMED,

200mM EDTA, 0.1% quercetin and crude enzyme extract. The reaction was monitored at 406nm for 1 min. Percentage inhibition of autoxidation of quercetin was determined.

2.5.2.2. Catalase (CAT) assay

The CAT activity was carried out according to Zeng et al. (2012). The reaction mixture contained of 50mM potassium phosphate buffer of pH 7, 8mM hydrogen peroxide (H_2O_2) (30%), and the crude enzyme extract. The decrease in the absorbance at 240nm was recorded for 2min. The activity was calculated and expressed as μ moles of H_2O_2 decomposed/min/mg of protein.

2.5.3. Anti-oxidant biomarker Assay

Reduced Glutathione (GSH) assay

The reduced GSH level in the samples was determined according to Ellman's method (Ellman, 1959) with slight modification. The cells were homogenized in extraction buffer consisting of 0.6% sulfosalicylic acid, 0.1% Tritonx100 in 0.1M potassium phosphate buffer pH 7.5 with 5mM EDTA. The cell lysate was collected by centrifuging at 8000rpm for 10min at 4⁰C. For the determination of reduced GSH content, to 100 μ l of about extract, Tris-HCl buffer of pH 8.2, 10mM DTNB was added and incubated for 10min at ambient temperature. The reaction concoction was spinned at 3,000g for 5min. The absorbance of clear supernatant was measured at 415nm. The amount of reduced GSH in the sample was determined by using standard reduced GSH curve. The amount of reduced GSH was expressed in the terms of μ M/mg of protein.

2.6. Secretory protein analysis by SDS-PAGE – A preliminary study

The cells were separated from the culture broth by centrifuging at 12,000rpm for 20min at 4⁰C. The filtrate was filtered through 0.22 μ m membrane to remove the cell debris. The proteins were precipitated by TCA-acetone precipitation method with modification (Wu et al. 2014 and Chen et al. 2015). Briefly, the filtrate was added with 10% TCA followed by overnight incubation at 4⁰C. The precipitated proteins were separated by centrifuging at 12,000rpm, 15min at 4⁰C. Then the proteins were suspended in 80% ice-cold acetone encompassed of 0.07% β -mercaptoethanol and incubated for 2hrs at 4⁰C. The contents were centrifuged; pellet was washed thrice with methanol and acetone respectively. Finally, the protein pellets were air-dried and solubilized in buffer containing urea (7M), thiourea (2M), CHAPS (1%) and 65mM DTT. The Bradford method (Bradford, 1976) was adapted for determination of protein concentration in the samples. The 60 μ g of protein was suspended in SDS-sample buffer (Tris-Hcl buffer of pH 6.8, SDS, β -mercaptoethanol, glycerol and bromophenol blue), heated at 95⁰C for 5min. An equal amount of protein (60 μ g/sample) was loaded along with the marker; the proteins were separated on 10% SDS-polyacrylamide mini gel on constant voltage of 70V for 30min, followed by 80V till the complete run. The gel was stained by CBB staining technique and differences in the banding pattern was noted (Laemmli, 1970).

Results And Discussion

3.1. Isolation of yeasts from industrial effluents and contaminated ground water.

The industrial effluent collected from the Bengaluru and contaminated ground water from Gajapathi area in Odisha, lead to the isolation, growth and axenic culturing of one and two yeast colonies respectively. The isolates were coded as BANG3, ODBG2, and ODBG4 for further studies and molecular identification. The microscopic observation exhibited the single cell structure ovoid in shapes with various budding stages along with clear and distinguished cell wall structure. Figure 01 represents the macroscopic and microscopic views of the yeast isolates. There were not much changes in the morphological and microscopic observations, thus the exact identification the isolates was not revealed, and indicated the necessity of molecular based identification.

3.2. Determination of MIC (Minimum inhibitory concentration) /Metal Tolerance Test.

The metal tolerance test/MIC revealed the capability of the isolate ODBG2 later identified as *Candida parapsilosis* strain ODBG2, to utilize the heavy metals for its growth and tolerated up to the concentration of 2.0mM Pb, 1.5mM Cd and 0.5mM Cr. The significant difference in the growth pattern was observed after 48hr of metal-isolate interaction with respect to Cd and Pb. While significant difference was recorded after 24hr under hexavalent Cr stress. In the second isolate BANG3; *Candida* sp. strain BANG3; exhibited the metal tolerance up to 2mM of Pb and 1.0mM Cd, with significant growth difference with respect to control (without HM amended) was observed after the incubation period of 72hr and 24hr respectively, whereas zero tolerance towards Cr(VI). The isolate ODBG4; *Candida vishwanathii* strain ODBG4 was adaptive to grow under Pb at 1.5mM followed by Cr and Cd at 1.0mM with a significant difference after the time interval of 48hr. The growth curve of yeast isolate ODBG2, BANG3 and ODBG4 with and without metal amended is shown in the Fig. 02. The recent study by Bansal et al. (2019) on *Candida parapsilosis*, an environmental isolate and its metal tolerance have indicated the survival capability and utilization under heavy metal such as Ni, Pb and Cd at the concentration of 12mM. Similarly, the various researches have shown the ability of *Candida* sp. to tolerate and to grow under various heavy metals which were isolated from contaminated sources like soil, water and industrial effluent system. The difference in metal tolerance might be ascribed to the level of HM contamination in the water samples, the contact or incubation time (Azcon et al. 2010; Munner et al. 2013; Moreno et al. 2019) as well as dependent on metal ion status and species (Balsalobre et al. 2003).The MICs of different isolates has been summarized in the **Table 01**.

3.3. Molecular Identification of heavy meal tolerant yeast isolates/strains.

The PCR amplicons of the isolates with the codes ODBG2, BANG3, and ODBG4 resulted in the product of ~ 500bp size (**Fig. 3**). Further the sequencing of the region ITS1, 5.8rRNA and ITS2 using ITS 1 (forward primer) of the amplicons ODBG2, BANG3, and ODBG4 resulted in 493bp, 438bp, and 447bp product. BLAST analysis showed > 98% identity with sequences of *Candida parapsilosis* (KX652405, JN989529, MNT33072) with respect to our isolate code ODBG2, *Candida* sp. (MH802509, MN124747, KJ706734)

with respective isolate code BANG3 and *Candida vishwanathii* (KU729067, MK394124, KC608220) with respect isolate code ODBG4. The obtained and identified sequences were deposited in GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) and Accession numbers were obtained. The code of isolates and accession no. has been represented in the table 02.

The evolutionary history was inferred by using the Maximum Likelihood method and Tamura-Nei model (Tamura and Nei, 1993). Evolutionary analyses was conducted in MEGA X (Felsenstein, 1985; Kumar et al., 2018), **Fig. 4** represent the phylogenetic tree.

3.4. Oxidative stress-induced enzymatic and non-enzymatic studies (SOD, CAT and GSH).

3.4.1 Effect of heavy metal-induced oxidative stress on SOD activity

The primary defense antioxidant enzymes in any biological system are attributed towards the activity of superoxide dismutase (SOD), catalase (CAT) and peroxidases (PO). The activity of SOD was detected based on the percentage inhibition of quercetin autoxidation in the control and treated samples. The SOD activity in *Candida parapsilosis* strain ODBG2 was found to be high in the range of 16% (Cd 0.5mM), 27.5% (Cd 1.0mM), 22.5% (Pb 0.5mM) and 25% (Pb 1.0mM) during first 24hr of stress, however the gradual decrease in the abundance at 48hr and thereafter (Fig. 5A). The SOD activity in *Candida viswanathii* strain ODBG4 was relatively high for Cd (0.5mM concentration) at 24hr, for Pb (1mM) it slightly increased at 48 and 72hr (Fig. 5B), indicating minimal effect on SOD. SOD is the primary defense enzyme found to be active during abiotic stress-induced intracellular ROS in biological systems, which interacts and converts into H₂O₂ and water molecule, leading to the sequential activation of other primary defense enzymes such as CAT and PO, subsequently the non-enzymatic small compounds like GSH (Bandyopadhyay et al., 1999). The varied response of SOD with respective different time intervals and concentrations has been observed in fungi such as *Pleurotus ostretus* HAU-2 upon Pb (Zhang et al., 2016), *Trichosporon asahii* upon different heavy metal and metalloid treatments (Ilyas et al., 2014). SOD was found to be not effective at the higher concentration rate and incubation time, which is indicating the role of CAT and PO in the system.

3.4.2. Effect of heavy metals induced oxidative stress enzyme – CAT

The CAT showed increased activity of 38 and 40 µmoles/mg of protein/min upon Pb stress at 0.5mM and 1.0mM concentration during at 48hr incubation, subsequently decreased on 72hr of incubation with 20 and 28 µmoles mg of protein/min in *Candida parapsilosis* strain ODBG2 (Fig. 6A). Similarly, CAT activity was higher at 48hr under Pb stress at both the concentrations with 15 and 16 µmoles/mg of protein/min in *Candida viswanathii* strain ODBG4, whereas the Cd stress at 0.5mM induced an steady increase with activity of about 16 µmoles and 18 µmoles/mg of protein/min at 48hr and 72hr respectively (Fig. 6 B). The activity of CAT was found to vary based on the metal ion concentration, time and the isolate. The high CAT activity in the basal cells might trigger other relative enzymes such as PO, which might be strategic in mitigating the toxicity of heavy metal ion such as Cd, thus exhibiting steady increase under Cd stress (Pradhan et al., 2017).

3.4.3 Effect of heavy metal-induced oxidative stress on non-enzymatic antioxidant – Reduced GSH

The intracellular GSH plays a key role for maintaining cell homeostasis under normal and stress conditions. In the current study, the levels of reduced GSH in the cells under heavy metal stress was found to be time and concentration-dependent with respect to the treated yeast isolates. In *Candida parapsilosis* strain ODBG2, the levels of reduced GSH under Cd and Pb stress at 24hr was found to be lesser to that of untreated cells, whereas at 48hr there was an increase by 1.9 fold (Cd 1.0mM) followed by 0.87 fold in Pb (1.0mM) at 48hr. In *Candida viswananthii* strain ODBG4, a maximum of 1.6 and 1.7 fold increased reduced GSH was found at Cd and Pb (1.0mM) concentration. Increased GSH content at 48hr under Pb stress is similar to the studies by Rehaman and Anjum (2011), with Cd stress in *Saccharomyces* spp. (Fauchon et al., 2002). Further, decrease in reduced GSH at 72h under metal treatment might be due to the interaction of SH-group of GSH with that of heavy metal, the protective ability towards the toxicity and detoxification of heavy metals (Gharieb and Gadd, 2004; Ilyas et al., 2017) or due to the increased intracellular ions induced activation of other pathways, which leads to imbalance riot in homeostasis (Zhang et al., 2016). The reduced GSH content under different metal stress and at different time intervals is represented in the Fig. 7(A) and Fig. 7(B) for the *Candida parapsilosis* strain ODBG2 and *Candida viswananthii* strain ODBG4 respectively.

3.5. Alterations in Secretory proteins upon different heavy metal stress in *Candida parapsilosis* strain ODBG2 and *Candida viswananthii* strain ODBG4 by SDS-PAGE.

Candida parapsilosis strain ODBG2 and *Candida viswananthii* strain ODBG4 extracted at 48hr of post-heavy metal stress (0.5mM concentration) of Pb(II), Cr(IV) and Cd (II) showed similar secretory protein profiles on 10% SDS-PAGE (Fig: 8.A and Fig. 8.B). In the present study, the SDS-PAGE profile showed two (2) over-expressed and intense protein bands in the molecular range of ~ 40-45kDa under Cd stress and faint polymorphic bands in the range of ~ 66-116kDa. The Pb stress also signposted the secretory protein prominently expressed approximately at ~ 43-44kDa. The similar profiling under Cd and Pb in these isolates might be attributed to similar kind of signal perception, gene activation, protein and 1⁰ and 2⁰ metabolite production at the species level. Since this work is been carried out for the first time, further studies are required for specifying these expressed protein bands at various molecular ranges.

Conclusion

The current study is emphasized on the differential ability of heavy metal tolerance among *Candida* sp. i.e *Candida parapsilosis* strain ODBG2 and *Candida viswananthii* strain ODBG2 isolated from industrial effluent-mediated contaminated ground water system. These isolates have an ability to be used as potent heavy metal detoxifiers in contaminated environment. It can be presumed that the potentiality of heavy metal tolerance towards Cd and Pb is due to communication in intra and extra-cellular secretory molecules modulated cell signaling during log phase growth under heavy metal stress. This contemporary work, has paved a way for further investigations on exploring exoproteome strategies in *Candida* sp. upon heavy metal stress.

Abbreviations

Pb –Lead

Cd –Cadmium

mM- Millimolar

SOD –Superoxide dismutase

CAT-Catalase

GSH-Glutathione

YPDA –Yeast peptone dextrose agar

HM-Heavy metal (s)

MIC – Minimal inhibitory concentration

ITS – Internal Transcribe Sequences

PCR –Polymerase chain reaction

TBE-Tris-borate EDTA

BLAST – Basic Local Alignment search tool

MEGA –Molecular Evolutionary Genetics Analysis

PMSF- Phenylmethylsulfonyl fluoride

H₂O₂ – Hydrogen peroxide

DTNB - 5, 5'-dithiobis(2-nitrobenzoic acid)

SDS-PAGE: Sodium dodecyl sulphate–polyacrylamide gel electrophoresis

CHAPS: 3-cholamidopropyl dimethylammonio 1-propanesulfonate

DTT – Dithiothreitol

Declarations

Author contributions

Supervision of research work: NG and SJ. Benchwork: GB. Data validation: GB, NG, KH and SJ. Figures and charts: GB, NG and SJ. Contributed reagents/chemicals: NG. Wrote the original manuscript: GB, NG and SJ. All the authors read and approved the manuscript.

Conflict of Interest

The authors do not have any competing interest in the submitted work.

Acknowledgements

Authors are thankful for Department of Studies in Biotechnology, University of Mysore, Mysuru for facilities provided. The authors also extend their thanks to facilities available at Laboratory of Plant Healthcare and Diagnostics, PG Department of Biotechnology and Microbiology, Karnatak University, Dharwad.

References

1. Azcón R, del Carmen Perálvarez M, Roldán A, Barea JM (2010) Arbuscular mycorrhizal fungi, *Bacillus cereus*, and *Candida parapsilosis* from a multi-contaminated soil alleviate metal toxicity in plants. *Microb Ecol* 59(4):668–677. DOI 10.1007/s00248-009-9618-5
2. Balsalobre L, De Silóniz MI, Valderrama MJ, Benito T, Larrea MT, Peinado JM (2003) Occurrence of yeasts in municipal wastes and their behaviour in presence of cadmium, copper and zinc. *J. Basic Microbiol: An International Journal on Biochemistry, Physiology, Genetics, Morphology, and Ecology of Microorganisms* 43(3):185–193. 10.1002/jobm.200390021
3. Bandyopadhyay U, Das D, Banerjee RK (1999) Reactive oxygen species: oxidative damage and pathogenesis. *Curr. Sci.* 658–666
4. Bhavya G, Kumar CRS, Swati K, Geetha N (2015) In search of industrial clean-up clients; evaluation of heavy metal tolerability of rhizospheric *Trichoderma*. *Int J Innov Sci Eng Technol* 2:948–952
5. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72(1–2):248–254. 10.1016/0003-2697(76)90527-3
6. Chen G, Zhou Y, Zeng G, Liu H, Yan M, Chen A, Guan S, Shang C, Li H, He J (2015) Alteration of culture fluid proteins by cadmium induction in *Phanerochaete chrysosporium*. *J Basic Microbiol* 55(2):141–147. 10.1002/jobm.201300398
7. Ellman GL (1959) Tissue sulfhydryl groups. *Arch Biochem Biophys* 82(1):70–77
8. Fauchon M, Lagniel G, Aude JC, Lombardia L, Soularue P, Petat C, Marguerie G, Sentenac A, Werner M, Labarre J (2002) Sulfur sparing in the yeast proteome in response to sulfur demand. *Mol cell* 9(4):713–723. 10.1016/S1097-2765(02)00500-2
9. Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evol* 39(4):783–791. 10.1111/j.1558-5646.1985.tb00420.x

10. Gharieb MM, Gadd GM (2004) Role of glutathione in detoxification of metal (loid) s by *Saccharomyces cerevisiae*. *Biomaterials* 17(2):183–188. 10.1023/B:BIOM.0000018402.22057.62
11. Ilyas S, Rehman A, Ilyas Q (2017) Heavy metals induced oxidative stress in multi-metal tolerant yeast, *Candida* sp. PS33 and its capability to uptake heavy metals from wastewater. *Pakistan Journal of Zoology*, 49(3)
12. Ilyas S, Rehman A, Coelho AV, Sheehan D (2016) Proteomic analysis of an environmental isolate of *Rhodotorula mucilaginosa* after arsenic and cadmium challenge: Identification of a protein expression signature for heavy metal exposure. *J Proteom* 141:47–56. 10.1016/j.jprot.2016.04.012
13. Ilyas S, Rehman A, Varela AC, Sheehan D (2014) Redox proteomics changes in the fungal pathogen *Trichosporon asahii* on arsenic exposure: identification of protein responses to metal-induced oxidative stress in an environmentally-sampled isolate. *PLoS One* 9(7):p.e102340. 10.1371/journal.pone.0102340
14. Kengaiah J, Nandish SKM, Ramachandraiah C, Shivaiah A, Vishalakshi GJ, Paul M, Santhosh MS, Shankar RL, Sannanigaiah D (2020) Protective effect of tamarind seed coat ethanol extract on eryptosis induced by oxidative stress. *Biochemistry* 85(1):119–129. 10.1134/S0006297920010113
15. Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35(6):1547. 10.1093/molbev/msy096
16. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227(5259):680–685. 10.1038/227680a0
17. Moreno A, Demitri N, Ruiz Baca E, Vega González A, Polentarutti M, Cuéllar Cruz M (2019) Bioreduction of precious and heavy metals by *Candida* species under oxidative stress conditions. *Microb Biotechnol* 12(6):1164–1179. 10.1111/1751-7915.13364
18. Muneer B, JIqbal M, Shakoori FR, Shakoori AR (2013) Tolerance and biosorption of mercury by microbial consortia: potential use in bioremediation of wastewater. *Pakistan Journal of Zoology*, 45(1)
19. Pradhan A, Herrero-de-Dios C, Belmonte R, Budge S, Lopez Garcia A, Kolmogorova A, Lee KK, Martin BD, Ribeiro A, Bebes A, Yucel R (2017) Elevated catalase expression in a fungal pathogen is a double-edged sword of iron. *PLoS pathogens*13(5):p.e1006405. 10.1371/journal.ppat.1006405
20. Rehman A, Anjum MS (2011) Multiple metal tolerance and biosorption of cadmium by *Candida tropicalis* isolated from industrial effluents: glutathione as detoxifying agent. *Environ Monit Assess* 174(1):585–595. 10.1007/s10661-010-1480-x
21. Tamura K, Nei M (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* 10(3):512–526. 10.1093/oxfordjournals.molbev.a040023
22. White TJ, Bruns T, Lee SJW, T, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods applications* 18(1):315–322

23. Wu X, Xiong E, Wang W, Scali M, Cresti M (2014) Universal sample preparation method integrating trichloroacetic acid/acetone precipitation with phenol extraction for crop proteomic analysis. *Nature protocols* 9(2):362–374. 10.1038/nprot.2014.022
24. Zeng GM, Chen AW, Chen GQ, Hu XJ, Guan S, Shang C, Lu LH, Zou ZJ (2012) Responses of *Phanerochaete chrysosporium* to toxic pollutants: physiological flux, oxidative stress, and detoxification. *Environ Sci Technol* 46(14):7818–7825. 10.1021/es301006j
25. Zhang S, Zhang X, Chang C, Yuan Z, Wang T, Zhao Y, Yang X, Zhang Y, La G, Wu K, Zhang Z (2016) Improvement of tolerance to lead by filamentous fungus *Pleurotus ostreatus* HAU-2 and its oxidative responses. *Chemosphere* 150:33–39. 10.1016/j.chemosphere.2016.02.003
26. Shaji E, Santosh M, Sarath KV, Prakash P, Deepchand V, Divya BV (2021) Arsenic contamination of groundwater: A global synopsis with focus on the Indian Peninsula. *Geosci Front* 12(3):101079. 10.1016/j.gsf.2020.08.015
27. Mohankumar K, Hariharan V, Rao NP (2016) Heavy metal contamination in groundwater around industrial estate vs residential areas in Coimbatore, India. *Journal of clinical diagnostic research: JCDR* 10(4):BC05
28. Kachiprath B, Solomon S, Jayanath G, Philip R (2019) Mangrove microflora as potential source of hydrolytic enzymes for commercial applications. *IJMS* 48(5):678–684
29. Geetha N, Bhavya G, Abhijith P, Shekhar R, Dayananda K, Jogaiah S (2021) Insights into nanomycoremediation: Secretomics and mycogenic biopolymer nanocomposites for heavy metal detoxification. *J Hazard Mater* 409:124541. 10.1016/j.jhazmat.2020.124541
30. Dey P, Malik A, Mishra A, Singh DK, von Bergen M, Jehmlich N (2020) Mechanistic insight to mycoremediation potential of a metal resistant fungal strain for removal of hazardous metals from multimetal pesticide matrix. *Environ Pollut* 262:114255
31. Santos DK, Resende AH, de Almeida DG, Soares da Silva RDCF, Rufino RD, Luna JM, Banat IM, Sarubbo LA (2017) *Candida lipolytica* UCP0988 biosurfactant: potential as a bioremediation agent and in formulating a commercial related product. *Front Microbiol* 8:767
32. Selvam RM, Nithya R, Devi PN, Shree RB, Nila MV, Demonte NL, Thangavel C, Maheshwari JJ, Lalitha P, Prajna NV, Dharmalingam K (2015) Exoproteome of *Aspergillus flavus* corneal isolates and saprophytes: identification of proteoforms of an over secreted alkaline protease. *J Proteom* 115:23–35

Tables

Table 1: Minimal Inhibitory concentration of yeast isolates to different heavy metal concentrations.

Isolates code	MIC in mM		
	Pb (II)	Cd (II)	Cr (VI)
ODBG2	2	1.5	0.5
BANG3	2	1	-
ODBG4	1.5	1	1

Table 2: Details of the identified isolates with their accession number.

Isolate code	Accession no.	Identification/deposition name
ODBG2	MZ841637	<i>Candida parapsilosis</i> strain ODBG2
BANG3	MZ841638	<i>Candida</i> sp. strain BANG3
ODBG4	MZ841639	<i>Candida viswanathii</i> strain ODBG4

Figures

Figure 1

Macroscopic view of the yeast isolates A: ODBG2; B: BANG3; C: ODBG4 on Yeast extract peptone agar media. The respective microscopic views at 40X in a, b and c showing the typical yeast cell structure.

Candida parapsilosis strain ODBG2 growth under Pb stress

Candida parapsilosis strain ODBG2 growth under Cd stress

Candida parapsilosis strain ODBG2 growth under Cr stress

Candida sp. strain BANG3 growth under Pb stress

Candida sp. strain BANG3 growth under Cd stress

Candida viswanathii strain ODBG4 growth under Cr stress

Candida viswanathii strain ODBG4 growth under Pb stress

Candida viswanathii strain ODBG4 growth under Cd stress

Figure 2

The growth of yeast isolates under different heavy metal stress and at different time intervals. Respective MICs is represented in the Table 1. Based on the results obtained further, the isolate code ODBG2 and ODBG4 were subjected for antioxidant enzymes and reduced GSH content studies upon Cd and Pb stress at different time intervals.

Figure 3

1.2% Agarose gel showing the colony PCR product of ~ 500bp size of the isolates ODBG2, BANG3 and ODBG4 in the Lane 4,5, and 6. Lane 1: Marker: 100bp. Along with other samples product in lane 2 and 3 (not discussed in this article).

Figure 4

The phylogenetic analysis by Maximum Likelihood method with 1000bootstrap replication in MEGA X, representing the evolutionary relationship of the isolates with other 5.8rRNA ITS1-5.8-ITS 2 sequences of related *Candida* species.

Figure 5

A: Inhibition of Quercetin by SOD heavy metal-treated cells in *Candida parapsilosis* strain ODBG2. The control protein samples were normalized to get 50% inhibition activity during the assay. Based on this inhibition activity, the test samples were assayed. The bar value represents the mean $\pm$ SD of triplicates, where $p < 0.0001$. Significant difference with respect to control samples is represented by Dunnett's multiple comparisons test. B: Inhibition of Quercetin by SOD in heavy metal-treated cells in *Candida viswanathii* strain ODBG4. The protein samples control was normalized to get the 50% inhibition activity during the assay. Based on this inhibition activity, the test samples were assayed. The bar value represents the mean $\pm$ SD of triplicates, where $p < 0.0001$. Significant difference with respect to control samples is represented with Dunnett's multiple comparisons test.

Figure 6

(A): Catalase activity in heavy metal-treated cells in *Candida parapsilosis* strain ODBG2. The bar value presences the mean $\pm$ SD of triplicates, where $p < 0.0001$. Significant difference with respect to control samples is represented with Dunnett's multiple comparisons test. (B): Reduced GSH levels in heavy metal treated cells in *Candida viswanantii* strain ODBG4. The bar value represents the mean $\pm$ SD of triplicates, where $p < 0.0001$. Significant difference with respect to control samples is represented by Dunnett's multiple comparisons test.

Figure 7

(A): Reduced GSH levels in heavy metal-treated cells in *Candida parapsilosis* strain ODBG2. The bar value represents the mean $\pm$ SD of triplicates, where $p < 0.0001$. Significant difference with respect to control samples is represented by Dunnett's multiple comparisons test. (B): Reduced GSH levels in heavy metal-treated cells in *Candida viswananthii* strain ODBG4. The bar value represents the mean $\pm$ SD of triplicates, where $p < 0.0001$. Significant difference with respect to control samples is represented by Dunnett's multiple comparisons test.

Figure 8

A: Secretory protein profile of *Candida parapsilosis* strain ODBG2. Secretory protein (60 μ g) separated on 10% SDS-PAGE and stained with CBB R-250. The image was documented in invert image mode in Chemdoc (Biorad). Lane 1- Protein Molecular Weight Marker; 2- Control (without metal treatment); 3-Pb; 4 – Cr and 5-Cd; 6- marker (low molecular weight). Red arrow marks indicates the protein bands. B: Secretory protein profile of *Candida viswananthii* strain ODBG2. Secretory protein (60 μ g) separated on 10% SDS-PAGE and stained with CBB R-250. The image was documented in invert image mode in Chemdoc

(Biorad). Lane 1- Protein Molecular Weight Marker; 2- Control (without metal treatment); 3-Pb; 4 – Cr and 5-Cd; 6- marker (low molecular weight). Red arrow marks indicates the protein bands.