First evidence showing that Pepper vein yellows virus P4 protein is a movement protein

Sangsang Li

Hunan University

Xianyang Su

Anhui Academy of Agricultural Sciences

Xiangwen Luo

Hunan Academy of Agricultural Sciences

Yu Zhang

Hunan University

Deyong Zhang

Hunan University

Jiao Du

Hunan Academy of Agricultural Sciences

Zhanhong Zhang

Hunan Academy of Agricultural Sciences

Xian OuYang

Hunan Academy of Agricultural Sciences

Songbai Zhang (✉ zsongb@hotmail.com)

Hunan Plant Protection Institute <https://orcid.org/0000-0001-8883-4525>

Yong Liu

Hunan University

Research article

Keywords: Pepper vein yellows virus, P4 protein, Plasmodesmata, movement protein, cell-to-cell movement

Posted Date: December 5th, 2019

DOI: <https://doi.org/10.21203/rs.2.18235/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Plant viruses move through plasmodesmata (PD) to infect new cells. To overcome the PD barrier, plant viruses have developed specific protein(s) to guide their genomic RNAs or DNAs to path through the PD.

Results In the present study, we analyzed the function of Pepper vein yellows virus P4 protein. Our bioinformatic analysis showed that the P4 protein contains an transmembrane domain, encompassing the amino acid residue 117-138. The P4 protein was found to target PD and form small punctates near walls. The P4 deletion mutant or the substitution mutant lost their function to produce punctates near the walls inside the fluorescent loci. The P4-YFP fusion was found to move from cell to cell in infiltrated leaves, and P4 could complement Cucumber mosaic virus movement protein deficiency mutant to move between cells.

Conclusion Taking together, we consider that the P4 protein is a movement protein of Pepper vein yellows virus.

Background

Plasmodesmata-mediated macromolecular trafficking is critical for plant growth and development [1]. Plant viruses have been shown to traffic between host cells through plasmodesmata (PD), and such trafficking is crucial for viral systemic infection [2].

PD is considered as a bottleneck for plant virus infection in plant, due mainly to its size exclusion limit (SEL) and/or the intricate and dynamic regulations controlled by the host defense mechanism [3]. To overcome this bottleneck, plant viruses have evolved to encode movement protein(s) (MP) to facilitate their intracellular trafficking, in a form of viral replication complexes or viral particle [4]. The MPs produced by plant viruses are strikingly different [5]. The pioneer report of viral MP is the 30 kDa MP of Tobacco mosaic virus, this MP was considered to guide TMV virion to move between cells [6]. The second type of MPs have two or three specialized MPs, and are referred to as double or triple gene block proteins (DGBps and TGBps) [7]. The third type of MPs are low molecule MPs, such as NSm encoded by Tomato spotted wilt tospovirus (TSWV) [8].

Pepper vein yellows virus (PeVYV), a new member in the genus Polerovirus, family Luteoviridae, was reported to infect multiple important solanaceous crops in Europe [9], Asia [10] and Africa [11]. Although the P4 protein of PeVYV was recently reported as a viral movement protein, based on the bioinformatic comparison to the genome structure of the movement protein of the type Potato leafroll virus species (PLRV) [12], the biological function and the key domain of P4 protein needed for PD trafficking remained unknown.

In the present study, we tried to determine the key domain needed for the interaction between the P4 protein and PD. Our result revealed that PeVYV P4 protein localized in PDs in cell walls and a domain covering amino acid (aa) residues 117 to 138 played an important role in PeVYV P4 PD localization.

Additionally, PeVYV could facilitate yellow fluorescent protein (YFP) cell-to-cell movement, and P4 could complement Cucumber mosaic virus movement protein deficiency mutant to move between cells.

Methods

Plants

Nicotiana benthamiana plants were grown inside a culture room set at 25°C with a 16 h light/8 h dark illumination. The seeds of *N. benthamiana*, which is a laboratory cultivar were obtained from The Plant Virology Laboratory of Ningbo University, Ningbo, China and stored in our lab. When the plants reached six-eight weeks old, they were infiltrated individually with one of the agrobacterium cultures made in this study.

P4 protein construction prediction

Transmembrane (TM) domain inside the P4 protein was predicted using five commonly used methods available on Internet: DAS (<http://www.sbc.su.se/miklos/DAS>), HMMTOP (<http://www.enzim.hu/hmmtop/>), TMPred (https://embnet.vital-it.ch/software/TMPRED_form.html), Split (<http://split.pmfst.hr/split/4>), and Topcons (<http://topcons.cbr.su.se>).

Plasmid constructs and agroinfiltration

Full length PeVYV *P4* gene and its alanine substitution mutant or deletion mutant aa position (117-138) were constructed individually through overlap PCR using primers listed in Table S1. The wild type (WT) *P4* and its mutants were cloned into expression vector pGWB441(YFP) or pGWB505(GFP) using the Gateway technology [13]. Plasmid pPDLP8-YFP and pPDLP8-RFP (marker protein localized in PD) was from a previously described source and was used to localize plasmodesmata in cell walls [14]. These plasmids were introduced individually into *Agrobacterium tumefaciens* strain GV3101. *Agrobacterium* cultures containing the corresponding plasmid were grown overnight in a YEP medium (10 g yeast extract, 10 g Bacto peptone and 5 g NaCl in one liter H₂O, pH 7.0) supplemented with 100 mg/L kanamycin and 50 mg/L rifampicin. *Agrobacterium* cells were pelleted and then incubated for three hours in an infiltration buffer (10 mM MgCl₂, 10 mM MES, pH 5.9, and 150 µM acetosyringone). The cultures were further diluted to OD₆₀₀ = 0.2 and then infiltrated individually into the abaxial side of *N. benthamiana* leaves. The infiltrated plants were again grown at 25°C with a 16 h light/8 h dark illumination.

Western blotting

Western blotting was operated as previously described [5]. Total protein was separated by electrophoresis in 10% SDS-PAGE and transferred onto a PVDF membrane. The antigens on the PVDF membrane were detected with polyanitbody against PeVYV *P4*, then incubated by AP-coupled goat anti-mouse IgG (1:5 000 dilution; Sigma) and 5-bromo-4-chloro-3-indolylphosphate/nitroblue tetrazolium (NBT/BCIP) staining (Sangon Biotech, Shanghai, China).

YFP, GFP and RFP images

Agro-infiltrated leaves were detached from the plants and examined for fluorescence from different YFP, GFP or RFP fusion proteins under a Nikon TI-E+C2 confocal laser-scanning microscope (Nikon Microsystems, Watford, United Kingdom). The excitation wavelength was set at 514 nm for YFP and GFP, and 555 nm for RFP, and the emission wavelength was set at 520-550 nm for YFP and GFP, and 583 nm for RFP. The images presented in the figures are either single image or multi-layered images to achieve the maximum signal intensity.

Results

P4 protein construction

Membrane association and plasmodesmata targeting are crucial functions of viral movement proteins. To investigate domain(s) in the P4 protein that are responsible for membrane association, we first predicted transmembrane (TM) domain(s) in the P4 protein using several bioinformatics algorithms. Results shown in Figure S1 revealed two potential TM domains in the P4 protein. Both DAS and TMpred algorithms showed that the P4 protein was an integral membrane protein and the two predicted TM domains were located at a region encompassing aa residue 127-129 (by DAS algorithm) and aa residue 117-138 (by TMpred algorithm), respectively.

P4 protein targets PD of host cells

To determine whether the P4 protein could target plasmodesmata in cell walls, we first infiltrated *N. benthamiana* leaves with *A. tumefaciens* cultures carrying the pP4-YFP or pPDL8-YFP plasmid. The infiltrated leaves were harvested at 48 hpi and then examined for the intracellular localization patterns of the two fusion proteins by Confocal Microscopy. The result showed that the P4-YFP fusion protein accumulated as yellow fluorescence punctates in the cytoplasm and near the cell walls (Figure 1, upper panel, red arrows showed). The PD-YFP fusion protein was found to produce small yellow fluorescence punctates near the cell walls, but not large punctates in the cytoplasm like the P4-YFP (Figure 1, lower panel).

To confirm if the computer-predicted TM domains in the P4 protein had the ability to target plasmodesmata, we fused the P4 mutants $\Delta P4^{117-138}$ and $\Delta P4^{AAA117-138}$ to YFP, respectively, and expressed them in *N. benthamiana* leaves. By 48 hpi, the infiltrated leaves were harvested and examined under the confocal microscope. The results showed that the yellow fluorescence from the $\Delta P4^{117-138}$ -YFP fusion or from the $\Delta P4^{AAA117-138}$ -YFP fusion was near cell walls, but no yellow punctates were observed in the cell walls, reminiscent of PD localization (Figure 2). The co-localization of P4 or its mutants with PDL8 (marker protein localized in PD) showed that only P4 could target PD (Figure 3).

P4 protein functions as movement protein

To determine whether the P4 protein can move through PD, we infiltrated *N. benthamiana* leaves with *A. tumefaciens* culture carrying the pP4-YFP plasmid. As a control, we infiltrated *N. benthamiana* leaves with a different culture carrying the pPDLP8-YFP plasmid. A total of 35 fluorescent loci expressing PD-YFP were examined under the confocal microscope at 24 hpi and the PD-YFP fusion was found in single cells only (Figure S2 and Table S2). At the same time point, the P4-YFP fusion in 21 of the 23 fluorescent foci had trafficked through the PD and entered the neighboring cells.

To further determine whether the P4 protein functions as movement protein to drive virus cell-to cell moving through PD, we infiltrated *N. benthamiana* leaves with *A. tumefaciens* culture carrying the plasmid of P4 or its mutants and CMV infectious clone with its movement protein 3a replaced by GFP . As a control, we infiltrated *N. benthamiana* leaves with a different culture carrying the plasmid of expressing P4 and its mutants. As figure 4 shown, P4, not P4 mutants could drive *Cucumber mosaic virus* movement protein deficiency mutant cell-to cell moving through PD. A total of 46 of the 50 fluorescent loci expressing P4-GFP and CMV movement protein deficiency mutant were examined under the confocal microscope at 6 dpi, and the P4 mutants was found in single cells only or less than 8% virus entered the neighboring cells (Figure 4 and Table 1).

Table 1 Movement of PeVYV P4 between epidermal cells in *N. benthamiana* leaves at 6 dpi

Constructs	No. of loci examined	No. of loci with a single cell (%)	No. of loci with more than 2 cells (%)	ρ -value
Vector	50	50 ^a (100%)	0	
CMV 3a	50	0	50(100%)	
P4	50	4(8%)	46(92%)	<0.05 ^b
ΔP4 117-138	50	50(100%)	0	
ΔP4 AAA117-138	50	42(84%)	8(16%)	<0.05

^a Loci with signal cells expressing GFP fluorescence.

^b ρ -value was determined using the unpaired two-tailed Student *t*-test.

Discussion

It is well documented that different plant viruses encode different movement proteins (MPs) to traffic through PD between two cells [6–8]. For example, the 17 kDa movement protein of Potato leafroll virus (PLRV, genus Polenovirus, family Luteoviridae), was reported as a low molecular weight movement protein of PLRV [15]. In the present study, the PeVYV P4 protein (17.5 kDa), which was previously

predicted as a virus movement protein, was analyzed and found to localize to PD in cell walls. A domain encompassing amino acid residues 117–138 in the P4 protein was determined to have an ability to target PDs.

The P4-YFP fusion protein was found to traffic between epidermal cells, and P4 could complement heterological Cucumber mosaic virus movement protein deficiency mutant to move between cells. These results provided the first evidence showing that the PeVYV 17.5 kDa protein is a viral MP.

Interestingly, in the *N. benthamiana* leaf epidermal cells expressing P4-YFP fusion protein, large and small yellow fluorescent punctates were observed (Fig. 2). Previous studies have shown that many viruses can produce aggregates during their infection in plant and these aggregates may have multiple functions including preventions of degradations by host cellular machinery [16] and/or providing viral and host factors for virus replication [17]. So the large punctates were found in cytoplasm and the small punctates were found near the cell walls.

Conclusion

The results presented here demonstrated that the 17.5 kDa protein of PeVYV is a movement protein of the virus. Whether this protein also has a role in virus replication and/or counteraction of host defense machinery requires further investigation.

Abbreviations

aa: amino acid; DGBps: double gene block proteins; PeVYV: Pepper vein yellows virus; PD: plasmodesmata; PLRV: *Potato leafroll virus*; TGBps: triple gene block proteins; TM: Transmembrane; TSWV: *Tomato spotted wilt tospovirus*

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analyzed during this study are included in this published article and its supplementary information files.

Competing Interests

The authors have declared that no competing interest exists.

Funding

This study was financially supported by the National Natural Sciences Foundation of China (31672003 directed to ZZH, 31772133 directed to ZSB). Funding bodies had no role in study design, collected data, analysis, or writing.

Author Contributions

LSS, SXY, ZZH, OYX and LXW - Carried the experimental work. ZY, ZDY and DJ - Collected and analyzed the data, ZSB and LY, Designed study, guided data interpretation, and wrote the manuscript. All authors approved the manuscript before it was submitted by the corresponding author.

Acknowledgments

I would like to thank Jian Yang for technical assistance of experiments of subcellular localization of P4 protein.

Author details

1. Longping Branch, Hunan University, Key Laboratory of Pest Management of Horticultural Crop of Hunan Province, Changsha 410125, China;
2. Hunan Plant Protection Institute, Hunan Academy of Agricultural Science, Changsha 410125, China
3. Plant Protection Institute of Anhui Academy of Agricultural Science, Hefei 230001, China

References

1. Kim I, Zambryski PC Cell-to-cell communication via plasmodesmata during *Arabidopsis* Curr Opin Plant Biol. 2005; 8: 593-599.
2. Heinlein M. Plasmodesmata: channels for viruses on the move. Methods Mol Biol. 2015; 12: 25-52.
3. Waigmann E, Ueki S, Trutnyeva K, Citovsky V. The ins and outs of nondestructive cell-to-cell and systemic movement of plant viruses. Crit Rev Plant Sci. 2004; 23: 195-250
4. Boevink P, Oparka KJ. Virus-host interactions during movement processes. Plant Physiol. 2005; 138: 1815-1821.
5. Lazarowitz SG, Beachy RN. Viral movement proteins as probes for intracellular and intercellular trafficking in plants. Plant Cell. 1999; 11: 535-548.
6. Liu C, Nelson RS. The cell biology of Tobacco mosaic virus replication and movement. Front Plant Sci. 2013; 4: 12
7. Morozov SY, Solovyev AG. Triple gene block: modular design of a multifunctional machine for plant virus movement. J Gen Virol. 2003; 84, 1351-1366

8. Feng Z, Xue F, Xu M, Chen X, Zhao W, Garcia-Murria MJ. The ER-membrane transport system is critical for intercellular trafficking of the NSm movement protein and Tomato spotted wilt tospovirus. *PLoS Pathog.* 2016; 12: e1005443.
9. Rast ATB. Occurrence of *Pepper yellow vein virus* in the Netherlands. *Neth J Plant Pathol* 1988; 94: 311-313.
10. Zhang SB, Zhao ZB, Zhang DY, Liu Y, Luo XW, Liu J, Wu LF, Peng J. First report of pepper vein yellows virus infecting red pepper in mainland China. *Plant Dis.* 2015; 99: 1190.
11. Knierim D, Tsai WS, Kenyon L. Analysis of sequences from field samples reveals the presence of the recently described pepper vein yellows virus (genus *Polerovirus*) in six additional countries. *Arch Virol.* 2013; 158: 1337-1341.
12. DeBlasio SL, Xu Y, Johnson RS, Rebelo AR, MacCoss MJ, Gray SM, Heck M. The interaction dynamics of two potato leafroll virus movement proteins affects their localization to the outer membranes of mitochondria and plastids. *Viruses*; 2018; 10: E585.
13. Bryksin AV, Matsumura I. Overlap extension PCR cloning: a simple and reliable way to create recombinant plasmids. *Biotechniques.* 2010; 48(6): 463-465.
14. Ye ZW, Chen QF, Chye ML. Arabidopsis thaliana Acyl-CoA-binding protein ACBP6 interacts with plasmodesmata-located protein PDLP8. *Plant Signal Behav.* 2017; 12(8): e1359365.
15. Schmitz J, Stussi-Garaud C, Tacke E, Prüfer D, Rohde W, Rohfritsch O. In situ localization of the putative movement protein (pr17) from *potato leafroll luteovirus* (PLRV) in infected and transgenic potato plants. *Virology.* 1997; 235(2): 311-322.
16. Moshe A, Gorovits R. Virus-induced aggregates in infected cells. *Viruses.* 2012; 4(10): 2218-2232.
17. Kaido M, Abe K, Mine A, Hyoudou K, Taniguchi T, Taniguchi H, Mise K, Okuno T. GAPDH-A recruits a plant virus movement protein to cortical virus replication complexes to facilitate viral cell-to-cell movement. *PLoS Pathogen.* 2014; 10(11): e1004505.